

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,900

Open access books available

185,000

International authors and editors

200M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



Variation of Greenhouse Gases in Urban Areas-Case Study: CO₂, CO and CH₄ in Three Romanian Cities

Iovanca Haiduc and Mihail Simion Beldean-Galea
*Babeş-Bolyai University, Faculty of Environmental Science,
 Cluj-Napoca,
 Romania*

1. Introduction

The natural equilibrium of atmospheric gases has been maintained for millions of years, but with the beginning of the industrial age, it became more fragile due to human activity. In the Intergovernmental Panel on Climate Change Report (IPCC) named "Climate Change 2007" (IPCC-AR4, 2007) it is specified that "the keep going emissions of the greenhouse gases (GHG) at/over current rate, will cause in the future global warming and will induce more global climate changes in the 21st century than those of 20th century". More than , in the coming IPCC Report named "Carbon cycle including ocean acidification (CCT)" (IPCC-AR5, 2010) is stipulated that ocean acidification will be a further critical and direct consequence of increasing atmospheric GHG concentrations.

In 1886, the chemist Svante Arrhenius (Nobel prize for Chemistry in 1903) calculated for the first time the CO₂ contribution (from fossil fuel combustion) to climatic changes and used for the first time the term of "greenhouse effect". Almost 100 years were necessary for the confirmation of Arrhenius predictions about the evolution of global climatic factors, and the fact that CO₂ is the main greenhouse gas, with a contribution of 55% to Global Warming Effect. The first IPCC Report (IPCC-FAR, 1990) draws the conclusion about the possible existence of a global warming phenomenon. The second IPCC Report (IPCC-SAR, 1995) shows the contribution of humans to global warming and predicts a major warming in the 21st century. The third IPCC Report (IPCC-TAR, 2001) affirms a very probable (60% - 90%) warming for the next century. In the IPCC-AR4 Report (IPCC-AR4, 2007) adds for the understanding of the impact of climate changes over the vulnerability and the adaptation of the environment, the most relevant scientific, technical and socio-economical information from more than 1500 scientific papers. This report accepts with a probability of over 90% that the emission of greenhouse gases and not the environmental conditions gives the global warming effect.

The IPCC Guide from 2006 makes an inventory of gases from atmosphere and distinguishes between:

- a. gases with GWP (Global Warming Potential) listed in IPCC 2001: CO₂, CH₄, N₂O, hydro fluorocarbons, per fluorocarbons, SF₆, NF₃, SF₅CF₃, halogenated ether, C₄F₉OC₂H₅, CHF₂OCF₂OC₂F₄OCHF₂, CHF₂OCF₂OCHF₂ and other halocarbons CF₃I, CH₂Br₂, CHCl₃, CH₃Cl, CH₂Cl₂.

b. gases without GWP: $C_3F_7C(O)C_2F_5$, C_7F_{16} , C_4F_6 , C_5F_8 and C_4F_8O .

As a follow-up of these reports, the scientific community had started a cycle of research programs having as scientific goal the complex study (emission sources, consumption sources, the balance of changes between the components of environment etc.) of these gases and the effect produced by them (Projects CARBOEUROPE, AEROCARD, CHIOTTO, Global Carbon Project, IGOS, NACP etc).

The CO_2 it is an important green-house gas and its level in the atmosphere has significantly increased from 280 ppm in the pre-industrial era to current 380 ppm. The first increase of 50 ppm occurred during a period of ca. 200 years, starting with the beginning of the industrial revolution until 1973. Between 1973 and 2006 the concentration of CO_2 increased with another 50 ppm.

The Global Warming Potential (GWP) for CO_2 is conventionally choose as 1, i.e. the atmospheric residence time between 50 and 200 years, and its contribution to the greenhouse effect is ca. 52 %.

According to IPCC Report (IPCC-AR4, 2007), the contribution of anthropic CO_2 is predominant, and the main anthropic sources are:

- the energetic sector 30 %
- industrial processes 20 %
- fuels used in transportation 20 %
- burning of biomass 9.1 %
- processing and distribution of fossil fuels 8.4 %
- other sources 12.5%

According to IPCC estimations, the increase of CO_2 concentration in the atmosphere leads to climate changes and will produce a global warming of the planet through the greenhouse effect.

The monitoring of the global levels of CO_2 is the concern of American government since about 30 years. The National Oceanic & Atmospheric Administration - Earth System Research Laboratory (NOAA-ESRL) performs measurements for main greenhouse gases in about 68 locations spread all over the world. According to NOAA data, the concentration of CO_2 has an ascending trend. Figure 1 shows the CO_2 variations in Mauna Loa for the period of 2003-2008 according to NOAA-ESRL source.




Fig. 1. Variation of the monthly average of CO_2 in Mauna Loa

At the European level, the CarboEurope-Clusters Program (CarboEurope) carries out measurements of atmospheric CO₂ concentrations in 61 locations in 17 countries. This program contains eight different projects working together to contribute to the understanding of the carbon cycle at the European level. The projects involved in this program are: AEROCARB, CAMELS, CARBOAGE, CARBODATA, CARBOEUROFLUX, CARBOEUROPE GHG; CARBOINVENT, CHIOTTO, CARBODATA, TACOS, EUROSIBERIAN CARBONFLUX, FORCAST, GREENGRASS, RECAB, TACOS-INFRASTRUCTURE, and TCOS SIBERIA. Two other projects, CARBOMONT and SILVISTRAT are associated to this program.

In 1998 Idso and co-workers (Idso et al., 2001) introduced the term “urban DOME” for the persistence of CO₂ over the urban cities as a result of anthropogenic contribution to CO₂ budget. After that, several individual studies regarding to CO₂ variation in urban areas have been reported (Day et al., 2002; Idso et al., 2002). These studies showed that the concentration of CO₂ in urban area is higher than the CO₂ concentration in rural area and this fact is a consequence of human activities. A literature review about these results is presented in this chapter.

Methane (CH₄) is another important greenhouse gas, with GWP = 25 and a residence time of more than 100 years. It is produced both naturally and through human activities. The global mixing ratios of CH₄ in the atmosphere have more than doubled since the pre-industrial period, rising from around 750 ppb (parts per billion) in 1800 (Simpson et al., 2002; Dlugokencky et al., 2003) to the current level of around 1770 ppb (NOAA-ESRL). The global trend of the methane concentration in the air is ascending, with a rate of increase of 5–10 ppb/year, and for the period 1984–2004 this tendency is shown in Figure 2.




Fig. 2. The variation of the global average concentration of CH₄. (source NOAA-ESRL)

The main natural source of methane are dominated by wetlands. The primary way for CH₄ transformation is its destruction in the atmosphere by hydroxyl radicals (Prinn et al., 1995, 2001). Some CH₄ is also oxidized by microorganisms (called methanotrophs), which use CH₄ as a source of carbon and energy. Tropospheric CH₄ is eventually oxidized to carbon dioxide; its atmospheric lifetime is estimated to be 8–12 years (NOAA-ESRL; Cunnold et al., 2002; IPCC-AR4, 2007).

It is estimated that the contribution of methane to the greenhouse effect is 18%, and the most important sources are:

- Residual agricultural products 40%
- Processing and distribution of fossil fuels 29.6%

- Storage and processing of domestic waste 18.1%
- Burning of biomass and grazing 6.6%
- Other sources 4.8%

In the IPCC Report (IPCC-TAR, 2001) is suggested that the natural sources account for ca. 40% of total methane sources. Aikawa (Aikawa, 2006) indicates that the transportation contributes 1.1% of CH₄ emissions in Japan and suggests that there is a small influence of methane emissions from mobile sources on the concentrations in ambient air. The possible evolution of anthropogenic methane emissions at global level has been discussed by Cofala (Cofala et al., 2007) who predicts an increase of CH₄ emissions from 250 Tg/year in 1990 to 420 Tg/year in 2030.

Carbon monoxide (CO) is the most significant pollutant. It has a short life time in the atmosphere due to its reaction with other atmospheric components, such as hydroxyl radicals. It has an indirect radiative effect by increasing the concentration of methane and tropospheric ozone. In urban areas, CO reacts photochemically with aldehydes, to produce peroxy radicals. These radicals react with nitrogen oxide, to form nitrogen dioxide, which is the main responsible for the formation of the fotochemical smog.

Through natural processes, the CO can be oxidized to CO₂, thus contributing to the increase of the later in the atmosphere. Thus, through resulting products the anthropogenic CO can indirectly contribute to the greenhouse effect and to the global warming.

Anyway, even if CO₂ and CH₄ are the main greenhouse gases, with a contribution of more than 70% (CO₂ 55%, CH₄ 15%) on global warming (IPCC-AR4, 2007), for a good estimation of these two gases to the Global Warming Potential (GWP) in urban areas it is necessary to take into account the indirect contribution of CO which is not a greenhouse gas but changes the atmospheric chemistry and the abundance of other greenhouse gases. CO is a key air pollutant which can be utilized like tracer in the separation of CO₂ and CH₄ from biogenic and anthropogenic sources (Daniel & Solomon, 1998).

Different studies (Daniel & Solomon, 1998; Fuglesvedt et al., 1996; Prather, 1996 cited in IPCC-TAR, Chapter 4, 2001) estimate the indirect GWP of the CO due to O₃ production and to feedbacks on the CH₄. This approach was made using a box model and estimate the indirect GWP of CO for time horizons of 20, 100, and 500 years. The indirect GWP value due to CO is gave in table 1.

| Authors/Models | Indirect Global Warming Potentials of CO, Time horizon | | |
|---|--|-----------|-----------|
| | 20 years | 100 years | 500 years |
| Daniel and Solomon (1998): box model considering CH ₄ feedbacks only | 2.8 | 1.0 | 0.3 |
| Fuglestvedt et al. (1996): two-dimensional model including CH ₄ feedbacks and tropospheric O ₃ production by CO itself | 10 | 3.0 | 1.0 |
| Johnson and Derwent (1996): two-dimensional model including CH ₄ feedbacks and tropospheric O ₃ production by CO itself | - | 2.1 | - |

Table 1. Estimated Indirect Global Warming Potentials of CO for time horizons of 20, 100, and 500 years, (source IPCC-TAR, 2001)

Regarding to long time measurement of CO₂ concentration in urban area as well as the variation of meteorological parameters allow to understanding the rule of the ecosystem functioning and meteorological parameters over inter-annual variation in carbon fluxes. The inter-annual variation of CO₂ fluxes has been typically studied either by modeling approaches (Higuchi et al., 2005; Ito et al., 2005; Bergeron & Strachan, 2011) or by correlation coefficient analyses together with meteorological parameters (Aubinet et al., 2002; Aurela et al., 2004; Wohlfahrt et al., 2008). These studies showed that, the CO₂ flux is strongly influenced by biological vegetation cycle and the variation of meteorological parameters. Thus the maximum value of CO₂ is registered during the cold season while the minimum value of CO₂ was registered during the summer. Another study (Sottocornola & Kiely, 2010), show that the wet conditions favored the CO₂ uptake by the ecosystem in autumn and in winter, while the warmer and dryer weather reduce the sequestration of CO₂ in the ecosystem. A study performed in urban and sub-urban area of Montreal (Bergeron & Strachan, 2011) showed that the CO₂ flux is also influenced by the anthropic activity. According to this study, *“Lower emissions at the suburban site are attributed to the large biological uptake in summer and to its relatively low population density inducing low anthropogenic emissions. Higher emissions at the urban site are partly associated with its greater population and building density, promoting higher emissions from vehicular traffic and heating fuel combustion. Vehicular traffic CO₂ emissions influenced the diurnal cycle of CO₂ fluxes throughout the year at the urban site. At the suburban site, summer CO₂ fluxes were dominated by vegetation sources and sinks as daytime CO₂ uptake occurred and CO₂ fluxes responded to incoming light levels and air temperature in a fashion similar to natural ecosystems. To a lesser extent, the vegetation component also helped offset CO₂ emissions from other sources in summer at the urban site.”*

This chapter will present the results of a case study of CO₂, CH₄ and CO variations during one year in three selected cities from Romania with different anthropic activity. In order to identify the influence of biogenic and anthropogenic sources to the budget of mentioned greenhouse gases the ¹³CO₂ and ¹³CH₄ isotopic composition have been determinate. Experimental results were finally correlated with meteorological parameters.

2. CO₂ in urban area

2.1 Trends of CO₂ variation in urban areas. A literature review

The problem of urban carbon dioxide came into the attention of scientists in the year 1998, with the discovery and characterization of the urban CO₂ dome of Phoenix, Arizona, USA by Idso and col. (Idso et al., 2001). Early work found that under certain meteorological conditions, urban CO₂ concentrations could be as high as 550-600 ppm (some 200 ppm higher than the surrounding countryside) (Idso et al., 2002). Soegaard and Moller-Jensen (Soegaard & Moller-Jensen, 2003) studied the urban CO₂ dome of Copenhagen and indicated that "traffic is the largest single CO₂ source in the city," and demonstrate that "emission rates range from less than 0.8 g CO₂ m⁻² h⁻¹ in the residential areas up to a maximum of 16 g CO₂ m⁻² h⁻¹ along the major entrance roads in the city center."

Following these studies, research regarding the urban CO₂ domes has been performed in many other parts of the world (Table 2). The results obtained from studies conducted in several cities from all over the world, show several commonalities. Thus, anthropogenic CO₂ emissions are the primary source of the urban CO₂ dome; the dome is generally stronger in city centers, in winter, on weekdays, at night, under conditions of heavy traffic, close to the ground, with little to no wind, and in the presence of strong temperature inversions. These

conclusions are in agreement with the data provided by Commonwealth Scientific and Industrial Research Organisation (CSIRO) which indicate that typical concentrations of CO₂ in urban areas is situated between 350 and 600 ppm and depend on meteorological parameters and urban agglomeration.

| Authors | Place of measurements | Period of measurement | CO ₂ range concentrations |
|-----------------------|------------------------------------|--------------------------|---|
| Coutts A. M. | Melbourne, Australia | February - July, 2004 | 355 - 380 ppm (daily mean concentration). |
| Day T. A. et al. | Phoenix, USA | March - April, 2000 | 377 - 396 ppm (daily mean concentration) |
| George K. et al. | Baltimore, USA | 2007 | 488 in urban area, 442 in sub-urban area, 422 in rural area |
| Ghuri B. | Six cities, Pakistan | 2003 - 2004 | 270 - 325 ppm in Islamabad, 289 - 389 ppm in Quetta, 316.5 - 360 ppm in Karachi, 324.1 - 380 ppm in Lahore 295.2 - 356 ppm in Rawalpindi, 312 - 382 ppm in Peshawar. |
| Gratani L. et al. | Rome, Italy | 1995 - 2004 | 367 ± 29 ppm in 1995 (monthly mean variation) 477 ± 30 ppm in 2004 (monthly mean variation) 414 ± 25 ppm green zone 505 ± 28 centrale zone |
| Grimmond et al. | Chicago, USA | July 11- August 14, 1995 | 338 - 370 ppm (diurnal variation) 405 - 441 ppm (nightly variation) |
| Idso S. B. et al. | Phoenix, USA | 2000 | 390.2 ± 0.2 ppm (minimum daily concentration) 424.3 - 490.6 ppm (maximum daily concentration) 619.3 ppm (maximum daily concentration in cold season) |
| Kuc T. et al. | Kasprowy Wierch and Krakow, Poland | 2000 | 370 ppm (monthly mean variation in Kasprowy Wierch) 370 - 430 ppm (monthly mean variation in Krakow) |
| Moriwaki R. et al. | Tokyo, Japan | October-Nov., 2005 | 380 - 580 ppm (daily mean concentration). |
| Nasrallah H.A. et al. | Kuwait City, Kuwait | 1996 - 2001 | 368 - 371 ppm (daily mean concentration at 7 metter high). |
| Velasco E. et al. | Mexico city, USA | June 11- August 14, 1995 | 398 - 444 ppm (daily variation) 421 ppm (daily mean) |

Table 2. Overview of urban CO₂ measurements

Taking in to account the concentration of CO₂ from the cities, some of researches were focused on the impact of local CO₂ emissions over local temperature. Thus, Balling et al. running the CO₂ concentration through a radiation model calculated that local CO₂ emissions modify the local temperature with more than one-tenth of one degree Celsius. In fact, Balling

et al. suggest that this increasing of temperature is insignificant by comparing it to the overall urban heat island in Phoenix which typically adds 5 to 10 degrees C. (Balling, Jr., et al., 2001). Recently, Jacobson (Jacobson, 2010) found that domes form above cities more than a decade ago, cause local temperature increases that in turn increase the amounts of local air pollutants, raising concentrations of health-damaging ground-level ozone, as well as particles in urban air. Also, this study has shown that "CO₂ dome" that develops over urban areas is a local problem, which creates much more health problems than in rural areas.

The conclusions of Jacobson about the human health effects of CO₂ created many controversies; therefore, more research is necessary on the measurement of CO₂ variation over the urban areas.

2.2 CO₂ variation in three Romanian cities. Case study

The available literature contains no information about the variation of CO₂ concentrations in Romania in urban areas. However, the American system of global monitoring of CO₂ (NOAA-ESRL) has a station of continuous measurement of the main parameters of air quality placed in Constanta, which also measures CO₂ concentrations.

According to NOAA-ESRL data, the variation of CO₂ concentrations at the Constanța measurement station has an ascending trend; the yearly average values are between 365 ppm in 1995 and 395 ppm in 2007. According to the same source the concentration of CO₂ has a seasonal variation with maxima in the cold season and minima in the warm season.

In order to study the influence of anthropic activity upon the CO₂ level the study was performed in three different Romanian cities from Cluj County, as follows:

- a. Cluj-Napoca city, 400 000 inhabitants,
- b. Turda city, 59 600 inhabitants
- c. Huedin town, 10 000 inhabitants

The case study has been carried out during one year (four seasons) from July 2008 to June 2009.

The measurements were performed monthly (during 8 hours from 8.30 am to 15.30 pm) in all the three selected locations using a NDIR CO₂ analyzer model EMG-4. The measurements of CO₂ levels were performed in a portable meteorological shelter.

For the estimation of the anthropogenic contribution over the CO₂ budget in all three studied areas a reference point situated outside of the cities has been chosen (Roba et al, 2009).

For the Cluj-Napoca city, three measurement points have been selected: one situated in a zone with intense traffic (Piața Mărăști), one with moderate traffic (Cartier Grigorescu) and a reference point located in a periphery location (meteorological station in Cartier Gruia).

The results of the measurements recorded in the three locations in the city of Cluj-Napoca show that during the year the CO₂ level is strongly influenced by the amplitude of the anthropic activities. The highest levels were recorded in the location with intense anthropic activity (Piata Marasti) and the lowest levels in the reference point located outside of the city. The recorded values are comprised between 380 and 530 ppm (Mărăști), 376-456 (Grigorescu) and 373-444 (reference point).

It was also found that the highest values were recorded during the months of October and November (during the final period of the biological cycle of plants) and during the winter. It was also found that starting with the months of March; the CO₂ concentrations decrease and become comparable with the summer values (Figure 3).

For the Turda city two measurement points have been selected: one in the city (Potaisa School) and another one located outside of the city (Meteo Station).

The results of the measurements in the two locations indicate a concentration difference between 15 and 20 ppm, depending of the period of measurements. The largest values of CO₂ concentration were obtained in the months of October and November, at the end of the biological cycle of plants and in the winter. The concentrations measured are in the range 380-502 ppm at Potaisa site and 371-450 ppm at the reference point.

In the town of Huedin the measurements were carried out simultaneously in two locations: one in the interior of the city (Liceul Octavian Goga) and a reference point in the peripheral area (Meteo Station). The results show a slight difference between the two locations. It was also observed again that the largest values are recorded in the months of October and November, i.e. at the end of the biological cycle of plants, and during the winter.

Starting with the month of March, the CO₂ concentrations decrease to normal values, comparable to those of the summer. The values recorded are in the range 355- 433 ppm (Octavian Goga site) and 350-411 ppm (reference point).

A comparison of the measured values carried out in different locations shows that the CO₂ concentrations depend on the size of the town, the highest values being recorded in Cluj-Napoca city, followed by Turda and Huedin (Figure 3).




Fig. 3. Annual variation of CO₂ in urban studied areas

Diurnal variations were also observed; the highest values were measured in the morning and the lowest values at the astronomic midday, both in summer and in winter seasons (Figure 4).




Fig. 4. Diurnal variation of CO₂ in urban studied areas

Taking into account the results of this case study we can conclude that the variation of CO₂ in studied urban areas is in agreement with other results reported in the scientific literature, which report that in urban areas the CO₂ levels are situated between 350 and 600 ppm, depending on meteorological parameters and urban agglomeration, with the observation that in the absence of rainfall the CO₂ level increases both in urban areas and outside.

3. CH₄ in urban area

3.1 Trends of CH₄ variation in urban areas. A literature review

Regarding to CH₄ variation in urban areas there are only a few studies about the level of methane in urban atmosphere, and these results are reported starting after the year 1980. For Romania such data are available only after 1995.

According to data from CSIRO the common concentrations of CH₄ in urban areas have values between 1700-2500 ppb and are influenced by meteorological parameters and urban agglomeration. Kuc and co-workers (Kuc et al., 2003) show that, the CH₄ level in urban areas is comprised between the natural level (1650 ppb) and 4200 ppb. Ito and co-workers (Ito et al. 2000) compared the atmospheric CH₄ concentrations recorded in Nagoya with the values measured at Mauna Loa Observatory in Hawaii (USA) and estimated that the excess concentration of CH₄ in the urban atmosphere of Nagoya was 170 ppb in 1988 and 150 ppb in 1997. A selective bibliography on the subject is presented in Table 3.

3.2 CH₄ variation in three Romanian cities. Case study

In Romania, the available literature provides no information about the variation of methane concentrations in urban areas and no such studies are reported. However, the American System of Global Monitoring of Air Quality (NOAA-ESRL), has a station of continuous measurements of atmospheric methane concentrations at Constanța. According to NOAA data, the concentration of CH₄ has an ascending trend, with average values comprised between 1880 ppb in 1995 and 1980 ppb in 2006. According to the same source, the methane concentration shows a seasonal variation, with maxima in the summer months and minima in the autumn and spring.

| Authors | Place of measurements | Period of measurement | CH ₄ range concentrations |
|------------------------|---------------------------------|------------------------|--|
| Aikawa M. et al. | Urban, Sub-urban, Nagoya, Japan | 2004 | 1.80 - 1.84 ppm -urban area 1.78 - 1.80 ppm-suburban area |
| Derwent R.G. et al. | Island | 1990 - 2003 | 1.75 - 2.00 ppm |
| Ghuri B. et al. | Pakistan | 2003 - 2004 | 0.5 - 1.7 ppm |
| Hsu Y.K. et al. | California, USA | April 2007 - Feb. 2008 | 1.75 - 2.16 ppm |
| Ito A. et al. | Nagoya, Japan | 1983 - 1997 | 1.85 ppm in 1998, 1.91 ppm in 1995, 1.90 ppm in 1997. 1983 - 1997 increase 13 ppb/year |
| Kuc T. et al. | Kasprowy Wierch Krakow, Poland | 2000 | 1650 ppb Kasprowy Wierch (monthly mean concentration) 2000 - 2800 ppb Krakow (monthly mean concentration) |
| Sikar E. & La Scala N. | Urban area, Brasil | 1998 - 1999 | 1.80 ppm |
| Smith F.A. et al. | Mexico City | March, 1993 | 1.8 ppm during the night 7.971 ppm in the morning 2.001 - 2.999 middle of the day |
| Thi Nguyen H et al. | Seul, Korea | 1996 - 2006 | 2.24 ± 0.42 ppm urban road-side 2.06 ± 0.31 ppm urban background |
| Veenhuysen D. et al. | Amsterdam, Netherlands | 1994 | 1.75 - 3.00 ppm |
| Wang J.L. et al. | Sub-urban area, Taiwan | 1-27 April, 2000 | 1.9 - 3.7 ppm |

Table 3. Overview of urban CH₄ measurements

Our case study has been focused on the measurement of the CH₄ variation in three urban areas from Cluj county as described in sub-chapter 2. The study was carried out during one year (four seasons) from July 2008 to June 2009. The measurements were performed monthly in all selected areas at the astronomic midday (in Romania at 12.30 h). In all three areas a measurement point located in the city and a reference point located outside has been selected. The samples were collected in Cluj-Napoca, in Marasti location in the city and as reference point at Gruia location. In Turda the measurements in the city were made at Liceul Potaisa, and the reference point at the Meteo station. In Huedin the measurements were made at Liceul Octavian Goga in the town and the reference point was the meteo station.

For atmospheric CH₄ measurements, the air samples were collected by the flask sampling method and analysed by gas chromatography technique (GC) coupled with a flame ionisation detector (FID) (Cristea et al., 2009).

The results show a significant variation of atmospheric methane, depending on the season and the urban agglomeration degree. Thus, the highest values were recorded in the city of Cluj-Napoca (11.5 ppm in April 2009) and the lowest values were recorded in the month of August 2008 (2.5 ppm).

In Turda the concentrations of atmospheric methane were comprised between 2.2 and 8.6 ppm, with the lowest values measured in August 2008 (2.2 ppm) and the highest values recorded in April 2009 (8.6 ppm).

In Huedin the concentrations of methane were measured between 1.4 and 7 ppm, with the lowest values recorded in July 2008 (1.4 ppm) and the highest ones in April 2009 (7 ppm).

Significant differences are also observed between the methane concentrations in the interior of the cities and the reference points located outside. These differences were recorded throughout the experiments, which leads to the conclusion that the anthropic activities, the automobile traffic in particular, are an important source of methane in the urban atmosphere.

The analysis of the methane concentrations in the three areas investigated indicates a similar profile for the measurements in the interior of the cities, with minima in the summer months and maxima during the spring. This may be attributed to the absence of rains in the spring (March-April). In May 2009, when the precipitations started, the concentration of atmospheric methane became closer to the values measured at the reference points.

For the reference points, the values of the atmospheric methane concentrations are in the range 2.1-4.2 ppm in Cluj-Napoca, 1.7-3.5 in Turda and 1.4-2.9 ppm in Huedin. As for the measurements in the city, a slight increase of the concentrations were observed in the spring period of 2009, due to the lack of precipitations.




Fig. 5. The variation of methane concentration in the urban areas of Cluj county locations.

The results of our measurements indicate that the atmospheric CH₄ level in urban areas is strongly influenced by the size of the urban agglomeration as well as by the meteorological parameters. These results are in agreement with other results from scientific literatures.

4. CO in urban areas

4.1 Trends of CO variation in urban areas. A literature review

According to World Health Organization data (WHO, 2000), in the main European cities the average atmospheric CO concentration is situated under 2 mg/m³ air, with a maximum concentration lower than 6.0 mg/m³ air. At global level, the concentration of CO is composed between 0.05 and 0.12 ppm in the air. This concentration is an average between the values measured in urban and rural areas. In rural areas the CO concentration is due mainly to natural processes, but in urban areas it is strongly influenced by anthropic activities.

The CO concentration in the air of urban zones depends upon the density of combustion sources, the topography of the measurements location, the meteorological conditions and from the distance between the measurement point and the auto traffic routes.

The monitoring of CO in USA is carried out since 1980; currently, there are 243 measurement stations distributed all over the USA territory. According to EPA Reports (EPA, 2009), the concentration of CO decreased in the period 1980-2006 from 14 ppm to 3 ppm.

In Europe, the monitoring of CO in urban areas is 20 years old. Several European projects were in action, to evaluate the exposure of the population to CO, and the measurements were carried out both with fixed and mobile stations. According to WHO data (WHO, 2000) in large European cities the CO concentrations (during 8 hours measurements) are situated below 20 mg/m³ in the air, and the maxima are not higher than 10 mg/m³ in the air.

The first network for the measurement of pollutants resulted from anthropic activities was created in France in 1979 under the name AIRPARIF and measures the daily, monthly and annual concentrations of NO_x, SO₂, O₃, PM, CO and of some organic compounds. According to this source, the CO concentration in the Paris region decreased from 4000 µg/m³ air in 1994 to 1200 µg/m³ air in 2006. In 2004 started measuring background measurements. The variation of annual average decreased from 500 µg/m³ air in 2003 to 400 µg/m³ air in 2006.

In Great Britain, the measurement of the concentrations of atmospheric pollutants dates from 1973; currently, there are more than 100 stations in urban zones for continuous monitoring of the air quality parameters. In London, the quality of air is monitored by as many as 30 stations. The network was created in 1993 under the name London Air Quality Network (LAQN), and since 1997 this network also measures the evolution of daily CO concentrations.

At the European level functions the European Environment Agency (EEA) with 32 members: all the 27 EU member countries, also Island, Liechtenstein, Norway, Switzerland and Turkey. Under the coordination of EEA was created the European Environment Information and Observation Network (EIONET), with the role of processing and validating the data from the stations of the member countries connected to this network. The information is available as Reports to interested users. Among the workstations connected to EIONET, 163 measure the concentrations of CO. According to EEA data, the concentration of CO at the European level decreased from 1 mg/m³ air in 1995 to 0.5 mg/m³ air in 2005.

In addition to the data from the monitoring stations there are numerous studies about the determination of CO concentrations in urban zones all over the world. A synthesis of these results is given in Table 4.

| Authors | Place of measurements | Period measurement | CO range concentrations [ppm] |
|--------------------------|-----------------------|--------------------|--|
| Chatterton T. et al. | Norwich, UK | 1997 - 1998 | 0.4 - 10.9 |
| Chelani A.B. et al. | Delhi, India | 2000 - 2003 | 1.66 - 8.4 |
| Cheng C. S. et al. | Canada | 1974 - 2000 | Montreal: 0.5 - 2.1, Toronto: 0.7 - 3.7 |
| Corti A. et al. | Salerno, Italy | Not specified | 0.55 - 0.85 |
| DEQ-Oregon | Portland-SUA | 1980 - 1998 | 13.0 - 4.7 |
| Emmerson K. et al. | Birmingham, UK | 1999 - 2000 | 0.17 - 0.66 |
| EPA | USA | 1990 - 2006 | Washington (decrease from 9 to 2), New-York (8.6 - 1.8), Los Angeles (14 - 4) , |
| Ghauri B. et al. | Pakistan | 2003 - 2004 | Islamabad : 6 - 13; Quetta: 1.9 - 14 , Karachi: 1.6 - 8.0; Lahore: 1.3 - 12; Rawalpindi: 1.6 - 8 |
| Ghose M. K. et al. | Calcutta, India | 2003 | 2.6 - 5.1 |
| Jones S. G. et al. | Paris, France | 1997 | 0.38 - 1.45 |
| Kim S.Y. et al. | Seoul , Korea | 2002 | 0.8 - 44.0 |
| Kukkonen J. et al. | Helsinki, Sweden | 1997 | 0.1 - 4.5 |
| Lijteroff R. et al. | San Luis, Argentina | 1994 - 1995 | 3.43 - 9.17 |
| Linden J. et al. | Burkina Faso, Africa | 2004 - 2005 | Background: 1 - 9, Traffic: 6.5 - 6.0 |
| Makra L. et al. | Szeged, Hungary | 1997 - 2001 | 0.24 - 0.93 |
| Manning A.J. et al. | Leek, UK | 1997 | 0.25 - 4.0 |
| Martín M.L. et al. | Bay of Algeciras | 1999 - 2001 | 0.5 - 2.9 |
| Milton R. et al. | Londra, UK | 2004 - 2005 | 0.9 - 14.9 |
| Muttamara S. et al. | Bangkok | 1997 | 8.23 - 26.89 |
| Ni-Bin Chang et al. | Kaohsiung, Taiwan | 1995 | 0.1 - 2.0 |
| Park S. S. et al. | Seoul, Korea | 1998 - 1999 | 1.74 - 2.81 |
| Reich S. et al. | Buenos Aires | 2001 | 0.60 - 2.44 |
| Rubio M. A. et al. | Santiago City, Chile | 2005 - 2006 | 0.31 - 3.06 |
| Sanchez-Coyllo O. et al. | Sao-Paulo, Brazilia | 1999 | 1.20 - 4.00 |
| Sathitkunarath S. et al. | Chiang Mai, China | 2002 | 0.9 - 1.5 |
| Shiva Nagendra S.M. | Delhi, India | 1997 - 1999 | 0.1 - 18 |
| Turias I.J. et al. | Campo de Gibraltar | 1999 - 2001 | 0.4 - 4.5 |
| Venegas L.E. et al. | Buenos Aires | 1994 - 1996 | Autumn:10,0, Winter: 9.80, Spring :10,7 |

Table 4. Overview of urban CO measurements

4.2 Case study Cluj-Napoca city

In Romania, the monitoring of air quality is done by the Agencies for Environment Protection and follows the concentrations of nitrogen oxides, sulfur dioxide, ozone, BTEX, material particles, etc. Of these, 53 monitoring stations are connected to the European EIONET System and 12 stations also measure the concentrations of CO in urban zones.

In the city of Cluj-Napoca, there are four stations for continuous monitoring of air quality, and beginning with August 2005 the Agency measures the CO concentrations in two locations in the city of Cluj-Napoca and one in the city of Dej. The measurements in Cluj county show that the concentration of CO in the atmosphere is much below the admitted level (10 mg/m^3) and varies between 0.09 and 0.4 mg/m^3 air.

In this case study the measurements were performed daily in Cluj-Napoca at the astronomic midday (in Romania at 12.30 h) using a NDIR CO analyzer Horiba model APMA-360. The results showed that the CO level in Cluj-Napoca is less than 1 mg/m^3 with a tendency of accumulation during the winter season. It is also observed a trend of accumulation during the spring months, due to the lack of precipitations. Beginning with the end of May, when the rain regime becomes normal, the values of CO concentrations are around 0.1 mg/m^3 air (Figure 6).




Fig. 6. Variation of CO concentrations in Cluj-Napoca in the period July 2008-June 2009

5. Anthropogenic contribution in CO₂ and CH₄ budgets

5.1 Isotopic ¹³C CO₂ measurements

Knowledge of the terrestrial CO₂ cycle will help to understand the climate change phenomena and to predicting the future atmospheric CO₂ concentrations and global temperatures. By estimating the terrestrial CO₂ cycle, including such factors as emissions, storages and fluxes and combining this with the isotope compositions of atmospheric CO₂ will help to identify the contribution of different factors to the atmospheric CO₂ budget. More than the observation of the CO₂ isotopic composition provides important information about sources such as fossil fuel combustion and biogenic respiration. The determination of isotopic concentrations of ¹³C enforces the analyze of different species of atmospheric CO₂ and CH₄ collected in situ in glass recipients of different measures (flask sampling) by mass spectrometry. Thus, Takahashi et al. (2001, 2002) using CO₂ isotope compositions method

for investigating the sources of atmospheric CO₂ observed carbon isotope compositions of $\Delta^{14}\text{C}$ and $\delta^{13}\text{C}$ in atmospheric CO₂ and estimated the contributions of fossil fuels and biogenic respiration, while Pataki (Pataki et al., 2003, 2006a,b) observing $\delta^{13}\text{C}$ and $\delta^{18}\text{O}$ isotope compositions in atmospheric CO₂ has reported the contribution of natural gas combustion, gasoline combustion and biogenic respiration over the total atmospheric CO₂ budget.

To identify sources of carbon and to quantify these sources it is used the Keeling plot (Pataki et al., 2003). The equation used in the Keeling plot is derived from the basic assumption that the atmospheric concentration of a substance (CO₂, CH₄) in air reflects the combination of some background amount of the substance that is already present in the atmosphere and some amount of substance that is added or removed by sources or sinks:

$$C_T = C_A + C_S \quad (1)$$

where C_T , C_A , and C_S are the concentrations of the substance in air, in the background of atmosphere, and that contributed by sources, respectively. Isotope ratios of these different components can be expressed by a simple mass balance equation:

$$C_T \delta_T = C_A \delta_A + C_S \delta_S \quad (2)$$

where δ_T , δ_A , and δ_S represent the isotopic composition of the substance in the atmosphere, in the background, and of the sources, respectively. By combine Eqs. 1 and 2 it is obtained equation 3:

$$\delta_T = C_A (\delta_A - \delta_S) / C_T + \delta_S \quad (3)$$

This is a linear relationship with a slope of $C_A (\delta_A - \delta_S)$ and an intercept at the δ_S value of the net sources/sinks in the atmosphere.

According to Pataki (Pataki et al., 2006) the mixing ratios originating from local sources (C_S) is composed from the CO₂ mixing of natural gas combustion (C_N) and CO₂ mixing of gasoline combustion (C_G):

$$C_T = C_A + C_N + C_G \quad (4)$$

Using the values of measured concentrations of CO₂ (C_T) in the same time and in the same air as the measurements of δ_T , and know the δ_N , δ_G and C_N it is possible to estimate C_G . The isotopic mass balance equation in this case is:

$$\delta_T C_T = \delta_B C_B + \delta_N C_N + \delta_G C_G \quad (5)$$

where δ_T and δ_G represent the isotopic composition of the CO₂ results from natural gas combustion and δ_G is the isotopic composition of the CO₂ results from gasoline combustion. In order to study the effects of the emission and diffusion of CO₂ from fossil fuel combustion most of the recent studies have focused on urban areas where CO₂ is mainly the product of power plants and transportation. The results of these measurements were correlated with variations in carbon isotopic composition and its show that while the natural level of $\delta^{13}\text{C}$ value is -8.02‰ , in urban areas the $\delta^{13}\text{C}$ values is down to -12‰ for atmospheric CO₂. This difference is given by the increasing input of CO₂ derived from fossil fuel (Clark- Thorne and Yapp, 2003, Lichtfouse et al., 2003; Widory and Javoy, 2003, Newman S et al. 2008, Wada et al 2010).

5.2 Isotopic ^{13}C measurements

The carbon isotopic composition (^{12}C , ^{13}C and ^{14}C) of atmospheric methane is used to estimate the local CH_4 sources contribution over the CH_4 budget in a local areas (Moriizumi et al., 1998)

According to (Miller et al., 2003 cited by Cuna et al. 2008) the methane mixing ratio in air, $[\text{CH}_4]$, and its isotopic ratio, $\delta^{13}\text{C}_{\text{CH}_4}$, may be derived from three main sources: methane produced by microbial, $[\text{CH}_4]_{\text{micr}}$, fossil methane, $[\text{CH}_4]_{\text{ff}}$, and methane produced from biomass burning, $[\text{CH}_4]_{\text{bmb}}$

$$[\text{CH}_4]_{\text{T}} = [\text{CH}_4]_{\text{micr}} + [\text{CH}_4]_{\text{ff}} + [\text{CH}_4]_{\text{bmb}} + [\text{CH}_4]_{\text{bg}} \quad (1)$$

In equation (1) $[\text{CH}_4]_{\text{bg}}$ is defined as the smoothed marine boundary layer (MBL) at the latitude of interest (Dlugokencky et al., 1994 cited by Cuna et al. 2008).

Each of these emissions has a more-or-less distinct isotopic signature with bacterial methane $\delta^{13}\text{C}_{\text{micr}} \approx 60\text{‰}$, thermogenic methane $\delta^{13}\text{C}_{\text{ff}} \approx 40\text{‰}$, and biomass burning methane $\delta^{13}\text{C}_{\text{bmb}} \approx 25\text{‰}$ (Quay et al., 1999 cited by Cuna et al. 2008).

$$\delta^{13}\text{C}[\text{CH}_4]_{\text{T}} = \delta^{13}\text{C}[\text{CH}_4]_{\text{micr}} + \delta^{13}\text{C}[\text{CH}_4]_{\text{ff}} + \delta^{13}\text{C}[\text{CH}_4]_{\text{bmb}} + \delta^{13}\text{C}[\text{CH}_4]_{\text{bg}} \quad (2)$$

Separating CH_4 sources using isotopic signatures is complicated by enrichment during uptake processes such as bacterial CH_4 oxidation, or methanotrophy (Chanton et al., 2005 cited by Cuna et al. 2008). Thus, the methane mixing ratio changes over time according to Eq. (3), where $[\text{CH}_4]_{\text{S}}$ is the sum of all sources and τ is the lifetime of methane with respect to its destruction by OH and addition from other processes (Montzka et al., 2000; Hein et al., 1997 cited by Cuna et al. 2008):

$$d[\text{CH}_4] / dt = [\text{CH}_4]_{\text{S}} - ([\text{CH}_4] / \tau) \quad (3)$$

The $\delta^{13}\text{C}$ of methane measured in an air sample results from several different sources, such that

$$\delta^{13}\text{C}_{\text{S}} = \delta^{13}\text{C}_{\text{bg}} + \varepsilon \quad (4)$$

In Eq. (4) $\delta^{13}\text{C}_{\text{S}}$ is the flux-weighted isotopic ratio of all sources expressed in δ notation, $\delta^{13}\text{C}_{\text{bg}}$ is the atmospheric background isotopic ratio and ε is the average isotopic fractionation associated with these processes (Cantrell et al., 1990 cited by Cuna et al. 2008). Using the values of measured concentrations of methane $[\text{CH}_4]_{\text{T}}$ in the same time and in the same air as the measurements of $\delta^{13}\text{C}$, it is possible to estimate the contribution of all sources at the atmospheric methane budget. Thus, Moriizumi (Moriizumi et al., 1998) analyzing the CH_4 in Nagoya, Japan found that “the contribution of fossil CH_4 to local CH_4 released from the urban area was calculated to be $102 \pm 8\%$, and its $\delta^{13}\text{C}$ was $-40.8 \pm 3.0\text{‰}$. In a suburban area of Nagoya fossil, CH_4 contributed to less than 10% of local release and the calculated value of $\delta^{13}\text{C}$ for non-fossil CH_4 was approximately -65‰ , which is within the range of reported values of $\delta^{13}\text{C}$ for CH_4 derived from bacterial CH_4 sources such as irrigated rice paddies”. Kuc (Kuc et al. 2003), measuring the CH_4 in Krakow found that “The linear regression of $\delta^{13}\text{C}$ values of methane plotted versus reciprocal concentration yields the average $\delta^{13}\text{C}$ signature of the local source of methane as being equal to -54.2‰ . This value agrees very well with the measured isotope signature of natural gas being used in Krakow ($-54.4 \pm 0.6\text{‰}$) and points to leakages in the distribution network of this gas as the main anthropogenic source of CH_4 in the local atmosphere”. Nakagawa (Nakagawa et al., 2005), using the stable carbon and hydrogen isotopic compositions ($\delta^{13}\text{C}$

and δD) of methane quantified the contribution of automobile exhaust to local CH₄ budget. The authors estimated that for local sources, automobile exhaust in Nagoya, Japan, contribute significant amounts (up to 30%) of CH₄ to the troposphere in the studied area. Studies performed in wetlands showed that the isotopic signature $\delta^{13}C$ of methane is situated between -67.4 and -53.3‰ with lower values in the summer and higher values in the winter (Cuna et al., 2008; Tarasova et. al., 2006). These values confirm that in the wetlands, the biogenic CH₄ is the main source of atmospheric CH₄.

5.3 Isotopic ¹³CO₂ measurements in Cluj county. Case study

In order to study the role of CO₂ resulted from anthropic activities in the urban atmosphere of Cluj county, the variation of CO₂ concentrations and the corresponding $\delta^{13}C$ values, in samples collected in the three areas (Cluj-Napoca, Turda, Huedin) were measured (by flask sampling) during a whole year (July 2008-June 2009). For each area two points of measurements were selected, one in the city and one reference outside of the city. The determination of CO₂ concentrations was done with an infrared gas analyzer, and the isotopic ratios were measured with a DELTA V Advantage, Thermo Finnigan mass spectrometer. A graphic representation of the isotopic ratios as a function of 1/ [CO₂] gives a Keeling plot and the value of the intercept of Keeling slope provide information about the isotopic signature of the source. Depending on the climatic conditions and the size of the urban agglomeration, correlations between $\delta^{13}C$ values and corresponding CO₂ concentrations were between -11 ‰ for Cluj-Napoca, -10.0‰ for Turda and -9.0‰ for Huedin (Tables 5-7). If we consider $\delta^{13}C = -8‰$ the isotopic composition of natural CO₂, the anthropogenic contribution for CO₂ budget is higher for Cluj-Napoca and near the natural level for small town Huedin. As the data in tables show, for the locations in the interior of the cities, the isotopic values are displaced from the average values by 0.5-1.5 ‰ compared with the reference points, which suggests that the CO₂ source in the urban location is composed from the zone with $\delta^{13}C = -8‰$, with clean air, and an anthropic source which can be CO₂ resulted from burning fossil fuels (mainly gasoline and methane) plus a biogenic source of CO₂ resulted from the respiration of the local vegetation. The largest difference occurred in the municipality of Cluj-Napoca (1.376), where the average values of $\delta^{13}C = -$

| Measurement month | Time | City Point (Mărăști) | | Reference Point | |
|-------------------|-------|-----------------------|--------------------|-----------------------|--------------------|
| | | CO ₂ (ppm) | $\delta^{13}C$ (‰) | CO ₂ (ppm) | $\delta^{13}C$ (‰) |
| July 2008 | 12.30 | 386 | -8.773 | 373 | -8.670 |
| August 2008 | 12.30 | 433 | -10.529 | 409 | -8.928 |
| September 2008 | 12.30 | 435 | -10.683 | 400 | -8.901 |
| October 2008 | 12.30 | 530 | -10.926 | 416 | -8.936 |
| November 2008 | 12.30 | 526 | -10.836 | 444 | -8.940 |
| January 2009 | 12.30 | 538 | -10.200 | 450 | -8.949 |
| February 2009 | 12.30 | 773 | -11.012 | 438 | -8.939 |
| March 2009 | 12.30 | 665 | -10.543 | 455 | -8.943 |
| April 2009 | 12.30 | 682 | -10.657 | 428 | -8.921 |
| May 2009 | 12.30 | 458 | -9.542 | 420 | -8.901 |
| June 2009 | 12.30 | 425 | -9.230 | 395 | -8.763 |

Table 5. Values of CO₂ (ppm) and $\delta^{13}C_{PDB}$ (‰) in Cluj-Napoca city

| Measurement month | Time | City Point (Potaisa) | | Reference Point | |
|-------------------|-------|-----------------------|-----------------------|-----------------------|-----------------------|
| | | CO ₂ (ppm) | δ ¹³ C (‰) | CO ₂ (ppm) | δ ¹³ C (‰) |
| July 2008 | 12.30 | 380 | -8.797 | 371 | -8.720 |
| August 2008 | 12.30 | 403 | -8.826 | 382 | -8.762 |
| September 2008 | 12.30 | 416 | -8.802 | 391 | -8.851 |
| October 2008 | 12.30 | 450 | -8.973 | 406 | -8.878 |
| November 2008 | 12.30 | 502 | -10.620 | 450 | -8.953 |
| January 2009 | 12.30 | 463 | -9.274 | 425 | -8.900 |
| February 2009 | 12.30 | 483 | -9.560 | 450 | -8.940 |
| March 2009 | 12.30 | 450 | -9.200 | 420 | -8.760 |
| April 2009 | 12.30 | 490 | -10.146 | 435 | -9.120 |
| May 2009 | 12.30 | 399 | -8.870 | 371 | -8.832 |
| June 2009 | 12.30 | 425 | -9.132 | 355 | -8.900 |

Table 6. Values of CO₂ (ppm) and δ¹³C_{PDB} (‰) in Turda city

| Measurement month | Time | City Point (O.Goga) | | Reference Point | |
|-------------------|-------|-----------------------|-----------------------|-----------------------|-----------------------|
| | | CO ₂ (ppm) | δ ¹³ C (‰) | CO ₂ (ppm) | δ ¹³ C (‰) |
| July 2008 | 12.30 | 376 | -8.180 | 370 | -8.168 |
| August 2008 | 12.30 | 408 | -8.438 | 379 | -8.187 |
| September 2008 | 12.30 | 387 | -8.175 | 379 | -8.141 |
| October 2008 | 12.30 | 405 | -8.382 | 376 | -8.138 |
| November 2008 | 12.30 | 404 | -8.390 | 387 | -8.221 |
| January 2009 | 12.30 | 422 | -9.455 | 411 | -8.324 |
| February 2009 | 12.30 | 416 | -9.342 | 385 | -8.122 |
| March 2009 | 12.30 | 406 | -9.142 | 400 | -8.786 |
| April 2009 | 12.30 | 409 | -10.620 | 390 | -9.200 |
| May 2009 | 12.30 | 402 | -8.761 | 379 | -8.100 |
| June 2009 | 12.30 | 355 | -8.956 | 350 | -8.212 |

Table 7. Values of CO₂ (ppm) and δ¹³C_{PDB} (‰) in Huedin town

10.266‰ were obtained in the city center, compared with the δ¹³C = -8.890‰ for the reference point. For the other locations studied (Turda and Huedin) the isotopic concentrations are close to the atmospheric background, namely -9,291‰ for Turda and -8,895‰ for Huedin, comparable with the average values for the reference points (-8.874‰ for Turda and -8.327‰ and Huedin) suggesting that anthropic CO₂ is not contributing to the pollution.

5.4 Isotopic ¹³CH₄ measurements in Cluj county. Case study

For the evaluation of the role of CH₄ resulted from anthropic activities in the urban atmosphere in the Cluj county, the variation of CH₄ concentrations and the corresponding δ¹³C values were measured in air samples collected in three areas: Cluj-Napoca (Piața

Mărăști), Turda (Potaisa School) and Huedin (O. Goga High School) during the period between January-June 2009, using the flask sampling. Again, the $\delta^{13}\text{C}$ values were measured with a ThermoFinnigan Delta V Advantage mass spectrometer. The methane concentrations in the same samples were measured with a gas chromatograph equipped with a FID detector. The observed variation of methane concentrations (Table 8) is rather large, between 4.7 and 11.5 ppm. The values measured are above the atmospheric background, which suggests that there is an anthropic source of CH₄ in all investigated areas. The average value of $\delta^{13}\text{C} = -40$ ‰ suggests that the source of methane in the atmosphere is the gas fuel network of the urbane zone investigated.

| Measurement month | Cluj-Napoca | | Turda | | Huedin | |
|-------------------|-----------------------|---------------------------|-----------------------|---------------------------|-----------------------|---------------------------|
| | CH ₄ (ppm) | $\delta^{13}\text{C}$ (‰) | CH ₄ (ppm) | $\delta^{13}\text{C}$ (‰) | CH ₄ (ppm) | $\delta^{13}\text{C}$ (‰) |
| January 2009 | 11.5 | -39.21 | 10.4 | -39.45 | 10.6 | -39.78 |
| March 2009 | 8.0 | -37.90 | 6.3 | -38.75 | 5.5 | -38.97 |
| April 2009 | 11.5 | -39.80 | 11.0 | -38.92 | 10.5 | -38.98 |
| May 2009 | 8.4 | -38.02 | 7.4 | -38.80 | 4.7 | -38.85 |

Table 8. Simultaneous value of CH₄ (ppm) and $\delta^{13}\text{C}$ (‰) in the Cluj district.

6. Correlation between CO₂ trend variation in urban area and variation of meteorological parameters

6.1 Case study Cluj-Napoca city

For this study a daily measurements of the CO₂ concentration and the main meteorological parameters (temperature, relative humidity and wind velocity) were recorded for a whole calendar year, beginning with July 2008 until June 2009. The measurements were carried out in the centre of Cluj-Napoca city, the time of midday (12.30), at the selected latitude. The results of measurements revealed a daily variation of CO₂ concentrations correlated with the meteorological factors and biological cycles of plants. Thus, the largest values of CO₂ were recorded in the fall and winter in the absence of vegetation, and the lowest values in the summer months, when the biologic cycle of plants is at the maximum.

The graphic representation of the CO₂ values as a function of meteorological parameters indicates a direct correlation with temperature and an inverse correlation with the wind velocity and relative humidity.

A computation of linear regression slopes of CO₂ versus air temperature for two months from winter (January and February) and two months from summer (July and August) gives positive slopes for both seasons with a highest correlation factor (0.666) in winter in the absence of photosynthesis (figure 7).

The same representation for CO₂ and for relative humidity shows a negative slope in summer and a positive slope in the winter. The computation of linear regression slopes for CO₂ versus wind velocity show a negative slopes both in summer and in winter (figure 7).

The results of this case study show that in urban area it is difficult to estimate by correlation coefficient analyses the influence of meteorological parameters over the CO₂ variation. In order to correlate the variation of CO₂ concentrations with the variation of meteorological parameters a statistic analysis of data is necessary. For the statistical approach the regression analysis and principal component analysis (PCA) has been used.




Fig. 7. Correlation between CO₂ variation and meteorological parameters in two seasons

6.1.1 The regression analysis

The regression analysis was performed with the aid of Curve estimation model of SPSS statistics program and the regression coefficients were calculated, having the CO₂ concentration as independent variable and the meteorological parameters as dependent variables. For analysis the squares of regression coefficients and the regression curves were used. The regression coefficients R² were computed for the most frequently used types of regression, namely linear, logarithmic, polynomial and exponential (Table 8).

The analysis of regression coefficients leads to some important conclusions.

The only meteorological parameter which correlated with the CO₂ concentration over the 0.6 threshold is the air temperature. Both the air humidity and the wind velocity have very low regression coefficients, suggesting that there is a low probability that the variation of the CO₂ is influenced by these meteorological factors. However, there are singular situations, when the correlation coefficients are close to the 0.6 threshold. Thus, for the relative

| Month | Rt | Ta °C | RH % | V m/s |
|----------------|----|-------|------|-------|
| I (January) | Li | .842 | .042 | .014 |
| | Lo | .843 | .040 | .013 |
| | Po | . | .040 | . |
| | Ex | . | .041 | . |
| II (February) | Li | .802 | .000 | .426 |
| | Lo | .789 | .000 | .430 |
| | Po | . | .001 | . |
| | Ex | . | .000 | . |
| III (March) | Li | .461 | .011 | .166 |
| | Lo | .456 | .011 | .168 |
| | Po | .550 | .003 | .090 |
| | Ex | .549 | .003 | .087 |
| IV (April) | Li | .392 | .034 | .064 |
| | Lo | .392 | .033 | .064 |
| | Po | .368 | .037 | .012 |
| | Ex | .368 | .039 | .012 |
| V (May) | Li | .750 | .462 | .546 |
| | Lo | .741 | .462 | .544 |
| | Po | .865 | .454 | .380 |
| | Ex | .869 | .455 | .381 |
| VI (June) | Li | .632 | .485 | .000 |
| | Lo | .632 | .489 | .000 |
| | Po | .673 | .520 | .020 |
| | Ex | .673 | .516 | .019 |
| VII (July) | Li | .117 | .158 | .003 |
| | Lo | .121 | .159 | .004 |
| | Po | .116 | .161 | .019 |
| | Ex | .113 | .160 | .018 |
| VIII (August) | Li | .028 | .271 | .088 |
| | Lo | .026 | .278 | .082 |
| | Po | .025 | .166 | .120 |
| | Ex | .027 | .160 | .126 |
| IX (September) | Li | .368 | .033 | .337 |
| | Lo | .376 | .035 | .342 |
| | Po | .481 | .004 | .287 |
| | Ex | .471 | .003 | .283 |

| | | | | |
|----------------|----|------|------|------|
| X (October) | Li | .121 | .045 | .088 |
| | Lo | .123 | .047 | .092 |
| | Po | .206 | . | .065 |
| | Ex | .202 | . | .061 |
| XI (November) | Li | .270 | .189 | .143 |
| | Lo | .269 | .180 | .144 |
| | Po | . | .152 | . |
| | Ex | . | .160 | . |
| XII (December) | Li | .571 | .172 | .028 |
| | Lo | .581 | .176 | .027 |
| | Po | .667 | .196 | .024 |
| | Ex | .656 | .193 | .023 |

Table 9. The Estimation of regression coefficients (R^2). Note: The types of regression R are: Li - linear, Lo - logarithmic, Po - polynomial, Ex - exponential

humidity the highest regression coefficient was 0.52, for the type polynomial in June 2009. For the wind velocity the largest coefficient was 0.56 in May 2009, for the linear regression. The regression between CO_2 concentration and temperature reveals two interesting aspects. The 0.6 threshold of the correlation coefficient was overfull filled in the three months of winter, at the end of spring and beginning of the summer, in May and June. The lowest value of the correlation coefficient was observed in August when the measurements were made at high temperatures over 30 Celsius degree. Large values, above 0.8 were observed in January and May. To illustrate the correlations we present in table 9 all the situations when the correlation coefficient was higher than 0.6.

6.1.2 The analysis of main components (PCA)

This type of analysis was necessary, because we wanted to see, which is the weight of meteorological parameters and CO_2 concentrations in the explanation of total variations. The PCA (Principal Component Analysis) method without factor rotation was used and the results are shown in Table 10.

| Component | Initial Eigenvalues | | | Extraction Sums of Squared Loadings | | |
|-----------|---------------------|---------------|--------------|-------------------------------------|---------------|--------------|
| | Total | % of Variance | Cumulative % | Total | % of Variance | Cumulative % |
| 1 | 1.760 | 44.002 | 44.002 | 1.760 | 44.002 | 44.002 |
| 2 | 1.066 | 26.646 | 70.648 | 1.066 | 26.646 | 70.648 |
| 3 | .900 | 22.491 | 93.139 | .900 | 22.491 | 93.139 |
| 4 | .274 | 6.861 | 100.000 | | | |

Table 10. Explanation of total variations. Extraction Method: Principal Component Analysis

After the value of 0.5 was selected for the Eigenvalue, three components were extracted which together explain 93.139% of the total variation. The first component explains 44.002 %

of the variation, the second 26.649 and the third 22.5 %. The difference to 100 % is due to a fourth component, which was eliminated because it had an Eigenvalue of only 0.274. The matrix of components (Table 11) allows the identification of each factor. Thus, the first factor is very well correlated with the air temperature (inverse correlation) and with the air humidity (direct correlation). The second component is very well correlated with the wind velocity. The concentration of CO₂ is very well correlated with the third component.

| | Component | | |
|-----------------|-----------|-------|-------|
| | 1 | 2 | 3 |
| CO ₂ | .480 | -.283 | .824 |
| Ta | -.897 | -.245 | .045 |
| V | -.045 | .961 | .234 |
| RH | .850 | -.048 | -.406 |

Table 11. Matrix of components

7. Conclusions

The monitoring of the CO₂ levels in urban area could estimate the contribution of anthropic activities over the global CO₂ level. This contribution is essential in order to establish the presence of CO₂ dome over the cities. The results of presented case study confirm the presence of a CO₂ dome over the urban studied area. More than that, it confirms that the anthropogenic CO₂ emissions are the primary source of the urban CO₂. Taking into account the obtained results it can be observed that the level of CO₂ in urban areas is influenced by the size of the city and by the amplitude of anthropic activities. Thus the highest values of CO₂ were obtained in the biggest city Cluj-Napoca (between 380 and 530 ppm at Mărăști Square, 376-456 at Grigorescu and 373-444 at reference point) followed by Turda (380-502 ppm at Potaisa School and 371-450 ppm at the reference point) and Huedin (355- 433 ppm at O. Goga High School and 350-411 ppm at the reference point. It is also observed that the concentration of urban CO₂ has an annual variation with the lower value in the summer and the highest value in the autumn and spring. Regarding the daily CO₂ variation it also observed that it is dominated by the photosynthesis.

The results of the atmospheric methane measurements show a significant variation depending on the season and the urban agglomeration degree. Thus, the methane concentrations in the three investigated areas indicate a similar profile for the measurements carried out in the cities, with minima in the summer months and maxima during the spring. The highest values were recorded in the city of Cluj-Napoca (11.5 ppm in April 2009) and the lowest values were recorded in the month of August 2008 (2.5 ppm). In Turda the concentrations of atmospheric methane were comprised between 2.2 and 8.6 ppm, with the lowest values measured in August 2008 (2.2 ppm) and the highest values recorded in April 2009 (8.6 ppm) while in Huedin the concentrations of methane varied between 1.4 and 7 ppm, with the lowest values recorded in July 2008 (1.4 ppm) and the highest ones in April 2009 (7 ppm). Significant differences are also observed between the methane concentrations in the interior of the cities and the reference points located outside. These differences prove that the anthropic activities, in particular the automobile traffic, are an important source of methane in the urban atmosphere.

The carbon isotopic composition measurement of CO₂ and CH₄ is the best way to establish the biogenic and anthropic contribution at CO₂ and CH₄ budget in urban areas. Regarding the case study performed in three Romanian cities the results of ¹³CO₂ show that the value of δ¹³C is depending on the size of the urban agglomeration. Thus, the lower value -11 ‰ were obtained for Cluj-Napoca, followed by Turda (-10.0‰) and Huedin (-9.0‰). If we consider δ¹³C = -8‰ the isotopic composition of natural CO₂, the anthropogenic contribution for CO₂ budget is higher for Cluj-Napoca and near the natural level for small town Huedin. More than, the recorded data show a difference of 0.5-1.5 ‰ between the measurements city points and reference points which suggests that the CO₂ source in the urban location is composed from the zone with δ¹³C = -8‰, with clean air, and an anthropic source which can be CO₂ resulted from burning fossil fuels (mainly gasoline and methane) plus a biogenic source of CO₂ resulted from the respiration of the local vegetation. The largest difference occurred in the municipality of Cluj-Napoca (1.376), where the average values of δ¹³C = -10.266‰ were obtained in the city center, compared with the δ¹³C = -8.890‰ for the reference point. For the other locations studied (Turda and Huedin) the isotopic concentrations are close to the atmospheric background, namely -9,291‰ for Turda and -8,895‰ for Huedin, comparable with the average values for the reference points (-8.874 for Turda and -8.327 and Huedin).

Regarding the ¹³CH₄ measurements the results obtained in the case study are above -40 ‰ which suggests that there is an anthropic source of CH₄ in all investigated areas. We think that these values are a consequence of methane resulted from gas fuel network of the urbane investigated areas.

Regarding the correlation between CO₂ variation in urban area and variation of meteorological parameters the results of the case study indicate a direct correlation of CO₂ level with temperature and an inverse correlation with the wind velocity and relative humidity. Although, these correlations are poor and the analysis of regression coefficients showed that only the air temperature is correlated with the CO₂ concentration over the 0.6 threshold. The 0.6 threshold of the correlation coefficient was overfull filed in the three months of winter, at the end of spring and beginning of the summer, in May and June. The lowest value of the correlation coefficient was observed in August when the measurements were made at high temperatures, over 30 degrees Celsius. Large values, above 0.8 were observed in January and May. The air humidity and the wind velocity have very low regression coefficients, suggesting that in the urban areas studied, there is a low probability that the variation of the CO₂ is influenced by these meteorological factors. Thus, for the relative humidity the highest regression coefficient was 0.52, for the type polynomial in June 2009 while for the wind velocity the largest coefficient was 0.56 in May 2009, for the linear regression.

Taking into account the results obtained, the present case study shows that the variation of CO₂ in urban area is in agreement with other results reported in the scientific literature. Thus, according to the carbon isotopic composition measurements of CO₂ the anthropogenic CO₂ emissions are the primary source of the urban CO₂ dome; the dome is generally stronger in city centers, in winter, under conditions of heavy traffic, with little or no wind, and in the presence of strong temperature inversions.

8. Acknowledgements

The case study has been supported by the Romanian National Authority for Scientific Research, Project no. 213-1/2007

9. References

- Aikawa, M.; Hiraki, T. & Eiho, J. (2006). Vertical atmospheric structure estimated by heat island intensity and temporal variations of methane concentrations in ambient air in an urban area in Japan. *Atmospheric Environment*, Vol. 40, Iss. 23, pp. 4308-4315.
- AIRPARIF: Available from: <http://www.airparif.asso.fr/index.php?setlang=en>
- Aubinet, M.; Heinesch, B. & Longdoz, B. 2002. Estimation of the carbon sequestration by a heterogeneous forest: night flux corrections, heterogeneity of the site and inter-annual variability. *Global Change Biology*, Vol. 8, pp. 1053-1071.
- Balling Jr.; R.C., Cervený, R.S. & Idso, C. D. (2001). Does the urban CO₂ dome of Phoenix, Arizona contribute to its heat island? *Geophysical Research Letters*, Vol. 28, pp. 4599-4601.
- Bergeron, O. & Strachan I.B. (2011). CO₂ sources and sinks in urban and suburban areas of a northern mid-latitude city. *Atmospheric Environment*, Vol. 45, pp. 1564-1573.
- CarboEurope, Available from: <http://carboeurope.org/>
- Chang, N.-B. & Tseng, C.C. (1999). Optimal design of a multi-pollutant air quality monitoring network in a metropolitan region using Kaohsiung, Taiwan as an example. *Environmental Monitoring and Assessment*, Vol. 57, pp. 121-148.
- Chantterton, T.; Dorling, S.; Lovett, A. & Stephenson, (2000). Air quality in Norwich, UK, Multi-scale modeling to assess the significance of city, county and regional pollution sources. *Environmental Monitoring and Assessment*, Vol. 65, pp. 425-433.
- Chelani, A.B. & Devotta, S. (2007). Air Quality Assessment in Delhi: Before and After CNG as fuel. *Environmental Monitoring and Assessment*, Vol. 125, pp. 257-263.
- Cheng, C.S.; Campbell, M.; Li, Q.; Li, G.; Auld, H.; Day, N.; Pengelly, D.; Gingrich, S. & Yap, D. (2007). A Synoptic Climatological Approach to Assess Climatic Impact on Air Quality in South-central Canada. Part I: Historical Analysis. *Water Air Soil Pollution*, Vol. 182, pp. 131-148.
- Clark-Thorne, S.T. & Yapp, J.C. (2003). Stable carbon isotope constraints on mixing and mass balance of CO₂ in an urban atmosphere: Dallas metropolitan area, Texas, USA. *Applied Geochemistry*, Vol. 18, pp. 75-95.
- Cofala, J.; Amann, M.; Klimont, Z.; Kupiainen, K. & Isaksson L.H. (2007). Scenarios of global anthropogenic emissions of air pollutants and methane until 2030. *Atmospheric Environment*, Vol. 41, Iss.38, pp. 8486-8499.
- Corti, A. & Senatore A. (2000). Project of an Air Quality Monitoring Network for Industrial Site in Italy. *Environmental Monitoring and Assessment*, Vol. 65, pp. 10-9117.
- Coutts A.M.; Beringer, J. & Tapper, N.J. (2007). Characteristics influencing the variability of urban CO₂ fluxes in Melbourne, Australia. *Atmospheric Environment*, Vol. 41, No.1, pp. 51-62.
- Cristea, G.; Haiduc, Iv.; Beldean-Galea, M.S. & Cuna, S. (2009). The variation of CH₄ in the urban areas from Cluj county. *Studia Universitatis Babeş-Bolyai, series Geographia*, Iss. 3, pp. 91-94
- CSIRO, Available from: <http://www.csiro.au/science/Climate-Change.html>
- Cuna, S.; Pendall, E.; Miller, J.B.; Tans, P.P.; Dlugokencky, E. & White, J.W.C. (2008). Separating contributions from natural and anthropogenic sources in atmospheric methane from the Black Sea region, Romania. *Applied Geochemistry*, Vol. 23, pp. 2871-2879.

- Daniel J.S. & Solomon S. (1998). On the climate forcing of carbon monoxide. *Journal of Geophysical Research-Atmospheres*, Vol. 103, pp. 13249-13260.
- Daniel, J.S. & Solomon, S. (1998). On the climate forcing of carbon monoxide. *Journal of Geophysical Research*, Vol. 103, No.D11, pp13249-13260: in IPCC-TAR, 2001, Chapter 4.
- Day, T.A.; Gober, P.; Xiong, F.S., & Wentz E.A. (2002). Temporal patterns in near-surface CO₂ concentration over contrasting vegetation types in the Phoenix metropolitan area. *Agricultural and Forest Meteorology*, Vol. 110, Nr. 3, pp. 229-245.
- DEQ-Oregon, 2007: Available from: <http://www.deq.state.or.us/aq/toxics/index.htm>
- Derwent, R.G.; Simmonds, P.G.; O'Doherty, S.; Stevenson, D.S.; Collins, W.J.; Sanderson, M.G.; Johnson, C.E.; Dentener, F.; Cofala, J.; Mechler, R. & Amann, M. (2006). External influences on Europe's air quality: Baseline methane, carbon monoxide and ozone from 1990 to 2030 at Mace Head, Ireland. *Atmospheric Environment*, Vol. 40, pp. 844-855.
- EEA: Available from: <http://www.eea.europa.eu>
- EIONET: Available from: <http://www.eionet.europa.eu>
- Emmerson, K.M.; Carslaw, N.; Carpenter, L.J.; Heard, D.E.; Lee, J.D. & Pilling, M. J.(2005) Urban Atmospheric Chemistry During the PUMA Campaign 1: Comparison of Modelled OH and HO₂ Concentrations with Measurements. *Journal of Atmospheric Chemistry*, Vol. 52, pp. 143-164.
- EPA, 2009: Available from: <http://www.epa.gov/oar/airtrends/carbon.html>
- Fuglestvedt, J.S.; Isaksen, I.S.A. & Wang, W.-C. (1996). Estimates of indirect global warming potentials for CH₄, CO, and NO_x. *Climate Change*, Vol. 34, pp. 405-437: in IPCC-TAR, 2001, Chapter 4.
- George, K.; Ziska, L.H.; Bunce, J.A. & Quebedeaux, B. (2007). Elevated atmospheric CO₂ concentration and temperature across an urban-rural transect. *Atmospheric Environment*, Vol. 41, No.35, pp.7654-7665.
- Ghauri, B.; Lodhi, A. & Mansha, M. (2007). Development of baseline (air quality) data in Pakistan. *Environmental Monitoring Assessment*, Vol. 127, pp. 237-252.
- Ghose, M.K.; Paul, R. & Banerjee, R. K. (2005). Assessment of the status of urban air pollution and its impact of human health in the city of Kolkata. *Environmental Monitoring and Assessment*, Vol. 108, pp. 151-167.
- Gratani, L. & Varone L. (2005). Daily and seasonal variation of CO₂ in the city of Rome in relationship with the traffic volume. *Atmospheric Environment*, Vol. 39, No.14, pp. 2619-2624.
- Grimmond, C.S.B.; King, T.S.; Cropley, F.D.; Nowakb, D.J., & Souch, C. (2002). Local-scale fluxes of carbon dioxide in urban environments: methodological challenges and results from Chicago, *Environmental Pollution*, Vol. 116, No.1, pp. S243-S254.
- Higuchi, K.; Shashkov, A.; Chan, D.; Saigusa, N.; Murayama, S.; Yamamoto, S.; Kondo, H.; Chen, J.; Liu, J. & Chen B. (2005). Simulations of seasonal and inter-annual variability of gross primary productivity at Takayama with BEPS ecosystem model. *Agricultural and Forest Meteorology*, Vol. 134, Iss.1-4, Pp. 143-150.
- Hsu, Y.K.; VanCuren, T.; Park, S.; Jakober, C.; Herner, J.; FitzGibbon, M.; Blake, D.R. & Parrish, D.D. (2010). Methane emissions inventory verification in southern California. *Atmospheric Environment*, Vol. 44, pp. 1-7.


- Idso, C.D.; Idso, S.B. & Balling, R.C. (2001). An intensive two-week study of an urban CO₂ dome. *Atmospheric Environment*, Vol. 35, Iss.6, pp. 995-1000.
- Idso, S.B.; Idso, C.D. & Balling, R.C. (2002). Seasonal and diurnal variations of near-surface atmospheric CO₂ concentrations within a residential sector of the urban CO₂ dome of Phoenix, AZ, USA. *Atmospheric Environment*, Vol. 36, pp. 1655-1660.
- IPCC-AR1 , 1992, Available from: http://www.ipcc.ch/ipccreports/far/IPCC_1990_and_1992_Assessments/English/ipcc-90-92-assessments-overview.pdf
- IPCC-AR4 , 2007, Available from: http://www.ipcc.ch/pdf/assessment-report/ar4/syr/ar4_syr.pdf
- IPCC-AR5 , 2010, Available from: <http://www.ipcc-wg2.gov/AR5/ar5.html>
- IPCC-SAR, 1995, Available from: <http://www.ipcc.ch/pdf/climate-changes-1995/ipcc-2nd-assessment/2nd-assessment-en.pdf>
- IPCC-TAR, 2001, Available from: http://www.grida.no/publications/other/ipcc_tar/
- Ito, A.; Takahashi, I.; Nagata, Y.; Chiba, K. & Haraguchi, H. (2000). The long-term evolutions and the regional characteristics of atmospheric methane concentrations in Nagoya, 1983–1997. *Science of the Total Environment*, vol. 263, pp. 37-45.
- Ito, A.; Saigusa, N.; Murayama, S.; Yamamoto, S. (2005). Modeling of gross and net carbon dioxide exchange over a cool-temperate deciduous broad-leaved forest in Japan: Analysis of seasonal and interannual change. *Agricultural and Forest Meteorology*, Vol. 134, Iss.1–4, pp. 122-134.
- Jacobson, M. Z. (2010). Enhancement of local air pollution by urban CO₂ domes. *Environmental Science and Technology*, Vol. 44, pp. 2497-2502.
- Johnson, C.E.; Collins, W.J.; Stevenson, D.S. & Derwent, R.G. (1999). The relative roles of climate and emissions changes on future oxidant concentrations. *Journal of Geophysical Research*, Vol 104, pp. 18631-18645; in IPCC-TAR, 2001, Chapter 4.
- Jones, S.G.; Fisher, B.E.A; Gonzales-Flesca, N. & Sokhi, R. (2000). The use of measurement programmes and models to assess concentrations next to major roads in urban areas. *Environmental Monitoring and Assessment*, Vol. 64, pp. 531-547.
- Kèlomé, N.C.; Lévêque, J.; Andreux, F.; Milloux, M.-J. & Oyédé, L.-M. (2006). C₄ plant isotopic composition ($\delta^{13}\text{C}$) evidence for urban CO₂ pollution in the city of Cotonou, Benin (West Africa). *Science of the Total Environment*, Vol. 366, pp. 439-447.
- Kim, S.-Y.; O'Neill, M.S.; Lee, J.-T.; Cho, Y.; Kim, J. & Kim, H. (2007). Totlul. *Intetnational Archives Occupational Environmental Health*, Vol. 80, pp. 701-710.
- Kuc, T.; Rozanski, K.; Zimnoch, M.; Necki, J. M. & Korus A. (2003). Anthropogenic emissions of CO₂ and CH₄ in an urban environment. *Applied Energy*, Vol. 75, Iss. 3-4, pp. 193-203.
- Kukkonen, J.; Valkonen, E.; Walden, J.; Koskentalo, T.; Karppinen, A.; Berkowicz, R. & Kartastenpaa, R. (2000). Measurements and modeling of air pollution in a street canyon in Helsinki. *Environmental Monitoring and Assessment*, Vol. 65, pp. 371-379.
- LAQN: Available from: <http://www.londonair.org.uk>
- Lichtfouse, E.; Lichtfouse, M. & Jaffrezic, A. (2003). $\delta^{13}\text{C}$ values of grasses as a novel indicator of pollution by fossil-fuel-derived greenhouse gas CO₂ in urban area. *Environmental Science and Technology*, Vol. 37, pp. 87-90.
- Lijteroff, R.; Cortinez, V. & Raba, J. (1999). Urban development and air quality in San Luis City, Argentina. *Environmental Monitoring and Assessment*, Vol. 57, pp. 169-182.

- Lindén, J.; Thorsson, S. & Eliasson, I. (2007). Carbon Monoxide in Ouagadougou, Burkina Faso - A Comparison between Urban Background, Roadside and In-traffic Measurements. *Water Air Soil Pollution*, DOI 10.1007/s11270-007-9538-2.
- Makra, L.; Mika, J.; Bartzokas, A.; Beczi, R.; Borsos, E. & Sumeghy, Z. (2006). An objective classification system of air mass types for Szeged, Hungary, with special interest in air pollution levels. *Meteorological Atmosphere Physics*, Vol. 92, pp. 115-137.
- Manning, A.J.; Nicholson, K.J.; Middleton, D.R. & Rafferty, S. C. (2000). Field study of wind and traffic to test a street canyon pollution model. *Environmental Monitoring and Assessment*, Vol. 60, pp. 283-313.
- Martín, M.L.; Turias, I.J.; González, F.J.; Galindo, P.L.; Trujillo, F.J.; Puntonet, C.G. & Gorriz, J.M. (2008). Prediction of CO maximum ground level concentrations in the Bay of Algeciras, Spain using artificial neural networks. *Chemosphere*, Vol. 70, Iss.7, pp. 1190-1195.
- Milton, R. & Steed, A. (2007). Mapping carbon monoxide using GPS tracked sensors. *Environmental Monitoring and Assessment*, Vol. 124, pp. 1-19.
- Moriwaki, R.; Kanda, M. & Nitta H. (2006). Carbon dioxide build-up within a suburban canopy layer in winter night. *Atmospheric Environment*, Vol. 40, No.8, pp.1394-1407.
- Muttamara, S.; Leong, S.T. (2000). Titlul. *Environmental Monitoring and Assessment*, Vol. 60, pp. 163-180.
- Nakagawa, F.; Tsunogai, U.; Komatsu, D.D.; Yamada, K.; Yoshida, N.; Moriizumi, J.; Nagamine, K.; Iida, T. & Ikebe, Y. (2005). Automobile exhaust as a source of ¹³C- and D-enriched atmospheric methane in urban areas. *Organic Geochemistry*, Vol. 36, pp. 727-738.
- Nasrallah, H.A.; Balling Jr. R.C.; Madi, S.M. & Al-Ansari L. (2003). Temporal variations in atmospheric CO₂ concentrations in Kuwait City, Kuwait with comparisons to Phoenix, Arizona, USA. *Environmental Pollution*, Vol. 121, No.2, pp. 301-305.
- Newman, S.; Xu, X.; Affek, H.P.; Stolper, E. & Epstein, S. (2008). Change in mixing ratio and isotopic composition of CO₂ in urban air from the Los Angeles basin, California, between 1972 and 2003. *Journal of Geophysical Research*, vol. 113, pp. D23304.
- NOAA-ESRL, Available from: <http://www.esrl.noaa.gov/>
- Park, S.S. & Kim, Y.J. (2007). Titlul. *Environmental Monitoring and Assessment*, Vol. 128, pp.231-240.
- Pataki, D.E.; Alig, R.J.; Fung, A.S.; Golubiewski, N.E.; Kennedy, C.A.; Mcpherson, E.G.; Nowak, D.J.; Pouyat, R.V. & Romero-Lankao, P. (2006a). Urban ecosystems and the North American carbon cycle. *Global Change Biology*, Vol. 12, pp. 2092-2102.
- Pataki, D.E.; Bowling, D.R. & Ehleringer, J.R., (2003). Seasonal cycle of carbon dioxide and its isotopic composition in an urban atmosphere: anthropogenic and biogenic effects. *Journal of Geophysical Research*, Vol. 108, NO.D23, pp. 4735.
- Pataki, D.E.; Bowling, D.R.; Ehleringer, J.R. & Zobitz, J.M. (2006b). High resolution atmospheric monitoring of urban carbon dioxide sources. *Geophysical Research Letters*, Vol. 33, pp. L03813.
- Reich, S.; Magallanes, J.; Dawidowski, L.; Gomez, D.; Groselj, N & Zupan, J. (2006). An analysis of secondary pollutants in Buenos Aires city. *Environmental Monitoring and Assessment*, Vol. 119, pp. 441-457.

- Roba, C., Haiduc, Iv. & Beldean-Galea, M.S.(2009). Diurnal and seasonal variation of CO₂ in urban areas from Cluj county. *Studia Universitatis Babeş-Bolyai, seria Geographia*, Iss, 3, pp. 49-55
- Rubio, M.A.; Lissi, E.; Jorquera, H.; Salinas, E.; Castro, J. & Cádiz, M. (2007). Carbon monoxide concentrations in Santiago City at street levels and their vertical gradient. *Environmental Monitoring and Assessment*, DOI 10.1007/s10661-007-9857-1.
- Sanchez-Ccoyllo, O.R.; Silva Dias, P.L.; de Fatima Andrade, M. & Freitas, S. R. (2006). Determination of O₃-, CO- and PM10-transport metropolitan area of São Paulo, Brazil synoptic-scale analysis of back trajectories. *Meteorology and Atmospheric Physics*, Vol. 92, pp. 83-93.
- Sathitkunnarat, S.; Wongwises, P.; Pan-Aram, R. & Zhang M. (2006). Titlul. *Advances. in Atmospheric Sciences*, Vol. 26, No.6, pp. 901-908.
- Shiva Nagendra, S. M. & Khare, M. (2005). Modelling urban air quality using artificial neural network. *Clean Technology and Environment Policy*, Vol. 7, pp.116-126.
- Sikar, E. & La Scala, N. (2004). Methane and Carbon Dioxide Seasonal Cycles at Urban Brazilian Inland Sites. *Journal of Atmospheric Chemistry* , Vol. 47, No. 2, pp. 101-106.
- Smith, F.A.; Elliott, S.; Blake, D.R. & Sherwood Rowland, F. (2002). Spatiotemporal variation of methane and other trace hydrocarbon concentrations in the Valley of Mexico. *Environmental Science & Policy*, Vol. 249, pp. 1-13.
- Soegaard, H.& Moller-Jensen, L. (2003). Towards a spatial CO₂ budget of a metropolitan region based on textural image classification and flux measurements. *Remote Sensing of Environment*, vol. 87, pp. 283-294.
- Sottocornola, H. & Kiely G. (2010).Hydro-meteorological controls on the CO₂ exchange variation in an Irish blanket bog. *Agricultural and Forest Meteorology*, Vol. 150, pp. 287-297.
- Takahashi, H.; Hiyama, T.; Konohira, E.; Takahashi, A.; Yoshida, N. & Nakamura, T. (2001). Balance and behavior of carbon dioxide at an urban forest inferred from the isotopic and meteorological approaches. *Radiocarbon*, Vol. 43, pp. 659-669.
- Takahashi, H.A.; Konohira, E.; Hiyama, T.; Minami, M.; Nakamura, T. & Yoshida, N. (2002). Diurnal variation of CO₂ concentration, D¹⁴C and d¹³C in an urban forest: estimate of the anthropogenic and biogenic CO₂ contributions. *Tellus*, Vol. 54B, pp. 97-109.
- Thi Nguyen, H.; Kim, K.H.; Ma, C.J.; Cho, S.J. & Ryeul Sohn, (2010). A dramatic shift in CO and CH₄ levels at urban locations in Korea after the implementation of the Natural Gas Vehicle Supply (NGVS) program. *Journal of Environmental Research*, Vol. 110, No.4, pp. 396-409.
- Turias, I.J.; González, F.J.; Martin, M.L. & Galindo, P.L. (2006). A competitive neural network approach for meteorological situation clustering. *Atmospheric Environment*, Vol. 40, Iss. 3, pp. 532-541.
- Veenhuysen, A.; Vermeulen, T.; Hofschreuder P. & Van Den Bulk W.C.M. (1998). Methane emission of the Amsterdam urban area. *Water, Air, and Soil Pollution* , Vol. 107, pp. 321-333.
- Velasco, E.; Pressley, S.; Allwine, E.; Westberg H. & Lamb, B. (2005). Measurements of CO₂ fluxes from the Mexico City urban landscape. *Atmospheric Environment*, Vol. 39, No.38, pp. 7433-7446.

- Venegas, L.E. & Mazzeo, N.A. (2000). Carbon monoxide concentration in a street canyon of Buenos Aires City (Argentina). *Environmental Monitoring and Assessment*, Vol. 65, pp. 417-424.
- Wada, R.; Pearce, J.K.; Nakayama, T.; Matsumi, Y.; Hiyama, T.; Inoue, G. & Shibata, T. (2010). Observation of carbon and oxygen isotopic compositions of CO₂ at an urban site in Nagoya using Mid-IR laser absorption spectroscopy. *Atmospheric Environment*, pp.1-7.
- Wang, J.L.; Kuo, S.R.; Maa, S. & Chen, T.Y. (2001). Construction of a low-cost automated chromatographic system for the measurement of ambient methane. *Analytica Chimica Acta*, Vol. 448, pp. 187-193.
- WHO, 2000: Available from:
http://www.euro.who.int/__data/assets/pdf_file/0005/74732/E71922.pdf
- Widory, D. & Javoy, M. (2003). The carbon isotopic composition of atmospheric CO₂ in Paris. *Earth Planet Science Letter*, Vol. 215, pp. 289-298.
- Wohlfahrt, G.; Hammerle, A.; Haslwanter, A.; Bahn, M.; Tappeiner, U. & Cernusca, A., (2008). Seasonal and inter-annual variability of the net ecosystem CO₂ exchange of a temperate mountain grassland: effects of weather and management. *Journal of Geophysical Research*, Vol. 113, pp. D08110.
- Yamada, K.; Ozaki, Y.; Nakagawa, F.; Tanaka, M. & Yoshida, N. (2003) An improved method for measurement of the hydrogen isotope ratio of atmospheric methane and its application to a Japanese urban atmosphere, *Atmospheric Environment*, Vol. 37, Iss.14, pp. 1975-1982.

IntechOpen



Air Quality-Models and Applications

Edited by Prof. Dragana Popovic

ISBN 978-953-307-307-1

Hard cover, 364 pages

Publisher InTech

Published online 09, June, 2011

Published in print edition June, 2011

Air pollution has been a major transboundary problem and a matter of global concern for decades. High concentrations of different air pollutants are particularly harmful to large cities residents, where numerous anthropogenic activities strongly influence the quality of air. Although there are many books on the subject, the one in front of you will hopefully fulfill some of the gaps in the area of air quality monitoring and modeling, and be of help to graduate students, professionals and researchers. The book is divided in five sections, dealing with mathematical models and computing techniques used in air pollution monitoring and forecasting; air pollution models and application; measuring methodologies in air pollution monitoring and control; experimental data on urban air pollution in China, Egypt, Northeastern U.S, Brazil and Romania; and finally, the health effects due to exposure to benzene, and on the influence of air pollutants on the acute respiratory diseases in children in Mexico.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Iovanca Haiduc and Mihail Simion Beldean-Galea (2011). Variation of Greenhouse Gases in Urban Areas- Case Study: CO₂, CO and CH₄ in Three Romanian Cities, Air Quality-Models and Applications, Prof. Dragana Popovic (Ed.), ISBN: 978-953-307-307-1, InTech, Available from: <http://www.intechopen.com/books/air-quality-models-and-applications/variation-of-greenhouse-gases-in-urban-areas-case-study-co2-co-and-ch4-in-three-romanian-cities>

INTECH
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2011 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the [Creative Commons Attribution-NonCommercial-ShareAlike-3.0 License](#), which permits use, distribution and reproduction for non-commercial purposes, provided the original is properly cited and derivative works building on this content are distributed under the same license.

IntechOpen

IntechOpen