

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,900

Open access books available

186,000

International authors and editors

200M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.

For more information visit www.intechopen.com



Inhibition of Soybean Lipoxygenases – Structural and Activity Models for the Lipoxygenase Isoenzymes Family

Veronica Sanda Chedea¹ and Mitsuo Jisaka²

¹Laboratory of Animal Biology, National Research Development Institute
for Animal Biology and Nutrition (IBNA),

²Faculty of Life Sciences and Biotechnology, Shimane University,

¹Romania

²Japan

1. Introduction

Lipoxygenases (EC 1.13.11.12, linoleate:oxygen, oxidoreductases, LOXs) which are widely found in plants, fungi, and animals, are a large monomeric protein family with non-heme, non-sulphur, iron cofactor containing dioxygenases that catalyze the oxidation of polyunsaturated fatty acids (PUFA) as substrate with at least one 1Z, 4Z-pentadiene moiety such as linoleic, linolenic and arachidonic acid to yield hydroperoxides (Gardner, 1991).




Fig. 1. Lipoxygenase substrates, linoleic, α- linolenic and arachidonic acid.

Theorell et al. (1947) succeeded in crystallizing and characterizing lipoxygenase (LOX) from soybeans and since then among plant LOXs, soybean lipoxygenase isozyme 1 (LOX-1) can be regarded as the mechanistic paradigm for these nonheme iron dioxygenases (Coffa et al., 2005; Minor et al., 1996; Fiorucci et al., 2008).

Designing agents to modulate activities of the variety of so closely homologous enzymes, such as different LOXs, require an intimate knowledge of their 3D structures, as well as information about metabolism of the potential xeno- or endobiotics. So far only the structures of soybean isozymes LOX-1 and LOX-3 have been determined for native enzymes, and several structures of their and rabbit 15-LOX (from reticulocytes) molecular complexes with inhibitors are known. Due to lack of sufficiently purified human enzymes most of the structural research has been done on soybean LOX (Skrzypczak-Jankun et al., 2003).

Understanding the mechanism of inhibition of LOXs can have profound effect in the development of many anti-cancer and anti-inflammatory drugs. On the basis of the available LOX data it was suggested that a combination of LOX modulators might be needed to shift the balance of LOX activities from procarcinogenic to anticancerogenic as a novel strategy for cancer chemoprevention (Skrzypczak-Jankun et al., 2003).

The aim of the present study is to present knowledge on different lipoxygenases having the soybean lipoxygenases as a structural and activity template for their inhibition by natural antioxidant compounds as theoretical approach for food biochemistry and medical applications.

2. Lipoxygenase structure and activity

The three-dimensional structure of soybean lipoxygenase-1 has been determined to 2.3 Å resolution by single crystal X-ray diffraction methods (Boyington et al., 1993). It is a two-domain, single-chain prolate ellipsoid of dimensions 90 x 65 x 60 Å with a molecular mass of 95 kDa. The 839 residues are organised in two domains: one 146 residue N-terminal domain (domain I), and a major, 693 residue C-terminal domain (domain II) (Prigge et al., 1997). Overall, the three-dimensional structure of lipoxygenase-1 shows a helical content of 38.0% and a β -sheet content of 13.9%. The structure of another crystal form of soybean lipoxygenase-1 determined to 1.4 Å resolution (Minor et al., 1996) showed very similar results. The structure of lipoxygenase-3, another soybean lipoxygenase isozyme (Skrzypczak-Jankun, 1997) shows that the lipoxygenase-3 isozyme is very similar in structure despite significant differences in sequence: 857 residues vs 839, deletions at 7 positions, insertions at 25 positions, and substitutions at 224 residues (72% identity).

Soybean seed isoenzymes are 94–97 kDa monomeric proteins with distinct isoelectric points ranging from about 5.7 to 6.4, and can be distinguished by optimum pH, substrate specificity, product formation and stability (Siedow, 1991; Mack et al., 1987). LOX-1 is the smallest in size (838 amino acids; 94 kDa), exhibits maximal activity at pH 9.0 and converts linoleic acid preferentially into the 13-hydroperoxide derivative. LOX-2 is characterized by a larger size (865 amino acids; 97 kDa), by a peak of activity at pH 6.8, and forms equal amounts of the 13- and 9-hydroperoxide compounds (Loiseau et al., 2001). LOX-2 oxygenates the esterified unsaturated fatty acid moieties in membranes in contrast to LOX-1 which only uses free fatty acids as substrates (Maccarrone et al., 1994). LOX-3 (857 amino acids; 96.5 kDa) exhibits its maximal activity over a broad pH range centred around pH 7.0 and displays a moderate preference for producing a 9-hydroperoxide product. It is the most active isoenzyme with respect to both carotenoid cooxidation and production of oxodienoic acids (Ramadoss, 1978).

3. Lipoxygenase reaction

The initial step of LOX reaction is removal of a hydrogen atom from a methylene unit between double bonds in substrate fatty acids (Fig. 2A). The resulting carbon radical is stabilized by electron delocalization through the double bonds. Then, a molecular oxygen is added to the carbon atom at +2 or -2 position from the original radical carbon, forming a peroxy radical as well as a conjugated *trans,cis*-diene chromophore. The peroxy radical is then hydrogenated to form a hydroperoxide. The initial hydrogen removal and the following oxygen addition occur in opposite (or antarafacial) sides related to the plane

formed by the 1*Z*,4*Z*-pentadiene unit. In most LOX reactions, particularly those in plants, the resulting hydroperoxy groups are in *S*-configuration, while one mammalian LOX and some marine invertebrate LOXs produce *R*-hydroperoxides. Even in the reactions of such “*R*-LOXs”, the antarafacial rule of hydrogen removal and oxygen addition is conserved. In cases of plant LOXs, including soybean LOXs, the usual substrates are C18-polyunsaturated fatty acids (linoleic and α -linolenic acids), and the products are their 9*S*- or 13*S*-hydroperoxides (Fig. 2B). Most plant LOXs react with either one of the regio-specificity, while some with both. Therefore, based on the regio-specificity, plant LOXs are classified into 9-LOXs, 13-LOXs, or 9/13-LOXs.




Fig. 2. LOX reaction showing the principal steps of LOX reaction (Panel A), and the actual reactions of plant LOXs and α -linolenic acid (Panel B). HPOTE: hydroperoxyoctadecatrienoic acid.

4. Biological and metabolic functions

4.1 In plants

Lipid peroxidation is common to all biological systems, both appearing in developmentally and environmentally regulated processes of plants (Feussner & Wasternack, 2002). The hydroperoxy polyunsaturated fatty acids, synthesized by the action of various highly specialised forms of lipoxygenases, are substrates of at least seven different enzyme families (Feussner & Wasternack, 2002). Signaling compounds such as jasmonates, antimicrobial and antifungal compounds such as leaf aldehydes or divinyl ethers, and a plant-specific blend of volatiles including leaf alcohols are among the numerous products. Thus, the lipoxygenase pathway becomes an initial step in the interaction of plants with pathogens, insects, or abiotic stress and at distinct stages of development (Feussner & Wasternack, 2002).

4.2 In humans

Besides polyunsaturated fatty acids, H_2O_2 , fatty acid hydroperoxides, and synthetic organic hydroperoxides support the lipoxygenase-catalyzed xenobiotic oxidation the major reactions documented thus far including oxidation, epoxidation, hydroxylation, sulfoxidation, desulfuration, dearylation, and N-dealkylation (Kulkarni, 2001). It is noteworthy that lipoxygenases are also capable of glutathione conjugation of certain xenobiotics (Kulkarni, 2001). Available data suggest that lipoxygenases contribute to in vivo metabolism of endobiotics and xenobiotics in mammals (Kulkarni, 2001).

Recent reviews describe the role of lipoxygenase in cancer (Bhattacharya et al., 2009; Pidgeon et al., 2007; Moreno, 2009), inflammation (Duroudier et al., 2009; Hersberger, 2010) and vascular biology (Chawengsub et al., 2009; Mochizuki & Kwon, 2008) and for an extensive presentation of the role of eicosanoids in prevention and management of diseases the reader is referred to the review of Szeffel et al. (2011).

5. Interaction of lipoxygenase with inhibitors as theoretical approach for food industry and medical applications

In terms of the structure and function, LOXs are unique, because their metal cofactor is a single ion bound by the side chains of the surrounding amino acids and the carboxylic group of the C-terminus, and their inhibitors bind to or near the Fe co-factor (Skrzypczak-Jankun et al., 2007). Lipoxygenases are inhibited by a large number of chemicals, some of which also serve as co-substrates (Kulkarni, 2001).

5.1 Importance of lipoxygenase inhibition for food industry

Besides their physiological role, plant lipoxygenases are of significant importance to the food industry, since these enzymes have been implicated in the generation of the flavour and aroma in many plant products. For instance, they are responsible for the undesirable 'beany', 'green' and 'grassy' flavours produced during processing and storage of protein products derived from legume seeds (Fukushima, 1994; Robinson et al., 1995) and the development of the stale flavour in beer during storage (Kobayashi et al., 1993). Lipoxygenases also play an important role in the baking industry. They are quite effective as bleaching agents, increase mixing tolerance and improve dough rheology (Nicolas & Potus, 1994; Larreta-Garde, 1995; Cumbee et al., 1997; Borelli^b et al., 1999).

Freshly refined soybean oil is practically odourless and bland, but "green, grassy, fishy" off-flavors may develop quickly if the oil is heated or stored under conditions that expose it to light and oxygen or by contamination with pro-oxidant metals such as copper and iron (Berk, 1992). "Beany" flavour is the principal inconvenience of traditional soymilk and its products (e.g., tofu) and is caused by some ketones and aldehydes, particularly hexanals and heptanals, produced through LOX catalyzed oxidation (Berk, 1992).

Fish lipids are susceptible to oxidation owing to the high levels of polyunsaturated fatty acids (PUFA), even in frozen storage, and this can affect the flavour, texture, taste, aroma and shelf life of fish (Ke & Ackman, 1976). Since the direct interaction between oxygen and highly unsaturated lipids is kinetically hindered (Kanner et al., 1987), the enzymatic initiation of oxidation by enzymes such as lipoxygenase, peroxidases and microsomal enzymes has been gaining favour.

Green tea glazing was shown to improve the storage quality of frozen bonito fillets (Lin & Lin, 2005). In addition, hot water tea extract was shown to suppress the pro-oxidant activities of the dark meat and skin of blue sprat (Seto et al., 2005). Banerjee (2006) proposes that the improvement in the shelf life of fish by green tea polyphenols is at least in part due to inhibition of LOX resulting in delaying oxidation of fish lipids and because of that impregnation of muscle fillets in tea extract by itself or in combination with other natural inhibitors may improve the shelf-life and storage quality of fish fillets.

Besides its function of oxidizing the polyunsaturated fatty acids (linoleic, linolenic and arachidonic), the enzyme may also catalyse the co-oxidation of carotenoids, resulting in the loss of natural colorants and essential nutrients (Robinson et al., 1995). LOX have been

implicated in the generation of the flavour and aroma in many plant products, in the decolourisation of pigments and in the potential of compromising the anti-oxidant status (Casey, 1999). In pasta the involvement of LOX in colour loss is demonstrated by positive correlation between the decrease of β -carotene content after pastification and LOX activities in semolina. In addition to this, the hydroperoxidation and bleaching activities of LOX are highly correlated demonstrating that the bleaching might be ascribable to a co-oxidative action by LOX (Borrelli^a et al., 1999).

During pasta processing in which the maximal pigment degradation by LOX activity occurs (Borrelli^b et al., 1999), it is shown that externally added β -carotene can act as inhibitor of the LOX-catalysed linoleate hydroperoxidation and an inverse relation between the % of carotenoid loss and the initial carotenoid content in semolina from durum varieties, showing similar LOX activity, was found (Trono et al., 1999).

The complete characterisation of lipoxygenase from pea seeds (*Pisum sativum* var. *Telephone L.*) gives possibility to avoid destructive influence during food processing and storage (Szymanowska et al., 2009) by the action of this enzyme.

5.2 LOX inhibition in cancer

Molecular studies of the well-known relationship between polyunsaturated fatty acid metabolism and carcinogenesis have revealed novel molecular targets for cancer chemoprevention and treatment (Lipkin et al., 1999; Willett, 1997; Klurfeld & Bull, 1997; Guthrie & Carroll, 1999).

The role of lipoxygenase in the development and progression of cancer is complex due to the variety of lipoxygenase genes that have been identified in humans, in addition to different profiles of lipoxygenase observed between studies on human tumor biopsies and experimentally induced animal tumor models (Pidgeon et al., 2007). The literature emerging on the role of lipoxygenases in tumor growth, for the most part, suggests that distinct lipoxygenase isoforms, whose expression are lost during the progression of cancer, may exhibit anti-tumor activity, while other isoforms may exert pro-tumorigenic effects and are preferentially expressed during the development of various cancers.

The involvement of 5-lipoxygenase and 12-lipoxygenase in human cancer progression is now supported by a growing body of literature. The involvement of 15-lipoxygenase-1 in colorectal cancer involves its implication in carcinogenesis having pro-carcinogenic as well as anti-carcinogenic roles (Bhattacharya et al., 2009). The co-localization of these enzymes and the similarities of their bioactions on cancer cell growth suggest that the simultaneous inhibition of these enzymes may represent novel and promising therapeutic approaches in selected cancer types (Pidgeon et al., 2007). Therefore, when targeting the regulation of arachidonic acid metabolism, blocking 5-lipoxygenase, 12-lipoxygenase and 15-lipoxygenase-1 without altering the expression of the anti-carcinogenic 15-lipoxygenase-2 may be the most effective, however at present no drug recapitulates these capabilities (Pidgeon et al., 2007).

5.3 Mechanisms of lipoxygenase inhibition

In general, lipoxygenase inhibitors can bind covalently to iron or form the molecular complexes blocking access to iron (Skrzypczak-Jankun et al., 2007). It was pointed out by Walther et al., that a course of inhibition, by the drug ebselen, (noncompetitive vs competitive) and its reversibility depend on the oxidation state of iron, i.e. whether the enzyme is catalytically silent with Fe^{2+} when it binds covalently, causing irreversible

inhibition or preoxidized and active with Fe^{3+} in the presence of the fatty acid substrate (Walther et al., 1999). In both cases the enzyme's performance can be illustrated by a classic Lineweaver-Burk plot. Many inhibitors do not follow such a linear relation between velocity and the inhibitor's concentration showing a hyperbolic curve instead as observed by Skrzypczak-Jankun et al. (2002) for polyphenolic inhibitors (curcumin, quercetin, epigallocatechin gallate and epigallocatechin) interacting with soybean lipoxygenase-3. In general, the kinetic data are seldom reported (Skrzypczak-Jankun et al., 2007). Xenobiotic oxidation by soy lipoxygenase has been investigated and described, while human enzymes lack such thorough studies (Skrzypczak-Jankun et al., 2007). The *in vivo* susceptibility of lipoxygenases' inhibitors may depend not only on the source of lipoxygenase and its isozyme (Pham et al., 1998; Schewe et al., 1986) but also on the oxidation state of iron and the competition between peroxidase and co-oxidase activities of enzyme (Borbulevych et al., 2004).

The first mode to inhibit the lipoxygenase would be a direct reduction of iron to its inactive form. For soybean lipoxygenase, it has been demonstrated that nordihydroguaiaretic acid rapidly reduces the active ferric species of the enzyme to its inactive ferrous form, thus causing interruption of the catalytic cycle (Kemal et al., 1987). For the polyphenol inhibition of lipoxygenase it was firstly suggested that this molecules strongly complex of the ferric iron moiety of the lipoxygenase, thus preventing its reduction *via* the catalytic cycle as proposed for the action of 4-nitrocatechol on the soybean lipoxygenase-1 (Spaapen et al., 1980). The second observation that complexation of the flavanols with Fe^{3+} did not abolish the inhibitory effect may rule out a direct complexation of the iron moiety in ferric lipoxygenase by these catechol compounds. The X-ray analysis shows 4-nitrocatechol near iron with partial occupancy, blocking access to Fe but not covalently bound to it (Skrzypczak-Jankun et al., 2004). If a similar mode of action holds for the interaction of flavanols with mammalian lipoxygenases, the corresponding iron polyphenol complexes may retain their lipoxygenase-inhibitory effect.

A third conceivable mode of action of polyphenols is the effective reduction of hydroperoxides that are essential activators of lipoxygenase *via* conversion of the enzymatically silent ferrous species to the active ferric form. The observation of Schewe et al. (2001) that lowering of the hydroperoxide tone by glutathione plus glutathione peroxidase did not modulate the inhibitory effects of flavanols on 15-lipoxygenase-1 does not support the latter possibility.

In case of carotenoids, more specifically, β -carotene, the lipoxygenase was inhibited by keeping it in the inactive form of Fe(II) (Serpen & Gökmen, 2006). These authors suggest that β -carotene reacts with linoleyl radical ($\text{L}\cdot$) at the beginning of the chain reaction, so it prevents the accumulation of conjugated diene forms ($\text{LOO}\cdot$, LOO^- and LOOH). Since $\text{L}\cdot$ transforms back to its original form of LH, the enzyme cannot complete the chain reaction and thus remains in the inactive Fe(II) form, which is not capable of catalyzing linoleic acid hydroperoxidation (Serpen & Gökmen, 2006).

Wu et al. (1999) have reported that β -carotene scavenges the linoleyl peroxy radical ($\text{LOO}\cdot$) by a hydrogen transfer mechanism and the oxidation of β -carotene occurs during this action. In these conditions, it is absolutely clear that the amount of inactivated enzyme depends on the concentration of β -carotene present in the medium (Serpen & Gökmen, 2006).

According to Mahesha et al. (2007) the lipoxygenase inhibition by isoflavones follows the next mechanism: an electron donated by isoflavones is accepted by the ferric form (Fe^{3+}) of lipoxygenase, which is reduced to resting ferrous form (Fe^{2+}), thus inhibiting lipoxygenase.

Genistein is neither consumed nor does state change during the course of the reaction of lipoxygenase (Mahesha et al., 2007), while quercetin entrapped within lipoxygenase undergoes degradation (Borbulevych et al., 2004).

5.4 Inhibition of soybean lipoxygenase by different classes of polyphenols

Gillmor et al. (1997) obtained the structure of the rabbit reticulocyte enzyme as a complex with the inhibitor RS75091. Located in one of the hydrophobic channels of the enzyme, the inhibitor was found to be close but not binding the iron atom of the catalytic site. These observations provided the first indications of how the native enzyme can interact with potential ligands (Pham et al., 1998).

Natural flavonoids don't affect only the lipoxygenase oxidation of its classical substrates but also the co-oxidation of xenobiotics by this enzyme. Epigallocatechin-gallate, quercetin and rutin proved to reduce the co-oxidation rate of guaiacol, benzidine, paraphenylenediamine and dimethoxybenzidine by soybean lipoxygenase-1 (Hu et al., 2006). This data suggest that flavonoids may have anticarcinogenic and antitoxic effect through inhibition of oxidative activation generated by lipoxygenase (Hu et al., 2006). Green tea polyphenols have potent free radical quenching and antioxidant activities (Wiseman et al., 1997) and have structural features that may specifically interfere with the arachidonic acid cascade, including the lipoxygenase pathway (Hong & Yang, 2003; Hussain et al., 2005). In addition, with growing concerns regarding the safety of synthetic antioxidants such as BHT and BHA, alternative mechanisms of antioxidant protection by the use of natural antioxidants have been in review over the years (Barlow, 1990).

Polyphenols, mainly flavonoids and phenolic acids, are abundant in a number of dietary sources such as certain cocoas, tea, wine, fruits and vegetables. More than 8000 different flavonoids of natural origin are known (Schewe & Sies, 2003). The flavonoids exist in nature as aglycons (free form) or conjugated (with O-glucosides or methylated). The aglycons can be subdivided in different subclasses (flavanols, flavanones, flavones, izoflavones, flavonols, anthocyanidines, aurones, chalcones) in function of how the B ring from their structure is linked to the heterocycle C, of the oxidation state and of the functional groups linked to the C ring (Beecher, 2003).




Fig. 3. The basic structure of flavonoids.

The basic structure of flavonoids is represented by the flavan nucleus containing 15 carbons structured in 2 benzene rings, named A and B and linked by a C₃ unit, which together with an oxygen atom forms the γ -pyronic or γ -pyranic ring, named the C ring as shown in Fig.3.

A number of *in vitro* and *in vivo* studies as well as clinical trials suggest beneficial effects of flavonoids for health, counteracting the development of cardiovascular diseases, cancer and obesity. Bors et al. (1990) were the first to claim three partial structures contributing to the

radical-scavenging activity of flavonoids: (a) an *o*-dihydroxyl structure in the B ring (catechol structure) as a radical target site providing good electron delocalization and stabilization of the phenoxy radical; (b) a 2,3-double bond with conjugation to the 4-oxo group which is necessary for delocalization of an unpaired electron from the B ring, (c) hydroxyl groups at the 3- and 5-positions, which are necessary for enhancement of radical scavenging activity, increasing the delocalisation of electrons across the flavonoid scaffold. The catechol group is essential for the radical-scavenging activity of flavan-3-ols and flavanones lacking 2,3-double bonds (Bors et al., 1990).

In parallel to the free radical-scavenging properties the following structural features were found to enhance the inhibitory potency: (i) presence of a catechol arrangement in the B or A ring, (ii) a carbonyl group together with a 2,3-double bond in the C ring (Schewe & Sies, 2003). Other structural features were opposite to the free radical scavenging potencies: (i) presence of a 3-OH group in the C ring diminished than reinforced the inhibition of lipoxygenases, (ii) in the absence of a catechol arrangement there was an inverse correlation to the total number of OH groups in the flavonoid molecule (Schewe & Sies, 2003). Although either reducing or ferric iron chelating properties are prerequisites for a lipoxygenase-inhibitory compound, and both of them are also inherent to flavonoids, the inhibitory effects cannot be ascribed solely to one of these mechanisms (Schewe & Sies, 2003). The inhibition of lipoxygenases by flavonoids appears to be of more complex nature (Schewe & Sies, 2003).

Inhibitors studies of lipoxygenase from pea showed that phenolic antioxidant components were effective and can be used to protect food lipids against oxidation (Szymanowska et al., 2009). The conducted research proved that activity of lipoxygenase from pea seeds could be effectively inhibited by some phenolic compounds. The most effective inhibitor is caffeic acid (about 57% of inhibition). Flavonoids like catechin and quercetin considerably inhibit the lipoxygenase activity. Inhibitors used for investigation in this study were placed in the following order: caffeic acid > quercetin > catechin > benzoic acid > ferulic acid > kaempferol (Szymanowska et al., 2009).

5.5 Lipoxygenase inhibition by quercetin

Quercetin is the most abundant among the flavonoid molecules and can be found in the fruits, vegetables, seeds, nuts, and flowers of many plants. Its documented impact on human health includes cardiovascular protection, anticancer, antiviral, anti-inflammatory activities, antiulcer effects and cataract prevention. Like other flavonoids, quercetin appears to combine both lipoxygenase-inhibitory activities and free radical-scavenging properties in one agent and thus belongs to a family of very effective natural antioxidants (Sadik et al., 2003). Quercetin is a flavonol that can be easily oxidized in an aqueous environment, and in the presence of iron and hydroxyl free radicals (Borbulevych et al., 2004).

The inhibition of rabbit 15-lipoxygenase-1 and of soybean lipoxygenase-1 by quercetin was studied in detail (Sadik et al., 2003). Quercetin modulates the time course of the lipoxygenase reaction in a complex manner by exerting three distinct effects: (i) prolongation of the kinetic lag period, (ii) instant decrease in the initial rate after the lag phase being overcome, (iii) time-dependent inactivation of the enzyme during reaction, but not in the absence of substrate (Schewe & Sies, 2003). The literature data obviously indicate that quercetin represents one of the most potent inhibitors of different LOXs (Schneider & Bucar^a, 2005; Schneider & Bucar^b, 2005).

The study of Banerjee (2006) shows that green tea polyphenols are very potent inhibitors of mackerel muscle LOX, with EGCG (epigallocatechin gallate) as the most effective inhibitor (IC_{50} 0.13 nM) followed by ECG (epicatechin gallate) (IC_{50} 0.8 nM), EC (epicatechin) (IC_{50} 6.0 nM), EGC (epigallocatechin) (IC_{50} 9.0 nM) and C (catechin) (IC_{50} 22.4 nM). Chocolate and cocoa are also sources of catechins. Epigallocatechin gallate isolated from the seeds of *Theobroma cacao* had the best inhibitory activity on rabbit 15-lipoxygenase-1, with an IC_{50} =4M, epicatechin gallate had IC_{50} =5M and epicatechin an IC_{50} =60M (Schewe et al., 2002). A better inhibition of epicatechin (IC_{50} approx. 15M) was registered in the case of recombinant human platelet 12-lipoxygenase.

Obtained from X-ray analysis, the 3D structure of the resulting complex of (-)-epigallocatechin gallate (EGCG) interacting with soybean lipoxygenase-3 reveals the inhibitor depicting (-)-epigallocatechin that lacks the galloyl moiety (Skrzypczak-Jancun et al., 2003). The A-ring is near the iron co-factor, attached by the hydrogen bond to the C-terminus of the enzyme, and the B-ring hydroxyl groups participate in the hydrogen bonds and the van der Waals interactions formed by the surrounding amino acids and water molecules (Skrzypczak-Jancun et al., 2003).




Fig. 5. Lipoxygenase-3 in complex with epigallocatechin gallate as an inhibitor determines the degradation of natural flavonoid to epigallocatechin (Skrzypczak-Jancun et al., 2003).

X-ray analysis of soybean lipoxygenase-3 crystals soaked with EGCG shows the molecular complex of LOX-3 with (-)-epigallocatechin molecule.

5.7 Inhibitory effects of soybean isoflavones on lipoxygenase activity

Soybeans are important sources of isoflavone levels (Song et al., 1998), present as 12 derivatives, including free genistin, daidzin, glycitin and their acetyl, malonyl or glycosylated forms. Isoflavones are composed of 2 benzene rings (A and B) linked through a heterocyclic pyrane C ring. The position of the B ring discriminate flavonoid flavones (C2-position) from isoflavones (C3-position).




Fig. 6. Dietary isoflavones inhibitors of lipoxygenase.

The impact of dietary isoflavones, daidzein and genistein, on the health of adults and infants is well documented, an increasing interest for these compounds being registered due to their biological effects including: estrogen-like activity, prevention of breast (Warri et al., 2008), prostate (Matsumura et al., 2008) and colon cancer (Mac Donald et al., 2005), antioxidant activity (Malencić et al., 2007; Sakthivelu et al., 2008), prevention of menopausal symptoms and osteoporosis (Ma et al., 2008), and heart disease (Xiao, 2008).

In the work of Mahesha et al. (2007) the inhibition of soy lipoxygenase-1 and 5-lipoxygenase from human polymorph nuclear lymphocyte by isoflavones, genistein and daidzein as glycosilated and unglycosilated compounds was studied. Soybean isoflavones inhibit LOX either as aglycons, or as glucosides. Isoflavones exert combined dual actions as inhibitors: they compete with the hydroperoxide formation to prevent the generation of LOX active ferric state (1) and also are capable of reducing the ferric enzyme to its inactive ferrous form (2) (Mahesha et al., 2007).

Vicaş et al. (2011) showed that genistein was almost twice more potent inhibitor than daidzein at similar concentration with concentration that induces 50% soybean lipoxygenase-1 inhibition values of 5.33 mM versus 11.53 mM. Genistein and daidzein proved to be noncompetitive with inhibition constants K_i of 33.65 and 43.45 mM, respectively (Vicaş et al., 2011). The inhibitory efficiency of the genistein and daidzein depended both on their concentration and on the substrate's concentration (Vicaş et al., 2011).

5.8 Inhibition of soybean lipoxygenase by carotenoids

There are over 600 fully characterized, naturally occurring molecular species belonging to the class of carotenoids. In humans, some carotenoids (the provitamin A carotenoids: α -carotene β -carotene, γ -carotene and the xanthophyll, β -cryptoxanthin) are best known for converting enzymatically into vitamin A; diseases resulting from vitamin A deficiency remain among the most significant nutritional challenges worldwide. Also, the role that carotenoids play in protecting those tissues that are the most heavily exposed to light (e.g. photo protection of the skin, protection of the central retina) is perhaps most evident, while other potential roles for carotenoids in the prevention of chronic diseases (cancer, cardiovascular disease) are still being investigated. Because carotenoids are widely consumed and their consumption is a modifiable health behaviour (via diets or supplements), health benefits for chronic disease prevention, if real, could be very significant for public health (Mayne, 2010). Carotenoids are isoprenoid molecules which contain a polyene chains, with or without cyclisation at the ends.




Fig. 7. Carotenoids serving as lipoxygenase co-substrates

The existence of an enzyme "carotene oxidase" in soybeans, which catalyzes the oxidative destruction of carotene was reported by Bohn and Haas in 1928 (Bohn & Haas, 1928). Four years later, Andre and Hou found that soybeans contained an enzyme, lipoxygenase (linoleate oxygen oxidoreductase), which they termed "*lipoxidase*", catalyzing the peroxidation of certain unsaturated fatty acids (Andre & Hou, 1932).

In 1940 the observation that "*lipoxydase*" is identical to "*carotene oxidase*" was published (Sumner & Sumner, 1940). These early findings of lipoxygenase peroxidizing the unsaturated fats and bleaches the carotene were reported as the result of studies on the oxidation of crystalline carotene or carotene dissolved in unsaturated oil. Surprisingly it was found that the carotene oxidase had an almost negligible bleaching action upon the crystalline carotene. On the contrary, when one employs carotene dissolved in a small quantity of fat, the bleaching is extremely rapid. With excessive quantities of fat, the rate of bleaching of the carotene diminishes, and it was concluded that the effect of added fat upon the rate of bleaching of carotene is probably due to a coupled oxidation (Sumner & Sumner, 1940).

Studying the soya-lipoxygenase-catalyzed degradation of carotenoids from tomato Biacs and Daood (2000) found that β -carotene was the most sensitive component, followed by lycoxanthin and lycopene. Their results also implied that β -carotene can actively perform its antioxidant function during the course of lipid oxidation. It seems that oxidative degradation and, accordingly, antioxidant activity of each carotenoid depends on the rate of its interaction with the peroxy radical produced through the lipoxygenase pathway (Biacs & Daood, 2000) and thus is able to inhibit lipoxygenase. The inhibition of the hydroperoxide formation by carotenoids has been attributed to their lipid peroxy radical-trapping ability (Burton & Ingold, 1984).

In vitro, lycopene is a substrate of soybean lipoxygenase. The presence of this enzyme also significantly increased the production of lycopene oxidative metabolites (dos Anjos Ferreira et al., 2004; Biacs & Daood, 2000). It was reported that during the co-oxidation of β -carotene

by LOX-mediated hydroperoxidation reactions, inhibition of LOX activity takes place also (Lomnitski et al., 1993; Trono et al., 1999; Pastore et al., 2000). The activity of soybean lipoxygenase-1 was inhibited by β -carotene which breaks the chain reaction at the beginning stage of linoleic acid hydroperoxidation (Serpen & Gökmen, 2006). Besides soybean lipoxygenase (Ikedioby & Snyder, 1977; Hildebrand & Hymowitz, 1982) carotene oxidation during lipoxygenase-mediated linoleic acid oxidation has been reported in various studies for the enzymes extracted from potato (Aziz et al., 1999), pea (Yoon & Klein, 1979; Gökmen et al., 2002), wheat (Pastore et al., 2000), olive (Jaren-Galan et al., 1999) and pepper (Jaren-Galan & Minguez-Mosquera, 1997). Soybean lipoxygenase-1 and recombinant pea lipoxygenase-2 and lipoxygenase-3, oxidizing β -carotene, yield apocarotenal, epoxy-carotenal, apocarotenone and epoxy-carotenone (Wu et al., 1999).

Through molecular modeling Hazai et al. (2006) predicted that lycopene and lycophyll bind with high affinity in the superficial cleft at the interface of the β -barrel and the catalytic domain of 5-LOX (the “cleavage site”) suggesting potential direct competitive inhibition of 5-LOX activity by these molecules after in vivo supplementation, particularly in the case of the dial metabolite.

5.9 Quinone and semiquinone formation during the lipoxygenase inhibition reaction

In his excellent review from 2001, Kulkarni presents the studies up to that date indicating the semiquinone and quinone formation in different lipoxygenase catalyzed reactions of xenobiotics oxidation. Diethylstilbestrol (DES) is a human transplacental carcinogen. DES-quinone, one of the metabolites of DES, binds to DNA and is presumed to be the ultimate toxicant. Although DES-quinone formation by human tissue lipoxygenase has yet to be examined, soybean lipoxygenase has been shown to initiate one-electron oxidation to DES semiquinone in the presence of H_2O_2 (Nunez-Delicado et al., 1997). Subsequent dismutation of two molecules of DES semiquinone yields one molecule each of DES-quinone and DES.

Although phenol is oxidized slowly by different lipoxygenase isoenzymes, potato 5-lipoxygenase (Cucurou et al., 1991) and soybean lipoxygenase-1 (Cucurou et al., 1991; Mansuy et al., 1988), substituted phenols and catechols undergo extensive one-electron oxidation and yield the corresponding reactive phenoxyl radicals or semiquinones. These free radicals polymerize to yield a mixture of complex metabolites.

However, it seems that quercetin may act, in most of cases, after being metabolically activated (Metodiewa et al., 1999), and despite a constant increase of knowledge on both positive and negative biological effects of this natural product, it remains often unclear which activated form should play a role in a given process (Fiorucci et al., 2007). Indeed, semiquinone and quinone forms of quercetin, deriving from the abstraction of respectively one or two $H\bullet$, are involved in many oxidative processes (Metodiewa et al., 1999; Gliszczynska-Swiglo et al., 2003; Hirakawa et al., 2002). For instance, quercetin reduces peroxy radicals involved in lipid peroxidation, and through this reaction, a semiquinone species is produced, which then undergoes a disproportionation to generate a quinone form (Fiorucci et al., 2007). Three semiquinone forms for quercetin have been considered by Fiorucci et al. (2008) in order to study the quercetin binding to lipoxygenase-3 by molecular modeling simulations. In the case of lipoxygenase-catechol complexes, the formation of the catechol-iron(III) complex of soybean lipoxygenase 1 gradually results in reduction of the cofactor and release of the semiquinone but no evidence of quinone formation in the UV-visible spectra of samples of the native enzyme treated with catechol was obtained (Spaapen et al., 1980; Nelson, 1988; Pham et al., 1998).

Besides their antioxidant properties, catechins have been described to display pro-oxidant activity having the potential to oxidize the quinones or semiquinones resulting in redox cycling and reactive oxygen species production as well as in thiol, DNA and protein alkylation (Galati & O'Brien, 2004; van der Woude et al., 2006).

Our previous study shows that the oxidation products of catechins are formed within the cellular matrix but also in the extracellular medium (Chedea et al., 2010). We have demonstrated by UV-Vis spectroscopy, that the quinones are involved in the modulation of lipoxygenase activity in the presence of catechins within the cells (Chedea et al., 2010). This conclusion is in agreement with that of Sadik et al. (2003) and Banerjee (2006). An irreversible covalent modification of soybean LOX by flavonoids has been suggested by Sadik et al. (2003) whereby during the formation of fatty acid peroxy radical in the LOX pathway, the flavonoids are co-oxidized to a semi-quinone or quinone, which in turn may bind to sulfhydryl or amino groups of the enzyme causing inhibition (Banerjee, 2006).

6. Conclusions

The knowledge presented in this study addressed the lipoxygenase pathway inhibition by antioxidant polyphenols at two levels: human diet and human health or to a larger extent human disease prevention and treatment. Review articles (Jachak, 2006; Schneider & Bucar^a, 2005; Schneider & Bucar^b, 2005) summarize natural products with inhibitory properties toward LOX (Skrzypczak-Jankun et al., 2007). Natural remedies almost never consist of a single ingredient and usually are a mixture of many in proper proportions, with a synergistic effect of their simultaneous action being absolutely necessary for beneficial medicinal results. Thus, one should proceed with caution, since the action of a selected single compound may not be the same (Skrzypczak-Jankun et al., 2007). The X-ray studies of soybean complexes with quercetin, curcumin, EGCG, EGC indicated conversion of these inhibitors into their metabolites (Skrzypczak-Jankun et al., 2007), which is not surprising considering the co-oxidative activity of LOXs (Kulkarni, 2001). As already presented a question arises concerning the lipoxygenase inhibition: "What is really inhibiting LOX, a given chemical or its LOX metabolite?" (Skrzypczak-Jankun et al., 2007). The results presented so far indicate a complex mode of inhibition involving the inhibitor itself but also its reaction product with lipoxygenase.

The lipoxygenase researcher faces the next antinomy: despite the difference in the number of amino acids between plant and mammalian LOXs, these proteins are amazingly similar in topology with high similarities in the active site of these enzymes. It is believed that all LOXs follow the same catalytic mechanism; however, it is probably the vicinity of the iron site that determines the regio and stereospecificity of the particular enzyme (Skrzypczak-Jankun et al., 2003). In this contradictorily state of facts soybean lipoxygenases stands as a control point in terms of structure, activity and thus inhibition.

7. Acknowledgements

V.S. Chedea is a Japan Society for the Promotion of Science (JSPS) postdoctoral fellow. The authors wish to thank Dr. Ewa Skrzypczak-Jankun for the enlightening discussions during this manuscript writing and Ms. Nana Henmi for helping with the structures drawing.

8. References

- André, E. & Hou, K.-W. (1932). La présence d'une oxydase des lipides ou lipoxydase dans graine de soja, glycine soja lieh, *Comptes Rendus*, Vol.194, pp. 645–647.
- Aziz, S., Wu, Z. & Robinson, D.S. (1999). Potato lipoxygenase catalyzed co-oxidation of β -carotene. *Food Chemistry* Vol.64, pp. 227–230.
- Banerjee, S. (2006). Inhibition of mackerel (*Scomber scombrus*) muscle lipoxygenase by green tea polyphenols, *Food Research International*, Vol.39, pp. 486–491.
- Barlow, S. M. (1990). Toxicological aspects of antioxidants used as food additives, In B. J. F. Hudson (Ed.). *Food antioxidants*, London: Elsevier Applied Science, pp. 253–308.
- Bednar, W., Holzmann, K. & Marian, B. (2007). Assessing 12(S)-lipoxygenase inhibitory activity using colorectal cancer cells overexpressing the enzyme, *Food and Chemical Toxicology*, Vol.45, pp. 508–514.
- Beecher, G.R. (2003). Overview of dietary flavonoids: Nomenclature, occurrence and intake, *Journal of Nutrition*, Vol. 133, pp. 3248S–3254S.
- Bergström^a, S., Krabisch, L., Samuelsson, B., & Sjövall, J. (1962). Preparation of prostaglandin F from prostaglandin E, *Acta Chemica Scandinavica*, Vol.16, pp.969–974.
- Bergström^b, S., Dressler F., Ryhage R., Samuelsson B. & Sjövall, J. (1962). The isolation of two further prostaglandins from sheep prostate glands, *Arkiv foer Kemi*, Vol.19, pp. 563–567.
- Berk, Z. (1992). Technology of production of edible flours and protein products from soybeans, In: *FAO Agricultural Services Bulletin*. Vol. 97. Date of access: 19.01.2010, Available from: <http://www.fao.org/docrep/t0532e/t0532e01.htm>.
- Bhattacharya, S., Mathew, G., Jayne, D.G., Pelengaris, S. & Khan, M. (2009). 15-Lipoxygenase-1 in colorectal cancer: A review, *Tumor Biology*, Vol. 30, pp. 185–199.
- Biacs, P.A. & Daood, H.G. (2000). Lipoxygenase-catalysed degradation of carotenoids from tomato in the presence of antioxidant vitamins. *Biochemical Society Transactions*, Vol. 28, pp. 839.
- Bohn, R.M. & Haas, L.W. (1928). Chemistry and methods of enzymes, in *Chemistry and methods of enzymes*, ed by Sumner JB and Somers GF, Academic Press, New York.
- Borbulevych, O.Y., Jankun, J., Selman, S. & Skrzypezak-Jankun, E. (2004). Lipoxygenase interactions with natural flavonoid, quercetin, reveal a complex with protocatechuic acid in its X-Ray structure at 2.1 Å resolution. *Proteins, Structure, Function and Bioinformatics*, Vol. 54, pp.13–19.
- Borrelli^a, G.M., Troccoli, A., Fares, C., Trono, D., De Leonardis, A.M., Padalino, L., Pastore, D., Del Giudice, L. & Di Fonzo, N. (1999). Lipoxygenase in durum wheat: What is the role in pasta colour?, *Options Mediterraneennes*, pp. 497–499, Date of access 12.02.2011, Available from: <http://ressources.ciheam.org/om/pdf/a40/00600082.pdf>
- Borrelli^b, G.M., Troccoli, A., Di Fonzo, N. & Fares, C. (1999). Durum wheat lipoxygenase activity and other quality parameters that affect pasta colour, *Cereal Chemistry*, Vol. 76, pp. 335–340.
- Bors, W., Heller, W., Michael, C. & Saran, M. (1990). Flavonoids as antioxidants: Determination of radical-scavenging efficiencies. In L. Packer & A. N. Glazer (Eds.). *Methods in enzymology* (Vol. 186, pp. 343–355). San Diego: Academic Press.
- Boyington, J. C., Gaffney, B. J. & Amzel L. M. (1993). Structure of soybean lipoxygenase-1. *Biochemical Society Transactions*, Vol. 21, pp. 744–748.

- Burton, G.W. & Ingold, K.U. (1984). β -carotene: An unusual type of lipid antioxidant. *Science*, Vol. 224, pp. 569-573.
- Casey, R. (1999). *Lipoxygenases in seed proteins*, pp. 685-708. Shewry, P.R. and Casey, R. eds. Kluwer Academic Publishers.
- Chawengsub, Y., Gauthier, K. M. & Campbell, W. B. (2009). Role of arachidonic acid lipoxygenase metabolites in the regulation of vascular tone, *American Journal of Physiology- Heart Circulatory Physiology*, Vol. 297, pp. H495-H507, ISSN: 0363-6135, eISSN: 1522-1539.
- Chedea, V. S., Vicaș, S.I., Oprea, C. & Socaciu, C. (2006). Lipoxygenase kinetic in relation with lipoxygenase's inhibition by quercetin, *Proceedings of the XXXVI Annual Meeting of the European Society for New Methods in Agricultural Research (ESNA)*, 10 – 14 September 2006, Iasi-Romania, pp. 449-456, ISBN (10) 973-7921-81-X ; ISBN (13) 978-973-7921-81-9.
- Chedea, V.S., Braicu, C. & Socaciu, C. (2010). Antioxidant/prooxidant activity of a polyphenolic grape seed extract, *Food Chemistry*, Vol. 121, pp. 132-139.
- Coffa, G., Schneider, C. & Brash, A.R. (2005). A comprehensive model of positional and stereo control in lipoxygenases. *Biochemical and Biophysical Research Communications*, Vol. 338, pp. 87-92, ISSN: 0006-291X.
- Cucurou, C., Battioni, J. P., Daniel, R. & Mansuy, D. (1991). Peroxidase-like activity of lipoxygenase: different substrate specificity of potato 5-lipoxygenase and soybean 15-lipoxygenase and particular affinity of vitamin E derivatives for the 5-lipoxygenase. *Biochimica et Biophysica Acta*, Vol. 1081, pp. 99-105.
- Cumbee, B., Hildebrand, D.F. & Addo, K. (1997). Soybean flour lipoxygenase isozymes effects on wheat flour dough rheological and breadmaking properties. *Journal of Food Science*, Vol. 62, pp. 281-283, ISSN:1750-3841.
- dos Anjos Ferreira, A. L., Yeum, K. J., Russell, R. M., Krinsky, N. I. & Tang, G. (2004). Enzymatic and oxidative metabolites of lycopene. *The Journal of Nutritional Biochemistry*, Vol. 15, pp. 493-502, ISSN: 0955-2863.
- Duroudier, N.P., Tulah, A.S. & Sayers, I. (2009). Leukotriene pathway genetics and pharmacogenetics in allergy. *Allergy*, Vol. 64, pp. 823-839.
- Feussner, I. & Wasternack, C. (2002). The Lipoxygenase Pathway. *Annual Review in Plant Biology*, Vol. 53, pp. 275-297.
- Fiorucci, S., Golebiowski, J., Cabrol-Bass, D. & Antonczak, S. (2007). DFT Study of quercetin activated forms involved in antiradical, antioxidant, and prooxidant biological processes, *Journal of Agricultural and Food Chemistry*, Vol. 55, pp. 903-911.
- Fiorucci, S., Golebiowski, J., Cabrol-Bass, D. & Antonczak, S. (2008). Molecular simulations enlighten the binding mode of quercetin to lipoxygenase-3. *Proteins*, Vol. 73, pp. 290-298.
- Fukushima, D. (1994). Recent progress on biotechnology of soybean proteins and soybean protein food products. *Food Biotechnology*, Vol. 8, pp.83-135.
- Galati, G. & O'Brien, P. J. (2004). Potential toxicity of flavonoids and other dietary phenolics: Significance for their chemopreventive and anticancer properties. *Free Radical Biology & Medicine*, Vol. 37, pp. 287-303, ISSN: 0891-5849.
- Gardner, H.W. (1991). Recent investigations into the lipoxygenase pathway in plants. *Biochimica Biophysica Acta*, Vol. 1084, No. 3, pp. 221-239.

- Gillmor, S. A., Villasenor, A., Fletterick, R., Sigal, E. & Browner, M.F. (1997). The structure of mammalian 15-Lipoxygenase reveals similarity to the lipases and the determinants of substrate specificity. *Nature Structural & Molecular Biology*, Vol. 4, pp. 1003–1009, ISSN: 1545-9993.
- Gliszczynska-Swiglo, A., van der Woude, H., de Haan, L., Tyrakowska, B., Aarts, J. M. M. J. G. & Rietjens, I. M. C. M. (2003). The role of quinone reductase (NQO1) and quinone chemistry in quercetin cytotoxicity. *Toxicology in Vitro*, Vol. 17, pp. 423-431, ISSN: 0887-2333.
- Gökmen, V., Bahçeci, S. & Acar, J. (2002). Characterization of crude lipoxygenase extract from green pea using a modified spectrophotometric method. *European Food Research and Technology*, Vol. 215, pp. 42–45, ISSN:1438-2377.
- Guthrie, N. & Carroll, K. K. (1999). Specific versus non-specific effects of dietary fat on carcinogenesis. *Progress in Lipid Research*, Vol. 38(3), pp. 261-271, ISSN: 0163-7827.
- Hazai, E., Bikádi, Z., Zsila, F. & Lockwood, S. F. (2006). Molecular modeling of the non-covalent binding of the dietary tomato carotenoids lycopene and lycophyll, and selected oxidative metabolites with 5-lipoxygenase. *Bioorganic and Medicinal Chemistry*, Vol. 14, pp. 6859–6867.
- Hersberger, M. (2010). Potential role of the lipoxygenase derived lipid mediators in atherosclerosis: leukotrienes, lipoxins and resolvins. *Clinical Chemistry and Laboratory Medicine*, Vol. 48(8), pp.1063–1073, ISSN: 1434-6621.
- Hildebrand, D.F. & Hymowitz, T. (1982). Carotene and chlorophyll bleaching by soybeans with and without seed lipoxygenase-1. *Journal of Agricultural and Food Chemistry*, Vol. 30, pp. 705–708, ISSN: 0021-8561.
- Hirakawa, K., Oikawa, S., Hiraku, Y., Hirosawa, I. & Kawanishi, S. (2002). Catechol and hydroquinone have different redox properties responsible for their differential DNA-damaging ability. *Chemical Research in Toxicology*, Vol. 15, pp.76-82, ISSN: 0893-228X.
- Ho, C. T., Chen, Q., Shi, H. & Rosen, R. T. (1992). Antioxidative effect of polyphenol extract prepared from various Chinese teas. *Preventive Medicine*, Vol. 21, pp. 520-525, ISSN: 0091-7435.
- Hong, J.G. & Yang, C.S. (2003). Effects of tea polyphenols on arachidonic acid metabolism in human colon. In F. Shahidi et al. (Eds.), ACS symposium series: Vol. 851. *Food factors in health promotion and disease prevention* (pp. 27–38). Washington, DC: American Chemical Society.
- Hu, J., Huang, Y., Xiong, M., Luo, S. & Chen, Y. (2006). The effects of natural flavonoids on lipoxygenase-mediated oxidation of compounds with benzene ring structure—a new possible mechanism of flavonoid anti-chemical carcinogenesis and other toxicities. *International Journal of Toxicology*, Vol. 25, pp. 295-301.
- Hussain, T., Gupta, S., Adhami, V. M. & Mukhtar, H. (2005). Green tea constituent epigallocatechin-3-gallate selectively inhibits COX-2 without affecting COX-1 expression in human prostate carcinoma cells, *International Journal of Cancer*, Vol. 113, pp. 660–669.
- Ikediyobi, C.O. & Snyder, H.E. (1977). Co-oxidation of β -carotene by an isoenzyme of soybean lipoxygenase. *Journal of Agricultural and Food Chemistry*, Vol. 25, pp. 124–127, ISSN: 0021-8561.

- Jachak, S. M. (2006). Cyclooxygenase inhibitory natural products: current status. *Current Medicinal Chemistry*, Vol. 13, pp. 659-678, ISSN: 0929-8673.
- Jaren-Galan, M. & Minguez-Mosquera, M.I. (1997). β -Carotene and capsanthin co-oxidation by lipoxygenase: kinetic and thermodynamic aspects of reaction. *Journal of Agricultural and Food Chemistry*, Vol. 45, pp. 4814-4820, ISSN: 0021-8561.
- Jaren-Galan, M., Carmona-Ramon, C. & Minguez-Mosquera, M.I. (1999). Interaction between chloroplast pigments and lipoxygenase enzymatic extract of olives. *Journal of Agricultural and Food Chemistry*, Vol. 47, pp. 2671-2677, ISSN: 0021-8561.
- Kanner, J., German, J. B. & Kinsella, J. E. (1987). Initiation of lipid peroxidation in biological systems. *CRC Critical Reviews in Food Science and Nutrition*, Vol. 25, pp. 317-364.
- Ke, P. J. & Ackman, R. G. (1976). Metal catalyzed oxidation in mackerel skin and meat lipids. *Journal of the American Oil Chemists Society*, Vol. 53, pp. 636-640.
- Kemal, C., Louis-Flamberg, P., Krupinski-Olsen, R. & Shorter, A.L. (1987). Reductive inactivation of soybean lipoxygenase 1 by catechols: a possible mechanism for regulation of lipoxygenase activity. *Biochemistry*, Vol. 26, No. 22, pp. 7064-7072.
- Klurfeld, D.M. & Bull, A.W. (1997). Fatty acids and colon cancer in experimental models, *The American Journal of Clinical Nutrition*, Vol. 66, pp. 1530-1538, ISSN: 0002-9165.
- Kobayashi, N., Kaneda, H., Kano, Y. & Koshino, S. (1993). The production of linoleic and linolenic acid hydroperoxides during mashing. *Journal of Fermentation and Bioengineering*. Vol. 76, pp. 371-375.
- Kulkarni, A.P. (2001). Lipoxygenase - a versatile biocatalyst for biotransformation of endobiotics and xenobiotics. *Cellular and Molecular Life Sciences*, Vol. 58, pp. 1805-1825.
- Larreta-Garde, V. (1995). Lipoxygenase in making breads, cookies and crackers. *Oléagineux Corps Gras Lipides*, Vol. 2, pp. 363-365.
- Lekli, I., Ray, D. & Das, D.K. (2010). Longevity nutrients resveratrol, wines and grapes. *Genes and Nutrition*, Vol. 5, pp. 55-60, ISSN: 1555-8932.
- Lin, C. C. & Lin, C. S. (2005). Enhancement of the storage quality of frozen bonito fillets by glazing with tea extracts, *Food Control*, Vol. 16, pp. 169-175.
- Lipkin, M., Yang, K., Edelman, W., Xue, L., Fan, K., Risio, M., Newmark, H. & Kucherlapati R. (1999). Preclinical mouse models for cancer chemoprevention studies. *Annals of the New York Academy of Sciences*, Vol. 889, pp. 14-19.
- Loiseau, J., Benoît, L. V., Macherel, M.-H. & Le Deunff, Y. (2001). Seed lipoxygenases: occurrence and functions. *Seed Science Research*, Vol. 11, pp. 199-211.
- Lomnitski, L., Bar-Natan, R., Sklan, D. & Grossman, S. (1993). The interactions between β -carotene and lipoxygenase in plant and animal systems. *Biochimica Biophysica Acta*, Vol. 1167, pp. 331-338.
- Ma, D.F., Qin, L.Q., Wang, P.Y. & Katoh, R. (2008). Soy isoflavone intake inhibits bone resorption and stimulates bone formation in menopausal women: meta-analysis of randomized controlled trials. *European Journal of Clinical Nutrition*, Vol. 62, pp. 155-161, ISSN: 0954-3007.
- Maccarrone, M., van Aarie, P.G.M., Veldink, G.A. & Vliegthart, J.F.G. (1994). In vitro oxygenation of soybean biomembranes by lipoxygenase-2. *Biochimica et Biophysica Acta*, Vol. 1190, pp. 164-169.


- Macdonald, R.S., Guo, J., Copeland, J., Browning, J.D. JR., Sleper, D., Rottinghaus, G.E. & Berhow, M.A. (2005). Environmental influences on isoflavones and saponins in soybeans and their role in colon cancer. *Journal of Nutrition*, Vol. 135, pp. 1239-1242.
- Mahesha, H.G., Singh, S.A. & Appu Rao, A.G. (2007). Inhibition of lipoxygenase by soy isoflavones: Evidence of isoflavones as redox inhibitors. *Archives of Biochemistry and Biophysics*, Vol. 461, pp. 176-185.
- Malencic, D., Popovic, M. & Miladinovic, J. (2007). Phenolic content and antioxidant properties of soybean (*Glycine max* (L.) Merr.) seeds, *Molecules*, Vol. 12, pp. 576-581.
- Mansuy, D., Cucurou, C., Biatry, B. & Battioni, J. P. (1988). Soybean lipoxygenase-catalyzed oxidations by linoleic acid hydroperoxide: different reducing substrates and dehydrogenation of phenidone and BW755C. *Biochemical and Biophysical Research Communications*, Vol. 151, pp. 339-346.
- Matsumura, K., Tanaka, T. & Kawashima, H. (2008). Involvement of the estrogen receptor beta in genistein induced expression of p21 (waf1/cip1) in PC-3 prostate cancer cells. *Anticancer Research*, Vol. 28, pp. 709-714.
- Mayne, S.T. (2010). Carotenoids a colorful and timely research field. In *Carotenoids, physical, chemical and biological functions and properties* edited by Landrum T., CRC Press Taylor and Francis Group, USA.
- Metodiewa, D., Jaiswal, A. K., Cenas, N., Dickancaite, E. & Segura-Aguilar, J. (1999). Quercetin may act as a cytotoxic prooxidant after its metabolic activation to semiquinone and quinoidal product. *Free Radical Biology and Medicine*, Vol. 26, pp. 107-116.
- Minor, W., Steczko, J., Stec, B., Otwinowski, Z., Bolin, J.T., Walter, R. & Axelrod, B. (1996). Crystal structure of soybean lipoxygenase L-1 at 1.4 Å resolution. *Biochemistry*, Vol. 35 pp. 10687-10701.
- Mochizuki, N. & Kwon, Y.-G. (2008). 15-Lipoxygenase-1 in the vasculature: Expanding roles in angiogenesis, *Circulation Research*, Vol.102, pp. 143-145.
- Moreno, J.J. (2009). New aspects of the role of hydroxyeicosatetraenoic acids in cell growth and cancer development. *Biochemical Pharmacology*, Vol. 77, pp. 1-10, ISSN: 0006-2952.
- Nelson, M. J. (1988). Catecholate complexes of ferric soybean lipoxygenase 1. *Biochemistry*, Vol. 27, pp. 4273-4278.
- Nicolas, J. & Potus, J. (1994). Enzymatic oxidation phenomena and coupled oxidations – Effects of lipoxygenase in bread making and of polyphenol oxidase in fruit technology. *Sciences des Aliments*, Vol. 14, pp. 627-642.
- Nunez-Delicado, E., Sanchez-Ferrer, A. & Garcia-Carmona, F. (1997). Hydroperoxidative oxidation of diethylstilbestrol by lipoxygenase. *Archives of Biochemistry and Biophysics*, Vol. 348, pp. 411-414, ISSN: 0003-9861.
- Pastore, D., Trono, D., Padalino, L., Simone, S., Valenti, D., Fronzo, N.D. & Passarella, S. (2000). Inhibition by α -tocopherol and L-ascorbate of linoleate hydroperoxidation and β -carotene bleaching activities in durum wheat semolina. *Journal of Cereal Science*, Vol. 31, pp. 41-54, ISSN: 0733-5210.
- Pham, C., Jankun, J., Skrzypczak-Jankun, E., Flowers, R.A. & Funk, M.O. (1998). Structural and thermochemical characterization of lipoxygenase-catechol complexes. *Biochemistry*, Vol. 37, pp. 17952-17957.

- Pidgeon, G. P., Lysaght, J., Krishnamoorthy, S., Reynolds, J. V., O'Byrne, K., Nie, D. & Honn, K. V. (2007). Lipoxygenase metabolism: roles in tumor progression and survival. *Cancer and Metastasis Reviews*, Vol. 26, pp. 503–524, ISSN: 0167-7659.
- Prigge, S.T., Boyington, J.C., Faig M., Doctor, K.S., Gaffney, B.J. & Amzel, L.M. (1997). Structure and mechanism of lipoxygenases. *Biochimie*, Vol. 79, pp. 629-636.
- Ramadoss, C.S., Pistorius, E.K. & Axelrod, B. (1978). Coupled oxidation of carotene by lipoxygenase requires two isoenzymes. *Archives of Biochemistry and Biophysics*, Vol. 190, pp. 549–552.
- Robinson, D.S., Wu, Z.C., Domoney, C. & Casey, R. (1995). Lipoxygenases and the quality of foods. *Food Chemistry*, Vol. 54, pp. 33–43.
- Sadik, C.D., Sies, H. & Schewe, T. (2003). Inhibition of 15-lipoxygenases by flavonoids: structure–activity relations and mode of action. *Biochemical Pharmacology*, Vol. 65, pp. 773-781.
- Sakthivelu, G., A. Devi, M.K., Giridhar, P., Rajasekaran, T., Ravishankar, G.A., Nikolova, M.T., Angelov, G.B., Todorova, R.M. & Kosturkova, G.P. (2008). Isoflavone composition, phenol content, and antioxidant activity of soybean seeds from India and Bulgaria. *Journal of Agricultural and Food Chemistry*, Vol. 56, pp. 2090-2095, ISSN: 0021-8561.
- Schewe, T. & Sies, H. (2003). Flavonoids as protectants against prooxidant enzymes, *Research monographs*, Institut für Physiologische Chemie I; Heinrich-Heine-Universität Düsseldorf– Date of access: 7.07.2008. Available from: <http://www.uni-duesseldorf.de/WWW/MedFak/PhysiolChem/index.html>.
- Schewe, T., Kühn, H. & Rapoport, S. (1986). Positional specificity of lipoxygenases and their suitability for testing potential drugs. *Prostaglandins, Leukotrienes & Medicine*, Vol. 23, pp. 155-160, ISSN: 0952-3278.
- Schewe, T., Kühn, H. & Sies H. (2002). Flavonoids of cocoa inhibit recombinant human 5-Lipoxygenase. *The Journal of Nutrition*, Vol. 56, pp. 1825-1829.
- Schewe, T., Sadik, C., Klotz, L.-O., Yoshimoto, T., Kühn, H. & Sies, H. (2002). Polyphenols of cocoa: inhibition of mammalian 15-Lipoxygenase. *Biological Chemistry*, Vol. 382, pp. 1687– 1696.
- Schneider, I. & Bucar^a F. (2005). Lipoxygenase inhibitors from natural plant sources. Part I Medicinal plants with action on arachidonate 5-lipoxygenase. *Phytotherapy Research*, Vol. 19, pp. 81-102.
- Schneider, I. and Bucar^b F. (2005). Lipoxygenase inhibitors from natural plant sources. Part II Medicinal plants with action on arachidonate 12-lipoxygenase, 15-lipoxygenase and leukotriene receptor antagonists. *Phytotherapy Research*, Vol. 19, pp. 263-272.
- Serpen, A. & Gökmen, V. (2006). A proposed mechanism for the inhibition of soybean lipoxygenase by β -carotene. *Journal of the Science of Food and Agriculture*, Vol. 86, pp. 401–406.
- Seto, Y., Lin, C. C., Endo, Y. & Fujimoto, K. (2005). Retardation of lipid oxidation in blue sprat by hot water tea extracts, *Journal of the Science of Food and Agriculture*, Vol. 85, pp. 1119–1124.
- Siedow, J.N. (1991). Plant lipoxygenases. Structure and function. *Annual Review of Plant Physiology and Plant Molecular Biology*, Vol. 42, pp. 145-188.

- Skrzypczak-Jankun, E., Amzel, L.M., Kroa, B.A. & Funk, M.O. (1997). Structure of soybean lipoxygenase L3 and a comparison with its L1 isoenzyme. *Proteins*, Vol. 29, pp. 15-31.
- Skrzypczak-Jankun, E., Zhou, K. & Jankun J. (2003). Inhibition of lipoxygenase by (-)-epigallocatechin gallate: X-ray analysis at 2.1 Å reveals degradation of EGCG and shows soybean LOX-3 complex with EGC instead. *International Journal of Molecular Medicine*, Vol. 12, pp. 415-420.
- Skrzypczak-Jankun, E., Chorostowska-Wynimko, J., Selman, S.H., & Jankun, J. (2007). Lipoxygenases – A challenging problem in enzyme inhibition and drug development. *Current Enzyme Inhibition*, Vol. 3, pp. 119-132.
- Skrzypczak-Jankun, E., Zhou, K., McCabe, N.P., Kernstock, R., Selman, S.H., Funk, M.O., Jr. & Jankun, J. (2002). In *Natural Polyphenols in Complexes with Lipoxygenase - Structural Studies and their Relevance to Cancer*, pp. 205. 93rd Annual Meeting AACR, San Francisco, 2002; AACR: San Francisco.
- Song, T.B., Buseman, K. & Murphy, G. (1998). Soy isoflavone analysis: quality control and a new internal standard. *American Journal of Clinical Nutrition*, Vol. 68, pp. 1474-1479.
- Spaapen, L. J., Verhagen, J., Veldink, G. A. & Vliegthart, J. F. (1980). Properties of a complex of Fe(III)-soybean lipoxygenase-1 and 4-nitrocatechol. *Biochimica and Biophysica Acta*, Vol. 617, No.1, pp. 132-140.
- Sumner, J.B. & Sumner, R.J. (1940). The coupled oxidation of carotene and fat by carotene oxidase. *Journal of Biological Chemistry*, Vol. 134, pp. 531-533.
- Szefel, J., Piotrowska, M., Kruszewski, W.J., Jankun, J., Lysiak-Szydłowska, W. & Skrzypczak-Jankun, E. (2011). Eicosanoids in prevention and management of diseases, *Current Molecular Medicine*, Vol. 11, pp. 13-25.
- Szymanowska, U., Jakubczyk, A., Baraniak, B. & Kur, A. (2009). Characterisation of lipoxygenase from pea seeds (*Pisum sativum* var. Telephone L.). *Food Chemistry*, Vol. 116, pp. 906-910.
- Theorell, H., Holman, R. T. & Åkeson, Å. (1947). Crystalline lipoxidase. *Acta Chemica Scandinavica*, Vol. 1, pp. 571-576.
- Trono, D., Pastore, D. & Fonzo, N.D. (1999). Carotenoid dependent inhibition of durum wheat lipoxygenase. *Journal of Cereal Science*, Vol. 29, pp. 99-102.
- van der Woude, Boersma, H., Alink, M. G., Vervoort, G. M., J. & Rietjens, I. M. C. M. (2006). Consequences of quercetin methylation for its covalent glutathione and DNA adduct formation. *Chemico-Biological Interactions*, Vol. 160, pp. 193-203.
- Vicaș, S. I., Chedea, V. S. & Socaciu, C. (2011). Inhibitory effects of isoflavones on soybean lipoxygenase-1 activity, *Journal of Food Biochemistry*, Vol. 35, pp. 613-627.
- Walther, M., Holzhutter, H.G., Kuban, R.J., Wiesner, R., Rathmann, J. & Kuhn, H. (1999). The inhibition of mammalian 15-lipoxygenases by the anti-inflammatory drug ebselen: dual-type mechanism involving covalent linkage and alteration of the iron ligand sphere *Molecular Pharmacology*, Vol. 56, pp. 196-203.
- Warri, A., Saarinen, N.M. & Makela, S. (2008). The role of early life genistein exposures in modifying breast cancer risk. *British Journal of Cancer*, Vol. 98, pp. 1485-1493, ISSN 0007-0920.
- Willett, W.C. (1997). Nutrition and cancer. *Salud Publica Mexican*, Vol. 39, No. 4, pp. 298-309.

- Wu, Z., Robinson, D. S., Hughes, R. K., Casey, R., Hardy, D. & West, S. I. (1999). Co-oxidation of β -carotene catalysed by soybean and recombinant pea lipoxygenases. *Journal of Agricultural and Food Chemistry*, Vol. 47, pp. 4899-4906.
- Xiao, C. W. (2008). Health effects of soy protein and isoflavones in humans. *Journal of Nutrition*, Vol. 138, No.6, pp. 1244 - 1249.
- Yoon, S. & Klein, B.P. (1979). Some properties of pea lipoxygenase isoenzymes. *Journal of Agricultural and Food Chemistry*, Vol. 27, pp. 955-962, ISSN: 0021-8561.
- Zimeri, J. & Tong, C.H. (1999). Degradation kinetics of (-)epigallocatechin gallate as a function of pH and dissolved oxygen in a liquid model system. *Journal of Food Science*, Vol. 64, pp. 753-758.

IntechOpen



Recent Trends for Enhancing the Diversity and Quality of Soybean Products

Edited by Prof. Dora Krezhova

ISBN 978-953-307-533-4

Hard cover, 536 pages

Publisher InTech

Published online 28, October, 2011

Published in print edition October, 2011

This book presents new aspects and technologies for the applicability of soybean and soybean products in industry (human food, livestock feed, oil and biodiesel production, textile, medicine) as well as for future uses of some soybean sub-products. The contributions are organized in two sections considering soybean in aspects of food, nutrition and health and modern processing technologies. Each of the sections covers a wide range of topics. The authors are from many countries all over the world and this clearly shows that the soybean research and applications are of global significance.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Veronica Sanda Chedea and Mitsuo Jisaka (2011). Inhibition of Soybean Lipoxygenases – Structural and Activity Models for the Lipoxygenase Isoenzymes Family, Recent Trends for Enhancing the Diversity and Quality of Soybean Products, Prof. Dora Krezhova (Ed.), ISBN: 978-953-307-533-4, InTech, Available from: <http://www.intechopen.com/books/recent-trends-for-enhancing-the-diversity-and-quality-of-soybean-products/inhibition-of-soybean-lipoxygenases-structural-and-activity-models-for-the-lipoxygenase-isoenzymes-f>

INTECH
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2011 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen