

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,900

Open access books available

186,000

International authors and editors

200M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



Heat-Transfer-Model Analysis of the Thermal Effect of Intrusive Sills on Organic-Rich Host Rocks in Sedimentary Basins

Dayong Wang, Minglong Zhao and Tian Qi

*Key Laboratory of Ocean Energy Utilization and Energy Conservation of Ministry of Education, School of Energy and Power Engineering, Dalian University of Technology
China*

1. Introduction

Numerous geological explorations demonstrate that magmatic intrusions may increase the geothermal gradient in sedimentary basins, accelerating the thermal maturation of organic matter in strata, and promoting the hydrocarbon generation (Fjeldskaar et al., 2008; Jones et al., 2007). They may also be beneficial to the migration and accumulation of oil and gas by providing them with pathways, reservoirs, covering conditions and trapping constructions (Feng and Tang, 1997; Li, 2000; Othman and Ward, 2002; Othman et al., 2001; Wang et al., 1990). Therefore, it is of great significance to study the thermal effect of igneous intrusions on organic-rich host rocks.

A lot of the organic-rich host rocks are argillaceous rocks, e.g., shales in the DSDP 41-368 hole near Cape Verde Rise in eastern Atlantic and mudstones in Xia 38 well block in the Huimin Sag of Bohai Bay, which generally have the relatively low permeability (e.g., $<10^{-16}$ m²). Under such circumstances, the hydrothermal convection in host rocks can be reasonably ignored, and heat conduction models can be used to approximately describe the heat transfer in host rocks (Hanson, 1995; Hayba and Ingebritsen, 1997). Thus, these models are often used as a geothermometer to indicate the temperature range in which the thermal metamorphism of these host rocks takes place (Barker et al., 1998; Santos et al., 2009; Stewarta, 2005; Turcotte and Schubert, 1982; Wang et al., 2007, 2008). Several types of heat conduction models have been constructed and used in some geological researches (Galushkin, 1997; Wang et al., 2007, 2011). However, only a small portion of these researches specially compare and quantify the difference in the prediction results of different heat conduction models (Jeager, 1959; Galushkin, 1997; Wang et al., 2011). Due to the apparent importance of the accuracy of heat conduction models in these researches, it is still required to further explore and distinguish the applicable conditions of these models based on some geological cases.

In this study, we accordingly investigate the difference in the prediction results of three types of commonly used heat conduction models by taking an intrusive sill in the Bellata-1 Well in the Gunnedah Basin, Australia as an example. These models assume different intrusion mechanisms of magma and different evolution states of pore water during cooling of magma. By comparing the prediction results of these models with the measured vitrinite-reflectance (R_o) geothermometer, we also discuss the potential intrusion mechanism of the sill and the state of pore water in host rocks during cooling of the sill.

2. Geology of the Bellata-1 Well in the Gunnedah Basin, Australia

The Bellata-1 Well is located in the Gunnedah Basin of northern Australia and intersects Permian, Triassic, Jurassic and Cretaceous strata in turn (Fig 1). The total thickness of the Permian and Triassic strata reaches 451 m, overlain by the 640m thick Jurassic and Cretaceous sediments. A 15.68m thick mafic basaltic sill was found in the lower part of Triassic Napperby Formation. The intrusion of the sill took place between the Late Triassic and Early Jurassic with a current burial depth of 847.60 m (Othman and Ward, 2002; Othman et al., 2001). The Triassic strata are mainly composed of organic-rich mudstones. The R_o profile adjacent to the sill shows the effect of the significant local heating: the R_o value can be as high as up to 2.43% within the contact aureole, whereas the R_o value in the unaffected parts is only 0.57-0.74%. The oil generated by these organic-rich rocks due to the thermal effect of the intrusive sill is found in the Jurassic Pilliga Sandstone (Othman et al., 2001). Therefore, the igneous sill of the Bellata-1 Well and its host rocks constitute an ideal geological example for numerically investigating the thermal effects of igneous intrusions on organic-rich host rocks.




Fig. 1. Stratigraphic characteristics and vitrinite reflectance profile around an igneous sill in Well Bellata-1 (Othman and Ward, 2002; Othman et al., 2001)

3. Method

3.1 Heat conduction models

Some general assumptions are required in constructing heat conduction models: 1) The shape of the intrusion is regular, dike-like or sill-like; 2) Convection motion in the intrusion is not considered; 3) Heat loss due to the escape of volatiles is neglected. Thus, the basic heat conduction equations in one dimension which can be used to describe the heat transfer

between intrusive magmas and host rocks are expressed (Barker *et al.*, 1998; Shan *et al.*, 1998; Stewart *et al.*, 2005):

For magma intrusions :

$$\frac{\partial}{\partial Z} \left(K_{\text{magma}} \cdot \frac{\partial T}{\partial Z} \right) = \rho_{\text{magma}} \cdot \frac{H}{L_2 - L_1} \cdot \frac{\partial T}{\partial t} + \frac{\partial(\rho_{\text{magma}} \cdot C_{\text{magma}} \cdot T)}{\partial t} \quad (1)$$

For host rocks :

$$\frac{\partial}{\partial Z} \left([1 + [Nu]] \cdot K_{\text{host}} \right) \cdot \frac{\partial T}{\partial Z} = \frac{\partial}{\partial t} (\rho_{\text{host}} \cdot C_{\text{host}} \cdot T) + A_1 + [A_2] \quad (2)$$

Where T is the temperature; t represents the time; K means the thermal conductivity; C is the specific heat; ρ denotes the density; A_1 and A_2 represent the latent heat consumed by the dehydration and decarbonation reactions and pore-water volatilization per unit volume of host rocks and per unit time; the subscripts, i.e., magma and host, denote magma and host rocks, respectively. H represents the latent crystallization heat of melted magma; $L_1 - L_2$ is the crystallization temperature range of intrusive magma. Nu represents the Nusselt number and can be used to implement approximately the hydrothermal convection in host rocks (Galushkin, 1997). In this study, a finite difference method is used to obtain the numerical solution of Eqns. (1) and (2).

3.2 Model parameters

As the host rocks were located on or near the ground surface at the intrusion moment of the sill (between Late Triassic and Early Jurassic), we approximately assume that the strata with the current depth of 640 m is located on the ground surface during the intruding of magma. The surface temperature and the geothermal gradient are assumed to be approximately equal to 25 °C and 30 °C/Km, respectively. The temperature of the melted mafic magma is usually about 1250 °C (Barker *et al.*, 1998; Wohletz, 1999). Its thermal conductivity and density are usually equal to 2.1 J m⁻¹ s⁻¹ °C⁻¹ and 2700 Kg/m³ (Barker *et al.*, 1998; Wohletz, 1999), respectively. We specify the specific heat of the sill to be equal to 1200 J/Kg (Galushkin 1997; Wang *et al.*, 2010). The latent heat of crystallization of melted magma equals 400 KJ/Kg, and the corresponding crystallization temperature range is 1150 °C - 1250 °C. According to Wang *et al.* (2007), Organic-rich mudstones generally have the relatively low thermal conductivity. For example, the thermal conductivity of mudstones in the region of England is 1.4-1.6 W/mK (Midttømme *et al.*, 1998), and the thermal conductivity of mudstones at the depth of 850 m in the Huimin Sag of Bohai Bay is about 1.4 W/mK (Wang *et al.*, 2007). The specific heat and density of mudstone matrix can be specified to be equal to 820 J/Kg and 2700 Kg/m³, respectively. We specify 1.9 J m⁻¹ s⁻¹ °C⁻¹ as its thermal conductivity. This value is the same with that of the host rocks of the diabase sill of Well Xia38 in the Huimin Sag of Bohai Bay, as the latter has the same lithology and the similar intrusion depth with our example. At the buried depth of about 200 m, the boiling point of pore water may reach 200 °C, and its latent volatilization heat is about 1939.73 KJ/Kg. The porosity of the host rock is about 0.5 in terms of the depth - porosity relationship of mudstones (Allen and Allen, 2005). We calculate the total specific heat and total thermal conductivity of host rocks based on the computational equations of Galushkin (1997), Travis *et al.* (1991), Wang *et al.* (2007) and Wohletz *et al.* (1999).

3.3 Simulated cases

Three types of one-dimensional heat conduction models are built to simulate the heat transfer between the sill and its host rocks (Table 1). We adopt the method of Galushkin (1997) to implement the finite-time intrusion mechanism of magma: The temperature at the axis of the sill is set as 300 °C when the sill begins to form; and the time of the pre-cooled shell formation is equal to 2.2 hours; the total time of the sill formation is about 4.4 hours.

In order to verify the applicability of these three heat conduction models to the modeled sill, we need to compare the prediction results of the models with the measured vitrinite-reflectance (R_o) geothermometer. We adopt the vitrinite reflectance - the peak-temperature (T_{peak}) relation (i.e. $T_{peak} = (\ln R_o + 1.19) / 0.00782$) of *Barker et al.* (1998) to calculate the T_{peak} of the overlying host rocks based on the measured R_o values and then compare it with the predictions of the models.

Case No.	Intrusion mechanism of magma	Pore-water volatilization	Hydrothermal convection in overlying host rocks	Dehydration and decarbonation of host rocks
1	instantaneous	not considered	not considered	considered
2	instantaneous	considered	not considered	considered
3	finite-time	not considered	not considered	considered

Table 1. Three cases for simulation

4. Results and discussion

The T_{peak} profiles of host rocks predicted by three types of heat conduction models are shown in Fig. 2. The contact temperature (T_c) predicted by Case 2 reaches 852 °C, and is higher than that predicted by the other cases. Actually, pore-water volatilization can decrease the thermal conductivity of host rocks. As a result, the diffusion of the heat of the sill in host rocks is depressed, and near the contact, heat from the sill congregates and rapidly increases the contact temperature. Comparably, the T_c predicted by Case 3 is lowest and only reaches 706 °C. This is apparently due to the heat loss caused by the pre-cooled shell of the sill compared to the instantaneous intrusion mechanism. In addition, the computation based on Case 1 deduces the highest degree of the thermal effect of the sill on its host rocks, whereas the prediction from Case 3 results in the lowest one. This indicates that the intrusion mechanism of magma may play a more important role in lowering the thermal effect of the intrusion than the heat sinks in host rocks.

By comparing the predicted T_{peak} with the measured R_o geothermometer, it is obviously observed that the T_{peak} predicted by all of these three models is much lower than the R_o geothermometer in the region where it is 75 m away from the margin of the sill (i.e. $X/D \approx 5$). This demonstrates that the increase in the temperature of strata due to the subsequent sedimentation after cooling of the sill have covered up the thermal influence of the intrusion on host rocks in this region. The heat conduction model assuming the instantaneous intrusion mechanism and ignoring pore-water volatilization matches well with the measured R_o geothermometer. Othman et al. (2001, 2002) once reported that the Napperby Formation is mainly composed

of low-permeability mudstones (shale) and has the abnormal high pressure. Consequently, volatilization and escape of pore water can likely be restricted. This is consistent with the prediction of Case 1. All of these indicate that the instantaneous intrusion mechanism likely represents natural conditions and that the effect of pore-water volatilization is insignificant.




Fig. 2. Comparison between vitrinite-reflectance geothermometer of Baker et al. (1998) and peak temperature of host rocks predicted by three types of heat conduction models, assuming different intrusion mechanisms of magma and the state of pore water during cooling of magma

5. Conclusions

The following conclusions can be made based on the heat-conduction-model analysis of the T_{peak} of the host rocks of a mafic sill of the Bellata-1 Well from the Gunnedah Basin, Australia:

1. The consideration of pore-water volatilization can increase the T_c prediction, while it is converse for the finite-time intrusion mechanism. The computation based on the heat conduction model assuming the instantaneous intrusion mechanism and considering

- pore-water volatilization deduces the highest T_c among three types of heat conduction models, whereas the computation based on the model assuming the finite-time intrusion mechanism and ignoring pore-water volatilization results in the lowest T_c .
2. The degree of thermal effect deduced by the heat conduction model assuming the instantaneous intrusion and ignoring pore-water volatilization is highest, while that deduced by the model assuming the finite-time intrusion and ignoring pore-water volatilization is lowest. This indicates that the intrusion mechanism of magma may play a more important role in lowering the thermal effect of the intrusion than the heat sinks in host rocks.
 3. The heat conduction model assuming the instantaneous intrusion mechanism and ignoring pore-water volatilization matches well with the measured vitrinite-reflectance geothermometer. Considering the real geological characteristics of the host rocks, it can be concluded that the instantaneous intrusion mechanism likely represents natural conditions and that the effect of pore-water volatilization is insignificant.

6. Acknowledgement

This work was financially supported by the National Science Foundation of China (No. 41004031 and 50736001), the 863 program (No. 2008AA062303 and 2006AA09209), the 973 program (No. 2009CB219507), the Ph.D. Programs Foundation of Ministry of Education of China (No. 20100041120039), the Research Foundation of Dalian University of Technology (No. 852003, 893210, 893316) and the Fundamental Research Funds for the Central Universities.

7. References


- Allen P A, Allen J R. Basin Analysis: Principles and Applications [M]. Oxford: Blackwell Scientific Publications, 2005, pp549.
- Barker C E, Boney Y, Lewan M D. Fluid inclusion and vitrinite-reflectance geothermometry compared to heat-flow models of maximum paleotemperature next to dikes, western onshore Gippsland Basin, Australia [J]. *International Journal of Coal Geology*, 1998, 37: 73~111.
- Feng Qiao, Tang Xiyuan. Magma activity's influence on conditions forming oil and gas pools[J]. *Geological Science and Technology Information (In Chinese with English abstract)*, 1997, 16(4): 59~65.
- Fjeldskaar W, Helset H M, Johansen H, et al. Thermal modeling of magmatic intrusions in the Gjallar Ridge, Norwegian Sea: implications for vitrinite reflectance and hydrocarbon maturation [J]. *Basin Research*, 2008, 20: 143-159.
- Galushkin Y I. Thermal effects of igneous intrusions on maturity of organic matter: A possible mechanism of intrusion [J]. *Organic Geochemistry*, 1997, 26(11-12): 645-658.
- Hanson R B. The hydrodynamics of contact metamorphism [J]. *Geological Society of America Bulletin*, 1995, 107: 595-611.
- Hayba D O., Ingebritsen S E. Multiphase groundwater flow near cooling plutons [J]. *Journal of Geophysical Research*, 1997, 102: 2235-12252.
- Jaeger J C. Temperatures outside a cooling intrusive sheet [J]. *American Journal of Science*, 1959, 257: 44~54.

- Jones S F, Wielens H, Williamson M C, Zentilli M. Impact of magmatism on petroleum systems in the Sverdrup Basin, Canadian Arctic Islands, Nunavut: A numerical modeling study [J]. *Journal of Petroleum Geology*, 2007, 30(3): 237-255.
- Li Yahui. Diabase and hydrocarbon reservoir formation on the nor-thern slope of Gaoyou sag[J]. *Journal of Geomechanics (In Chinese with English abstract)*, 2000, 6(2): 17~22.
- Midttømme K, Roaldset E, Aagaard P. Thermal conductivity of selected claystones and mudstones from England [J]. *Clay Minerals*, 1998, 33: 131~145.
- Othman R, Arouri K R, Ward C R, et al. Oil generation by igneous intrusions in the northern Gunnedah Basin, Australia[J]. *Organic Geochemistry*, 2001, 32: 1219~1232.
- Othman R, Ward C R. Thermal maturation pattern in the southern Bowen, northern Gunnedah and Surat Basins, northern New South Wales, Australia[J]. *International Journal of Coal Geology*, 2002, 51: 145~167.
- Santos R V, Dantas E L, de Oliveira C G, de Alvarenga C J, dos Anjos C W, Guimaraes E M, Oliveira F B. Geochemical and thermal effects of a basic sill on black shales and limestones of the Permian Irati Formation [J]. *Journal of South American Earth Sciences*, 2009, 28 (1): 14-24.
- Shan Yehua, Yuan Ding, Li Zhian. Numerical simulation of magmatic metamorphism [J]. *Coal Geology of China (In Chinese with English abstract)*, 1998, 10(2): 19~26.
- Stewart A K, Maseya M, Padgett P L, et al. Influence of a basic intrusion on the vitrinite reflectance and chemistry of the Springfield (No. 5) coal, Harrisburg, Illinois[J]. *International Journal of Coal Geology*, 2005, 63: 58~67.
- Travis B J, Janecky D R, Rosenberg N D. Three-dimensional simulation of hydrothermal circulation at mid-ocean ridges[J]. *Geophysical Research Letters*, 1991, 18(8): 1441~1444.
- Turcotte D L, Schubert G.. *Geodynamics Applications of Continuum Physics to Geological Problems* [M]. New York: Wiley, 1982, 1~450.
- Wang D., Lu X., Song Y., et al. Influence of the Temperature Dependence of Thermal Parameters of Heat Conduction Models on the Reconstruction of Thermal History of Igneous-intrusion-bearing Basins [J]. *Computers & Geosciences*, 2010, 36, 1339-1344, doi:10.1016/j.cageo.2010.03.014.
- Wang D., Lu X., Xu S., Hu W. Comment on "Influence of a basic intrusion on the vitrinite reflectance and chemistry of the Springfield (No. 5) coal, Harrisburg, Illinois" by Stewart et al. (2005) [J]. *International Journal of Coal Geology*, 2008, 73: 196-199.
- Wang D, Lu X, Zhang X, Xu S, Hu W, Wang L. Heat-model analysis of wall rocks below a diabase sill in Huimin Sag, China compared with thermal alteration of mudstone to carbargilite and hornfels and with increase of vitrinite reflectance [J]. *Geophysical Research Letters*, 2007, 34, L16312, doi:10.1029/2007GL030314.
- Wang D., Song Y., Liu W., Zhao M., Oi T. Numerical investigation of the effect of volatilization and the supercritical state of pore water on maturation of organic matter in the vicinity of igneous intrusions[J]. *International Journal of Coal Geology*, 2011, 87: 33-40.
- Wang Youxiao, Fan Pu, Cheng Xuehui, et al. Abnormal geothermal influence on hydrocarbon genesis of sedimentary organics[J], *Oil & Gas Geology (In Chinese with English abstract)*, 1990, 11(1): 73~77.

Wohletz K H, Orsi G, Civetta L. Thermal evolution of the Phlegraean magmatic system [J].
Journal of Volcanology and Geothermal Research, 1999, 91: 381~414.

IntechOpen

IntechOpen



Earth Sciences

Edited by Dr. Imran Ahmad Dar

ISBN 978-953-307-861-8

Hard cover, 648 pages

Publisher InTech

Published online 03, February, 2012

Published in print edition February, 2012

The studies of Earth's history and of the physical and chemical properties of the substances that make up our planet, are of great significance to our understanding both of its past and its future. The geological and other environmental processes on Earth and the composition of the planet are of vital importance in locating and harnessing its resources. This book is primarily written for research scholars, geologists, civil engineers, mining engineers, and environmentalists. Hopefully the text will be used by students, and it will continue to be of value to them throughout their subsequent professional and research careers. This does not mean to infer that the book was written solely or mainly with the student in mind. Indeed from the point of view of the researcher in Earth and Environmental Science it could be argued that this text contains more detail than he will require in his initial studies or research.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Dayong Wang, Minglong Zhao and Tian Qi (2012). Heat-Transfer-Model Analysis of the Thermal Effect of Intrusive Sills on Organic-Rich Host Rocks in Sedimentary Basins, Earth Sciences, Dr. Imran Ahmad Dar (Ed.), ISBN: 978-953-307-861-8, InTech, Available from: <http://www.intechopen.com/books/earth-sciences/heat-transfer-model-analysis-of-the-thermal-effect-of-intrusive-sills-on-organic-rich-host-rocks-in->

INTECH
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2012 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen