

# We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,900

Open access books available

185,000

International authors and editors

200M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index  
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?  
Contact [book.department@intechopen.com](mailto:book.department@intechopen.com)

Numbers displayed above are based on latest data collected.  
For more information visit [www.intechopen.com](http://www.intechopen.com)



# Newly-Recognized Small Molecule Receptors on Human Breast Cancer Cell Integrin $\alpha v \beta 3$ that Affect Tumor Cell Behavior

Hung-Yun Lin<sup>1</sup>, Faith B. Davis<sup>1</sup>, Mary K. Luidens<sup>1,2</sup>, Aleck Hercbergs<sup>3</sup>, Shaker A. Mousa<sup>4</sup>, Dhruva J. Bharali<sup>4</sup> and Paul J. Davis<sup>1,2</sup>

<sup>1</sup>Signal Transduction Laboratory, Ordway Research Institute, Albany, NY,

<sup>2</sup>Albany Medical College, Albany, NY,

<sup>3</sup>Cleveland Clinic, Cleveland, OH,

<sup>4</sup>Pharmaceutical Research Institute, Albany College of Pharmacy, Albany, NY, USA

## 1. Introduction

Hormonal regulation of the growth of breast cancer cells has been largely seen to result from interactions of estrogen and progestins with nuclear receptors for these steroids that may reside, unliganded, in cytoplasm or be transcriptionally active as steroid-protein nuclear receptor complexes. Receptors for nonpeptide hormones exist in the plasma membrane, however, and when activated may stimulate breast cancer cell proliferation. For example, the functional classical estrogen receptor- $\alpha$  (ER $\alpha$ ) is found in the cell membrane. Recently, the identification has been made of a novel receptor for thyroid hormone and for dihydrotestosterone (DHT) on the plasma membrane of cells; this receptor is the  $\alpha v \beta 3$  integrin, which promotes proliferation of human breast cancer cells by these two hormones. Integrins are heterodimeric structural proteins of the plasma membrane whose primary functions are to interact with extracellular matrix proteins and growth factors. One integrin,  $\alpha v \beta 3$ , bears discrete receptors for thyroid hormone (L-thyroxine, T<sub>4</sub>; 3, 5, 3'-triiodo-L-thyronine, T<sub>3</sub>) and for DHT. A receptor for the polyalcohol, resveratrol, also exists on this integrin in breast cancer cells, mediating the anti-proliferative, pro-apoptotic action of this compound. Resveratrol has certain structural features that are estrogen-like. Disparate actions of T<sub>4</sub>, T<sub>3</sub>, DHT and resveratrol that are initiated at the integrin depend downstream upon stimulation of the activity of mitogen-activated protein kinase (MAPK), suggesting the existence of distinct, function-specific pools of MAPK within the cell. Tetraiodothyroacetic acid (tetrac) is a model specific inhibitor of hormone actions on the thyroid hormone integrin receptor. Tetrac and a nanoparticulate formulation of tetrac block stimulation by thyroid hormone analogues of cancer cell proliferation and of angiogenesis. Interestingly, tetrac also acts in the absence of T<sub>4</sub> and T<sub>3</sub> to block the tumor-relevant angiogenic responses to vascular growth factors, e.g., vascular endothelial growth factor (VEGF), basic fibroblast growth factor (bFGF) and other growth factors. Tetrac and nanoparticulate tetrac also

disable expressions of families of genes important to cancer cell survival pathways. This chapter reviews the functions of the several nonpeptide hormone receptors on integrin  $\alpha v \beta 3$ .

The possibility that the hormone-directed biology of the breast cancer cell might be in part regulated from the cell surface was first suggested by the identification of estrogen receptor protein in the plasma membrane (Levin, 1999). Discrete receptors on the plasma membrane  $\alpha v \beta 3$  integrin of breast cancer cells have also recently been described for thyroid hormone (Bergh *et al.*, 2005; Cheng *et al.*, 2010; Davis, P. *et al.*, 2011), for dihydrotestosterone (DHT) (Lin, H. *et al.*, 2009b) and for resveratrol (Lin, H. *et al.*, 2006). The several functions of these membrane receptors include modulation of cancer cell proliferation and, in the case of thyroid hormone, of tumor-relevant angiogenesis (Cheng *et al.*, 2010; Mousa *et al.*, 2009). Expression of integrin  $\alpha v \beta 3$  is concentrated in tumor cells and rapidly-dividing endothelial and vascular smooth muscle cells, so that receptors for these hormones – and resveratrol is a polyalcohol with estrogen-like structural features – may be considered targets for manipulation of breast cancer. In the present review we will describe the features of these receptors in the breast cancer cell, and will also propose clinical therapeutic applications that are based on inhibition of these small molecule plasma membrane receptors.

## 2. Thyroid hormone stimulates human breast cancer cell proliferation via plasma membrane integrin $\alpha v \beta 3$

In the absence of estrogen, thyroid hormone (L-thyroxine [ $T_4$ ]) was shown in 2004 to enhance proliferation of human estrogen receptor- $\alpha$  (ER $\alpha$ )-positive breast cancer cells (Tang *et al.*, 2004). The thyroid hormone effect required extracellular-regulated kinases 1 and 2 (ERK1/2)-dependent phosphorylation of Ser-118 of ER $\alpha$ , precisely mimicking the action of estradiol (Kato *et al.*, 1995) on breast cancer cell proliferation. The nuclear estrogen receptor inhibitor ICI 182,780 (Fulvestrant<sup>®</sup>) blocked this action of thyroid hormone (Lin H. , unpublished), as did tetraiodothyroacetic acid (tetrac), an analogue of  $T_4$  in which the alanine side chain of  $T_4$  is converted to acetic acid (Tang *et al.*, 2004). Tetrac is an inhibitor of actions of  $T_4$  and  $T_3$  that are initiated at the thyroid hormone receptor site on integrin  $\alpha v \beta 3$  (Bergh *et al.*, 2005; Yalcin *et al.*, 2010a; Yalcin *et al.*, 2010b; Davis, F. *et al.*, 2006). Studied *in vitro*,  $T_4$  and  $T_3$  are anti-apoptotic in breast cancer cells (Tang *et al.*, 2004) and other tumor cells (Lin *et al.*, 2008a), at least in part via a mechanism that blocks effectiveness of the oncogene suppressor protein and pro-apoptotic factor, p53 (Lin *et al.*, 2011).

Acting via the plasma membrane integrin receptor, agonist thyroid hormone analogues  $T_4$  and  $T_3$  are also pro-angiogenic (Mousa *et al.*, 2009; Davis, P. *et al.*, 2009; Luidens *et al.*, 2010). In the absence of tumor cells and in the setting of tissue ischemia, stimulation of neovascularization by  $T_4$  and  $T_3$  may be desirable (Tomanek *et al.*, 1998; Chen *et al.*, 2010). In the setting of cancer, however, these agents appear to enhance tumor-related angiogenesis. The mechanism of angiogenesis is complex and involves the release by endothelial cells of vascular growth factors (Tomanek *et al.*, 1998) and the resulting autocrine effects of such factors.  $T_4$  and  $T_3$  may also enhance the actions of vascular growth factors (Davis, F. *et al.*, 2004).

The effectiveness of tetrac as an inhibitor of actions of  $T_4$  and  $T_3$  on the plasma membrane of cancer cells caused us to study tetrac as an anti-cancer and anti-angiogenic agent. We found

that unmodified tetrac as well as tetrac re-formulated as a nanoparticulate, in which it is covalently bound to poly-lactic-co-glycolic acid (PLGA), have anti-proliferative effects in tumor cells. These actions reflect the ability of tetrac and nanoparticulate tetrac to 1) antagonize the pro-proliferative, anti-apoptotic actions of  $T_4$  and  $T_3$ ; 2) to disable, in the absence or presence of  $T_4$  and  $T_3$ , the expression of a number of survival pathway genes (Glinsky *et al.*, 2004); and 3) to suppress the death-from-cancer gene signature of a number of cancer cell lines (Glinsky *et al.*, 2005). This 11-gene signature is a predictor of aggressiveness of cancer cells as demonstrated by shortened "time-to-tumor recurrence", the presence of distant metastasis, and death after tumor therapy. The PLGA formulation of tetrac acts exclusively at the integrin receptor for thyroid hormone; in contrast, unmodified tetrac acts at the integrin, but also gains access to the interior of the cell and may mimic actions of  $T_4$  (Moreno *et al.*, 2008).

Actions of nanoparticulate tetrac on gene expression in human breast cancer cells are coherent (Glinskii *et al.*, 2009). While the expression of a number of cyclin genes is suppressed and anti-apoptotic gene expression is decreased, pro-apoptotic gene expression is enhanced. Fairly remarkably, the expression of *thrombospondin 1* (*TBSP1*) is increased. The *TBSP1* protein is anti-angiogenic and the protein rarely accumulates in cancer cells. Nanoparticulate tetrac can also suppress expression of the *epidermal growth factor receptor* (*EGFR*) whose gene product supports cancer growth and angiogenesis and whose receptor is an oncologic target (Glinskii *et al.*, 2009). In the same study, however, while unmodified tetrac blocked expression of cyclin genes and certain genes relevant to apoptosis, this unmodified form of tetrac did not affect *EGFR* expression. Thus, the fit of the nanoparticulate formulation into its receptor on integrin  $\alpha\beta 3$  appears to be distinct from that of tetrac, itself. Nanoparticulate tetrac is also 10- to 100-fold more potent than unmodified tetrac, depending on the particular cell line studied (Glinskii *et al.*, 2009; Yalcin *et al.*, 2010a; Yalcin *et al.*, 2010b).

These observations on the two formulations of tetrac are consistent with the complexity of the receptor for thyroid hormone on the integrin (Cody *et al.*, 2007) and with the ability of the integrin to generate a spectrum of intracellular actions via several signal-transducing kinase pathways, including MAPK (extracellular-regulated kinases 1/2, or ERK1/2) and phosphatidylinositol 3-kinase (PI3K). Studied in human glioma cells by mathematical modeling of the kinetics of binding, the receptor site appears to contain two thyroid hormone-binding domains, one that recognizes only  $T_3$  and a second that transduces both the  $T_4$  and  $T_3$  signals (Lin *et al.*, 2009c). Recent computer graphics analysis of the interactions of  $T_3$  and  $T_4$  with the integrin (V. Cody, unpublished observations) is consistent with the existence of two hormone-binding domains in the receptor. While tetrac and nanotetrac interact with both domains, they do so differently, as the resulting gene expression profiles obtained in breast cancer cells indicate (Glinskii *et al.*, 2009).

It is important to note that the integrin-mediated anti-angiogenic activity of tetrac and nanotetrac transcend the inhibition of the pro-angiogenic thyroid hormone agonists,  $T_4$  and  $T_3$ . That is, in the absence of agonist thyroid hormone, tetrac can block the actions of vascular endothelial growth factor (VEGF) (Mousa *et al.*, 2008), bFGF (FGF2) (Mousa *et al.*, 2008), platelet-derived growth factor (PDGF) (SA Mousa: unpublished observations) and, as mentioned above, EGF. Because tumor cells can secrete multiple vascular growth factors to support their needs, it is desirable to identify potential therapeutic agents that antagonize the actions of more than one such growth factor.

It is clear from the foregoing that the actions of tetrac formulations at the cell surface integrin and of  $T_4$  and  $T_3$  at  $\alpha v\beta 3$  generate complex downstream events. However, it is also apparent that the hormone receptor on the integrin can also engage in crosstalk with growth factors on the plasma membrane. Interference by tetrac formulations with the actions of VEGF, bFGF and other growth factors imply this, since such growth factors include Arg-Gly-Asp (RGD) sequences that are validated by the RGD recognition site on the integrin before each growth factor can activate its growth factor-specific receptor clustered with the integrin (Davis, P. *et al.*, 2011). The  $\alpha v\beta 3$ -vascular growth factor receptor interaction, e.g., with VEGF (Sarmishtha *et al.*, 2005) and bFGF (Sahni and Francis, 2004), have been described in other studies that did not include thyroid hormone. The integrin receptor for agonist thyroid hormone analogues and for tetrac, however, influences other local cell membrane events. For example, activity of the  $Na^+/H^+$  exchanger (NHE1) is regulated by thyroid hormone analogues and this action is blocked by tetrac (Incerpi *et al.*, 2003). The insertion of Na, K-ATPase protein into the plasma membrane and activity of the ATPase (sodium pump) are also affected nongenomically by a MAPK- and PI3K-dependent mechanism (Lei *et al.*, 2007).

Such observations caused us to examine in doxorubicin-resistant human breast cancer cells (MCF-7) whether the state of resistance, in part determined by the expression and activity of the P-glycoprotein (MDR) pump (Dönmez *et al.*, 2010), may be affected by tetrac. Treatment of such breast cancer cells with tetrac reversed chemoresistance of the cells and resulted in increased intracellular residence time of radiolabeled doxorubicin (Rebbaa *et al.*, 2008). Such *in vitro* studies suggest that the combination of tetrac and doxorubicin should be tested in xenografts of doxorubicin-resistant MCF-7 cells. The mechanism of this particular action of tetrac has not been established. While it is possible that the function of the P-glycoprotein is affected by crosstalk with the tetrac-occupied hormone receptor on the integrin, we have also suggested that the action of tetrac on the NHE may be involved (Incerpi *et al.*, 2003). That is, inhibition of the NHE and a resultant intracellular decrease in pH may affect P-glycoprotein function because of the pH optimum of the MDR pump.

We would also point out that tetrac radiosensitizes tumor cells (Hercbergs *et al.*, 2009) by interfering with repair of DNA double-strand breaks (Hercbergs *et al.*, 2011). However, this has been studied to-date only in murine and human glioma cells and whether the agent affects DNA repair in breast cancer cells is not yet known.

A final consideration with regard to thyroid hormone is the recently described effect of the hormone on the abundance of integrin  $\alpha v\beta 3$  on the tumor cell plasma membrane. Agonist thyroid hormone analogues have a modest effect on the expression of the  $\alpha v\beta 3$  gene (Yonkers *et al.*, 2009) and may also increase the internalization of the  $\alpha v$ , but not the  $\beta 3$  component (Lin *et al.*, 2009). The disparate distribution of the  $\alpha v$  and  $\beta 3$  monomers within the cell is also conditioned by thyroid hormone, including import of  $\alpha v$  into the cell nucleus (Lin *et al.*, 2007).

### 3. Actions of resveratrol and the integrin receptor on the biology of breast cancer cells

Resveratrol is an extensively-studied, naturally-occurring alcohol with desirable properties in several biologic models. These activities include extension of lifespan in *C. elegans* (Zarse *et al.*, 2010) and remarkable anti-cancer properties (Pezzuto *et al.*, 2011; Hsieh *et al.*, 2011; Lin, C *et al.*, 2011). Substantial attention has been devoted to the metabolism of this agent because of its

rapid disappearance from the circulation post-administration to intact animals and its cellular uptake and chemical modification (Delmas *et al.*, 2011). The half-life of the parent compound is sufficiently short to promote speculation about the nature of the active biologic material.

We recently described a cell surface receptor for resveratrol on breast cancer cells which, like the thyroid hormone receptor on tumor cells (Bergh *et al.*, 2005), is on integrin  $\alpha v \beta 3$  (Lin *et al.*, 2006). The existence of such a receptor and its ability to transduce the plasma membrane resveratrol signal into MAPK activity and downstream into pro-apoptotic action suggested that the parent compound has substantial bioactivity. Integrins have been widely-viewed to bear receptors or binding sites only for large molecules—extracellular matrix proteins and growth factors (Plow *et al.*, 2000)—and thus it was surprising to find apparent biologically relevant binding sites for two small molecules on this integrin. It was also remarkable that the receptors for thyroid hormone and for resveratrol did not appear to interact functionally with one another. That is, both agents activated intracellular pools of MAPK (ERK1/2), but resveratrol was pro-apoptotic (Lin H *et al.*, 2008a; Lin C *et al.*, 2011) whereas thyroid hormones ( $T_4$ ,  $T_3$ ) were anti-apoptotic (Lin H *et al.*, 2008b), as noted above. Such observations suggested that the results of any efficacy testing of resveratrol as a chemotherapeutic agent in the presence of physiologic concentrations of thyroid hormone *in vitro* or in the intact animal with a normal pituitary-thyroid axis may be difficult to interpret (see below). An Arg-Gly-Asp (RGD) peptide prevents both the pro-apoptotic activity of resveratrol and the anti-apoptotic activity of thyroid hormone, indicating that the receptor is near the RGD recognition site on the integrin that was mentioned above (see section on *Thyroid Hormone Action*). The interactions of tetrac with either  $T_4$  or resveratrol, however, indicate that the two integrin binding sites are distinct, in that while tetrac inhibits the proliferative action of  $T_4$  in cancer cells, tetrac does not inhibit the pro-apoptotic actions of resveratrol (Lin C *et al.*, 2011).

Our own studies of the mechanism by which resveratrol may act to induce apoptosis in tumor cells have revealed several unexpected findings. First, resveratrol is able to induce p53-related apoptosis in cancer cells expressing certain mutations in p53. What is required in p53 for expression of resveratrol's apoptotic activity is that the Ser-15 be intact, as it is a target of phosphorylation by resveratrol-activated MAPK (Lin H *et al.*, 2002). Second, one of the actions of resveratrol in tumor cells is to induce a nuclear pool of cyclooxygenase-2 (COX-2) (Lin C *et al.*, 2011; Lin H *et al.*, 2009a). Chronic accumulation of COX-2 in cytoplasm is a marker of tumor cell aggressiveness (Schmitz *et al.*, 2006; Perdiki *et al.*, 2007) and long-term pharmacologic inhibition of the enzymatic activity of COX-2—the product of which is prostaglandins—appears to improve clinical outcomes or prevent emergence of certain cancers, such as that of the colon (Galalmb *et al.*, 2010).

Inducible COX-2 in the nucleus, however, is a wholly different biologic product. It is pro-apoptotic, can interact with Ser-15-phosphorylated p53 and can even bind to DNA (Lin H *et al.*, 2008b). The latter observation raises the possibility that COX-2 can be a co-activator and, indeed, resveratrol-induced nuclear COX-2 co-localizes with p300, a coactivator for p53 (Song *et al.*, 2010), as well as for proteins in the superfamily of nuclear hormone receptors (Kalkhoven, 2004). Activation of MAPK is required for formation of the complex of p53, p300 and COX-2, since that complex is not obtained in resveratrol-treated cells in the presence of the MEK-MAPK inhibitor, PD 98059 (Tang *et al.*, 2006). In contrast, inhibition of the enzymatic activity of COX-2 with indomethacin does not affect the pro-apoptotic activity of COX-2 (Tang *et al.*, 2006).

Third, a thyroid hormone analogue such as  $T_4$  prevents or disrupts the formation of the nuclear p53-COX-2 complex in cells treated with resveratrol (Lin *et al.*, 2008a, b). It is thought that this hormonal effect explains the blunting of the pro-apoptotic action of resveratrol in the presence of thyroid hormone. Competition between thyroid hormone and resveratrol for binding to the  $\alpha\beta3$  integrin is thought to be responsible for this inhibition. Not unexpectedly, in a tumor cell system that includes resveratrol and a physiologic concentration of  $T_4$ , the addition of tetrac protects the pro-apoptotic action of resveratrol from the anti-apoptotic effect of thyroid hormone (Lin *et al.*, 2008b). All of these actions compete at the  $\alpha\beta3$  integrin receptor. In the absence of  $T_4$ , the induction of apoptosis by resveratrol is not affected by the addition of tetrac (Lin *et al.*, 2008b).

The antecedent review suggests that further testing of resveratrol as a cancer chemotherapeutic agent might be pursued in two ways. First, the combination of nanotetrac and resveratrol may be evaluated against breast tumor cells *in vitro* or *in vivo* in the nude mouse xenograft model. Second, the potential manufacture of a nanoparticulate formulation of resveratrol in which the polyalcohol is covalently bound to the nanoparticle may be desirable, so as to permit biologic activity but restrict that effect of resveratrol to the integrin receptor. This formulation would prevent cellular uptake and subsequent metabolism/degradation of the nanoparticulate compound.

There is another implication with regard to the dependence of the pro-apoptotic action of resveratrol on an inducible pool of nuclear COX-2. New pharmacologic inhibitors of the enzymatic function of cyclooxygenase-2 should be examined for their ability to block the action of resveratrol, and perhaps of other polyalcohols as model inducers of nuclear COX-2. For example, NS398 is an experimental inhibitor of COX-2 that has been shown to prevent induction of COX-2 by resveratrol (Tang *et al.*, 2006).

#### **4. Dihydrotestosterone (DHT) acts via integrin $\alpha\beta3$ to induce proliferation of human breast cancer cells**

Although androgens may have an inhibitory effect on the proliferation of breast cancer cells, the actions of these steroids on such cells are variable. When we examined the action of dihydrotestosterone (DHT) on ER $\alpha$ -positive MCF-7 and ER $\alpha$ -negative MDA-MB-231 human breast cancer cells, we found that the androgen promoted proliferation in both cell lines (Lin *et al.*, 2009b). Integrin  $\alpha\beta3$  antibody inhibited the action of DHT in MDA-MB-231 cells, but was ineffective in MCF-7 cells (Lin *et al.*, 2009b). On the other hand, ICI 182,780 treatment and siRNA knockdown of ER $\alpha$  blocked the proliferative effect of DHT in MCF-7 cells, but not in the MDA-MB-231 cells. Thus, the mechanisms of DHT action differ in the two cell lines, and only in the ER-negative cells was there evidence for the existence of a DHT receptor on integrin  $\alpha\beta3$ . In neither breast cancer cell line could participation of a classical androgen receptor be implicated in the action of DHT. Tetrac did not affect the action of DHT (HY Lin: unpublished observations), indicating that these two small molecule receptors on the integrin function independently of one another.

It is not yet clear what the clinical significance may be of the DHT receptor on the breast cancer cell surface. We speculate, however, that in the patient with a recurrent ER $\alpha$ -positive tumor and taking tamoxifen or an aromatase inhibitor, residual circulating androgen may be promoting cancer cell proliferation. Useful information relevant to this possibility will come from determination of the androgen analogue-specificity of the receptor for DHT, and of a

possible contribution of DHT to breast cancer cell growth. In the case of the ER-negative human breast cancer cell, we have already demonstrated its susceptibility to DHT-stimulation (Lin *et al.*, 2009b).

It will also be important to analyze solid tumors beyond breast cancer for the presence of DHT-induced growth stimulation, including cancer cell growth as well as angiogenesis. In a recent report, Sieveking *et al.* (2010) have reported that androgen treatment of male endothelial cells *in vitro* enhanced angiogenesis; in contrast, gene knockdown of the androgen receptor (AR) in these cells caused unresponsiveness of these cells to androgen treatment. Female endothelial cells lacking AR did not respond to androgen treatment, but overexpression of the androgen receptor in female endothelial cells caused angiogenesis to occur.

## 5. Discussion and conclusions

Modulation of the proliferation of breast cancer cells is largely viewed as a function of the presence or absence of estrogen in ER-positive tumor cells and the presence of polypeptide growth factors that are autocrine or systemic. The emphasis on ER-mediated actions has grown from a broad understanding, emerging from a number of laboratories, of the molecular functions of ER in the cell nucleus and genomic actions of estrogens. Management of breast cancer—beyond surgery, tumor irradiation and chemotherapeutic agents—specifically emphasizes suppression of the action of endogenous estrogen with tamoxifen or inhibition of estrogen synthesis with aromatase inhibition. That estrogen may support breast tumor growth nongenomically is now under consideration (Silva & Shupnik, 2007), perhaps involving nuclear ER $\alpha$  insinuated into the plasma membrane or ER-like proteins in the membrane (Levin, 2011).

The concept that thyroid hormone may be a growth factor for breast cancer has been advanced by various investigators (Goodman *et al.*, 1980; Borek *et al.*, 1983; Cristofanilli *et al.*, 2005). That thyroid hormone could nongenomically support ER-positive human breast cancer cell proliferation and to be estrogen-like was shown when T<sub>4</sub> caused MAPK-dependent specific serine phosphorylation of nuclear ER, mimicking estradiol upon which depended stimulation of proliferation by the iodothyronine. This work led to the subsequent identification of a thyroid hormone receptor on plasma membrane integrin  $\alpha v \beta 3$  at which a variety of effects of thyroid hormone are initiated that are nongenomic in mechanism (Tang *et al.*, 2004). The integrin (and, therefore, the receptor) is expressed primarily on cancer cells of various types and on rapidly-dividing blood vessel cells (Dijkgraaf *et al.*, 2009; Wei *et al.*, 2009; Dimastromatteo *et al.*, 2010).

An interesting inhibitor of this receptor target exists. This is tetraiodothyroacetic acid (tetrac), a deaminated analogue of T<sub>4</sub> that blocks binding of T<sub>4</sub> and T<sub>3</sub> to the integrin, but also has novel anti-cancer activities at the  $\alpha v \beta 3$  hormone-binding site in the absence of T<sub>4</sub> and T<sub>3</sub>. These actions, initiated at the cell surface, are on expression of specific genes and the actions are inimical to tumor cell survival and on angiogenesis (Glinskii, A. *et al.*, 2009), including that initiated by several vascular growth factors. The hormone receptor on the integrin also disables or garbles crosstalk between the integrin and growth factor receptors that are clustered with  $\alpha v \beta 3$  (Davis, F. *et al.*, 2004; Mousa S. *et al.*, 2008).

Tetrac has been re-formulated as a nanoparticle in which the outer ring hydroxyl group is stably covalently bonded through a linker to the nanoparticle. This formulation does not

enter the cell and its actions are limited to the tetrac/thyroid hormone receptor on integrin  $\alpha v \beta 3$  (Bergh J. *et al.*, 2005). The agent has been shown to suppress growth of a variety of human cancer cell xenografts and to be anti-angiogenic at the tumor site (Davis F. *et al.*, 2004; Davis F. *et al.*, 2006; Glinskii A. *et al.*, 2009; Davis P. *et al.*, 2011). This hormone target on the integrin is fairly remarkable, in that as a single target, it has a relatively large number of downstream actions on cancer cells that emphasize vulnerability of such cells.

Recognition and characterization of the thyroid hormone receptor on integrin  $\alpha v \beta 3$  were premonitory to the identification of other small molecule binding sites – now understood to be receptors in that, when occupied, they cause predictable downstream cellular events. These receptors appear to be quite independent of one another and of the thyroid hormone binding domain. Identification of the resveratrol receptor on the integrin of breast cancer cells and other solid tumor cells may provide useful insights into the actions of this stilbene. The rapid cellular uptake and metabolism of resveratrol (Kroon *et al.*, 2010) – leading to pharmacokinetic and pharmacodynamic speculation about the relative bioactivity of resveratrol analogues – is less puzzling if, in fact, the parent compound (resveratrol) can act on the outside of the cell. That is, resveratrol binds to the integrin and rapidly initiates a MAPK-requiring, p53-dependent apoptotic process, regardless of uptake of the compound and chemical processing. What was somewhat surprising was the observation that resveratrol-induced apoptosis was blocked by thyroid hormone, but by a mechanism that begins at a different domain on the integrin and utilizes MAPK, but is effected by interruption of a resveratrol-induced phosphorylation sequence in the cancer cell nucleus that involves p53 and other nucleoproteins. The outcomes of a recent pharmaceutical industry-sponsored clinical trial of resveratrol as an anti-cancer agent were disappointing (McBride, 2010). Where there may be multiple explanations for these results, we can speculate that in the setting of normal thyroid function that the anti-cancer activity of resveratrol may be wholly suppressed by the hormone.

A third small molecule receptor on the integrin of breast cancer cells was recently described and this is a site that responds to the androgen, dihydrotestosterone (DHT) (Lin, H. *et al.*, 2009b). It was not wholly unexpected that an authentic steroid was found to act at the integrin, since resveratrol, while not a steroid, has certain structural features and functional activities that are estrogen-like. Acting via integrin  $\alpha v \beta 3$ , DHT is a trophic agent for breast cancer cells. Of some interest is that the mechanisms by which DHT acts *in vitro* differ in ER $\alpha$ -negative and ER $\alpha$ -positive cells. The cell surface integrin receptor for DHT is required for the proliferative effect of the androgen in ER $\alpha$ -negative cells, but is irrelevant in ER-positive cells, where the estrogen receptor protein, itself, is needed for DHT action. In neither type of cell is the authentic nuclear androgen receptor a part of the mechanism by which DHT recognizes the existence of a receptor on integrin  $\alpha v \beta 3$  in ER $\alpha$ -negative breast cancer cells. As we have pointed out above, recognition of the existence of the androgen receptor on  $\alpha v \beta 3$  in ER $\alpha$ -negative breast cancer may help to explain recurrences of tumor in postmenopausal women in whom some effect of circulating androgen, now apparent in the absence of estrogen, may be seen.

## 6. Acknowledgment

An endowment at Ordway Research Institute supported by M. Frank Rudy and Margaret Domiter Rudy supported a significant portion of the work described here.


## 7. References

- Bergh, J.J., Lin, H.Y., Lansing, L., Mohamed, S.N., Davis, F.B., Mousa, S., & Davis, P.J. Integrin  $\alpha v \beta 3$  contains a cell surface receptor site for thyroid hormone that is linked to activation of mitogen-activated protein kinase and induction of angiogenesis. *Endocrinology*. 2005;146:2864-71.
- Borek, C., Guernsey, D.L., Ong, A., & Edelman, I.S. Critical role played by thyroid hormone in induction of neoplastic transformation by chemical carcinogens in tissue culture. *Proc. Natl. Acad. Sci. USA*. 1983; 80:5749-5752.
- Chen, Y.F., Redetzke, R.A., Said, S., Beyer, A.J., & Gerdes, A.M. Changes in left ventricular function and remodeling after myocardial infarction in hypothyroid rats. *Am J Physiol Heart Circ Physiol*. 2010 Jan; 298(1):H259-62.
- Cheng, S.Y., Leonard, J.L., & Davis, P.J. Molecular aspects of thyroid hormone actions. *Endocr. Rev.* 2010; 31:139-70.
- Cody, V., Davis, P.J., & Davis, F.B. Molecular modeling of the thyroid hormone interactions with  $\alpha v \beta 3$  integrin. *Steroids*. 2007; Feb;72(2):165-70.
- Cristofanilli, M., Yamamura, Y., Kau, S.W., Bevers, T., Strom, S., Patangan, M., Krishnamurthy, S., Theriault, R.L., & Hortobagyi, G.N. Thyroid hormone and breast carcinoma. Primary hypothyroidism is associated with a reduced incidence of breast carcinoma. *Cancer*. 2005; 103:1122-1128.
- Davis, F.B., Mousa, S.A., O'Connor, L., Mohamed, S., Lin, H.Y., Cao, H.J., & Davis, P.J. Proangiogenic action of thyroid hormone is fibroblast growth factor-dependent and is initiated at the cell surface. *Circ Res*. 2004 Jun; 94(11):1500-6.
- Davis, F.B., Tang, H.Y., Shih, A., Keating, T., Lansing, L., Hercbergs, A., Fenstermaker, R.A., Mousa, A., Mousa, S.A., Davis, P.J., & Lin, H.Y. Acting via a cell surface receptor, thyroid hormone is a growth factor for glioma cells. *Cancer Res*. 2006 Jul 15;66(14):7270-5.
- Davis, P.J., Davis, F.B., Mousa, S.A., Luidens, M.K., & Lin, H.Y. Membrane receptor for thyroid hormone: physiologic and pharmacologic implications. *Annu. Rev. Pharmacol. Toxicol.* 2011;51:99-115.
- Davis, P.J., Davis, F.B., & Mousa, S.A. Thyroid hormone-induced angiogenesis. *Curr Cardiol Rev*. 2009 Jan; 5(1):12-16.
- De, S, Razorenova, O, McCabe, NP, O'Toole, T, Qin, J & Byzova, TV. VEGF-integrin interplay controls tumor growth and vascularization. *Proc. Natl. Acad. Sci. USA* 2005; 102:7589-7594.
- Delmas, D., Aires, V., Limagne, E., Dutartre, P., Mazué, F., Ghiringhelli, F., & Latruffe, N. Transport, stability, and biological activity of resveratrol. *Ann. N Y Acad. Sci.* 2011 Jan; 1215(1):48-59.
- Dijkgraaf, I., Beer, A.J., Wester, H.J., Application of RGD-containing peptides as imaging probes for  $\alpha v \beta 3$  expression. *Front. Biosci.* 2009; 14:887-899.
- Dimastromatteo, J., Riou, L.M., Ahmadi, M., Pons, G., Pellegrini, E., Broisat, A., Sancey, L., Gavrilina, T., Boturyn, D., Dumy, P., Fagret, D., & Ghezzi, C. In vivo molecular imaging of myocardial angiogenesis using the  $\alpha(v)\beta 3$  integrin-targeted tracer  $^{99m}\text{Tc}$ -RAFT-RGD. *J. Nucl. Cardiol.* 2010; 17:435-443.
- Dönmez, Y., & Gündüz, U. Reversal of multidrug resistance by small interfering RNA (siRNA) in doxorubicin-resistant MCF-7 breast cancer cells. *Biomed Pharmacother.* 2011, in press.

- Galamb, O., Spisák, S., Sipos, F., Tóth, K., Solymosi, N., Wichmann, B., Krenács, T., Valcz, G., Tulassay, Z., & Molnár, B. Reversal of gene expression changes in the colorectal normal-adenoma pathway by NS398 selective COX2 inhibitor. *Br. J. Cancer*. 2010 Feb 16;102(4): 765-73.
- Glinskii, A.B., Glinsky, G.V., Lin, H.Y., Tang, H.Y., Sun, M., Davis, F.B., Luidens, M.K., Mousa, S.A., Hercbergs, A.H., & Davis, P.J. Modification of survival pathway gene expression in human breast cancer cells by tetraiodothyroacetic acid (tetrac). *Cell Cycle*. 2009 Nov; 8(21): 3554-62.
- Glinsky, G.V., Glinskii, A.B., Stephenson, A.J., Hoffman, R.M., & Gerald, W.L. Gene expression profiling predicts clinical outcome of prostate cancer. *J Clin Invest*. 2004 Mar;113(6): 913-23.
- Glinsky, G.V., Berezovska, O., & Glinskii, A.B. Microarray analysis identifies a death-from-cancer signature predicting therapy failure in patients with multiple types of cancer. *J Clin Invest*. 2005 Jun;115(6):1503-21.
- Goodman, A.D., Hoekstra, S.J., & Marsh, P.S. Effects of hypothyroidism on the induction and growth of mammary cancer induced by 7,12-dimethylbenz(a)anthracene in the rat. *Cancer Res*. 1980; 40:2336-2342.
- Hercbergs, A., Davis, P.J., Davis, F.B., Ciesielski, M.J., & Leith, J.T. Radiosensitization of GL261 glioma cells by tetraiodothyroacetic acid (tetrac). *Cell Cycle*. 2009 Aug 15;8(16):2586-91.
- Hercbergs, A.H., Lin, H.Y., Davis, F.B., Davis, P.J., & Leith, J.T. Radiosensitization and production of DNA double-strand breaks in U87MG brain tumor cells induced by tetraiodothyroacetic acid (tetrac). *Cell Cycle*. 2011 Jan 15;10(2):352-7.
- Hsieh, T.C., Wong, C., Bennett, D.J., & Wu, J.M. Regulation of p53 and cell proliferation by resveratrol and its derivatives in breast cancer cells: an *in silico* and biochemical approach targeting integrin  $\alpha\beta3$ . *Int. J. Cancer*. In press, 2011.
- Incerpi, S., D'Arezzo, S., Marino, M., Musanti, R., Pallottini, V., Pascolini, A., & Trentalance, A. Short-term activation by low 17 $\beta$ -estradiol concentrations of the Na<sup>+</sup>/H<sup>+</sup> exchanger in rat aortic smooth muscle cells: physiopathological implications. *Endocrinology*. 2003 Oct;144(10):4315-24.
- Kalkhoven E. CBP and p300: HATs for different occasions. *Biochem. Pharmacol*. 2004 Sep 15;68(6):1145-55.
- Kato, S., Endoh, H., Masuhiro, Y., Kitamoto, T., Uchiyama, S., Sasaki, H., Masushige, S., Gotoh, Y., Nishida, E., Kawashima, H., Metzger, D., & Chambon, P. Activation of the estrogen receptor through phosphorylation by mitogen-activated protein kinase. *Science* 1995; 270:1491-4.
- Kroon, P.A., Iyeer, A., Chunduri, P., Chan, V., & Brown, L. The cardiovascular nutraceutical of resveratrol: pharmacokinetics, molecular mechanisms and therapeutic potential. *Curr. Med. Chem*. 2010; 17:2442-2455.
- Lei, J., Wendt, C.H., Fan, D., Mariash, C.N., & Ingbar, D.H. Developmental acquisition of T3-sensitive Na-K-ATPase stimulation by rat alveolar epithelial cells. *Am. J. Physiol. Lung Cell Mol. Physiol*. 2007 Jan; 292(1):L6-14.
- Levin, E.R. Cellular Functions of the Plasma Membrane Estrogen Receptor. *Trends Endocrinol. Metab*. 1999 Nov;10(9): 374-377.
- Lin, C., Crawford, D.R., Lin, S., Hwang, J., Sebuyira, A., Meng, R., Westfall, J.E., Tang, H.Y., Lin, S., Yu, P.Y., Davis, P.J., & Lin, H.Y. Inducible COX-2-dependent apoptosis in human ovarian cancer cells. *Carcinogenesis*. 2011 Jan;32(1): 19-26.

- Lin, H.Y., Davis, P.J., Tang, H.Y., Mousa, S.A., Luidens, M.K., Hercbergs, A.H., & Davis, F.B. The pro-apoptotic action of stilbene-induced COX-2 in cancer cells: convergence with the anti-apoptotic effect of thyroid hormone. *Cell Cycle*. 2009a Jun 15;8(12): 1877-82.
- Lin, H.Y., Lansing, L., Merillon, J.M., Davis, F.B., Tang, H.Y., Shih, A., Vitrac, X., Krisa, S., Keating, T., Cao, H.J., Bergh, J., Quackenbush, S., & Davis, P.J. Integrin  $\alpha v \beta 3$  contains a receptor site for resveratrol. *FASEB J*. 2006; 20:1742-4.
- Lin, H.Y., Shih, A., Davis, F.B., Tang, H.Y., Martino, L.J., Bennett, J.A., & Davis, P.J. Resveratrol-induced serine phosphorylation of p53 causes apoptosis in a mutant p53 prostate cancer cell line. *J. Urol*. 2002 Aug; 168(2): 748-55.
- Lin, H.Y., Sun, M., Lin, C., Tang, H.Y., London, D., Shih, A., Davis, F.B., & Davis, P.J. Androgen-induced human breast cancer cell proliferation is mediated by discrete mechanisms in estrogen receptor- $\alpha$ -positive and -negative breast cancer cells. *J. Steroid Biochem. Mol. Biol*. 2009b Feb; 113(3-5): 182-8.
- Lin, H.Y., Sun, M., Tang, H.Y., Lin, C., Luidens, M.K., Mousa, S.A., Incerpi, S., Drusano, G.L., Davis, F.B., & Davis, P.J. L-Thyroxine vs. 3,5,3'-triiodo-L-thyronine and cell proliferation: activation of mitogen-activated protein kinase and phosphatidylinositol 3-kinase. *Am J Physiol Cell Physiol*. 2009c May;296(5): C980-91.
- Lin, H.Y., Sun, M., Tang, H.Y., Simone, T.M., Wu, Y.H., Grandis, J.R., Cao, H.J., Davis, P.J., & Davis, F.B. Resveratrol causes COX-2- and p53-dependent apoptosis in head and neck squamous cell cancer cells. *J. Cell. Biochem*. 2008a Aug 15;104(6): 2131-42.
- Lin, H.Y., Tang, H.Y., Davis, F.B., & Davis, P.J. Resveratrol and apoptosis. *Ann N Y Acad Sci*. 2011 Jan;1215(1): 79-88.
- Lin, H.Y., Tang, H.Y., Lin, C., Davis, F.B., & Davis, P.J. Thyroid hormone induces nuclear accumulation of monomeric integrin  $\alpha v$  and formation of integrin-nucleoprotein complexes. *Thyroid* 2007 Oct;17(Suppl. 1): abstr. 261.
- Lin, H.Y., Tang, H.Y., Keating, T., Wu, Y.H., Shih, A., Hammond, D., Sun, M., Hercbergs, A., Davis, F.B., & Davis, P.J. Resveratrol is pro-apoptotic and thyroid hormone is anti-apoptotic in glioma cells: both actions are integrin- and ERK-mediated. *Carcinogenesis* 2008b Jan; 29(1): 62-9.
- Luidens, M.K., Mousa, S.A., Davis, F.B., Lin, H.Y., & Davis, P.J. Thyroid hormone and angiogenesis. *Vascul. Pharmacol*. 2010 Mar-Apr; 52(3-4):142-5.
- McBride, R. Glaxo ends resveratrol drug study. *Xconomy/Boston* 2010; 12/01
- Moreno, M., de Lange, P., Lombardi, A., Silvestri, E., Lanni, A., & Goglia, F. Metabolic effects of thyroid hormone derivatives. *Thyroid*. 2008 Feb;18(2):239-53.
- Mousa, S.A., Bergh, J.J., Dier, E., Rebbaa, A., O'Connor, L.J., Yalcin, M., Aljada, A., Dyskin, E., Davis, F.B., Lin, H.Y., & Davis, P.J. Tetraiodothyroacetic acid, a small molecule integrin ligand, blocks angiogenesis induced by vascular endothelial growth factor and basic fibroblast growth factor. *Angiogenesis*. 2008;11(2):183-90.
- Mousa, S.A., Davis, F.B., & Davis, P.J. Hormone-Integrin Cross Talk and Angiogenesis Modulation. *Immunology, Endocrine & Metabolic Agents in Medicinal Chemistry* 2009; 9:75-83.
- Perdiki, M., Korkolopoulou, P., Thymara, I., Agrogiannis, G., Piperi, C., Boviatsis, E., Kotsiakis, X., Angelidakis, D., Diamantopoulou, K., Thomas-Tsagli, E., & Patsouris, E. Cyclooxygenase-2 expression in astrocytomas. Relationship with microvascular parameters, angiogenic factors expression and survival. *Mol. Cell. Biochem*. 2007 Jan; 295 (1-2):75-83.

- Pervaiz, S., & Holme, A.L. Resveratrol: its biological targets and functional activity. *Antioxidants & Redox Signaling*. 2009 Oct 15; 11(11):2851-97.
- Pezzuto, J.M. The phenomenon of resveratrol: redefining the virtues of promiscuity. *Ann. N.Y. Acad. Sci.* 2011 Jan; 1215(1):123-30.
- Plow, E.F., Haas, T.A., Zhang, L., Loftus, J., & Smith, J.W. Ligand binding to integrins. *J. Biol. Chem.* 2000 Jul 21; 275(29):21785-8.
- Rebbaa, A., Chu, F., Davis, F.B., Davis, P.J., & Mousa, S.A. Novel function of the thyroid hormone analog tetraiodothyroacetic acid: a cancer chemosensitizing and anti-cancer agent. *Angiogenesis*. 2008 Nov; 11(3):269-76.
- Sahni, A., & Francis, CW. Stimulation of endothelial cell proliferation by FGF-2 in the presence of fibrinogen requires  $\alpha v\beta 3$ . *Blood*. 2004; 104:3635-3641.
- Schmitz, K.J., Callies, R., Wohlschlaeger, J., Kimmig, R., Otterbach, F., Bohr, J., Lee, H.S., Takeda, A., Schmid, K.W., & Baba, H.A. Overexpression of cyclo-oxygenase-2 is an independent predictor of unfavourable outcome in node-negative breast cancer, but is not associated with protein kinase B (Akt) and mitogen-activated protein kinase (ERK1/2, p38) activation or with Her-2/neu signaling pathways. *J. Clin. Pathol.* 2006 Jul; 59(7):685-91.
- Sieveking, D.P., Lim, P., Chow, R.W., Dunn, L.L., Bao, S., McGrath, K.C., Heather, A.K., Handelsman, D.J., Celermajer, D.S., & Ng, M.K. A sex-specific role for androgens in angiogenesis. *J. Exp. Med.* 2010 Feb 15; 207(2):345-52.
- Song, L., Gao, M., Dong, W., Hu, M., Li, J., Shi, X., Hao, Y., Li, Y., & Huang, C. p85 $\alpha$  mediates p53 K370 acetylation by p300 and regulates its promoter-specific transactivity in the cellular UVB response. *Oncogene* 2011, in press.
- Tang, H.Y., Lin, H.Y., Zhang, S., Davis, F.B., & Davis, P.J. Thyroid hormone causes mitogen-activated protein kinase-dependent phosphorylation of the nuclear estrogen receptor. *Endocrinology* 2004 July; 145(7):3265-72.
- Tang, H.Y., Shih, A., Cao, H.J., Davis, F.B., Davis, P.J., & Lin, H.Y. Resveratrol-induced cyclooxygenase-2 facilitates p53-dependent apoptosis in human breast cancer cells. *Mol. Cancer Ther.* 2006 Aug; 5(8):2034-42.
- Tomanek, R.J., Doty, M.K., & Sandra, A. Early coronary angiogenesis in response to thyroxine: growth characteristics and upregulation of basic fibroblast growth factor. *Circ. Res.* 1998 Mar; 82(5):587-93.
- Wei, L, Ye, Y., Wadas, T.J., Lewis, J.S., Welch, M.J., Achilefu, S., & Anderson, C.J. (64)Cu-labeled CB-TE2A and diamsar-conjugated RGD peptide analogs for targeting angiogenesis: comparison of their biological activity. *Nucl. Med. Biol.* 2009; 36:277-285.
- Yalcin, M., Bharali, D.J., Lansing, L., Dyskin, E., Mousa, S.S., Davis, F.B., Davis, P.J., & Mousa, S.A. Tetraiodothyroacetic acid (tetrac) and tetrac nanoparticles inhibit growth of human renal cell carcinoma. *Anticancer Res.* 2009 Oct; 29(10):3825-31.
- Yalcin, M., Dyskin, E., Lansing, L., Bharali, D.J., Mousa, S.S., Bridoux, A., Herberg, A.H., Lin, H.Y., Davis, F.B., Glinsky, G.V., Glinskii, A., Ma, J., Davis, P.J., & Mousa, S.A. Tetraiodothyroacetic acid (tetrac) and nanoparticulate tetrac arrest growth of medullary carcinoma of the thyroid. *J. Clin. Endocrinol. Metab.* 2010b Apr; 95(4):1972-80.
- Yonkers, M.A., & Ribera, A.B. Molecular components underlying nongenomic thyroid hormone signaling in embryonic zebrafish neurons. *Neural Dev.* 2009 Jun; 4:20.
- Zarse, K., Schmeisser, S., Birringer, M., Falk, E., Schmoll, D., & Ristow, M. Differential effects of resveratrol and SRT1720 on lifespan of adult *Caenorhabditis elegans*. *Horm. Metab. Res.* 2010 Nov; 42(12):837-9.



## **Targeting New Pathways and Cell Death in Breast Cancer**

Edited by Dr. Rebecca Aft

ISBN 978-953-51-0145-1

Hard cover, 190 pages

**Publisher** InTech

**Published online** 29, February, 2012

**Published in print edition** February, 2012

This book presents novel and interesting findings by multiple accomplished investigators in breast cancer. These chapters elucidate new mechanisms of breast cancer cell death as well as discuss new pathways for therapeutic targeting.

### **How to reference**

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Hung-Yun Lin, Faith B. Davis, Mary K. Luidens, Aleck Hercbergs Shaker A. Mousa, Dhruva J. Bharali and Paul J. Davis (2012). Newly-Recognized Small Molecule Receptors on Human Breast Cancer Cell Integrin  $\alpha v \beta 3$  that Affect Tumor Cell Behavior, Targeting New Pathways and Cell Death in Breast Cancer, Dr. Rebecca Aft (Ed.), ISBN: 978-953-51-0145-1, InTech, Available from: <http://www.intechopen.com/books/targeting-new-pathways-and-cell-death-in-breast-cancer/newly-recognized-small-molecule-receptors-on-human-breast-cancer-cell-integrin-alpha-v-beta-3-that-affect-tumor-cell-behavior>

**INTECH**  
open science | open minds

### **InTech Europe**

University Campus STeP Ri  
Slavka Krautzeka 83/A  
51000 Rijeka, Croatia  
Phone: +385 (51) 770 447  
Fax: +385 (51) 686 166  
[www.intechopen.com](http://www.intechopen.com)

### **InTech China**

Unit 405, Office Block, Hotel Equatorial Shanghai  
No.65, Yan An Road (West), Shanghai, 200040, China  
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元  
Phone: +86-21-62489820  
Fax: +86-21-62489821

© 2012 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen