

PUBLISHED BY

INTECH

open science | open minds

World's largest Science,
Technology & Medicine
Open Access book publisher



3,550+
OPEN ACCESS BOOKS



112,000+
INTERNATIONAL
AUTHORS AND EDITORS



115+ MILLION
DOWNLOADS



BOOKS
DELIVERED TO
151 COUNTRIES

AUTHORS AMONG
TOP 1%
MOST CITED SCIENTIST



12.2%
AUTHORS AND EDITORS
FROM TOP 500 UNIVERSITIES



Selection of our books indexed in the
Book Citation Index in Web of Science™
Core Collection (BKCI)

WEB OF SCIENCE™

Chapter from the book *Current Topics in Ionizing Radiation Research*
Downloaded from: <http://www.intechopen.com/books/current-topics-in-ionizing-radiation-research>

Interested in publishing with IntechOpen?
Contact us at book.department@intechopen.com

Radiation-Sensitivity and Transcription Profiles in Various Mutant p53 Cells

Kumio Okaichi, Nanaka Izumi, Yuma Takamura,
Shoichi Fukui and Takashi Kudo

Department of Radioisotope Medicine,

Atomic Bomb Disease and Hibakusha Medicine Unit,

Atomic Bomb Disease Institute,

*Nagasaki University Graduate School of Biomedical Sciences, Nagasaki,
Japan*

1. Introduction

The tumor suppressor gene *p53* plays an important role in determining radiosensitivity. The normal *p53* gene product accumulates after exposure to ionizing radiation, and causes growth arrest or promotes cell death through the apoptosis pathway (Figure 1). Mutation of the *p53* gene is the most common genetic alteration observed in human cancers (Nigro et al. 1989). It has been widely reported that cells with mutant *p53* are more resistant to ionizing radiation or DNA-damaging agents (Fan et al. 1994; Wattel et al. 1994; Lee et al. 1993; Hamada et al. 1996). On the other hand, there have been reports of cells harboring mutant *p53* that are sensitive to ionizing radiation and anticancer drugs (Biard et al. 1994; Fan et al.




Fig. 1. Pathway of p53. From irradiation by ionizing radiation to growth arrest or apoptosis of the cell through p53.

1995), although specific details of the mutations were not discussed. Mutant forms of *p53* differ in their properties according to the points of the mutation. For example, Crook et al. using a large series of *p53* mutants, found that not all transcriptionally active mutants retained the ability to suppress transformation, and that some tumor-derived point mutations conferred both transforming and transactivating activity (Crook et al. 1994). Some mutant forms of the *p53* gene do not merely induce the functional equivalent of *p53* loss (Harvey et al. 1995). The radiosensitivity of cells may depend on the type of *p53* mutation they harbor. It is important to determine which mutations affect the radiosensitivity of tumor cells, because tumor cell radiosensitivity has substantial clinical relevance in the context of tumor radiotherapy.

2. Mutation in *p53*

We prepared 15 types of cells harboring mutant forms of *p53* (T123A, L130V, Q143A, V157F, H168R, R175H, I195T, C242F, G244C, G245S, R273H, C277F, R280T, R282W and E286K) to examine their radiosensitivity. First, we created various mutations in the *p53* gene using a QuikChange site-directed mutagenesis kit (Stratagene, La Jolla, CA) in accordance with the manufacturer's protocol and integrated them into the LacSwitch inducible mammalian expression system (Stratagene, La Jolla, CA) (Okaichi et al. 1999). These mutant *p53* genes were then transformed into the Saos-2 cell line, which is null for *p53*, and stable transformants were obtained.

3. Radiation sensitivity of cells harboring *p53* mutation

Radiation sensitivity may depend on the position at which mutation occurs in *p53*. We previously examined various *p53* mutants for radiation sensitivity (Okaichi et al. 2008). Cells were subjected to γ -ray irradiation and then plated onto dishes. Colonies were examined after about one month to calculate the surviving fraction. The cells with wild-type *p53* showed higher radiation sensitivity than Saos-2 (*p53*-null) cells. Some mutations also resulted in increased radiation sensitivity, but mutations including hot spot mutations (175H, 245S 273H and 282W) showed almost no alteration of radiation sensitivity compared with Saos-2. Other mutations conferred an intermediate level of radiation sensitivity (Okaichi et al. 2008).

We then compared the radiosensitivity of these mutants with the frequency of mutation at each point, which is correlated with the tendency for tumorigenesis. Figure 2 shows the relationship between the frequency of *p53* mutation in human cells and radiosensitivity of the *p53* mutants. We divided these mutants into three groups; R (resistant), M (medium) and S (sensitive). The 175H, 244C, 245S 273H and 282W transformants were placed in group R, which was radioresistant and included a high frequency of mutation at all hot spots. The 130V, 143A, 168R, 277F and 286K transformants were placed in group M, which showed medium radiosensitivity and a low frequency of mutation. The 123A, 157F, 195T, 242F and 280T transformants were placed in group S, which was radiosensitive and showed a relatively low frequency of mutation.

As the radiosensitivity of these cells may be related to the induced expression of various genes by each type of *p53* mutation, we investigated the genes whose expression appeared to be related to the radiosensitivity of cells bearing *p53* mutations.




Fig. 2. Relationship between radiosensitivity of *p53* mutant cell lines and the frequency of *p53* mutation. The mutant cell lines were divided into three groups; R (radioresistant), M (medium radiosensitivity) and S (radiosensitive).

4. Transcriptional control in cells harboring *p53* mutation

We examined the expression of genes in cells harboring *p53* mutation that were subjected to γ -ray irradiation. For this we employed a DNA microarray (Gene Chip, Human Genome U133 Plus 2.0 Array; Affymetrix, Santa Clara, CA), containing over 54,000 probe sets, in accordance with the manufacturer's instructions. We extracted mRNA from each cell type 24 hours after irradiation at 6 Gy, and synthesized the cDNA. After we had synthesized, in turn, cRNA from the cDNA, we labeled the former with biotin and hybridized it with the DNA microarray. After staining and washing, we read the fluorescence using a scanner. The expression value (signal) of each gene was calculated and normalized using GeneSpring (Agilent Technologies, Santa Clara, CA) to adjust for minor differences between the experiments. In order to obtain the mean basal expression level of each gene, the signal values for unirradiated Saos-2 cells were used as the standard for the analysis. The change in value (signal log ratio) for each gene was calculated using Comparison Analysis in the software.

As radiosensitivity is intrinsically related to the apoptosis pathway, we summarized the genes associated with apoptosis, and these are shown in Table 1, where the cells harboring mutant *p53* cells are arranged from left to right according to their radiosensitivity, the cells harboring wild-type *p53* are located on the far left, and the parent cells, Saos-2, on the far right. We picked up genes showing an increase in gene expression of more than 2-fold, and indicated them by colored column. After irradiation, the cells with wild-type *p53* showed more than a 2-fold increase in the expression of 15 genes, whereas Saos-2 did not show any increase in the expression of apoptosis-related genes. Cells harboring mutant *p53* lacked

expression of many genes that were induced in cells harboring wild-type *p53*, but showed induction of some genes that were not induced in the latter. The 245S mutant cell line showed a particularly marked increase in the expression of many TNF-associated genes upon irradiation. As the expression of TNF-associated genes inhibits apoptosis, this would explain the radioresistance of 245S cells. We also noticed that the expression of TNF-associated apoptosis-inducing genes, such as TNFSF9 (0.15), TNFSF10 (0.18) and TNFSF21 (0.22), was decreased by more than half, in the 245S mutant cell lines. We were unable to explain the radiosensitivity of other mutant cell lines upon induction of apoptosis-related genes.

	Sensitive						Medium					Resistant					Saos
	Wild	195	123	242	157	280	277	168	286	130	143	244	273	245	282	175	Saos
CRADD	24.85	18.45	9.56	63.18	13.73	19.03	12.01	6.61	21.15	11.02	6.72	18.42	7.45	1.58	37.82	13.27	2.23
NFKBIA	4.35	2.05	2.86	1.73	2.54	4.72	4.02	3.33	6.15	2.52	1.87	2.54	3.44	0.82	0.88	4.21	2.43
PIK3R1	4.07	2.59	2.37	3.95	2.84	2.65	1.90	4.69	1.27	1.77	4.31	2.92	4.69	1.06	1.04	4.30	2.96
CASP1	27.65	87.42	9.50	0.52	0.37	8.97	1.51	1.29	4.27	3.97	4.75	5.19	2.04	0.49	9.59	4.81	1.47
CFLAR	2.89	2.00	1.51	3.52	3.07	1.47	1.82	2.39	1.51	1.69	2.89	2.58	2.51	3.71	1.81	1.48	2.29
IGF1R	3.51	4.16	1.52	5.86	4.24	2.77	1.81	3.83	0.92	1.70	3.94	3.39	3.19	1.04	0.89	1.79	2.26
MDM2	15.55	2.78	1.27	2.99	3.35	0.59	1.58	4.89	0.40	0.77	3.56	0.98	0.47	1.88	6.68	4.16	4.62
BCL2L1	6.52	2.62	0.90	3.44	0.70	0.69	1.76	1.03	3.58	2.11	1.44	5.89	0.62	2.73	6.90	0.84	2.04
CASP2	27.03	0.74	0.92	4.57	3.13	0.28	0.65	5.00	0.51	0.60	3.48	4.50	1.64	0.69	4.99	0.59	0.38
IKBKB	3.39	2.51	0.31	1.99	1.13	0.51	0.95	0.88	3.14	0.96	1.07	3.22	1.06	0.20	3.66	0.55	1.14
MCL1	3.24	2.09	0.56	2.28	0.95	0.55	1.10	1.13	1.22	1.72	1.86	2.50	1.96	0.51	2.41	1.20	1.37
FAS	2.16	0.44	3.79	0.36	1.02	0.19	0.51	0.90	0.33	0.70	2.07	0.44	1.08	1.49	0.45	2.29	1.10
TNFRSF10B	4.66	1.09	0.59	2.02	0.46	0.86	0.74	0.78	1.74	1.39	0.97	1.27	0.52	2.43	1.82	1.08	0.59
BIRC4	2.96	2.43	0.34	1.31	0.91	0.61	1.11	0.56	0.58	0.61	0.75	1.02	0.79	0.73	0.76	0.69	0.69
BAX	3.47	0.78	0.47	1.28	0.94	0.26	0.83	0.98	0.85	0.79	1.43	1.30	1.23	0.40	1.32	0.89	1.17
CASP4	0.21	1.35	0.56	3.79	1.33	1.97	0.87	7.81	0.97	13.87	9.53	0.72	2.73	1.24	2.63	0.44	0.98
BCL2L2	1.55	2.08	0.79	0.78	0.83	1.25	0.78	1.72	1.14	0.83	2.21	1.08	1.12	0.49	1.05	1.11	1.06
BCL2L11	0.74	0.39	0.95	1.15	1.28	0.43	0.34	1.12	2.01	0.63	1.30	1.19	1.75	0.95	2.42	1.22	1.26
MYC	0.34	0.53	0.39	0.21	0.25	0.34	0.44	0.71	0.12	1.95	0.77	0.22	0.45	2.04	0.10	0.41	1.19
NFKBIE	0.88	1.07	1.16	0.61	0.68	0.78	0.89	1.15	0.59	0.83	1.24	0.61	0.69	5.08	0.65	1.40	0.81
TNFRSF1A	1.63	0.84	0.85	0.67	0.64	0.66	0.94	0.55	0.76	0.61	0.61	0.85	0.84	2.39	0.54	0.61	0.85
TNFRSF25	0.32	0.06	1.13	1.03	1.03	0.16	0.91	1.27	0.20	2.49	1.03	1.53	0.98	5.92	2.29	0.37	0.25
TRAF2	0.75	0.88	0.96	1.36	0.62	0.81	0.57	1.03	0.98	0.58	0.99	0.85	1.28	2.31	0.98	0.99	1.08
TRAF3	1.14	1.60	1.06	1.03	0.72	0.79	0.92	0.55	0.94	0.77	0.98	0.87	0.67	2.03	1.31	1.28	1.01
BNIP3L	1.23	0.80	0.49	0.81	0.35	0.52	1.40	0.32	0.98	0.78	0.40	1.01	0.42	4.75	0.83	0.49	1.02
BID	1.09	0.79	1.18	0.66	0.80	0.62	1.43	1.77	1.19	1.83	2.01	0.69	0.77	0.36	0.85	1.32	0.92
TNFSF10	1.01	0.20	1.67	0.11	0.76	0.90	0.71	0.17	0.01	0.01	1.29	0.12	0.10	0.02	0.02	2.86	0.76

Table 1. Induction of gene expression in the apoptosis pathway by irradiation at 6 Gy. We listed the apoptosis genes whose expression was increased more than 2-fold in mutant cells. The numbers indicate the gene expression value in comparison with unirradiated Saos-2 cells. The colored columns indicate more than a 2-fold increase.

We speculated that certain genes might play an important role in making some cells radioresistant. In this connection, we listed those genes whose expression was increased more than 2-fold in radioresistant cells. Table 2 shows a list of genes whose expression was increased in more than 4 of the mutant strain of radioresistant cells. We paid attention to the level of gene expression in the radiosensitive mutant cells, because genes that play an important role in conferring radioresistance would be show lower levels of expression in radiosensitive cells. Expression of the genes CADPS2, DNPEP, NKTR, OVOS2, PSENEN, RASSF4, RBM14 and WTAP was not increased more than 2-fold in almost all of the

radiosensitive cells. CADPS2 acts as a calcium sensor in constitutive vesicle trafficking and secretion (Cisternas et al. 2003), and DNPEP is an aspartyl aminopeptidase that catalyzes the sequential removal of amino acids from the unblocked N termini of peptides and proteins (Nakamura et al. 2011). NKTR plays an important role in NK-cell cytotoxicity (Anderson et al. 1993). OVOS2 is a member of the ovostatin family and possesses trypsin-inhibitory activity (Saxena and Tayyab, 1997). PSENEN (Presenilin enhancer-2) is a component of the γ -secretase complex which catalyzes the final cleavage of amyloid precursor protein to generate the toxic amyloid β protein, the major component of plaques in the brain of Alzheimer disease patients, and protects embryos from apoptosis (Zetterberg et al. 2006). RASSF4 binds directly to activated K-Ras in a GTP-dependent manner via the effector domain, thus exhibits the basic nature of a Ras effector and plays an important role in Ras-dependent apoptosis (Eckfeld et al. 2004). RBM14 (CoAA) is a nuclear receptor coactivator protein at the interface of transcriptional coactivation and RNA splicing (Auboeuf et al. 2004). WTAP (Wilms' tumor 1-associating protein) is essential for embryonic development, and appears to exert an antiproliferative effect, inhibiting G₁-to-S phase cell cycle transition and also promoting apoptosis (Small et al. 2007).

	Sensitive						Medium					Resistant					Saos
	Wild	195	123	242	157	280	277	168	286	130	143	244	273	245	282	175	Saos
ARHGDI1	22.6	1.78	2.33	5.98	0.91	1.93	4.59	3.27	6.46	0.76	4.02	31.3	2.02	1.79	11.3	3.56	11.2
ATP6V0E1	1.39	1.84	1.72	2.43	1.49	2.22	1.77	1.84	1.86	1.58	1.64	3.53	2.03	3.24	1.23	2.02	2.54
C9orf86	3.04	1.49	2.00	2.57	1.79	2.97	2.34	1.84	4.49	0.61	1.73	3.79	2.07	3.47	3.56	2.16	2.10
CADPS2	0.05	0.73	2.6	0.21	0.06	0.18	0.21	1.03	0.61	1.63	1.62	1.20	4.49	4.01	2.74	3.40	1.40
CUEDC1	4.57	0.70	1.82	2.22	2.26	1.67	2.63	2.34	3.00	1.12	2.54	4.73	5.14	2.60	1.99	3.47	3.00
DNPEP	1.78	2.91	1.52	1.98	1.55	1.19	3.02	3.62	1.79	4.78	3.19	3.33	2.61	1.38	2.73	3.50	2.62
EIF4G1	3.70	2.07	1.20	2.50	1.44	1.32	1.99	2.45	3.30	1.08	2.72	3.38	2.01	3.77	3.26	2.20	3.87
FAM100B	2.79	2.95	1.37	2.51	1.77	1.79	2.17	1.72	3.16	2.48	1.56	3.06	2.43	3.60	5.01	1.72	2.37
FAM129B	4.89	3.13	1.92	1.71	1.63	2.02	3.48	3.6	5.18	1.21	2.49	4.32	2.02	4.07	1.98	2.86	3.83
FUS	0.75	1.84	1.34	1.35	2.86	2.04	2.10	2.05	0.53	1.70	2.62	3.39	2.82	3.38	0.93	2.73	1.37
HERC6	15.2	3.56	1.55	4.15	1.39	1.71	2.29	2.30	2.39	2.27	3.26	2.98	2.11	8.37	1.86	2.35	1.09
HOMER3	5.32	5.82	1.41	3.90	1.97	1.83	3.51	1.80	7.16	1.44	2.05	6.08	2.48	3.82	9.04	2.11	1.83
HSF1	9.23	2.53	1.80	3.01	1.91	1.60	2.72	2.77	4.01	0.99	3.40	6.41	2.09	2.70	7.47	2.07	4.29
LDLR	4.34	2.4	0.32	2.15	0.21	0.95	3.78	0.24	1.60	0.07	0.60	4.36	2.66	3.58	3.31	0.34	3.60
MYEF2	2.90	1.32	1.99	5.60	1.44	3.09	3.25	2.19	3.28	0.92	2.00	8.72	7.58	6.93	4.75	3.25	2.99
NKTR	0.06	1.04	1.54	2.05	1.08	1.85	2.07	1.85	1.04	2.00	1.58	2.12	3.96	2.70	1.12	2.59	1.34
OVOS2	0.56	0.80	0.65	1.96	1.24	0.33	0.19	1.87	3.03	4.3	2.32	2.01	2.83	7.44	2.93	0.27	1.04
PSENEN	4.48	1.44	2.74	1.99	1.41	1.86	1.68	2.61	2.94	1.24	2.65	2.29	2.76	3.50	1.94	3.08	2.86
RASSF4	0.53	0.47	1.13	11.9	0.48	0.58	0.85	1.49	33.1	0.25	1.85	13.9	0.27	2.80	42.7	2.93	0.57
RBM14	10.0	1.35	1.26	3.90	1.81	0.76	1.52	2.42	6.92	0.99	3.61	7.01	2.60	0.93	7.75	2.09	4.50
RHOU	0.34	0.36	4.82	4.11	1.00	0.92	0.42	2.66	1.13	2.15	5.27	3.48	3.39	5.72	1.92	7.63	1.32
SIRT6	2.24	1.71	3.02	1.73	1.33	2.38	1.37	1.82	2.48	1.29	1.87	2.14	2.14	2.37	1.53	3.07	1.96
SPAG9	2.13	1.73	1.03	2.41	3.38	1.72	2.04	3.19	1.12	1.46	3.33	2.14	2.64	7.77	1.14	2.09	2.69
TNRC6B	1.41	1.51	1.96	4.74	1.83	3.66	1.54	3.09	0.86	1.09	3.07	2.08	3.72	3.91	1.14	2.38	2.63
WTAP	0.79	0.80	3.32	1.97	1.89	1.68	2.46	2.02	0.99	3.93	3.15	2.99	2.43	2.69	2.27	4.07	1.71

Table 2. A list of the genes induced in radioresistant cells by irradiation at 6 Gy. We listed the genes that showed more than a 2-fold increase in almost all of the radioresistant cells. The numbers indicate the gene expression value in comparison with unirradiated Saos-2 cells. The colored columns indicate more than a 2-fold increase.

Among these genes, RASSF4 and WTAP are related to apoptosis, but exert a negative effect on radioresistant. As PSENEN blocks apoptosis, this gene may play an important role in radioresistance.

We approached this issue from the opposite perspective, and searched for genes that played an important role in conferring radiosensitivity. We listed genes showing more than a 2-fold increase in expression in radiosensitive cells. Table 3 shows a list of genes whose expression was increased in more than 4 of the radiosensitive mutant cell lines. Expression of CBR4, FOXP1, KPNA3, MFAP5, NEK3, TRIM2 and TRIM38 was not increased more than 2-fold in almost of all radioresistant cell lines. CBR4 (carbonyl reductase 4) is a mitochondrial NADPH-dependent quinone reductase that may be involved in the induction of apoptosis by cytotoxic 9, 10-phenanthrenequinone (Endo et al. 2008). FOXP1 is a forkhead transcription factor with functions in tissue and cell-type specific gene expression, and its gene is a direct target of p53-induced microRNA miR-34a (Rao et al. 2010). KPNA

	Wild	Sensitive					Medium					Resistant					Saos
		195	123	242	157	280	277	168	286	130	143	244	273	245	282	175	
ARNT	4.25	2.82	1.85	2.18	2.33	2.36	1.20	2.04	0.80	0.99	1.66	1.18	2.82	1.02	1.17	2.26	1.64
CBR4	0.87	2.09	2.69	1.56	2.43	2.64	1.99	2.72	1.04	2.80	2.33	1.07	1.87	1.94	1.16	2.12	1.07
CHD2	2.21	2.60	1.44	2.51	3.42	2.16	1.23	2.51	0.45	1.65	3.04	2.13	3.46	1.83	0.70	1.95	0.98
CRIM1	1.82	2.80	0.87	2.30	2.08	2.17	1.98	1.59	0.47	0.77	1.47	1.13	0.87	0.94	0.42	2.13	2.02
CSNK1A1	0.83	2.28	2.21	1.73	3.25	2.88	1.70	2.08	1.27	3.78	2.56	1.86	3.01	3.02	1.55	1.46	2.25
EPDR1	0.79	2.66	2.87	2.08	3.57	3.49	3.79	3.57	0.24	4.30	3.62	1.56	4.28	1.95	1.48	3.52	1.63
FOXP1	0.93	3.57	2.67	2.51	4.88	1.36	1.78	4.94	0.72	2.84	5.96	1.64	3.03	0.13	0.98	1.86	1.92
GNAS	0.90	2.94	1.40	2.45	3.81	2.22	1.95	2.60	1.89	5.39	2.46	2.76	2.47	1.16	1.94	1.89	2.10
IGF1R	3.51	4.16	1.52	5.86	4.24	2.77	1.81	3.83	0.92	1.70	3.94	3.39	3.19	1.04	0.89	1.79	2.26
KPNA3	1.24	2.92	0.79	2.07	2.69	2.07	1.84	1.45	0.66	0.96	1.80	1.29	1.59	2.03	0.92	1.20	2.06
KRAS	1.92	3.26	2.40	2.24	3.27	2.01	2.07	4.08	1.37	2.41	4.41	1.51	2.81	3.15	0.87	1.60	2.20
LYST	3.47	5.79	1.81	2.26	3.05	3.19	3.30	3.53	1.83	1.73	2.83	1.77	2.88	1.16	1.23	2.61	1.75
MARCKS	1.19	3.60	2.90	1.43	3.59	2.70	1.44	3.94	1.35	0.79	3.62	1.08	3.48	0.61	0.48	2.19	2.46
MFAP5	4.26	11.1	0.16	12.6	7.07	3.13	1.70	0.84	0.36	0.22	0.42	11.2	0.01	0.02	0.62	0.06	1.12
NBPF1	1.59	3.03	2.22	2.51	3.43	1.94	1.75	2.64	1.28	2.39	2.09	1.92	3.07	1.00	1.23	2.01	2.38
NBPF10	1.33	2.68	2.78	2.02	3.11	2.82	2.19	1.78	0.62	2.26	1.54	1.86	2.39	1.79	1.57	2.77	1.99
NEK3	1.37	2.93	1.23	2.10	2.45	2.84	1.72	2.64	0.89	0.90	1.99	2.36	1.83	0.52	1.06	0.89	1.39
PGF	1.11	1.94	2.53	2.02	3.22	3.12	1.37	2.46	0.88	2.05	2.65	1.51	1.99	2.67	1.46	3.45	1.68
PPP1R12A	2.40	2.32	2.11	2.32	2.83	3.16	2.24	3.53	1.23	1.24	3.84	1.65	2.68	1.77	1.13	2.63	1.92
QKI	2.33	3.30	2.96	1.69	5.23	2.53	1.83	3.39	1.94	9.80	3.07	1.85	3.27	0.91	1.94	2.54	2.50
RASAL2	4.52	5.9	1.16	2.73	2.79	2.44	2.89	2.19	1.61	0.89	2.36	3.06	3.16	0.89	1.32	1.72	2.14
SFRS11	2.01	3.99	3.54	1.96	7.18	2.86	1.47	4.95	0.53	1.96	4.51	1.98	5.37	1.31	0.95	6.45	2.95
SLC35E1	1.10	1.99	2.50	2.26	2.18	2.48	1.08	1.97	0.87	1.96	2.08	1.72	1.80	2.21	1.18	2.69	1.24
SPG21	1.35	2.13	3.53	2.21	3.25	3.73	1.93	2.72	1.16	2.61	1.97	1.60	1.88	2.43	0.84	3.25	1.62
SUMO2	1.72	2.41	2.55	1.87	2.28	2.77	1.54	2.10	1.63	1.30	2.53	2.30	1.96	1.88	1.35	2.83	1.64
TRIM165	0.99	2.51	3.18	3.07	3.07	2.5	1.76	4.25	0.7	5.08	3.87	1.39	4.51	1.51	1.68	3.62	2.75
TRIM13	2.78	3.09	1.61	3.27	3.82	3.22	2.02	3.22	1.12	0.99	2.12	3.26	1.88	3.99	1.20	1.42	0.92
TRIM2	1.67	3.04	0.47	2.53	2.05	2.13	4.37	0.82	0.73	1.14	0.57	1.23	1.36	0.06	0.49	0.42	2.32
TRIM38	1.87	2.15	1.71	2.45	3.57	2.54	1.51	2.75	1.03	3.26	2.86	1.95	1.59	2.92	1.75	1.78	1.93
TSPAN9	1.76	2.27	2.33	3.34	2.28	1.30	1.35	3.33	2.14	2.69	2.81	2.50	3.27	0.77	1.59	1.76	1.62
VGLL4	1.45	2.44	2.44	2.63	4.58	1.83	1.86	4.86	1.41	2.20	5.39	1.38	3.37	1.54	0.97	2.54	2.55
VPS13B	2.65	3.85	1.25	2.17	2.95	3.62	2.34	2.30	1.45	1.13	2.21	1.99	3.33	9.85	1.48	1.79	2.64
WNK1	1.69	2.33	2.48	2.03	3.01	1.00	1.29	2.98	1.22	1.99	3.21	2.22	1.72	4.99	1.52	1.84	1.34
WSB2	2.73	4.50	1.59	3.16	2.09	2.48	1.38	3.77	1.40	0.89	3.74	2.20	3.78	1.47	0.70	1.52	1.55
ZNF818	2.88	3.65	1.08	2.95	3.32	2.20	1.34	2.85	1.92	0.35	1.01	2.47	1.29	1.26	2.00	1.65	2.37

Table 3. A list of the genes induced in radiosensitive cells by irradiation at 6 Gy. We listed the genes that showed more than a 2-fold increase of expression in almost all of the radiosensitive cells. The numbers indicate the gene expression value in comparison with unirradiated Saos-2 cells. The colored columns indicate more than a 2-fold increase.

(karyopherin- α) proteins are responsible for the transport of proteins into and out of the nucleus through the nuclear pore complex, and KPUNA3 contributes genetically to schizophrenia (Wei and Hemmings 2005). MFAP5 (microfibrillar associated protein 5), also known as a microfibril-associated protein (MAGP2), is a highly significant indicator of survival and chemosensitivity of the cells (Spivey and Banyard, 2010). NEK3 is a serine/threonine kinase that contributes to PRL-mediated breast cell cancer motility through mechanisms involving Rac1 activation and paxillin phosphorylation (Miller et al. 2007). TRIM (tripartite motif-containing) proteins are a family comprising more than 70 members in humans and contain conserved RING, G-box, coiled-coil, and SPRY domains, most of which are involved in protein ubiquitination, but only a few of them have been well studied. TRIM2 mediates the p42/p44 MAPK-dependent ubiquitination of Bim (Bcl-2-interacting mediator of cell death) in rapid ischemic tolerance, and suppression of TRIM2 expression stabilizes the level of Bim protein and blocks neuroprotection (Thompson et al. 2011). TRIM38 has E3 ubiquitin ligase activity and can be degraded during virus infection (Liu et al. 2011).

As CBR4 is involved in the induction of apoptosis, this gene may play an important role in radiosensitivity. However, the precise role of these genes in radiosensitivity remains unknown.

We have attempted to perform hierarchical clustering analysis of RNA expression in these mutant *p53* cell lines using Gene Tree software, but were unable to find any clear relationship between radiosensitivity and gene expression.

5. Conclusions

Ionizing radiation is used extensively in medical diagnostic and treatment protocols. With a better understanding of radiation induced molecular processes, it might become possible to identify the radiosensitivity of individuals before the start of radiation therapy, leading to individualization of radiation treatment. Radiation-induced transcriptional responses have been studied using DNA microarray (Kis et al. 2006; Jen and Cheung, 2006). Some previous studies have also examined cells harboring mutant *p53* using DNA microarray (Amandson et al. 2003; Scian et al. 2004), but they did not examine each type of mutation.

In the present study, we prepared 15 mutant *p53* cell lines, cells harboring wild-type *p53* and Saos-2 cells (null for *p53*). We examined the radiosensitivity of the mutant cell lines and classified them as R (resistant), M (medium) or S (sensitive). We then studied the radiation-induced transcriptional responses in these cell lines, and examined the relationship between their radiation-induced gene expression and radiosensitivity. We found some genes that appeared to have some correlation with radiosensitivity, for example PSENE1 and CBR4. However, none of the genes directly determined the radiosensitivity of the cells. Further study will be needed to determine which of these genes is the main determinant of radiosensitivity.

Radiosensitivity may be determined by several genes working in collaboration. Mutation of *p53* leads not only to loss of function, but also gain of function. If such functions are related to growth arrest or DNA repair, then loss of function would confer radiosensitivity, and gain of function to radioresistance. On the other hand, if such functions are related to apoptosis, then loss of function would confer radioresistance and gain of function to radiosensitivity. Each mutation of *p53* may thus lead to loss of function and gain of some other function at the same time. This makes it very difficult to determine whether a certain

mutation of *p53* leads to the radiosensitivity on the basis of transcriptional analysis alone. Recently it has been reported that many kinds of microRNAs related to tumorigenesis or apoptosis are regulated by *p53* (He et al. 2007; Suzuki et al. 2009). Thus *p53* regulates not only mRNA but also microRNA. The regulation of microRNA in each mutant *p53* cell line would vary the degree of cell radiosensitivity. The available data suggest the importance of determining the type of mutation of *p53* and examining the regulation of overall transcriptional control in individual tumor cells in the context of radiotherapy.

6. Acknowledgments

This work was partly supported by the Global Center of Excellence (GCOE) of the Ministry of Education, Culture, Sports, Science and Technology of Japan.


7. References

- [1] Amundson SA, Bittner M and Fornace Jr JF (2003) Functional genomics as a window on radiation stress signaling. *Oncogene* 22: 5828-5833.
- [2] Anderson SK, Gallinger S, Roder J, Frey J, Young HA and Ortaldo JR (1993) A cyclophilin-related protein involved in the function of natural killer cells. *Proc Natl Acad Sci USA* 90: 542-546.
- [3] Auboeuf D, Dowhan DH, Li X, Larkin K, Ko L, Berget SM and O'Malley BW (2004) CoAA, a nuclear receptor coactivator protein at the interface of transcriptional coactivation and RNA splicing. *Mol Cell Biol* 24: 442-453.
- [4] Biard DS, Martin M, Rhun YL, Duthu A, Lefaix JL, May E and May P (1994) Concomitant *p53* gene mutation and increased radiosensitivity in rat lung embryo epithelial cells during neoplastic development. *Cancer Res* 54: 3361-3364.
- [5] Cisternas FA, Vincent JB, Scherer SW and Ray PN (2003) Cloning and characterization of human CADPS and CADPS2, new members of the Ca²⁺-dependent activator for secretion protein family. *Genomics* 81: 279-291.
- [6] Crook T, Marston N J, Sara EA and Vousden KH (1994) Transcriptional activation by *p53* correlates with suppression of growth but not transformation. *Cell* 79: 817-827.
- [7] Eckfeld K, Hesson L, Vos MD, Bieche I, Latif F and Clark GJ (2004) RASSF4/AD037 is a potential Ras effector/tumor suppressor of the RASSF family. *Cancer Res* 64: 8688-8693.
- [8] Endo S, Matsunaga T, Kitade Y, Ohno S, Tajima K, El-Kabbani O and Hara A (2008) Human carbonyl reductase 4 is a mitochondrial NADPH-dependent quinine reductase. *BBRC* 377: 1326-1330.
- [9] Fan S, el-Deiry WS, Bae I, Freeman J, Jondle D, Bhatia K, Fornace AJ, Jr Magrath I, Kohn KW and O'Connor PM (1994) *p53* gene mutations are associated with decreased sensitivity of human lymphoma cells to DNA damaging agents. *Cancer Res* 54: 5824-5830.
- [10] Fan S, Smith ML, Rivet DJ, Duba D, Zhan Q, Kohn KW, Fornace Jr AJ and O'Connor PM (1995) Disruption of *p53* function sensitizes breast MCF-7 cells to cisplatin and pentoxifylline. *Cancer Res* 55: 1649-164954.
- [11] Hamada M, Fujiwara T, Hizuta A, Gochi A, Naomoto Y, Takakura N, Takahashi K, Roth JA, Tanaka N and Orita K (1996) The *p53* gene is a potent determinant of

- chemosensitivity and radiosensitivity in gastric and colorectal cancers. *J Cancer Res Clin* 122: 360-365.
- [12] Harvey M, Vogel H, Morris D, Bradley A, Bernstein A and Donehower LA (1995) A mutant *p53* transgene accelerates tumor development in heterozygous but not nullizygous *p53*-deficient mice. *Nat Genet* 9: 305-311.
- [13] He L, He X, Lim LP, de Stanchina E, Xuan Z, Ling Y, Xue W, Zender L, Magnus J, Ridzon D, Jakson AL, Linsley PS, Chen C, Lowe SW, Cleary MA and Hannan GJ (2007) A microRNA component of the *p53* tumor suppressor network. *Nature* 447: 1130-1134.
- [14] Jen K-Y and Cheung VG (2006) Transcriptional response of lymphoblastoid cells to ionizing radiation. *Genome Res* 13: 2092-2100.
- [15] Kis E, Szatmari T, Keszei M, Farkas R, Esik O, Lumniczky K, Falus A and Safrany G (2006) Microarray analysis of radiation response genes in primary human fibroblasts. *Int J Radiation Oncology Biol Phys* 66: 1506-1514.
- [16] Lee JM and Bernstein A (1993) *p53* mutations increase resistance to ionizing radiation. *Proc Nat Acad Sci USA* 90: 5742-5746.
- [17] Liu X, Lei X, Zhou Z, Sun Z, Xue Q, Wang J and Hung T (2011) Enterovirus 71 induces degradation of TRIM38, a potential E3 ubiquitin ligase. *Virology J* 8: 61-68.
- [18] Miller SL, Antico G, Raghunath PN, Tomaszewski JE and Clevenger CV (2007) Nek3 kinase regulates prolactin-mediated cytoskeletal reorganization and motility of breast cancer cells. *Oncogene* 26: 4668-4678.
- [19] Nakamura Y, Inloes JB, Katagiri T and Kobayashi T (2011) Chondrocyte-specific microRNA-140 regulates endochondral bone development and targets Dnpep to modulate bone morphogenetic protein signaling. *Mol Cell Biol* 31: 3019-3028.
- [20] Nigro JM, Baker SJ, Preisiger AC, Jessup JM, Hostetter R, Cleary K, Bigner SH, Davidson N, Baylin S, Devilee P, Glover T, Collins FS, Weston A, Modali R, Harris CC and Vogelstein B (1989) Mutations in the *p53* gene occur in diverse human tumor types. *Nature* 342: 705-708.
- [21] Okaichi K, Wang L, Sasaki J, Saya H, Tada M and Okumura Y (1999) A point mutation of human *p53*, which was not detected as a mutation by a yeast functional assay, led to apoptosis but not p21Waf-1/Cip1/Sdi1 expression in response to ionizing radiation in a human osteosarcoma cell line, Saos-2. *Int J Radiat Oncol Biol Phys* 45: 975-980.
- [22] Okaichi K, Ide-Kanematsu M, Izumi N, Morita N, Okumura Y and Ihara M (2008) Variations in sensitivity to ionizing radiation in relation to *p53* mutant point. *Anticancer Res* 28: 2687-2690.
- [23] Rao DS, O'Connell RM, Chaudhuri AA, Garcia-Flores Y, Geiger TL and Baltimore D (2010) MicroRNA-34a perturbs B lymphocyte development by repressing the forkhead box transcription factor Foxp1. *Immunity* 33: 48-59.
- [24] Saxena I and Tayyab S (1997) Protein proteinase inhibitors from avian egg whites. *Cell Mol Life Sci* 53: 13-23.
- [25] Scian MJ, Stagliano KER, Ellis MA, Hassan S, Bowman M, Miles MF, Deb SP and Deb S (2004) Modulation of gene expression by tumor-derived *p53* mutants. *Cancer Res* 64: 7447-7454.

- [26] Small TW, Penalva LO and Pickering JG (2007) Vascular biology and the sex of flies: regulation of vascular smooth muscle cell proliferation by Wilms' tumor 1-associated protein. *Trends Cardiovasc Med* 17: 230-234.
- [27] Spivey KA and Banyard J (2010) A prognostic gene signature in advanced ovarian cancer reveals a microfibril-associated protein (MAGP2) as a promoter of tumor cell survival and angiogenesis. *Cell Adhesion & Migration* 4: 169-171.
- [28] Suzuki H, Yamagata K, Sugimoto K, Iwamoto T, Kato S and Miyazono K (2009) Modulation of microRNA processing by p53. *Nature* 460: 529-533.
- [29] Thompson S, Pearson AN, Ashley MD, Jessick V, Murphy BM, Gafken P, Henshall DC, Morris KT, Simon RP and Meller R (2011) Identification of a novel Bcl-2-interacting mediator of cell death (Bim) E3 ligase, tripartite motif-containing protein 2 (TRIM2), and its role in rapid ischemic tolerance-induced neuroprotection. *J Biol Chem* 286: 19331-19339.
- [30] Wattel E, Preudhomme C, Hecquet B, Vanrumbeke M, Quesnel B, Dervite I, Morel P and Fenaux P (1994) *p53* mutations are associated with resistance to chemotherapy and short survival in hematologic malignancies. *Blood* 84: 3148-3157.
- [31] Wei J and Hemmings GP (2005) The KPNA3 gene may be a susceptibility candidate for schizophrenia. *Neurosci Res* 52: 342-346.
- [32] Zetterberg H, Cambell WA, Yang HW and Xia W (2006) The cytosolic loop of the γ -secretase component Presenilin enhancer 2 protects Zebrafish embryos from apoptosis. *J Biol Chem* 281: 11933-11939.

INTECH



Current Topics in Ionizing Radiation Research

Edited by Dr. Mitsuru Nenoï

ISBN 978-953-51-0196-3

Hard cover, 840 pages

Publisher InTech

Published online 12, February, 2012

Published in print edition February, 2012

Since the discovery of X rays by Roentgen in 1895, the ionizing radiation has been extensively utilized in a variety of medical and industrial applications. However people have shortly recognized its harmful aspects through inadvertent uses. Subsequently people experienced nuclear power plant accidents in Chernobyl and Fukushima, which taught us that the risk of ionizing radiation is closely and seriously involved in the modern society. In this circumstance, it becomes increasingly important that more scientists, engineers and students get familiar with ionizing radiation research regardless of the research field they are working. Based on this idea, the book "Current Topics in Ionizing Radiation Research" was designed to overview the recent achievements in ionizing radiation research including biological effects, medical uses and principles of radiation measurement.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Kumio Okaichi, Nanaka Izumi, Yuma Takamura, Shoichi Fukui and Takashi Kudo (2012). Radiation-Sensitivity and Transcription Profiles in Various Mutant p53 Cells, Current Topics in Ionizing Radiation Research, Dr. Mitsuru Nenoï (Ed.), ISBN: 978-953-51-0196-3, InTech, Available from:

<http://www.intechopen.com/books/current-topics-in-ionizing-radiation-research/radiation-sensitivity-and-transcription-profile-in-various-mutant-p53-cells>

INTECH

open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821