

# Modified Atmosphere Packaging for Perishable Plant Products

Leonora M. Mattos<sup>1</sup>, Celso L. Moretti<sup>1</sup> and Marcos D. Ferreira<sup>2</sup>

<sup>1</sup>*Embrapa Vegetables,*

<sup>2</sup>*Embrapa Instrumentation  
Brazil*

## 1. Introduction

Packaging perishable plant products is one of the more important steps in the long and complicated journey from grower to consumer. Millions of different types of packages are used for produce around the world and the number continues to increase as the industry introduces new packaging materials and concepts. Packing and packaging materials contribute a significant cost to the produce industry; therefore it is important that packers, shippers, buyers, and consumers have a clear understanding of the wide range of packaging options available (Boyette et al., 1996). This fact chapter describes some of the many types of packaging materials, including their functions, uses, and limitations. Within packaging plastics for plant products, if commodity and film permeability characteristics are properly matched, an appropriate atmosphere can evolve passively through consumption of O<sub>2</sub> and production of CO<sub>2</sub> during respiration (Mir & Beaudry, 2002). Gas exchange and respiration rate through the package material are the processes involved in creating a modified atmosphere inside a package that will extend shelf life of fresh fruits and vegetables. The major methods for measuring respiration rates, along with their advantages and limitations are discussed. Modified atmosphere technologies have great potential in a wide range of applications in plant products. The usual methods of respiration rate determination can be the static system, the flowing system and the permeable system (Fonseca et al., 2002). The respiration rate of fresh produce can be expressed as O<sub>2</sub> consumption rate and/or CO<sub>2</sub> production rate. Factors affecting the respiration rate and respiratory quotient are outlined, stressing the importance of temperature, O<sub>2</sub> and CO<sub>2</sub> concentrations, and storage time (Kader et al., 1989). Modified atmosphere packaging should always be considered as a supplement to proper temperature and relative humidity management. The differences between beneficial and harmful concentrations of oxygen and carbon dioxide for each kind of produce are relatively small, so great care must be taken when using these technologies. Temperature has been identified as the most important external factor influencing respiration (Tano et al., 2007). The internal factors affecting respiration are the type and maturity stage of the commodity. Vegetables include a great diversity of plant organs such as fruits, roots, tubers, seeds, bulbs, sprouts, leaves and stems that have different metabolic activities and consequently different respiration rates. Different varieties of the same product exhibit specific respiration rates. The success of modified atmosphere packaging greatly depends on the accuracy of the predictive respiration rate (Kader, 2002). The main objective of this chapter is to present different packaging materials using modified atmosphere for perishable plant products, focusing

particularly on aspects of the respiration process, usual methods of measuring respiration rates and factors can be affect the respiration rate.

## 2. Modified atmosphere packaging

Modified atmosphere packaging (MAP) of fresh fruits and vegetables is based on modifying the levels of O<sub>2</sub> and CO<sub>2</sub> in the atmosphere produced inside a package sealed with some type of polymer film. It is desirable that the natural interaction that occurs between the respiration of the product and the packaging generates an atmosphere with low levels of O<sub>2</sub> and / or a high concentration of CO<sub>2</sub>. The growth of organisms that cause decay is thereby reduced and the life of the product is thus extended. Additionally, the desired atmosphere can reduce the respiration rate, and ethylene production, physiological changes. For example, it can inhibit chemical, enzymatic and microbiological mechanisms associated with the decay of fresh products, thus avoiding the use of other chemical or thermal process such as freezing, dehydration, and sterilization (Kader et al. 1989; Gorris & Tauscher, 1999; Saltveit, 1997; Fonseca et al., 2002).

The use of modified and controlled atmospheres has grown over the past 50 years, contributing significantly to extend the postharvest life and maintain the quality of various fruits and vegetables. So that changing the atmosphere occurs must have a combination of factors that influence the permeability of the product packaging and respiration in order to achieve an atmosphere of great balance for the conservation of the product. This balance is achieved when the respiration of the product consumes the same amount of O<sub>2</sub> entering the packaging and the production of CO<sub>2</sub> by respiration is equal to the amount that leaves the packaging (Day, 1993).

The first studies on modified atmospheres used reduced levels of O<sub>2</sub> in apple packaging in order to slow the ripening of fruits. The first challenge was to control the levels of O<sub>2</sub> in the package. Since then, an enormous variety of polymers with different properties have been developed to offer a wide range of options in features such as gas permeability, tensile strength and flexibility, among others. Presently, diverse systems of modified atmosphere packaging have been developed and used with a wide range of fruits and vegetables in order to provide optimal storage conditions and product longevity. Table 1 summarizes the optimal conditions of temperature and gas composition of O<sub>2</sub> and CO<sub>2</sub> for the transport and / or storage of some fruits and vegetables.

### 2.1 Passive

Modified atmospheres can be obtained passively between plant material and sealed package or intentionally using determined concentrations of gases. Modified atmosphere is formed as a result of vegetable respiration, which consumes CO<sub>2</sub> and releases O<sub>2</sub> in sealed package. In passive modification, the respiring product is placed in a polymeric package and sealed hermetically. Only the respiration of the product and the gas permeability of the film influence the change in gaseous composition of the environment surrounding the product. If the product's respiration characteristics are properly matched to the film permeability values, then a beneficial modified atmosphere can be passively created within a package. The polymer itself variably restricts gas exchange between the internal and external environments due to its selective permeability to O<sub>2</sub> and CO<sub>2</sub>. After a period of time, the

system reaches an equilibrium atmosphere containing of lower concentrations of O<sub>2</sub> and higher concentrations of CO<sub>2</sub> than in atmospheric air.

| Product               | Temperature<br>range (°C) | Atmosphere      |                  |
|-----------------------|---------------------------|-----------------|------------------|
|                       |                           | %O <sub>2</sub> | %CO <sub>2</sub> |
| Apples                | 0-5                       | 1-2             | 0-3              |
| Banana                | 12-16                     | 2-5             | 2-5              |
| Blackberry            | 0-5                       | 5-10            | 15-20            |
| Blueberry             | 0-5                       | 2-5             | 12-20            |
| Cherry, sweet         | 0-5                       | 3-10            | 10-15            |
| Cranberry             | 2-5                       | 1-2             | 0-5              |
| Grape                 | 0-5                       | 2-5             | 1-3              |
| Kiwifruit             | 0-5                       | 1-2             | 3-5              |
| Lemon                 | 10-15                     | 5-10            | 0-10             |
| Lychee (Lichti)       | 5-12                      | 3-5             | 3-5              |
| Mango                 | 10-15                     | 3-7             | 5-8              |
| Nuts and dried fruits | 0-10                      | 0-1             | 0-100            |
| Orange                | 5-10                      | 5-10            | 0-5              |
| Papaya                | 10-15                     | 2-5             | 5-8              |
| Persimmon             | 0-15                      | 3-5             | 5-8              |
| Pineapple             | 8-13                      | 2-5             | 5-10             |
| Plum                  | 0-5                       | 1-2             | 0-5              |
| Raspberry             | 0-5                       | 5-10            | 15-20            |
| Strawberry            | 0-5                       | 5-10            | 15-20            |
| Artichoke             | 0-5                       | 2-3             | 2-3              |
| Asparagus             | 1-5                       | Air             | 10-14            |
| Beans                 | 5-10                      | 2-3             | 4-7              |
| Broccoli              | 0-5                       | 1-2             | 5-10             |
| Brussels sprouts      | 0-5                       | 1-2             | 5-7              |
| Cabbage               | 0-5                       | 2-3             | 3-6              |
| Cantaloupes           | 2-7                       | 3-5             | 10-20            |
| Cauliflower           | 0-5                       | 2-3             | 3-4              |
| Celery                | 0-5                       | 1-4             | 3-5              |
| Cucumbers             | 8-12                      | 1-4             | 0                |
| Herbs*                | 0-5                       | 5-10            | 4-6              |
| Lettuce               | 0-5                       | 1-3             | 0                |
| Onions                | 0-5                       | 1-2             | 0-10             |
| Parsley               | 0-5                       | 8-10            | 8-10             |
| Pepper                | 5-12                      | 2-5             | 2-5              |
| Radish                | 0-5                       | 1-2             | 2-3              |
| Spinach               | 0-5                       | 7-10            | 5-10             |
| Tomatoes              | 12-20                     | 3-5             | 2-3              |

\*Herbs: chervil, chives, coriander, dill, sorrel and watercress, Adapted from Kader (2002)

Table 1. Summary of optimal conditions for modified atmosphere and temperature during transport and / or storage of fruits and vegetables

## 2.2 Active

The concept of active packaging has been developed to adjust the deficiencies in passive packaging such as when a film is a good barrier to moisture, but not to oxygen, the film can still be used along with an oxygen scavenger to exclude oxygen from the pack. An intentionally or actively obtained modified atmosphere occurs when the desired gas mixture is introduced into the container before sealing. In this way, atmospheric balance inside the package is reached faster or almost immediately. Sometimes, certain additives are incorporated into the polymeric packaging film or within packaging containers to modify the headspace atmosphere and to extend shelf-life. Another process is the acceleration of atmospheric balance under partial vacuum packaging is the process of removing the air before sealing, reducing the free space. Although the active modification of the atmosphere within the package incurs additional costs, the advantage is that the desired atmosphere is securely achieved in considerably less time.

## 2.3 O<sub>2</sub> and CO<sub>2</sub> limits

Safe levels of O<sub>2</sub> and CO<sub>2</sub> are important for package design. A lower O<sub>2</sub> limit has been associated with onset of fermentation and accumulation of ethanol and acetaldehyde (Beaudry et al., 1993). Fermentation is linked to the development of off-flavors and/or tissue damage. Effect of temperature on lower O<sub>2</sub> limit has been measured for a number of commodities including whole apple, apple slices, blueberry, and raspberry. In each case, lower O<sub>2</sub> limit increased with temperature. Lower O<sub>2</sub> limits vary from 0.15% to 5% and are influenced by temperature, commodity and cultivar (Beaudry and Gran, 1993).

It is necessary to know the main effects of gases on fresh fruits and vegetables and the interactions between gas and produce on the one hand and between gas and packaging material on the other hand to achieve the goal. It is important to recognize that while atmosphere modification can improve the storability of some fruits and vegetables, it also has the potential to induce undesirable effects. Fermentation and off-flavors may develop if decreased O<sub>2</sub> levels cannot sustain aerobic respiration (Kays, 1997). Similarly, injury will occur if CO<sub>2</sub> exceeds tolerable levels. Ranges of non-damaging O<sub>2</sub> and CO<sub>2</sub> levels have been published for a numbers of fruits and vegetables (Kader, 1997; Kupferman, 1997; Richardson and Kupferman, 1997; Saltveit, 1997; Beaudry 1999, 2000), minimally processed products (Gorny, 1997), and flowers and ornamentals (Reid, 1997). Horticultural crops differ in their tolerance for O<sub>2</sub> (Table 2) and CO<sub>2</sub> (Table 3).

## 3. Types of plastic films used in MAP

In a modified atmosphere packaging, changes start to take place immediately after packing the fresh produce as a result of the respiration of the packaged produce. The gases of the contained atmosphere and the external ambient atmosphere try to equilibrate by permeation through the package walls at a rate dependant upon the differential pressures between the gases of the headspace and those of the ambient atmosphere. It is in this context that the barrier to gases and water vapor provided by the packaging material must be considered. Thus, the success of the modified atmosphere packaging depends upon the barrier material used. MAP for fresh produce must also allow entry of oxygen to maintain the aerobic

| O <sub>2</sub> (%) | Commodity  |
|--------------------|--|
| 2                  | Lettuce (crisphead), pear  |
| 3                  | Artichoke, tomato  |
| 5                  | Apple (most cultivars), apricot, cauliflower, cucumber, grape, nashi, olive, orange, peach (clingstone), potato, pepper (bell) |
| 7                  | Banana, bean (green snap), kiwi fruit  |
| 8                  | Papaya   |
| 10                 | Asparagus, brussels sprouts, cabbage, celery, lemon, mango, nectarine, peach, persimmon, pineapple, sweet corn                 |
| 15                 | Avocado, broccoli, lychee, plum, pomegranate, sweetsop   |
| 20                 | Cantaloupe (muskmelon), durian, mushroom, rambutan   |
| 25                 | Blackberry, blueberry, fig, raspberry, strawberry  |

\*Data are from Beaudry (2000), Gorny (1997), Kader (1997), Kupferman (1997), Richardson and Kupferman (1997), and Saltveit (1997)

Table 2. O<sub>2</sub> limits below which injury can occur for selected horticultural crops held at typical storage temperatures

| CO <sub>2</sub> (kPa) | Commodity   |
|-----------------------|---|
| < 0.5                 | Chopped greenleaf, Romaine and iceberg lettuce, spinach, sliced pear, broccoli  |
| 1.0                   | Broccoli florets, chopped butterhead lettuce, sliced apple, brussels sprouts, cantaloupe, cucumber, crisphead lettuce, onion, apricot, avocado, banana, cherimoya, sweet cherry, cranberry, grape, kiwifruit, litchi, peach, plum |
| 2.0                   | Shredded and cut carrots, artichoke, cabbage, cauliflower, celery, bell and chili pepper, sweet corn, tomato, blackberry, fig, mango, olive, papaya, pineapple, pomegranate, raspberry, strawberry                                |
| 2.5                   | Shredded cabbage, blueberry   |
| 3.0                   | Cubed or sliced cantaloupe, low permeability apples and pears, persimmon  |
| 4.0                   | Sliced mushrooms  |
| 5.0                   | Green snap beans, lemon, lime, orange   |
| 10.0                  | Asparagus   |
| 14.0                  | Orange sections   |

\* Adapted from Herner (1987), Kader (1997), and Saltveit (1997)

Table 3. CO<sub>2</sub> partial pressures above which injury will occur for selected horticultural crops

metabolism of the product. In addition, some carbon dioxide must exit from the package to avoid build up of injurious levels of the gas. These packages are made of plastic films with relatively high gas permeability.

Packaging films that provide a wide range of physical properties, many of these individual films are combined through processes like lamination and co-extrusion. There are several groupings in MAP films such as in the plural, Vinyl Polymers, Styrene Polymers,

Polyamides, Polyesters and other polymers. Polypropylene is part of the Polyolefin group and used largely in MAP, in both forms: continuous and perforated. Sanz et al. (1999) studied the quality of strawberries packaged with polypropylene film, with proper perforations, during commercial postharvest practices. They concluded that perforated-mediated MA packaging helped to preserve fruit ripeness degree better, maintaining its nutritional value, measured as ascorbic acid content.

Many type of plastic films for packaging are available, but relatively few have been used to pack fresh fruits and vegetables, and even fewer have a gas permeability that makes them suitable for modified atmosphere packaging. The permeability of CO<sub>2</sub> should be 3 to 5 times the permeability of O<sub>2</sub>. Many polymers used to formulate packaging films are within this criterion (Table 4).

| Film type                 | Permeabilities<br>(cc/m <sup>2</sup> /mil/dia a 1 atm) |                |                                       |
|---------------------------|--|----------------|---------------------------------------|
|                           | CO <sub>2</sub>  | O <sub>2</sub> | CO <sub>2</sub> :O <sub>2</sub> ratio |
| Polyester                 | 180-390  | 52-130         | 3.0-3.5                               |
| Polyethylene, low density | 7,700-77,000   | 3,900-13,000   | 2.0-5.9                               |
| Polypropylene             | 7,700-21,000   | 1,300-6,400    | 3.3-5.9                               |
| Polystyrene               | 10,000-26,000  | 2,600-7,700    | 3.4-3.8                               |
| Polyvinyl chloride        | 4,263-8,138  | 620-2,248      | 3.6-6.9                               |

\*Adapted from Kader (2002)

Table 4. Permeability for packaging fresh fruits and vegetables

#### 4. Package parameters

Modified atmosphere (MA) packaging systems designed to produce optimum O<sub>2</sub> and CO<sub>2</sub> concentrations at suitable temperatures have been mathematically modeled (Chinnan, 1989; Lee et al., 1991; Exama et al., 1993; Talasila et al., 1995; Cameron et al., 1995).

The composition of the atmosphere inside the packaging results from the interaction of several factors including the permeability characteristics of the packaging material, the respiratory behavior of the plant material and the environment. Packaging films are selected so that the package has specific permeability characteristics and so that changes to these characteristics (temperature and humidity) over time, following the laws of physics. The concentration of gases within the packaging can be controlled to provide specific conditions. In contrast to these known and controllable factors are the often unknown and uncontrollable responses of the plant material. The plant species, the cultivar, cultural practices, the development stage, harvest management, the tissue type and post harvest handling all contribute and influence the response of the material to the atmosphere generated.

The scope of plant responses can be modified by the initial flow of gas before sealing the packaging as well as by the inclusion of chemical treatments to slow undesirable processes or decrease contamination. Each component of the packaging process needs to be examined separately to improve the understanding of what it contributes to potential packaging strategies.

Mathematical models can integrate the film permeability to O<sub>2</sub>, CO<sub>2</sub> and H<sub>2</sub>O, and the respiratory response of the commodity to O<sub>2</sub>, in some cases, CO<sub>2</sub>, along with its lower O<sub>2</sub> limit and upper CO<sub>2</sub> limit (Beaudry et al., 1992; Cameron et al., 1994; Lakakul et al., 1999; Fishman et al., 1996; Hertog et al., 1998). These models permit the identification of limiting features of the film, package design, and product and environment conditions.

The major factors to be taken into account while selecting the packaging materials are the type of package, the barrier properties needed (permeabilities of individual gases and gas ratios when more than one gas is used), physical properties of machinability and strength, integrity of closure (heat sealing), fogging of the film as a result of product respiration, printability and others.

#### 4.1 Respiration rate

Respiration is a process in which chemical reactions oxidize lipids and carbohydrates to carbon dioxide and water to produce energy, while the organelle responsible for aerobic respiration known as mitochondria. Part of the released energy is stored as chemical energy adenosine triphosphate (ATP) and part is lost as heat. This complex process can be influenced by several intrinsic factors such as product size, variety, maturity, type of tissue and extrinsic factors such as temperature, concentration of O<sub>2</sub> and CO<sub>2</sub> and mechanical damage (Day, 1993).

Knowledge of the minimum required for O<sub>2</sub> aerobic respiration is very important to avoid that the anaerobic pathway is the predominant route of respiration, which causes the accelerated loss of product quality. In order to maintain food safety, it is important to know the potential hazards of each product, the permeability of the films and the rate of respiration of fruits and vegetables (Watada et al., 1996).

Since both the rate of respiration and the permeability of the film are sensitive to temperature variations, and respond to these changes differently, it is expected that the package under modified atmosphere remains determined only within an atmosphere given temperature range (Zagory, 2000).

The maximal rate of respiration for most fruit and vegetable products undergoes a 4- to 6-fold increase from 0 to 15 °C (Beaudry et al., 1992; Cameron et al., 1994, 1995; Lakakul et al., 1999). This means that product respiration increases at two or three times the rate of LDPE permeability, and thirty times the rate of perforation permeability with increasing temperature. When respiratory demand for O<sub>2</sub> increases faster than O<sub>2</sub> permeation as temperature increases, O<sub>2</sub> levels decline and may pose a risk to product quality. This limits the usefulness of MAP in some situations.

Variation in the respiration rate of the product and the variation in film or permeability can influence package design. Variation in product respiration and package permeability has been measured for broccoli and the effect on package O<sub>2</sub> modeled (Cameron et al., 1993). Cameron et al. (1993) concluded that there is an estimable risk of the package O<sub>2</sub> falling sufficiently low to promote fermentation in any product. Packages should be designed to generate O<sub>2</sub> levels well above the lower limit to ensure aerobic conditions. Products such as broccoli, mushrooms, leeks and others have very high rates of respiration, and most continuous films do not have the capacity to provide enough O<sub>2</sub> to avoid fermentation.

Accordingly, there is commercial interest to develop films with high gas transmission rates. Films that have improved rates of gas transmission by virtue of their polymeric nature are often blends of two or three different polymers, where each polymer performs a specific function such as strength, transparency and improved gas transmission. Similarly, films can be laminated to achieve needed properties.

#### 4.2 Temperature

Temperature is one of the most important factors in extending the shelf-life of perishable products. Optimum storage temperature must be established for every product. Permeability of polymeric packaging films is also a function of temperature and it generally increases with the increase in temperature.

The effects of temperature on chemical reactions, including respiratory rate, traditionally quantified by  $Q_{10}$ , which is a coefficient by which it is possible to calculate how many times increases the rate of a reaction for each increase in temperature of 10 °C. The effect of temperature can also be quantified by the Arrhenius model, where the effect of temperature increase is given by the activation energy ( $E_a$ ) (Cameron et al., 1995). The temperature quotient is useful because it allows us to calculate the respiration rates at one temperature from a known rate at another temperature. However, the respiration rate does not follow ideal behavior, and the  $Q_{10}$  can vary considerably with temperature. At higher temperatures, the  $Q_{10}$  is usually smaller than at lower temperatures.

The storage temperature is very important for the evolution of microbial and visual quality of fresh fruits and vegetables. Knowledge of weather conditions and temperature in the cold chain for fresh produce is needed to determine the influence of the cold chain in real loss of quality and shelf life of these products (Cortez et al., 2002).

The temperature of the produce in the package is managed by circulating cool air around the outside of the package. The film and the headspace atmosphere are barriers to heat movement, prevent rapid cooling, and reduce the effectiveness of refrigeration. A 'safe radius' for the distance from the center of the package to the circulated air can be calculated based on the heat of respiration and the rate at which heat can be removed by the cooler air (Sharp et al., 1993). For instance, the center of a package of broccoli must have a radius of less than 14 cm to keep it within 1 °C of the refrigerated air.

Temperature control, when combined with correct use of packaging and modified atmosphere technology, is effective in controlling metabolic processes described above. However, the ideal temperature handling, storage and marketing of fresh fruits and vegetable is generally not respected. Some examples of the ranges of temperature and relative humidity are to be respected in Table 1.

If it is necessary to choose between mild temperatures that cause symptoms of chilling injury and temperatures accelerate senescence and microbial growth, the first must be chosen (Watada and Qi, 1999).

#### 4.3 Permeability

The permeability of the packaging material determines the atmospheric conditions in the headspace and ultimately the shelf-life of the product. If an atmosphere higher in carbon



dioxide and / or lower in oxygen is required, the material should be impermeable to the gases. Vegetables and fruits require a certain amount of oxygen in the headspace for maintenance of quality, therefore, packaging material for these products should be quite permeable to the oxygen, to allow atmospheric oxygen to replenish the gas in the package.

Transparency of packaging material to light is also important.

The choice of packaging material is an extremely important part of the MAP operation. The materials must be cost effective, have low water vapor transmission rate, high gas barrier, mechanical strength to withstand machine handling and subsequent storage and distribution of the finished pack as well as have the capability of giving high integrity seals to ensure retention of gas within the pack until opened by consumer. Also, once a gas atmosphere is applied, the level and proportion of headspace gas/gases is controlled only by judicious selection of packaging material with specified permeability characteristics. . Thickness is also a factor controlling permeability.

There are two ways to create barriers using film. The first strategy employs continuous films that control the movement of gases in and out of the box. The second strategy uses film with small openings or microperforations.

With continuous films, the movement of  $O_2$  and  $CO_2$  is directly proportional to the differences in gas concentration across the film. Constant levels of gases in the package are achieved when the product's consumption of  $O_2$  and production of  $CO_2$  are equal. This situation exists only when the respiration rate is constant (Fishman et al., 1996).

In perforated films, the rate of movement of gas through the perforated film is the sum of gas diffusion and atmospheric air infiltration through the polymer film. Generally, total gas flow through the perforations is much greater than gas movement through the film. The rate of gas exchange through microperforations is much greater than through continuous films. Perforated packaging is more suitable for vegetables that have a high demand for  $O_2$  (Gonzalez et al., 2008).

## 5. Respiration rate measurement

The respiration rate of fresh fruits and vegetables can be expressed as the rate of  $O_2$  consumption and  $CO_2$  production rate. The usual methods for determining the concentration of respiration are static or closed system, continuous flow system, and permeable system. The respiration rate is measured in permeable system, in other words, the product is within the dimensions and permeability of packaging film known (Beaudry, 1993; Joles et al., 1994; Lakakul, Beaudry, & Hernandez, 1999; Lee, Song, & Yam, 1996; Piergiovanni, Fava, & Ceriani, 1999; Smyth et al., 1998). The concentrations of  $O_2$  and  $CO_2$  are determined and stable mass balance is done on the system in order to estimate respiration rates.

The concentrations of gases in the system depend on the permeability characteristics of permeability, package size and weight of the product that were discussed previously. The time to reach equilibrium can be seen as a limitation of this method to measure the respiration rate. Definition of the steady-state concentration values is another difficulty of the permeable method.

For all methods have limitations, but this involves a greater number of variables, since the size of the packet are involved other parameters such as free volume, surface area and thickness of the gas exchange and the permeability characteristics. None of methods is clearly preferable over the others. When choosing the respiration rate determination method for a specific study, the benefits and limitations of each method should be taken into consideration. Fonseca et al. (1992) studied three different methods (closed, flow through and permeable systems) to measure respiration rate and observed their advantages and limitations. The main characteristics of these methods are summarized in Table 5.

| Characteristics                                      | System      |              |  |
|--|-------------|--------------|--|
|  | Closed      | Flow through | Permeable  |
| Non-destructive                                      | ✓           | ✓            | ✓  |
| Complexity of experimental set-up                    | ✓           | ✓            | ✓  |
| Ability to test different combinations of gases      | Simple      | Complex      | Complex  |
| Concentration is kept constant during the experiment | x           | ✓            | ✓  |
| Suitable for low respiring products                  | ✓           | x            |  |
| Suitable for high respiring products                 | x           | ✓            | ✓  |
| Accuracy is very sensitive to determination of       | Free volume | Flow-rate    | Permeability package dimensions, steady-state concentrations |

\*Adapted from Fonseca et al. (1992)

Table 5. Main characteristics of the three methods of respiration rate measurement

### 5.1 Factors affecting respiration rate

The factors affecting respiration are type and maturity stage of the commodity. Vegetables include a great diversity of plant organs such as roots, tubers, seeds, bulbs, fruits, sprouts, stems and leaves that have different metabolic activities and consequently different respiration rates. Even different varieties of the same product can exhibit different respiration rates (Fidler & North, 1967; Gran & Beaudry, 1992; Song et al., 1992). In general, non-climacteric commodities have higher respiration rates in the early stages of development that steadily decline during maturation (Lopez-Galvez, El-Bassuoni, Nie, & Cantwell, 2004). Respiration can also be affected by a wide range of environmental factors that include light, chemical stress, radiation stress, water stress, growth regulators, and pathogen attack. The most important postharvest factors are temperature, atmospheric composition, and physical stress.

One method to improve these problems would be to choose a film with permeability changes for O<sub>2</sub> similar to that of the respiration of the product, so if temperature increases, respiration and permeability of the film increase an equivalent amount.

## 5.2 Influence of temperature

Indeed the most important factor affecting postharvest life is temperature. This is because temperature has a profound effect on the rates of biological reactions, specifically metabolism and respiration. Over the physiological range of most crops, 0 to 30 °C increased temperatures cause an exponential rise in respiration.

Temperature of the product affects storability more than any other factor. Pre-cooling and temperature maintenance during handling and shipping are critical in preserving quality. Temperature also significantly affects film permeability and thereby the O<sub>2</sub> and CO<sub>2</sub> content of the package.

The elevated rate of respiration at high temperature could be used to rapidly establish the desired package atmosphere, but this would only be useful in the few situations in which it would be more important to rapidly establish the atmosphere than to slow physiological processes, eg., to reduce cut-surface browning.

Another solution to the MAP temperature problem is to develop a package system that senses either the environment or the physiological status of the enclosed product and responds by increasing the permeability to O<sub>2</sub> (Cameron et al., 1993). Such 'sense-and-respond' packaging is technically difficult to develop, and progress has only been conceptual at this time (Smyth et al., 1999). A third approach is to design packages to function at the highest temperatures typically encountered in the distribution and retail cool chain and, as far as possible, maintain control over the temperature of the packaged product, thereby adapting to the limitations imposed by the film. Most companies using MAP have adopted this simple solution. Generally, the lowest temperature feasible is maintained, since temperature has a much more significant influence on preserving quality than the application of low O<sub>2</sub> (Kays, 1997).

Reações metabólicas, entre elas a respiração, são reduzidas em 2 a 3 vezes para cada decréscimo de 10 °C na temperatura, o que permite retardar a maturação e a senescência do produto (Brecht, 1995).

## 5.3 Influence of gas composition

In MAP, the pack is flushed with a gas or a combination of gases. The common gases used are oxygen, nitrogen and carbon dioxide. Traces of carbon monoxide, nitrous oxide, ozone, argon, ethanol vapour and sulphur dioxide are also used. Minimum oxygen levels are used to pack plant under MA because oxygen can react with the fruits and vegetables resulting in the oxidative breakdown of them into their constitutive parts. Nitrogen is an inert gas. Carbon dioxide is responsible for the bacteriostatic and fungistatic effect in MA packaged fruits and vegetables. It retards the growth of moulds and aerobic bacteria. The inhibitory effect of carbon dioxide to micro-organisms is increased as the temperature is lowered.

The tolerance of any plant tissue at low O<sub>2</sub> tension is smaller as the storage temperature increases, since the requirements for aerobic respiration of the tissue increases with increasing temperature. Depending on the product damage associated with CO<sub>2</sub> can both increase and decrease with rise in temperature. CO<sub>2</sub> production increases with temperature, but its solubility decreases. Thus, the CO<sub>2</sub> concentration in the tissue may decrease or increase with increasing temperature. In addition, the physiological effect of CO<sub>2</sub> could be temperature dependent.

The activation energy is a parameter that has been used to characterize plastic packaging. Knowledge of the activation energy of the breath of product and packaging serves as an important tool to predict the effects of temperature fluctuations on the concentration of gases inside the package (Cameron et al., 1995).

## 6. Plant responses of MAP

Some of the most important factors that affect shelf life of fresh horticultural products are ripening and/or senescence, decay, and cut surface browning. The effect of modified atmosphere on these factors has been studied for different crops. The application of MA to affect these limiting factors can be restricted for some crops by adverse and/or non-beneficial physiological responses. Additionally, the good temperature management should be associated to by atmosphere modification.

Low O<sub>2</sub> and elevated CO<sub>2</sub> concentrations can significantly reduce the rates of ripening and senescence primarily by reducing the synthesis and perception of ethylene (Burg and Burg, 1967; Abeles et al., 1992). Changes in respiration and starch, sugars, chlorophyll, and cell wall constituents during this period can be reduced, and in some cases nearly arrested, by eliminating ethylene action through the use of low O<sub>2</sub>/high CO<sub>2</sub> atmospheres.

Chlorophyll loss, a desirable trait for many climacteric fruits, results in quality loss for many vegetables. Chlorophyll degradation during the senescence of green vegetables can be reduced by low O<sub>2</sub> and elevated CO<sub>2</sub> (Ku and Wills, 1999).

Modified atmospheres are most effective at reducing ripening prior to the onset of ripening, rather than at a later stage. At the same time, packaged fruits and vegetables are usually intended for immediate consumption by the consumer and an unripe product is not immediately edible or is of reduced quality relative to the ripe product. Thus, the advantage of extend shelf-life by retarding ripening runs counter to the needs of the consumer when retail MAP systems are used. Nevertheless, MAP can reduce the rate of ripening of some commodities such as tomato even during its later stages (Yang and Chinnan, 1988).

Decay control is a particularly important problem for many crops. Levels of above 10% CO<sub>2</sub> effectively slow or stop the growth of numerous decay organisms (Brown, 1922). Low O<sub>2</sub> has a very limited effect on decay organism activity or survival at levels above the fermentation threshold of most commodities. Strawberry, blueberry, blackberry, raspberry and cherry, they are examples that can be stored at CO<sub>2</sub> atmosphere between 10 and 20%.

A negative plant responses to modified atmosphere packaging is when the respiration is reduced as O<sub>2</sub> becomes limiting. Although there is a limit to which O<sub>2</sub> can be reduced. The lower O<sub>2</sub> limit is frequently considered to be the level of O<sub>2</sub> that induces fermentation. However there are lower O<sub>2</sub> levels that may confer benefits that outweigh the loss in flavor or other quality parameters. Ethanol, acetaldehyde, ethyl acetate and lactate are products of fermentation that can contribute to the development of off-flavors (Kays, 1997; Mattheis and Fellman, 2000).

Synthesis of aroma compounds are generally suppressed by high CO<sub>2</sub> and low O<sub>2</sub> levels, in part by their action on ethylene sensitivity, but also via action of O<sub>2</sub> on oxidative processes, including respiration required for substrate production. But low O<sub>2</sub> MAP may suppress aroma production so consumers perceive reduced quality upon opening the container.

## 7. Conclusions

The benefits of MAP technology to the manufacturer, retailer as well as consumer far outweigh the drawbacks. Nevertheless some critical points should be considered in this technology. The following list some advantages and disadvantages of MAP.

- Advantages
  - Increased shelf-life allowing lesser frequency of loading of retail display shelves.
- Improved presentation of the product
  - Hygienic stackable pack sealed and free from product drip and odor
  - Shelf-life can be increase by 50 to 400%.
  - Reduction in production and storage costs due to better utilization of space and equipment.
- Disadvantages
  - Capital cost of gas packaging machinery
  - Increased pack volume increases transport costs and retail display space
  - Cost of gases and packaging materials
  - Temperature control is of critical importance and, by itself, has a greater impact than atmosphere modification for most products
  - Potential growth of food borne pathogens due to non-maintenance of required storage temperature by retailers and consumers.

## 8. References

- Abeles, F. B., Morgan, P. W., & Saltveit, M. E. *Ethylene in plant biology*. 2<sup>nd</sup> ed. San Diego: Academic Press, 1992. 414 p.
- Beaudry, R.M., & Gran, C. D. (1993). Using a modified-atmosphere packaging approach to answer some postharvest questions: Factors affecting the lower oxygen limit. *Acta Hort.*, Vol. 362, pp. 203-212.
- Beaudry, R. M. (2000). Responses of horticultural commodities to low oxygen: limits to the expanded use of modified atmosphere packaging. *HortTechnology*, Vol. 10, pp. 491-500.
- Beaudry, R. M. (1999). Effect of O<sub>2</sub> and CO<sub>2</sub> partial pressure on selected phenomena affecting fruit and vegetable quality. *Postharvest Biol. Technol.*, Vol 14, pp. 293-303.
- Beaudry, R. M., Cameron, A. C., Shirazi, A, & Dostal-Lange, D. L. (1992). Modified-atmosphere packaging of blueberry fruit: Effect of temperature on package O<sub>2</sub> and CO<sub>2</sub>. *Amer. Soc. Hort.* Vol. 117, pp. 436-441.
- Beaudry, R. M., & Gran, C. D. (1993). Using a modified-atmosphere packaging approach to answer some postharvest questions: Factors affecting the lower oxygen limit. *Acta Hort.* Vol. 362, pp. 203-212.
- Boyette, M.D., Sanders, D.C., & Rutledge, G.A. (1996). Package requirements for fresh fruits and vegetables. The North Carolina Agricultural Extension Service. North Carolina State University, Ralley, NC. USA. Publication no 9/96-3m-TWK-260373-AG-414-8.
- Brecht, J. K. (1995). Physiology of lightly processed fruits and vegetables. *HortScience*, Vol. 30, No. 1, pp. 18-21.


- Brown, W. (1922). On the germination and growth of fungi at various temperatures and in various concentrations of oxygen and carbon dioxide. *Ann. Bot.* Vol. 36, pp. 257-283.
- Burg, S. P., & Burg, E. A. (1967). Molecular requirements for the biological activity of ethylene. *Plant Physiol.* Vol.42, pp. 114-152.
- Cameron, A. C., Beaudry, R. M., Banks, N. H., & Yelanich, M. V. (1994). Modified atmosphere packaging of blueberry fruit: modeling respiration and package oxygen partial pressures as a function of temperature. *Journal of the American Society for Horticultural Science.* Vol. 119, no. 3, pp. 534-539.
- Cameron, A. C., Patterson, B. D., Talasila, P. C., & Joles, D. W. (1993). Modeling the risk in modified atmosphere packaging: A case for sense-and-respond packaging. In: Proc. 6<sup>th</sup> Intl. Controlled Atmosphere Res. Conf., Ithaca, NY, pp. 95-112.
- Cameron, A. C., Talasila, P. C., & Joles, D.J. (1995). Predicting the film permeability needs for modified atmosphere packaging of lightly processed fruits and vegetables. *HortScience* Vol. 30, pp. 25-34.
- Cantwell, M. (2004). *Fresh-cut vegetables*. USA: University of California, Davis. p. 78-85. (Postharvest Horticulture Series n. 10.)
- Chinnan, M. S. (1989). Modeling gaseous environment and physio-chemical changes in fresh fruits and vegetables in modified atmospheric storage. (In) "Quality factors of fruits and vegetables - Chemistry and Technology". J. J. Jen (ed.). ACS Symposium Series No. 405, p. 189-202. Washington, D.C.
- Day, B. P. F. (1996). High oxygen modified atmosphere packaging for fresh prepared produce. *Postharvest News and Information*, Wallingford, v. 7, n. 3, p. 1N-34N.
- Fidler, J. C., & North, C. J. (1967). The effect of conditions of storage on the respiration of apples. I. The effects of temperature and concentrations of carbon dioxide and oxygen on the production of carbon dioxide and uptake of oxygen. *Journal of Horticultural Science*, 42, 189-206.
- Fishman, S., Rodov, V., & Ben-Yehoshua, S. (1996). Mathematical Model for Perforation Effect on Oxygen and Water Vapor Dynamics in Modified-Atmosphere Packages. *Journal of Food Science*, Vol. 61, No. 5, pp. 956-961.
- Fonseca, S. C., Oliveira, F. A. R., & Brecht, J. K. (2002). Modelling respiration rate of fresh fruits and vegetables for modified atmosphere packages: a review. *Journal of Food Engineering*, v. 52, p. 99-119.
- Fonseca, S.C., Oliveira, F.A.R., & Brecht, J.K. (2002). Modelling respiration rate of fresh fruits and vegetables for modified atmosphere packages: a review. *Journal of Food Engineering*, Vol. 52, pp. 99-119.
- González, J., Ferrer, A., Oria, R., & Salvador, M. L. (2008). Determination of O<sub>2</sub> and CO<sub>2</sub> transmission rates through microperforated films for modified atmosphere packaging of fresh fruits and vegetables. *Journal of Food Engineering*, Vol. 86, No. 2, pp. 194-201.
- Gorny, J. R. (1997). A summary of CA and MA requirements and recommendations for fresh-cut (minimally-processed) fruits and vegetables. In: J. Gorny (ed) *Fresh-cut fruits and vegetables and MAP*. Postharvest Hort. Series No. 19, Univ. Calif., Davis CA, CA'97 Proc. 5:30-66.

- Gorris, L., & Tauscher, B. (1999). Quality and safety aspects of novel minimal processing technology. *Processing of foods: Quality optimization and process assessment*. CRC Press, USA, pp. 325-339.
- Herner, R.C. (1987). High CO<sub>2</sub> effects on plant organs. In: J. Weichman (ed) *Postharvest Physiology of Vegetables*, Marcel Dekker, NY, pp. 239-253.
- Hertog, M.L.A.T.M., Peppelenbos, H. W., Evelo, R.G., & Tijskens, L.M.M. (1998). A dynamic and generic model of gas exchange of respiring produce: the effects of oxygen, carbon dioxide and temperature. *Postharvest Biol. Technol.* Vol. 14, PP. 335-349.
- Joles, D. W., Cameron, A. C., Shirazi, A., Petracek, P. D., & Beaudry, R. M. (1994). Modified atmosphere packaging oh 'Heritage' red raspberry fruit: respiratory response to reduce oxygen, enhanced carbon dioxide and temperature. *Journal of the American Society for horticultural Science*. Vol. 119, No. 3, pp. 540-545.
- Kader, A.A. (2002). *Post-harvest technology of horticultural crops*. Oakland: University of California, Division of Agriculture and Natural Resources Publication 3311, 535 pp.
- Kader, A.A. (1997). A summary of CA requirements and recommendations for fruits other than apples and pears. In: A. Kader (ed) *Fruits other than apples and pears*. *Postharvest Hort. Series No. 17*, Univ. Calif., Davis CA, CA'97 Proc. 2:1-36.
- Kader, A. A., Zagory, D., & Kerbel, E. L. (1989). Modified atmosphere packaging of fruits and vegetables. *Rev. Food Science and Nutrition*, Vol. 28, No. 1, pp. 1-30.
- Kays, S. J. (1997). *Postharvest physiology of perishable plant products*. Van Nostrand Reinhold, NY.
- Ku, V. V. V., & Wills, R. B. H. (1999). Effect of 1-methylcyclopropene on the storage life of broccoli. *Postharvest Biol. Technol.* Vol. 17, pp. 127-132.
- Kupferman, E. (1997). Controlled atmosphere storage of apples. In: E.J. Mitcham (ed) *Apples and Pears*. *Postharvest Hort. Series No. 16*, Univ. Calif., Davis CA, CA'97 Proc. 3:1-30.
- Lakakul, R., Beaudry, R. M., & Hernandez, R. J. (1999). Modeling respiration of apple slices in modified-atmosphere packages. *Journal of Food Science*. Vol. 64, No. 1, pp. 105-110.
- Lee, D. S., Hagggar, P. E., Lee, J., & Yam, K. L. (1991). Model for fresh produce respiration in modified atmospheres based on principles of enzyme kinetics. *Journal of food Science*, Vol. 56, No. 6, pp. 1580-1585.
- Mattheis, J.P., Fellman, & J.P. (2000). Impact of modified atmosphere packaging and controlled atmosphere on aroma, flavor and quality of horticultural produce. *HortTechnology*, Vol. 10, pp. 507-510.
- Mir, N., & Beaudry, R. M. (1986). *Modified Atmosphere Packaging*. USDA Handbook 66. Washington, D.C. GPO. 7/11/2011, Available from: <<http://www.ba.ars.usda.gov/hb66/015map.pdf>>.
- Reid, M.S. (1997). A summary of CA and MA requirements and recommendations for ornamentals and cut flowers. In: M.E. Saltveit (ed) *Vegetables and ornamentals*. *Postharvest Hort. Series No. 18*, Univ. Calif., Davis CA, CA'97 Proc. 4:129-136.
- Richardson, D. G., & Kupferman, E. (1997). Controlled atmosphere storage of pears. In: E.J. Mitcham (ed) *Apples and pears*. *Postharvest Hort. Series No. 16*, Univ. Calif., Davis CA, CA'97 Proc. 2:31-35.
- Saltveit, M. E. (1997). A summary of CA and MA recommendations for harvested vegetables. In: M.E. Saltveit (ed) *Vegetables and ornamentals*. *Postharvest Hort. Series No. 18*, Univ. Calif., Davis CA, CA'97 Proc. 4:98-117.

- Sanz, C., Pérez, A.G., Olías, R., & Olías, J.M. (1999). Quality of Strawberries Packed with Perforated Polypropylene. *Journal of Food Science*, Vol. 64, No. 4, pp. 748-752.
- Sharp, A. K., Irving, A.R., & Morris, S.C. (1993). Does temperature variation limit the use of MA packaging during shipment in freight containers? In: G. Blanpied, J. Bartsch and J. Hicks (eds) Proc. 6<sup>th</sup> Intl Contr. Atmos. Res. Conf., Cornell Univ., Ithaca NY, pp. 238-251.
- Shirazi, A., & Cameron, A.C. (1992). Controlling relative humidity in modified-atmosphere packages of tomato fruit. *HortScience*, Vol. 27, pp. 336-339.
- Smyth, A. B., Song, J., & Cameron, A. C. (1998). Modified atmosphere packaged cut iceberg lettuce: effect of temperature and O<sub>2</sub> partial pressure on respiration and quality. *Journal of Agricultural and Food Chemistry*, Vol. 46, pp. 4556-4562
- Smyth, A. B., Talasila, P. C., & Cameron, A. C. (1999). An ethanol biosensor can detect low-oxygen injury in modified atmosphere packages of fresh-cut produce. *Postharvest Biol. Technol.*, Vol. 15, pp.127-134.
- Talasila, P. C., Chau, K. V., & Brecht, J. K. (1995). Modified atmosphere packaging under varying surrounding temperature. *Trans. ASAE*, Vol. 38, pp. 869-876.
- Tano, K., Oulé, M. K., Doyon, G., Lencki, R. W., & Arul, J. (2007). *Postharvest Biology and Technology*, Vol. 46, pp. 212-221.
- Yang, C. C., & Chinnan, M.S. (1988). Modeling the effect of O<sub>2</sub> and CO<sub>2</sub> on respiration and quality of stored tomatoes. *Trans. ASAE*, Vol. 31, pp. 920-925.

INTECH





## **Polypropylene**

Edited by Dr. Fatih Dogan

ISBN 978-953-51-0636-4

Hard cover, 500 pages

**Publisher** InTech

**Published online** 30, May, 2012

**Published in print edition** May, 2012

This book aims to bring together researchers and their papers on polypropylene, and to describe and illustrate the developmental stages polypropylene has gone through over the last 70 years. Besides, one can find papers not only on every application and practice of polypropylene but also on the latest polypropylene technologies. It is also intended in this compilation to present information on polypropylene in a medium readily accessible for any reader.

### **How to reference**

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Leonora M. Mattos, Celso L. Moretti and Marcos D. Ferreira (2012). Modified Atmosphere Packaging for Perishable Plant Products, Polypropylene, Dr. Fatih Dogan (Ed.), ISBN: 978-953-51-0636-4, InTech, Available from: <http://www.intechopen.com/books/polypropylene/modified-atmosphere-for-perishable-plant-products>

# **INTECH**

open science | open minds

### **InTech Europe**

University Campus STeP Ri  
Slavka Krautzeka 83/A  
51000 Rijeka, Croatia  
Phone: +385 (51) 770 447  
Fax: +385 (51) 686 166  
[www.intechopen.com](http://www.intechopen.com)

### **InTech China**

Unit 405, Office Block, Hotel Equatorial Shanghai  
No.65, Yan An Road (West), Shanghai, 200040, China  
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元  
Phone: +86-21-62489820  
Fax: +86-21-62489821