

Non Invasive Prenatal Diagnosis of Down Syndrome

Dimitra Kappou¹, Eleftheria Papadopoulou² and Stavros Sifakis¹

¹*Department of Obstetrics & Gynecology, University Hospital of Heraklion, Crete*

²*Department of Pediatrics, University Hospital of Heraklion, Crete
Greece*

1. Introduction

Down syndrome (trisomy 21), which has an incidence of 1 in 800 live births, is considered to be the most frequent etiology of mental retardation and it is the predominant reason for women seeking prenatal diagnosis [Driscoll & Gross, 2009]. Trisomy 21 is used as a benchmark because it is the most common aneuploidy compatible with life and is associated with mental retardation and serious congenital anomalies. Currently used screening tests for aneuploidy are based on the assessment of fetal sonographic markers and/or the evaluation of biochemical markers in the maternal circulation during the first and second trimester. Screening test based on the combination of nuchal translucency assessment and biochemical markers at 11⁺⁰-13⁺⁶ weeks of gestation may detect 90-94% of pregnancies affected by Down syndrome at a false positive rate of 5% [Kagan et al., 2008]. The current gold standard for diagnosis of trisomy 21 is provided by invasive sampling of fetal genetic material through chorionic villus sampling (CVS) or amniocentesis followed by conventional cytogenetic or DNA analysis; however, both procedures are associated with an increased risk of fetal loss of about 1% and therefore they are recommended for pregnancies considered to be at high risk of fetal trisomy 21 [Alfirevic et al., 2003].

Since 1997, when cell free fetal DNA in maternal circulation was discovered, the research interest has focused on the development of reliable techniques for non-invasive prenatal diagnosis (NIPD) that would allow the direct analysis of fetal genetic material based on the discovery of cell-free fetal (cff) DNA and RNA in the maternal circulation. Current investigation fields of NIPD include fetal Rhesus D genotype determination in RhD negative women, fetal sex determination for sex-linked disorders and the role of cffDNA in pregnancy disorders such as preeclampsia but the holy grail of NIPD remains the detection of fetal aneuploidies [Honda et al., 2002; Bianchi et al., 2005]. The direct analysis of circulating fetal DNA for the NIPD of chromosomal aneuploidies is mainly complicated by the presence of the coexisting background maternal DNA. NIPD will hopefully overcome the limitations of the currently used methods for diagnosing Down syndrome antenatally and make prenatal testing safer for pregnant women and their fetuses. However, irrespective of which strategy is selected for isolating or distinguishing fetal genetic material in maternal plasma, the small quantity of cffDNA and cff mRNA poses severe technical challenges; all these issues should be addressed before the clinical application of these methods as screening test with high sensitivity, specificity and reproducibility.

In this chapter, we focus on recent advances in the NIPD of Down syndrome via the use of fetal cells or cell-free nucleic acids, and provide an overview of the future perspectives in terms of improvement of enrichment technologies and assaying methods and possibilities for clinical applications as well.

2. Intact fetal cells and cell-free fetal DNA in the maternal circulation

It has been established for over a century that fetal cells circulate in the maternal blood throughout gestation although the rarity of these cells limits eventually the practicability of a diagnostic process based on their enumeration in maternal plasma. A considerable challenge for the researchers in this field is to enrich or isolate these rare fetal cells, either for cytogenetic analysis by fluorescence in situ hybridization (FISH) or for analysis of fetal cell DNA by other molecular techniques [Bianchi and Hanson, 2006; Mavrou et al., 2007]. Several studies have reported that the number of fetal cells in maternal blood is markedly increased up to six times in women bearing aneuploid fetuses [Falcidia et al., 2004]. A large-scale study for this cell-based approach, conducted by the National Institute of Child Health and Human Development in the USA, demonstrated that detection of trisomy in these fetal nucleated erythrocyte cells is difficult possibly due to the fact that the chromosomes in these cells disintegrate some time before the nucleus is eliminated from the cell, making FISH analysis of samples from maternal circulation unreliable [Bianchi et al., 2002; Babochkina et al., 2005]. The use of fetal cells other than fetal nucleated red blood cells also found in maternal circulation has been studied, however these cells are able to persist for years, or even decades, following previous pregnancies and this persistence limits their potential value for NIPD [Guetta et al., 2003]. Possible explanations include a simple presence of these fetal cells or the fact that the maternal hematopoietic system becomes engrafted with fetal stem cells during pregnancy [Puszyk et al., 2008].

The first experimental demonstration of cffDNA in the maternal plasma and serum of women carrying male fetuses by Lo et al., (1997), opened up new possibilities in NIPD. Lo et al. were inspired by previous reports that documented the presence of tumour-derived DNA in the plasma of women suffering from a variety of cancer types. Potential sources of cffDNA include the fetal nucleated red blood cells which undergo apoptosis in the maternal circulation but the most likely source of origin is the placenta [Alberry et al., 2007]. Paternally derived DNA sequences in cffDNA can be reliably identified in maternal plasma, from as early as 5 weeks after conception and there is a positive correlation with gestational age. In particular, the median values of the quantity of cffDNA are 15.9, 21.5 and 52.0 genome equivalents/ml of blood in the first, second and third trimester respectively; the accumulation of cffDNA as pregnancy progresses lends further support to the placental origin of cffDNA [Lo et al., 1998; Sekizawa et al., 2001; Birch et al., 2005]. In particular, cytotrophoblasts (CTBs) are a likely candidate as a source of cffDNA; however, the increased rate of hyperploidy in these cells and the yet unknown relationship between the ploidy status of these cells and the ploidy of cffDNA in maternal circulation could make a diagnostic test problematical [Weier et al., 2005]. Moreover, there is a link between hypoxia and an increasing release of cffDNA that led to the suggestion that it may be a useful biomarker to assess well-being of the placenta during pregnancy [Tjoa et al., 2006]. It is known that cffDNA represents a mean of 3-6% of the DNA that is present in maternal plasma while the bulk of the DNA is derived from the mother herself and a rapid clearance occurs post partum with a half-life in the order of 16 minutes despite the narrow conflicting

results [Invernizzi et al., 2002; Rijnders et al., 2004]. First applications of cffDNA included prenatal determination of fetal sexing based on paternally derived DNA sequences such as SRY, determination of Rhesus-D status of the fetus and detection of paternally inherited genetic abnormalities [Lo, 2006; Van der Schoot et al., 2006].

3. Detection of trisomy 21 from nucleic acids in the maternal plasma

There are two approaches for the detection of trisomy 21 based on the use of cffDNA in maternal circulation. The first is defined as the relative chromosome dosage (RCD) method and compares the quantity of a chromosome 21-derived DNA sequence in cffDNA with the amount of a reference DNA sequence in cffDNA derived from a chromosome other than chromosome 21 [Lo et al., 2007a]. In a normal pregnancy the RCD of chromosome 21 is 2:2, whereas in trisomy 21 the RCD is expected to be 3:2. The second method is defined as the allele ratio (AR) method and involves the allelic ratio of single nucleotide polymorphisms (SNPs) present in a fetal-specific nucleic acid marker [Tong et al., 2006]. In a normal pregnancy where the fetus is heterozygous for a particular gene sequence, the AR in cffDNA is expected to be 1:1 whereas in a case of trisomy 21 the AR of chromosome 21 would be 2:1. The main disadvantage of this approach is that it is applicable only to heterozygous fetuses for the analyzed SNP.

3.1 Fetal DNA enrichment methods

The detection of fetal chromosomal aneuploidies with the aid of cffDNA presents considerable technical challenge: first, to select a subset of nucleic acid in maternal plasma that is completely fetal specific and second, to determine the chromosomal dosage in this subset. The major technical challenge that makes NIPD a demanding task is that cffDNA makes up a low proportion in maternal plasma in a high background of maternal DNA. Currently, several assay procedures are developed in order to enrich and enhance the fractional concentration of fetal DNA or just to distinguish the cffDNA in maternal blood samples. One point of differentiation between cffDNA and cell-free maternal DNA (cfmDNA) is that the first has a shorter size distribution (the majority being 145 bp in length or shorter whereas cfmDNA is significantly longer) [Li et al., 2004]. Based on this observation, researchers try to apply methods of size fractionation with the aid of various kits and columns that rely on the inability of large molecular weight DNA to pass through or by retention of low molecular weight DNA in a gel or column [Legler et al., 2007]. Main disadvantages of this approach are: a) the currently used electrophoretic method is labor-intensive and probably prone to contaminations and b) it is unknown if the provided DNA enrichment is enough satisfactory for the prenatal diagnosis of chromosomal aneuploidies [Lo, 2008]. In 2004, Dhallan's group proposed a specific blood processing protocol in which the addition of formaldehyde in maternal blood samples before centrifugation dramatically increased the percentage of fetal DNA recovered with the concurrent suppression of the maternal DNA background [Dhallan et al., 2004]. There are two speculations about the role of formaldehyde in increased yield of fetal DNA: a) prevention of maternal cell lysis and subsequent reduction of the amount of cfmDNA, and b) prevention of the degradation of cffDNA via its nuclease inhibitory effect [Dhallan et al., 2004]. The same research group supported that the application of this technique resulted in a significant increase in the proportion of cffDNA present from a maximum of about 6% to mean values of 20.2-25% in samples collected during various stages of gestation [Dhallan et al., 2004]. In an attempt to

reproduce these results, other investigators confirmed the previous results and reported a similar or a less pronounced increase of cffDNA (1-3%) whereas other studies yielded inconsistent results [Costa et al., 2004; Chinnapapagari et al., 2005]. A possible reason for this discrepancy is that the sample processing time differs between the studies and it is known that the amount of time spent in the tube affects the concentration of total cell-free DNA [Zhang et al., 2008]. Zhang et al. (2008) proposed that the formaldehyde addition will offer a beneficial effect if there is a delay > 6 hours in sample processing as they demonstrated no maternal blood lysis or released extra maternal free DNA into plasma within the first six hours. Future studies should be conducted to clarify the contribution of elapsed time between blood-taking and processing on the recovery of cell-free DNA from maternal plasma and determine other confounding factors in the effect of formaldehyde. The quantification of cffDNA in maternal circulation from women carrying Down syndrome fetuses could also serve as a prognostic marker for trisomy 21 as quantitative aberrations in biochemical markers of placental origin that contribute to the aneuploidies screening tests. Previous studies present conflicting results as both a two-fold increase and no significant difference in maternal concentration of cffDNA have been reported [Lee et al., 2002; Spencer et al., 2003]. Possible explanations of the observed discrepancy between the reported results include the small number of samples examined, the variable degree of placental apoptosis, the broad ranges of cffDNA concentration at each stage of pregnancy and other sampling or methodological variables that might affect the level of circulating cffDNA. In a recent study, DNA from pre-CVS maternal samples was extracted from 72 trisomy 21 and 264 control pregnancies and authors concluded that there is no difference in first trimester cffDNA levels and the quantification of cffDNA (studied only in pregnancies with male fetuses) has no prognostic value at least in the early stages of pregnancy [Gerovassili et al., 2007]. However, quantification of cffDNA in maternal plasma might be a valuable second-trimester serum marker of Down syndrome pregnancy. Farina et al., found that the maternal serum fetal DNA concentrations were elevated in 15 Down syndrome cases during the second trimester and that fetal DNA could give a 21% detection rate at a 5% false positive rate; in addition, fetal DNA increased the estimated detection rate of quadruple test from 81% to 86% at a 5% false positive rate [Farina et al., 2003]. Main limitation of this approach is that its screening performance has been evaluated only in pregnancies with male fetuses with the aid of unique DNA sequences on the Y chromosome and when a reliable gender-independent fetal DNA marker will be assayed, its clinical utility should be reassessed.

3.2 The role of epigenetic markers in rapid detection of Down syndrome

The term epigenetics refers to the molecular processes that affect gene expression with the concurrent avoidance of any change in DNA sequence or content. The most studied epigenetic process is the DNA methylation, which involves the addition of a methyl group to the cytosine residues of a DNA sequence and when it occurs in the promoters of genes has an inhibitory effect on the gene expression. Epigenetic markers for cffDNA have been discovered for other aneuploidies; these are *SERPINB5* (serpin peptidase inhibitor, clade B, member 5; also known as maspin) on chromosome 18 and *RASSF1A* (Ras association [RaIGDS/AF-6] domain family 1) on chromosome 3 [Lun et al., 2007]. In addition, the allelic ratio for placental-derived hypomethylated *SERPINB5* molecules in maternal plasma was further shown to be valuable in the non-invasive detection of trisomy 18 [Tong et al., 2006]. Nowadays, there is intense interest to identify differentially methylated DNA patterns on chromosome 21 between the placenta and maternal blood cells in order to

develop a similar method for the NIPD of Down syndrome. Such epigenetic markers could be useful either via the analysis of the epigenetic allelic ratios or directly compared with a placenta-derived DNA methylation marker on a reference chromosome [Tong et al., 2006]. A potential issue for any epigenetic approach to NIPD is the interindividual epigenetic variation as it has been documented in monozygotic twins; moreover, this process is regulated in a dynamic manner as epigenetic differences seem to increase over time in a process described as "epigenetic drift". However, epigenetic biomarkers sequences whose methylation has a functional significance may be subject to less individual variation than others with no functional constraint. The discovery of a number of DNA sequences that are differentially methylated between maternal and fetal DNA could provide novel markers for cfDNA via the quantification of fetal-specific DNA sequences derived from chromosome 21. One previous study described the methylation status for chromosome 21 in placenta and blood samples after the selection of sequences in promoter and non-promoter regions but it relied on an assay that used a methylation-sensitive restriction enzyme, HpaII that enables the analysis of a small proportion of all the CpG sites in the human genome [Old et al., 2007]. Differentially methylated sequences located at 21q22.3 (AIRE, SIM2 and ERG genes), 1q32.1 (CD48 gene and FAIM3 gene), 2p14 (ARHGAP25 gene) and 12q24 (SELPLG gene) were identified. Moreover, it was demonstrated that the methylation status for the sequences tested was not altered between early and term pregnancy [Old et al., 2007]. Recently, Chim et al. have performed a systematic search of 114 studied genomic regions (CpG islands) on chromosome 21 in a search for loci that were differentially methylated in placental tissue and blood cells and identified 22 (19%) that showed epigenetic differences between the maternal and fetal tissues [Chim et al., 2008]. The next step was to propose two new fetal-DNA epigenetic markers, *U-PDE9A* and *U-CG1137* found in the maternal circulation only during pregnancy and rapidly cleared upon delivery of the fetus [Chim et al., 2008]. This research group used a high resolution approach via bisulphite sequencing that increased the number of applicable CpG sites by 5-fold compared with the above-mentioned HpaII-based approach. These promising results suggest that fetal-specific epigenetic markers on chromosome 21 may provide a rich source of markers for NIPD. A novel method of trisomy 21 detection measures the ratio of a fetal-specific epigenetic marker on chromosome 21 (the putative promoter of the holocarboxylase synthetase (HLCS) gene) that is hypermethylated in the placenta and a genetic marker (ZFY, zinc finger protein, Y-linked) to determine the chromosome-dosage comparison in 5 maternal plasma samples from women carrying a fetus with Down syndrome [Tong et al., 2010]. Instead of ZFY, any other Y-chromosomal markers or any fetal-specific genetic targets that will be applied in female fetuses could also be used. Also, the placenta-specific epigenetic signature could be combined with the RNA transcripts of placental origin. This epigenetic-genetic chromosome dosage approach appears to be more precise compared to an approach based purely on epigenetic markers that will be extensively affected by the variability in the level of DNA methylation of individual molecules. Another group of investigators presented an alternative approach using methylation-dependent immunoprecipitation (MeDiP) that captures methylated sites combined with real-time quantitative PCR and identified 14 trisomy 21 cases and 26 euploid controls from pregnancies of 11-14 weeks old [Papageorgiou et al., 2011]. The accurate diagnosis of fetuses with Down syndrome was based on the ratio of a subset of fetal-specific methylated regions located on chromosome 21 compared with normal cases and regarding the clinical performance of the method both the sensitivity and

specificity were 100% [Papageorgiou et al., 2011]. The main methods performed for the study of DNA methylation are methylation-dependent immunoprecipitation (MeDiP), bisulphite conversion of DNA, and methylation sensitive restriction endonuclease assay to digest away the maternal sequences. Main limitations of the most commonly used methods for DNA methylation analysis are that the use of bisulphite-based reagents results in DNA degradation (up to 96%) and thus in reduction of target DNA available for subsequent analysis and the methylation sensitive restriction endonuclease assay is limited to the differentially methylated regions that contain a restriction site [Grunau et al., 2001].

4. Chromosome 21-encoded mRNA of placental origin in maternal circulation

In 2000, Poon et al. showed that mRNA transcribed from the Y chromosome could be detected in the plasma of women carrying male fetuses [Poon et al., 2000]. Since then, a series of reports confirmed that cell-free fetal mRNA (cffRNA) circulates in the maternal plasma in a relatively protected form and is predominately placental in origin; therefore, it could be valuable in NIPD for Down syndrome [Tsui et al., 2002]. The underlying mechanisms by which mRNA appears in the maternal plasma remain unknown, although programmed cell death (apoptosis) seems to be involved. The transfer of cffRNA is unidirectional from the placenta to the maternal circulation and microarray-based studies of the placenta are conducted to investigate the global mRNA expression profiles in placenta, a tissue type that is only present in the fetus [Maron et al., 2007]. The mRNA transcripts of two genes expressed in the placenta, human placental lactogen (hPL) and human chorionic gonadotrophin (β HCG) have already been detected and quantified throughout gestation in maternal circulation [Chiu et al., 2006].

The potential utilization of cffRNA in detecting fetal trisomy is based on the assumption that the allelic ratio in mRNA matches the chromosomal AR; therefore, the research interest is focused on the discovery of single nucleotide polymorphisms (SNPs) as biomarkers that will exhibit the 2:1 ratio of alleles in trisomy 21 to ascertain the aneuploidy status [RNA-SNP allelic ratio approach]. Candidate mRNA markers should be encoded from genes located on chromosome 21 and be detectable in maternal plasma during early pregnancy. The first valuable cffRNA marker shown to be highly accurate in assessing trisomy 21 is PLAC4 (placenta-specific 4) mRNA transcribed from the PLAC4 gene on chromosome 21 and originating exclusively from fetal cells in the placenta and cleared following delivery of the fetus [Lo et al., 2007 b]. If the fetus is euploid, that is containing two copies of chromosome 21 and thus two copies of the PLAC4 gene, the ratio of the two candidate SNP alleles would be 1:1. Similarly, the ratio of placental mRNA in maternal plasma that is transcribed from each of these two alleles would also be 1:1. However, if the fetus has trisomy 21, then the RNA-SNP allelic ratio would become 1:2 or 2:1 [Lo, 2009]. Compared with the epigenetic approach, the evaluation of RNA-SNP allelic ratio has two advantages; first, the transcription of a gene in the placenta will produce multiple copies of mRNA and second, application of reverse transcriptase PCR to detect mRNA markers is less complicated technique than bisulfite conversion methods used for the identification of epigenetic markers. The main drawback of this RNA-SNP allelic ratio approach is that it relies upon the fetus inheriting two different SNP alleles in a region which is transcribed into mRNA and therefore only fetuses heterozygous for the analysed SNP can be successfully diagnosed. Another candidate gene for this purpose is LOC90625 within the Down syndrome critical region that is over expressed in trisomy 21 placentas even from the first

trimester [Oudejans et al., 2003]. RNA from this chromosome was found to be present in 60-100% of maternal samples depending on the volume of plasma sample analysed leading to the conclusion that the detection of encoded m RNA could be used in NIPD.

This approach seems to be quite promising as in a recent study Lo et al. recruited a sample of 119 pregnancies and through the use of a mass spectrometry-based method for measuring the RNA-SNP allelic ratio precisely, demonstrated that this strategy could achieve a high diagnostic sensitivity and specificity for trisomy 21 (90% and 96.5% respectively) [Lo et al., 2007b]. The application of novel molecular techniques as digital PCR in which individual target molecules are amplified will possibly improve the protocols for plasma RNA processing and extraction and further increase the diagnostic yield. This method could also be useful in the detection of other fetal-derived m RNA species in maternal plasma. Hopefully, the reproducibility of these success rates maybe with the addition of other markers of similar value to that of PLAC4 in large-scale clinical trials will open up new avenues in NIPD.

5. Novel techniques for the prenatal detection of Down syndrome

The urgent need for the widespread application of NIPD in the detection of trisomy 21 has created strong interest in rapid and accurate single-molecule counting methods [digital PCR, multiplexed maternal plasma sequencing] which could be used in routine clinical diagnosis in the form of automated platforms. These methods will be gender- and polymorphism-independent and will detect trisomy 21 cases based on the presence of an elevated amount of chromosome 21 sequences in maternal blood. The main disadvantage of these approaches is that they require the counting of an extremely large number of molecules for markers that are not fetal-specific (random sequences from chromosome 21) and their use demands expensive equipment and reagents and complex bioinformatics methods. We present an overview of the currently proposed techniques that have been associated with encouraging results in the detection of fetuses with Down syndrome and will hopefully be moved into the practical application.

5.1 Digital polymerase chain reaction (PCR) technology

The above-mentioned approaches (the fetal enrichment techniques, the epigenetic markers, the RNA-SNP allelic ratio method) try to resolve the issue of the low fractional concentration of fetal DNA in maternal circulation and the technical challenges that it poses in the direct detection of chromosomal aneuploidies with conventional methods, for example by real-time PCR [Lo et al., 1998]. Recent reports have indicated that digital PCR, a method that was initially applied in the determination of the allelic frequencies of oncogenic alterations in samples from patients with cancer, could be a valuable new tool in NIPD of trisomy 21 [Zimmermann et al., 2008]. In 2007, Lo et al., used digital PCR to discriminate trisomy 21 placental DNA samples from euploid ones after having applied this method in the measurement of the RNA-SNP allelic ratio for non-invasive detection of fetal aneuploidy in microwell plates [Lo et al., 2007]. In particular, they were able to distinguish four aneuploid fetuses from nine normal ones based on the PLAC4 m RNA SNP approach. In their second trial, the same research group applied a dosage approach and compared the dosage of a locus on chromosome 21 to a locus on the reference chromosome 1 and tried to detect fetal aneuploidy in artificial mixtures of euploid and aneuploidy DNA with as low as 25% trisomic material (a concentration that could be obtained in clinical samples using

enrichment strategies for cffDNA sequences). The innovation of digital PCR is that multiple PCRs are performed in parallel and each PCR will contain either a single or no target molecule. Subsequently, the counting of the number of the positive reactions at the end of amplification will lead to the estimation of the number of input target molecules. The same research group has also defined the number of molecules needed for trisomy detection in different fetal DNA concentrations. Digital PCR does not depend on allelic distribution or gender and is able to detect signals in the presence of mosaics or contaminating maternal DNA; the widespread application of this method is still limited by that the fact that a large number of digital PCRs are needed for each analysis [Fan and Quake, 2007]. Fan and Quake used for their assay material obtained from a cell line with trisomy 21 and genetic material from cells with a normal genomic complement on a microfluidic chip [Fan and Quake, 2007]. The next step was to compare the dosage of an amyloid gene sequence on chromosome 21 to that of the GAPDH [glyceraldehydes 3-phosphate dehydrogenase] on chromosome 21 which was used as reference sequence. Their preliminary results suggest that digital PCR should be indicated for the discrimination between aneuploid and normal samples. It is noteworthy that the discrimination was possible even when the aneuploidy material represented a low proportion (10%) of the total material being examined, indicating the clear advantage of this method over conventional techniques such as real-time PCR or fluorescent quantitative PCR (QF-PCR). A significant barrier for using digital PCR is the small fraction of cffDNA in maternal plasma but an approach like size-fractionation that enriches cffDNA could overcome it. So far the reported encouraging results come from few preliminary studies and the question whether this method could be introduced as a screening tool has not yet been answered. In the near future, it seems likely that the efficacy of novel applications as microfluidic digital PCR and emulsion PCR that allow the simultaneous performance of few thousands of reactions in a single PCR step will be assessed [Zimmermann et al., 2008].

5.2 Shotgun sequencing DNA

Shotgun sequencing DNA technology is based on the massively parallel sequencing of DNA that produces tens of millions of short sequence tags in a single run followed by mapping to the chromosome of origin and measurement of the over- and underrepresentation of chromosomes from an aneuploidy fetus [Mardis 2008]. Fan et al., used this method and successfully identified all nine cases of trisomy 21 in their study population at gestational ages as early as the 14th weeks [Fan et al., 2008]. Further studies are required to specify technical features as the sample-volume limitations and the variations in the counts of sequenced fragments from sample to sample; in addition, this technology will contribute to current knowledge about cell-free nucleic acids revealing unknown features about plasma mRNA distributions and epigenetic features of plasma DNA.

5.3 Multiplexed maternal plasma DNA sequencing

Multiplexed maternal plasma sequencing can overcome the difficulty that poses the small proportion of fetal DNA in maternal circulation as it can identify and quantify millions of DNA fragments in biological samples in a span of days [Schuster 2008]. The feasibility and the diagnostic performance of this alternative approach has already been explored in three cohort studies that recruited few Down syndrome cases with promising results [Chiu et al., 2008; Fan et al., 2008; Chiu et al., 2010]. A recent large-scale validity study used multiplexed maternal plasma DNA sequencing analysis in 753 pregnant women at high risk for fetal

trisomy 21 according to the results of conventional screening who underwent invasive procedures for full karyotyping [Chiu et al., 2011]. Two different protocols (2-plex protocol and 8-plex protocol) were used with different levels of sample throughput followed by the measurement of the proportion of DNA molecules that originated from chromosome 21. The 2-plex protocol achieved 100% sensitivity and 97.9% specificity to rule out trisomy 21 with a positive predictive value of 96.6% and negative predictive value of 100% while the 8-plex protocol with which less plasma DNA molecules were analyzed, exhibited a relatively moderate diagnostic performance. The researchers also concluded that if the referrals for amniocentesis or CVS were based on the sequencing tests results, invasive diagnostic procedures could be avoided in about 98% of the cases in a high-risk population [Chiu et al., 2011]. Further studies will confirm the suitability of this method as first trimester screening test in the general population and its cost-effectiveness as it is currently expensive and not easily accessible to diagnostic laboratories.

6. Prenatal detection of Down syndrome through detection of trophoblasts in cervical smears

Fetal cells are also present in the uterine cavity from 5 to 15 weeks of pregnancy and are most probably exfoliated extravillous trophoblasts (shed from the placenta) [Holzgreve and Hahn, 2000]. Trophoblast cells can be retrieved from the cervical canal using aspiration, cryobrush or cotton wool swabs, endocervical lavage, and intrauterine lavage. Initial approaches using endocervical samples obtained by mucus aspiration or by cryobrush resulted in higher success rates of fetal sex prediction [Griffith-Jones et al, 1992; Falcinelli et al, 1998]. However, direct PCR amplifications from unpurified transcervical cells are likely to result in maternal cell contamination. A more recent study using PCR and FISH analyses on transcervical cells resulted in poor detection of fetal cells [Cioni et al, 2003]. To distinguish trophoblast cells from the predominant maternal cell population in transcervical cell samples, antibodies directed against placental antigens were employed [Koumantaki et al, 2001; Bulmer et al, 2003]. These analyses resulted in an overall detection rate of trophoblasts of 25 to 93%.

Another suggestion is that fetal cell search can be improved through better and faster recognition of fetal cells with the aid of automated scanning (automated microscope systems). Theoretically, the automated microscope could work faster and continuously and thus process more cells or more samples than the fatigued human. Analysis of interphase nuclei by FISH, can be used to detect numerical chromosome aberrations (Evans et al., 1992; Ward et al., 1993). The attraction of FISH as a relatively simple approach is based on experience with peripheral blood, amniocentesis samples, and transcervical samples, which have large numbers of cells to examine so that occasional poor signals are only a nuisance. With the few fetal cells available in fetal cell work, FISH quickly shows limitations. It seems necessary to further develop automated microscope systems, which would robotically identify and analyse putative target fetal cells. A recent study tested the hypothesis that fetal cells retrieved from the distal endocervical canal during the first trimester (as early as 5 weeks) may be a source of fetal genetic material for NIPD of trisomy 21 [Sifakis et al., 2011]. The hybridization of fetal cells with chromosome 21 specific probes followed by analysis with an automated fluorescence microscope led to the successful detection in 5 out of 5 trisomy 21 pregnancies [Sifakis et al., 2011]. Examples of the trisomy 21 cells detected, one from a male and the other from a female trisomy 21 pregnancy are shown in the Figure 1.

Additional studies with larger sample size are required to verify the potential of the utilization of fetal cells obtained via cervical samples for NIPD.




Fig. 1. Identification of trisomy 21 cells in cervical samples from trisomy 21 pregnancies. Panel A: Male trisomy 21 nucleus exhibiting X signal (white arrowhead), Y signal (broken blue arrow) and chromosome 21 signals (arrows). Panel B: Female trisomy 21 nucleus exhibiting two X signals (white arrow heads) and three chromosome 21 signals (arrows).

7. Conclusion

The development of a non-invasive genetic test for Down syndrome that would provide true genetic information without carrying risk for the progress of the pregnancy will continue to be an actively researched area in prenatal diagnosis. The trials performed so far highlight the medical and commercial potential of NIPD but the proposed techniques are not yet applicable in clinical practice. A major obstacle in the widespread application of NIPD in clinical diagnostics is that fetal DNA constitutes a small percentage of total DNA in maternal blood and intact fetal cells are even rarer. In the previous years the researchers were trying to discover Y chromosomes sequences or paternally inherited polymorphisms as targeted fetal DNA markers in maternal plasma but still there is no such a single marker that can be applied in all fetal-maternal pairs. One promising alternative approach appears to be the development of gender- and polymorphism- independent fetal DNA markers with a unique methylation pattern that will characterize the placental-derived free DNA in the maternal circulation. In parallel, the refinement of novel sequencing methods will create a

universal test for fetal aneuploidy by using maternal plasma DNA that will not depend on the presence of specific genetic polymorphisms at specific loci but on the enrichment and quantification of cfDNA in maternal peripheral blood. Also, an important goal of the ongoing research is to develop laboratory protocols with the aid of bioinformatics algorithms that will allow their application in large sample numbers. Nevertheless, large-scale studies will need to be performed to confirm the diagnostic efficacy of these methods and subsequently lead to introduction of the experimentally validated strategies into the clinical practice of fetal medicine.

8. References

- Alberry M, Maddocks D, Jones M, Abdel Hadi M, Abdel-Fattah S, Avent N, et al. Free fetal DNA in maternal plasma in anembryonic pregnancies: confirmation that the origin is the trophoblast. *Prenat Diagn* 2007;27:415-8.
- Alfirevic Z, Sundberg K, Brigham S. Amniocentesis and chorionic villus sampling for prenatal diagnosis. *Cochrane Database Syst Rev* 2003;(3):CD003252.
- Babochkina T, Mergenthaler S, Dinges TM, Holzgreve W, Hahn S. Direct detection of fetal cells in maternal blood: a reappraisal using a combination of two different Y chromosome-specific FISH probes and a single X chromosome-specific probe. *Arch Gynecol Obstet* 2005;273:166-9.
- Bianchi DW, Hanson J. Sharpening the tools: a summary of a National Institutes of Health workshop on new technologies for detection of fetal cells in maternal blood for early prenatal diagnosis. *J Matern Fetal Neonatal Med* 2006;19:199-207.
- Bianchi DW, Simpson JL, Jackson LG, Elias S, Holzgreve W, Evans MI, et al. Fetal gender and aneuploidy detection using fetal cells in maternal blood: analysis of NIFTY I data. National Institute of Child Health and Development Fetal Cell Isolation Study. *Prenat Diagn* 2002;22: 609-15.
- Bianchi DW, Avent ND, Costa JM, van der Schoot CE. Noninvasive prenatal diagnosis of fetal Rhesus D: ready for Prime(r) Time. *Obstet Gynecol* 2005;106:841-4.
- Birch L, English CA, O'Donoghue K, Barigye O, Fisk NM, Keer JT. Accurate and robust quantification of circulating fetal and total DNA in maternal plasma from 5 to 41 weeks of gestation. *Clin Chem* 2005;51:312-20.
- Bulmer JN, Cioni R, Bussani C, Cirigliano V, Sole F, Costa C, et al. HLA-G positive trophoblastic cells in transcervical samples and their isolation and analysis by laser microdissection and QF-PCR. *Prenat Diagn* 2003;23:34-9.
- Chim SSC, Jin S, Lee TYH, Lun FMF, Lee WS, Chan LYS, et al. Systematic search for placental epigenetic markers on chromosome 21:towards noninvasive prenatal diagnosis of fetal trisomy 21. *Clin Chem* 2008;54:500-11.
- Chinnapapagari SK, Holzgreve W, Lapaire O, Zimmermann B, Hahn S. Treatment of maternal blood samples with formaldehyde does not alter the proportion of circulatory fetal nucleic acids (DNA and mRNA) in maternal plasma. *Clin Chem* 2005;51:652-5.
- Chiu RWK, Lui W, Cheung M, Kumta N, Farina A, Banzola I, et al. Time profile of appearance and disappearance of circulating placenta-derived mRNA in maternal plasma. *Clin Chem* 2006;52:313-6.
- Chiu RKW, Chan KCA, Gao Y, Lau VYM, Zheng W, Leung TY, et al. Noninvasive prenatal diagnosis of fetal chromosomal aneuploidy by massively parallel genomic sequencing of DNA in maternal plasma. *Proc Natl Acad Sci USA* 2008;105:20458-63.

- Chiu RWK, Sun H, Akolekar R, Clouser C, Lee C, McKernan K, et al. Maternal plasma DNA analysis with massively parallel sequencing by ligation for noninvasive prenatal diagnosis of trisomy 21. *Clin Chem* 2010;54:459-63.
- Chiu RW, Akolekar R, Zheng YW, Leung TY, Sun H, Chan KC, et al. Non-invasive prenatal assessment of trisomy 21 by multiplexed maternal plasma DNA sequencing: large scale validity study. *BMJ* 2011;342:c7401.
- Cioni R, Bussani C, Scarselli B, Bucciantini S, Barciulli F, Scarselli G. Fetal cells in cervical mucus in the first trimester of pregnancy. *Prenat Diagn* 2003; 23:168-71
- Costa JM, Gautier E, Benachi A. Genetic analysis of the fetus using maternal blood. *Gynecol Obstet Fertil* 2004;32:646-50.
- Dhallan R, Au WC, Mattagajasingh S, Emche S, Bayliss P, Damewood M, et al. Methods to increase the percentage of free fetal DNA recovered from the maternal circulation. *JAMA* 2004;291:1114-9.
- Driscoll DA, Gross S. Clinical practice. Prenatal screening for aneuploidy. *N Engl J Med* 2009;360:2556-62.
- Evans MI, Klinger KW, Isada NB, Shook D, Holzgreve W, McGuire N, et al. Rapid prenatal diagnosis by fluorescent in situ hybridization of chorionic villi: an adjunct to long term culture and karyotype. *Am J Obstet Gynecol* 1992;167:1522-5.
- Falcidia E, Parano E, Grillo A, Pavone P, Takabayashi H, Trifiletti RR, et al. Fetal cells in maternal blood: a six-fold increase in women who have undergone amniocentesis and carry a fetus with Down syndrome: a multicenter study. *Neuropediatrics*. 2004 ;35:321-4.
- Falcinelli C, Battafarano S, Neri C, Mazza V, Ranzi A, Forabosco A. Analysis of fetal sex in TCC sample DNA: a contribution to the validation of this approach. *Prenat Diagn* 1998;18:1109-16
- Fan HC, Quake SR. Detection of aneuploidy with digital polymerase chain reaction. *Anal Chem* 2007;79:7576-9.
- Fan HC, Blumenfeld YJ, Chitkara U, Hudgins L, Quake SR. Noninvasive diagnosis of fetal aneuploidy by shotgun sequencing DNA from maternal blood. *Proc Natl Acad Sci USA* 2008;105:16266-71.
- Farina A, LeShane ES, Lambert-Messerlian GM, Canick JA, Lee T, Neveux LM, et al. Evaluation of cell-free fetal DNA as a second-trimester maternal serum marker of Down syndrome pregnancy. *Clin Chem* 2003;49:239-42.
- Gerovassili A, Garner C, Nicolaides KH, Thein SL, Rees DC. Free fetal DNA in maternal circulation: a potential prognostic marker for chromosomal abnormalities? *Prenat Diagn* 2007;27:104-10.
- Griffith-Jones MD, Miller D, Lilford RJ, Scott J, Bulmer J. Detection of fetal DNA in trans-cervical swabs from first trimester pregnancies by gene amplification: a new route to prenatal diagnosis? *Br J Obstet Gynaecol* 1992; 99:508-11.
- Grunau C, Clark SJ, Rosenthal A. Bisulfite genomic sequencing: systematic investigation of critical experimental parameters. *Nucleic Acids Res* 2001;29:E65-5.
- Guetta E, Gordon D, Simchen MJ, Goldman B, Barkai G. Hematopoietic progenitor cells as targets for non-invasive prenatal diagnosis: detection of fetal CD34+ cells and assessment of postdelivery persistence in the maternal circulation. *Blood Cells Mol Dis* 2003;30:13-21.
- Holzgreve W, Hahn S. Fetal cells in cervical mucus and maternal blood. *Baillieres Best Pract Res Clin Obstet Gynaecol* 2000;14:709-22.
- Honda H, Miharu N, Ohashi Y, Samura O, Kinutani M, Hara T, et al. Fetal gender determination in early pregnancy through qualitative and quantitative analysis of fetal DNA in maternal serum. *Hum Genet* 2002;110:75-9

- Invernizzi P, Biondi ML, Battezzati PM, Perego F, Selmi C, Cecchini F, et al. Presence of fetal DNA in maternal plasma decades after pregnancy. *Hum Genet* 2002;110:587-91.
- Kagan KO, Wright D, Baker A, Sahota D, Nicolaides KH. Screening for trisomy 21 by maternal age, fetal nuchal translucency thickness, free beta-human chorionic gonadotropin and pregnancy-associated plasma protein-A. *Ultrasound Obstet Gynecol* 2008;31:618-24.
- Koumantaki Y, Sifakis S, Dragatis G, Matalliotakis I, Froudarakis G, Papadopoulou E, et al. Microsatellite analysis provides efficient confirmation of fetal trophoblast isolation from maternal circulation. *Prenat Diagn* 2001;21: 566-70
- Lee T, LeShane ES, Messerlian GM, Canick JA, Farina A, Heber WW, et al. Down syndrome and cell-free fetal DNA in archived maternal serum. *Am J Obstet Gynecol* 2002;187:1217-21.
- Legler TJ, Liu Z, Mavrou A, Finning K, Hromadnikova I, Galbiati S, et al. Workshop report on the extraction of foetal DNA from maternal plasma. *Prenat Diagn* 2007;27:824-9.
- Li Y, Holzgreve W, Page-Christiaens GC, Gille JJ, Hahn S. Improved prenatal detection of a fetal point mutation for achondroplasia by the use of size-fractionated circulatory DNA in maternal plasma—case report. *Prenat Diagn* 2004;24:896-98.
- Lo YM, Corbetta N, Chamberlain PF, Rai V, Sargent IL, Redman CW, et al. Presence of fetal DNA in maternal plasma and serum. *Lancet* 1997;350:485-7.
- Lo YM, Tein MS, Lau TK, Haines CJ, Leung TN, Poon PM, et al. Quantitative analysis of fetal DNA in maternal plasma and serum: implications for noninvasive prenatal diagnosis. *Am J Hum Genet* 1998;62:768-75.
- Lo YM. Recent developments in fetal nucleic acids in maternal plasma: implications to noninvasive prenatal fetal blood group genotyping. *Transfus Clin Biol* 2006;13:50-2.
- Lo YMD, Lun FMF, Chan KCA, Tsui NB, Chong KC, Lau TK, et al. Digital PCR for the molecular detection of fetal chromosomal aneuploidy. *Proc Natl Acad Sci USA* 2007a;104:13116-21.
- Lo YM, Tsui NB, Chiu RW, Lau TK, Leung TN, Heung MM, et al. Plasma placental RNA allelic ratio permits noninvasive prenatal chromosomal aneuploidy detection. *Nat Med* 2007b;13:218-23.
- Lo YM. Fetal nucleic acids in maternal plasma. Toward the development of noninvasive prenatal diagnosis of fetal chromosomal aneuploidies. *Ann NY Acad Sci* 2008;1137:140-3.
- Lo Y. Noninvasive prenatal detection of fetal chromosomal aneuploidies by maternal plasma nucleic acid analysis: a review of the current state of the art. *BJOG* 2009;116:152-7.
- Lun FM, Chiu RW, Leung TY, Leung TN, Lau TK, Lo YM. Epigenetic analysis of RASSF1A gene in cell-free DNA in amniotic fluid. *Clin Chem* 2007;53:796-8.
- Maron JL, Johnson KL, Slonim D, Lai CQ, Ramoni M, Alterovitz G, et al. Gene expression analysis in pregnant women and their infants identifies unique fetal biomarkers that circulate in maternal blood. *J Clin Invest* 2007;117:3007-19.
- Mavrou A, Kouvidi E, Antsaklis A, Souka A, Kitsiou Tzeli S, Kolialexi A. Identification of nucleated red blood cells in maternal circulation: a second step in screening for fetal aneuploidies and pregnancy complications. *Prenat Diagn* 2007;27:150-3.
- Mardis ER. Next-generation DNA sequencing methods. *Annu Rev Genomics Hum Genet* 2008;9:387-402.
- Old RW, Crea F, Puszyk W, Hultén MA. Candidate epigenetic biomarkers for non-invasive prenatal diagnosis of Down syndrome. *Reprod Biomed Online* 2007;15:227-35.

- Oudejans CB, Go AT, Visser A, Mulders MA, Westerman BA, Blankenstein MA, et al. Detection of chromosome 21-encoded mRNA of placental origin in maternal plasma. *Clin Chem* 2003;49:1445-9.
- Papageorgiou EA, Karagrigroriou A, Tsaliki E, Velissariou V, Carter NP, Patsalis PC. Fetal-specific DNA methylation ratio permits noninvasive prenatal diagnosis of trisomy 21. *Nat Med* 2011;17:510-3.
- Poon LL, Leung TN, Lau TK, Lo YM. Presence of fetal RNA in maternal plasma. *Clin Chem* 2000;46:1832-4.
- Puszyk WM, Crea F, Old RW. Noninvasive prenatal diagnosis of aneuploidy using cell-free nucleic acids in maternal blood: promises and unanswered questions. *Prenat Diagn* 2008;28:1-6.
- Rijnders RJ, Christiaens GC, Soussan AA, van der Schoot CE. Cell-free fetal DNA is not present in plasma of nonpregnant mothers. *Clin Chem* 2004;50: 679-81.
- Schuster SC. Next-generation sequencing transforms today's biology. *Nat Methods* 2008;5:16-8.
- Sekizawa A, Kondo T, Iwasaki M, Watanabe A, Jimbo M, Saito H, et al. Accuracy of fetal gender determination by analysis of DNA in maternal plasma. *Clin Chem* 2001;47:1856-8.
- Sifakis S, Ghatpande S, Seppo A, Kilpatrick MW, Tafas T, Tsipouras P, et al. Prenatal diagnosis of trisomy 21 through detection of trophoblasts in cervical smears. *Early Hum Dev* 2010;86:311-3.
- Spencer K, de Kok JB, Swinkels DW. Increased total cell-free DNA in the serum of pregnant women carrying a fetus affected by trisomy 21. *Prenat Diagn* 2003;23:580-3.
- Tjoa ML, Cindrova-Dvaies T, Spasic-Boskovic O, Bianchi DW, Burton GJ. Trophoblastic oxidative stress and the release of cell-free fetoplacental DNA. *Am J Pathol* 2006;169:400-4.
- Tong YK, Ding C, Chiu RWK, Gerovassili A, Chim SS, Leung TY, et al. Noninvasive prenatal detection of fetal trisomy 18 by epigenetic allelic ratio analysis in maternal plasma: theoretical and empirical considerations. *Clin Chem* 2006;52:2194-202.
- Tong YK, Jin S, Chiu RW, Ding C, Chan KC, Leung TY, et al. Noninvasive prenatal detection of trisomy 21 by an epigenetic-genetic chromosome-dosage approach. *Clin Chem* 2010;56:90-8.
- Tsui NB, Ng EK, Lo YM. Stability of endogenous and added RNA in blood specimens, serum, and plasma. *Clin Chem* 2002;48:1647-53.
- Van der Schoot CE, Soussan AA, Koelewijn J, Bonsel G, Paget-Christiaens LG, de Haas M. Non-invasive antenatal RHD typing. *Transfus Clin Biol* 2006;13:53-7.
- Ward BE, Gersen SL, Carelli MP, McGuire NM, Dackowski WR, Weinstein M, et al. Rapid prenatal diagnosis of chromosomal aneuploidies by fluorescence in situ hybridization: clinical experience with 4,500 specimens. *Am J Hum Genet* 1993;52:854-65.
- Weier JF, Weier HU, Jung CJ, Gormley M, Zhou Y, Chu LW, et al. Human cytotrophoblasts acquire aneuploidies as they differentiate to an invasive phenotype. *Dev Biol* 2005;279:420-32.
- Zhang Y, Li Q, Hui N, Fei M, Hu Z, Sun S. Effect of formaldehyde treatment on the recovery of cell-free fetal DNA from maternal plasma at different processing times. *Clin Chim Acta* 2008;397:60-4.
- Zimmermann BG, Grill S, Holzgreve W, Zhong XY, Jackson LG, Hahn S. Digital PCR: a powerful new tool for noninvasive prenatal diagnosis? *Prenat Diagn* 2008;28:1087-93.



Prenatal Diagnosis and Screening for Down Syndrome

Edited by Prof. Subrata Dey

ISBN 978-953-307-355-2

Hard cover, 232 pages

Publisher InTech

Published online 17, August, 2011

Published in print edition August, 2011

This book provides a concise yet comprehensive source of current information on Down syndrome. Research workers, scientists, medical graduates and paediatricians will find it an excellent source for reference and review. This book focuses on exciting areas of research on prenatal diagnosis - Down syndrome screening after assisted reproduction techniques, noninvasive techniques, genetic counselling and ethical issues. Whilst aimed primarily at research worker on Down syndrome, we hope that the appeal of this book will extend beyond the narrow confines of academic interest and be of interest to a wider audience, especially parents and relatives of Down syndrome patients.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Dimitra Kappou, Eleftheria Papadopoulou and Stavros Sifakis (2011). Non Invasive Prenatal Diagnosis of Down Syndrome, Prenatal Diagnosis and Screening for Down Syndrome, Prof. Subrata Dey (Ed.), ISBN: 978-953-307-355-2, InTech, Available from: <http://www.intechopen.com/books/prenatal-diagnosis-and-screening-for-down-syndrome/non-invasive-prenatal-diagnosis-of-down-syndrome>

INTECH

open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2011 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the [Creative Commons Attribution-NonCommercial-ShareAlike-3.0 License](#), which permits use, distribution and reproduction for non-commercial purposes, provided the original is properly cited and derivative works building on this content are distributed under the same license.