

NSAID Induction of p75^{NTR} in the Prostate: A Suppressor of Growth and Cell Migration Via the p38 MAPK Pathway

Daniel Djakiew
Georgetown University Medical School
USA

1. Introduction

Chronic consumption of Non-Steroidal Anti-Inflammatory Drugs (NSAIDs) has been associated with a reduced incidence of prostate cancer (Nelson & Harris, 2000). Pathologic inflammation can induce oxidative stress generating free radicals that can subsequently react with infectious agents and surrounding cells during induction of innate immunity. Different forms of reactive oxygen species can form DNA adducts by halogenation, nitration and oxygenation of bases (Kang & Sowers, 2008). DNA repair of reactive oxygen species damaged bases can seal in mutations that may eventually lead to transformation of cells and carcinogenesis (Burrows, 2009). NSAIDs inhibit inflammation (Masferrer et al., 1995, Tegeder et al., 2001) and therefore may reduce the incidence of carcinogenesis by preventing free radical transformation of cells. NSAID inhibition of the cyclo-oxygenases (COXs) reduces inflammation, however, the mechanism of action of NSAIDs associated with reduced risk of prostate cancer appears independent of COX inhibition (Quann et al., 2007a; Quann et al., 2007b). NSAIDs represent a diverse category of pharmacological compounds with the common biological activity to reduce inflammation, temperature and pain, but with diverse chemical structures that undoubtedly interact with multiple target molecules with multiple mechanisms of action. In the prostate, selected aryl propionic acid NSAIDs such as ibuprofen and r-flurbiprofen (profens) inhibit epithelial cell growth in a COX independent manner (Quann et al., 2007a; Quann et al., 2007b). A mechanism by which these profens inhibit prostate growth appears to be via re-induction of the p75 neurotrophin receptor (p75^{NTR}) which has been shown to exhibit both tumor suppressor and metastasis suppressor activity (Krygier & Djakiew, 2001a; Krygier & Djakiew, 2002). In pathologic human prostate cancer tissues, the p75^{NTR} protein exhibits focal loss of expression which is further lost with malignant progression of tissues (Perez et al., 1997). Significantly, treatment of prostate tumor cells with profen NSAIDs promotes re-expression of the p75^{NTR} protein that inhibits growth (Quann et al., 2007a; Quann et al., 2007b) consistent with p75^{NTR} dependent tumor and metastasis suppressor activity (Krygier & Djakiew, 2001;2002). The COX independent pathway by which profen NSAIDs induce p75^{NTR} re-expression involves rapid phosphorylation of p38 MAPK and down stream effectors leading to re-expression of p75^{NTR} protein and suppressor activity (Quann et al., 2007b). This review discusses the body of evidence for the inhibitory role of the p75^{NTR} in prostate growth; the pathologic loss of

p75^{NTR} expression during progression to prostate cancer, and the ability of profen NSAIDs to re-induce p75^{NTR} protein expression through the p38 MAPK pathway with concomitant tumor and metastasis suppressor activity which provides a basis for NSAID associated reduced risk of prostate cancer.




Fig. 1. Affinities of the neurotrophin ligands, nerve growth factor (NGF), brain derived neurotrophic factor (BDNF), neurotrophin-3 (NT-3) and neurotrophin-4 (NT-4) for the family of tropomyosin receptor kinases (Trks) and the p75 neurotrophin receptor (p75^{NTR}). Solid arrows show primary affinities, while dotted arrows show secondary affinities.

2. Expression of neurotrophins and their receptors (p75^{NTR} and Trks) in the prostate

The progression of prostate cancer is accompanied by modifications in the expression of growth factors and their receptors (Bostwick et al, 2004). Amongst these are nerve growth factor (NGF) and its receptors, p75^{NTR} and the Trk family (Djakiew, 2000). The neurotrophin family of ligands (Leibrock et al., 1989, Maisonpierre et al., 1990, Hallbook et al., 1991) consisting of nerve growth factor (NGF), brain derived neurotrophic factor (BDNF), neurotrophin-3 (NT-3) and neurotrophin-4 (NT-4) which is identical to NT-5, all of which can bind with similar affinity to the p75^{NTR} (Bothwell, 1995) and with differential affinities to the Trk family (TrkA, TrkB, TrkC) of high affinity receptors (Figure 1). TrkA preferentially binds with NGF, but also binds BDNF. TrkB preferentially binds BDNF (Soppert et al., 1991), but also binds NT-3 and NT-4 (Berkemeier et al., 1991) and TrkC preferentially binds NT-3 (Lamballe et al., 1991), but also binds NT-4 (Figure 1). NGF immunoreactive protein has been localized to normal epithelium (MacGrogan, 1992; Paul et al., 1992) and the stroma of normal, BPH (Djakiew et al., 1991) and cancer tissues of the human prostate (MacGrogan et al., 1992; Djakiew, 1991; Graham et al., 1992). Exogenous NGF β has been shown to stimulate proliferation *in vitro* (Delsite & Djakiew, 1999; Angelsen et al., 1998; Pflug & Djakiew, 1998), and anchorage independent growth of several prostate tumor cell lines (Chung et al., 1992).

BDNF also is also expressed by human prostate stromal cells (Dalal & Djakiew, 1997). Hence, the two neurotrophins, NGF and BDNF appear to function as paracrine factors for prostate epithelial cell growth (Dalal & Djakiew, (1997).

The p75^{NTR} is expressed by normal human prostate epithelial cells (MacGrogan et al., 1992; Graham et al., 1992). Immunoblot (Pflug et al., 1992), immunofluorescence (Graham et al., 1992), and immunohistochemical (Pflug et al., 1992) studies have shown that p75^{NTR} protein expression progressively declines in human prostate cancer (Perez et al., 1997; Pflug et al., 1995; Djakiew et al., 1996). The p75^{NTR} is expressed in PIN tissue (Perez et al., 1997), and shows a gradual decline in the percentage of cells that retain expression with increasing Gleason score of pathologic prostate tissues (Perez et al., 1997). Hence, loss of p75^{NTR} expression appears to be correlated with cancer grade in organ-confined disease (Perez et al., 1997). The p75^{NTR} is also absent in three human cancer cell lines derived from metastases (Pflug et al., 1992). Loss of p75^{NTR} expression in prostate cancer may be related to its role in the induction of programmed cell death (Pflug & Djakiew, 1998; Djakiew et al., 1996). In this context, the p75 neurotrophin receptor (p75^{NTR}) is a 75 kD cell surface receptor glycoprotein that shares both structural and sequence homology with the tumor necrosis factor receptor super-family of proteins (Chao, 1994; Chapman, 1995). Some of these proteins (e.g. p75^{NTR}, p55^{TNFR}, Fas, DRs3-5) have similar sequence motifs of defined elongated structure (Chao, 1994) designated "death domains" based upon their apoptosis inducing function (Chao, 1994). Hence, re-expression of p75^{NTR} by stable and transient transfection showed that the p75^{NTR} inhibits growth of prostate epithelium *in vitro*, at least in part, by induction of programmed cell death (Pflug & Djakiew, 1998). Hence, loss of p75^{NTR} expression appears to eliminate a potential programmed cell death pathway in prostate cancer cells (Figure 2), thereby facilitating the growth of these cancer cells during carcinogenesis (Perez et al., 1997; Djakiew et al., 1996).

Since expression of the p75^{NTR} is lost during malignant progression and transformation of the prostate (Figure 2), NGF mediated growth of cancer cells has been shown to occur via the family of high affinity Trk receptors (Pflug & Djakiew, 1998; Dalal & Djakiew, 1997; Pflug et al., 1995). Differential expression of TrkA, TrkB and TrkC occurs in normal, organ confined and metastatic prostate cancer tissue (Chapman, 1995) and cell lines (Dalal & Djakiew, 1997) suggesting the presence of a neurotrophin mediated proliferative stimulus via the Trk receptors (Figure 2). Expression of the TrkA receptor has been observed in normal prostate epithelial cells, organ confined prostate adenocarcinoma tissues (Djakiew et al., 1996; Pflug et al., 1995; Djakiew et al., 1996; Dionne et al., 1998), in prostate tumors that have metastasized to the bone (Dionne et al., 1998), as well as in several human prostate tumor cell lines derived from metastases (Djakiew et al., 1996). Interestingly, although normal prostate epithelial cells do not express either TrkB or TrkC (Dionne et al., 1998), these receptors are expressed in metastatic prostate cancer of the bone (Dionne et al., 1998). Hence, it appears that normal prostate epithelial cells express TrkA, and that cancer cells continue to express TrkA within primary tumors and at metastatic foci, while TrkB and TrkC expression is subsequently up-regulated within metastatic prostate tumors (Dionne et al., 1998). Pharmacological inhibition of the Trk family of receptors has provided a basis for their role in mediating a proliferative stimulus in tumor cells. In this context, the indolocarbazole kinase inhibitors selectively antagonize Trk receptors at nanomolar concentrations (Berg et al., 1992) and inhibit NGF-stimulated Trk phosphorylation in cancer cell lines (Delsite & Djakiew, 1996). Concurrently, Trk selective indolocarbazoles inhibit growth of cancer cell lines *in vitro* (Delsite & Djakiew, 1996), and *in vivo* (Dionne et al., 1998).

Interestingly, Trk family mutations within the human prostate have not been identified (George et al., 1998). However, the absence of mutations in otherwise genetically unstable prostate tumor DNA suggests that intact Trk family signaling pathways may be important in prostate cancer development (George et al., 1998) consistent with a role as a proto-oncogene.




Fig. 2. Schematic diagram showing expression of NGF and BDNF in prostatic stroma for the paracrine regulation of epithelial cell growth. Pre-malignant epithelial cells express both TrkA and the p75^{NTR}, whereas metastatic tumor cells have lost expression of the p75^{NTR} and gained expression of both TrkB and TrkC.

3. Ectopic re-expression of p75^{NTR} induces tumor suppressor, metastasis suppressor and differentiation phenotypes in prostate tumor cells

Whereas the gene encoding p75^{NTR} is intact in prostate cancer cells (Krygiel & Djakiew, 2001b), expression of the p75^{NTR} protein is suppressed (Perez et al., 1997; Pflug et al., 1992; Pflug et al., 1995). Moreover, transient transfection of two constructs of p75^{NTR} into prostate cell lines that do not express the protein, one construct containing the full 2-kb 3' untranslated region and another that contains just a few hundred bases of the 3' untranslated region, showed that only the truncated construct allowed expression of the p75^{NTR} protein (Krygiel & Djakiew, 2001b). This led to the conclusion that elements of the 3' untranslated region of p75^{NTR} contribute to mRNA stability and p75^{NTR} protein expression (Krygiel & Djakiew, 2001b). Utilizing the truncated p75^{NTR} expression vector that allows protein expression, a series of stable prostate cell lines were developed that express different levels of p75^{NTR} protein (Pflug et al., 1992, Pflug et al., 1995). These cancer cells that

ectopically expressed the p75^{NTR} protein exhibited a retardation of cell cycle progression characterized by accumulation of cells in G1 phase with a corresponding reduction of cells in the S phase of the cell cycle (Krygiel & Djakiew, 2001a). In rank order, prostate cancer cells that expressed increased levels of p75^{NTR} protein exhibited increased rates of apoptosis and reduced rates of proliferation (Krygiel & Djakiew, 2001a). When the same series of tumor cells were injected into the flanks of SCID mice the growth of prostate tumors was suppressed in proportion to increased p75^{NTR} expression levels (Krygiel & Djakiew, 2001a) thereby functionally demonstrating that the p75^{NTR} exhibits tumor suppressor activity (Krygiel & Djakiew, 2001a). Further evidence for the tumor suppressor function of p75^{NTR} was shown utilizing a gene therapy strategy based upon intra-tumoral injection into xenografts of PC-3 prostate tumor cells of a lipoplex containing the p75^{NTR} gene. Administration of the p75^{NTR} gene into subcutaneous PC-3 xenografts suppressed in a dose-dependent manner the growth of tumors (Allen et al., 2004). Within the gene therapy treated tumors re-expression of the p75^{NTR} gene product was associated with increased apoptosis and reduced proliferation of tumor cells (Allen et al., 2004), the net effect of which was to reduce overall growth and tumor volume. Utilizing the same prostate tumor cells that exhibited a rank order (dose-dependent) increase in p75^{NTR} expression for growth of subcutaneous tumors in SCID mice (Krygiel & Djakiew, 2001a), treatment of these tumors with NGF stimulated both proliferation as indicated by PCNA expression, and apoptosis as indicated by TUNEL assay, the net result of which was no change in the overall growth of the tumors (Krygiel & Djakiew, 2002). However, NGF was found to increase the formation of smaller secondary satellite tumors, both contiguous and non-contiguous with respect to the primary tumor mass, indicating dose-dependent induction of metastasis (Krygiel & Djakiew, 2002). Significantly, the formation of satellite tumors was suppressed by the expression of p75^{NTR} thus showing that p75^{NTR} is a tumor suppressor of growth and a metastasis suppressor of NGF stimulated migration (Djakiew et al, 1993) of human prostate tumor cells (Krygiel & Djakiew, 2002). To better understand the molecular mechanism of p75^{NTR} on tumor and metastasis suppression a cDNA microarray composed of approximately 6,000 human cancer-related genes was used to determine the gene expression pattern altered by re-introduction of p75^{NTR} into PC-3 prostate tumor cells (Nalbandian et al., 2005). Comparison of the transcripts in the *neo* control and p75^{NTR}-transfected cells revealed 52 differentially expressed genes, of which 21 were up-regulated and 31 were down-regulated in the presence of p75^{NTR}. The known biological functions of these p75^{NTR} regulated genes suggested a role in the regulation of differentiation as well as cell adhesion, signal transduction, apoptosis, tumor cell invasion and metastasis (Nalbandian et al., 2005). Quantitative real-time polymerase chain reaction and immunoblot analysis confirmed increased CRABPI and IGFBP5 protein levels and decreased level of PLAU protein with increasing p75^{NTR} protein expression. Indeed, CRABPI was elevated far more than any other genes (Nalbandian et al., 2005). In this context, the retinoids, ATRA and 9-*cis* RA, that bind CRABPI, promoted functional cell differentiation in p75^{NTR} PC-3 cells, but not in *neo* control PC-3 cells. Subsequent examination of the retinoic acid receptors expression levels demonstrated an absence of RAR- β in the *neo* control cells and re-expression in the p75^{NTR} expressing cells, consistent with previous findings where RAR- β is believed to play a critical role as a tumor suppressor gene which is lost during de-differentiation of prostate epithelial cells. Whereas the RAR- α and - γ protein levels remained unchanged, RXR- α and - β also exhibited increasing protein levels with re-expression of the p75^{NTR} protein (Nalbandian et al., 2005). Moreover, the ability of p75^{NTR} siRNA to knockdown levels of RAR- β , RXR- α , and

RXR- β support the specificity of the functional involvement of p75^{NTR} in differentiation. Hence, re-expression of the p75^{NTR} appears to partially reverse de-differentiation of prostate cancer cells by up-regulating expression of CRABPI for localized sequestration of retinoids that are available to newly up-regulated RAR- β , RXR- α , and RXR- β proteins (Nalbandian et al., 2005). Hence, the p75^{NTR} has been shown to exhibit tumor suppressor (Krygiel & Djakiew, 2001a), metastasis suppressor (Krygiel & Djakiew, 2002) and differentiation (Nalbandian et al., 2005) functions all of which contribute to an anti-cancer phenotype in the prostate.

4. Effectors of signal transduction, cell cycle and apoptosis following constitutive p75^{NTR} expression in the prostate

Ectopic re-expression of p75^{NTR} down regulates the NF κ B and JNK pathways leading to reduced survival of tumor cells (Allen et al., 2005). As a member of the TNF receptor super-family the p75^{NTR} has been shown to mediate signal transduction through its intracellular death domain. Expression of two adaptor proteins, TRAF2 and RIP are down-regulated following restoration of p75^{NTR} protein by stable transfection of prostate tumor cells (Allen et al., 2005). Significantly, TRAF2 has previously been implicated as an upstream signaling molecule of both the NF κ B and JNK pathways (MacEwan, 2002), both of which function as potent effectors of transcription. Similarly, RIP has also been shown to interact with death receptors that signal through both the NF κ B and JNK pathways (Harper et al, 2003). Moreover, deletion constructs that lack much of the intracellular death domain were shown to rescue p75^{NTR} down-regulation of RIP (Allen et al., 2005). This p75^{NTR}-dependent reduction in RIP protein appeared to be a consequence of caspase-8 cleavage of RIP, since a caspase-8 inhibitor rescued the p75^{NTR}-dependent reduction in the levels of RIP. This observation is consistent with previous reports where caspase-8 cleavage of RIP was shown to prevent activation of the NF κ B pathway (Lin et al., 1999; Martinon et al., 2000) and suggests that caspase-8 cleavage of RIP is capable of preventing the previously reported RIP activation of the JNK pathway (Kelliher et al., 1998). Hence, both the NF κ B and JNK pathways appear to be regulated by p75^{NTR} expression (Allen et al., 2005). With regard to the NF κ B signaling bifurcation, ectopic re-expression of p75^{NTR} reduces levels of IKKs, with consequent reduced phosphorylation levels of I κ B α , stabilized levels of unphosphorylated I κ B α , and reduced levels of RelA (Allen et al., 2005). Moreover, the activity of the IKKs is regulated through the serine/threonine kinase activity of upstream full length RIP (Hur et al., 2003). Hence, the p75^{NTR}-dependent reduction in the levels of RIP, via caspase-8 cleavage of RIP, leads to an overall reduction in signal transduction components required for activation of NF κ B (Allen et al., 2005). Several of these p75^{NTR} dependent changes in signaling components are rescued with dominant negative death domain deletion constructs, thereby further establishing a link between the NF κ B pathway downstream of p75^{NTR} (Allen et al., 2005). With regard to the JNK signaling bifurcation, ectopic restoration of p75^{NTR} levels is associated with a reduction in the expression of the MKK4 kinase and a reduction in the phosphorylated (active) form of JNK. Since MKK4 phosphorylation of JNK has been shown to promote nuclear translocation (Cobb, 1999; Gonzalez et al., 2000) where activated JNK may function as an effector of transcription for growth (Gee et al., 2000; Herr & Debatin, 2001), p75^{NTR} associated suppression of JNK phosphorylation and suppression of translocation is consistent with the suppression of the JNK signaling pathway. Moreover,

deletion constructs that lack much of the intracellular death domain were shown to rescue p75^{NTR} down-regulation of MKK4 and phosphorylated JNK (Allen et al., 2005) thereby further establishing a link between the JNK pathway downstream of p75^{NTR} (Allen et al., 2005). At the functional level, ectopic re-expression of p75^{NTR} reduces cell survival (Allen et al., 2005), and dominant negative antagonism of IKK β or MKK4 partially rescues survival (Allen et al., 2005). Hence, it appears that both the NF κ B and JNK pathways promote cell survival and that p75^{NTR} down regulation of these pathways inhibits survival in prostate cancer cells, thereby providing a signal transduction pathway for the observed suppressor activities of the p75^{NTR} protein (Krygiel and Djakiew, 2001a; Krygiel and Djakiew, 2002). The observation that both signaling pathways bifurcate from the p75^{NTR} suggests a redundancy whereby robust down regulation of either pathway may function independently to promote suppressor activity.

Ectopic re-expression of p75^{NTR} alters the cell cycle kinetics of prostate tumor cells (Krygiel & Djakiew, 2001a). Cell cycle initiation and progression is cooperatively regulated by several classes of cyclin-dependent kinases (cdks). The expression of cyclin D1-cdk6 (Boonstra, 2003; Sherr, & Roberts, 1995) complexed with PCNA has been shown to promote hyperphosphorylation of Rb during progression through early to mid-G1 (Satyanarayana & Rudolph, 2004). Conversely, p16^{INK4a} binds and induces an allosteric conformational change in cdk6 that inhibits the binding of ATP thereby disrupting the formation of the cdk6-cyclin D1 complex (Satyanarayana & Rudolph, 2004; Golsteyn, 2005). Prostate cancer cells that express rank-order (dose-dependent) increased levels of p75^{NTR} show suppression of cdk6, PCNA and hypophosphorylated Rb, and up-regulation of p16^{INK4a} levels indicates that p75^{NTR} selectively regulates specific components of the holoenzyme complex (Khwaja et al., 2006) associated with retarded progression through early to mid-G1 of the cell cycle (Meyer et al., 2002). The rescue of cdk6, PCNA and phosphorylated Rb levels, and conversely suppression of p16^{INK4a} levels, by both a death domain deleted dominant-negative antagonist of p75^{NTR} and by NGF ligand, show a p75^{NTR}-dependent regulation of early to mid-G1 in prostate tumor cells (Khwaja, et al., 2006). Beyond mid-G1, near the G1/S restriction point, the expression of cyclin E complexed with cdk2 has been shown to promote hyper-phosphorylation of Rb (Boonstra, 2003; Shintani et al., 2002).

A rank-order increase in p75^{NTR} suppression of cyclin E and cdk2 in prostate tumor cells (Khwaja et al., 2006) indicate that progression through the G1/S restriction point is regulated by p75^{NTR} protein expression. (Figure 3). Since the Rb protein is a major effector of cell proliferation through its ability to regulate entry into the S phase (Sellers & Kaelin, 1996; , Huang et al., 2002), and the cyclin E/cdk2 holoenzyme complex hyper-phosphorylates Rb during the G1/S transition, the observation that p75^{NTR} expression can stabilize hypophosphorylated Rb and diminish the phosphorylation of Rb is consistent with a mechanism by which p75^{NTR} retards progression through the G1/S restriction point of the cell cycle (Khwaja et al., 2006). Furthermore, hypophosphorylated Rb has been shown to bind the E2F1 transcription factor (Harbour & Dean, 2000), so that the Rb/E2F1 complex can no longer promote transcription of PCNA preventing progression into the S phase of the cell cycle. Hence, the observation that p75^{NTR} associated hypo-phosphorylation of Rb in conjunction with suppression of E2F1 and PCNA expression (Westwood et al., 2002) further supports a role of p75^{NTR} in G1/S restriction point cyto-stasis (Khwaja et al., 2006). Beyond the G1/S restriction point, expression of the cyclin A/cdk2 holoenzyme complex has been shown to maintain hyper-phosphorylation of Rb during the S phase of the cell cycle (Sherr, 1996). Hence,

observations that p75^{NTR}-potentiated suppression of cdk2 and cyclin A support a selective effect of p75^{NTR} expression on maintaining hypo-phosphorylation of Rb (Khwaja et al., 2006), thereby retarding progression through the S phase of the cell cycle (Figure 3). Significantly, the rescue of cyclin E, cdk2, PCNA, phosphorylated Rb, E2F1 and PCNA by both a death domain deleted dominant-negative antagonist of p75^{NTR} and also by NGF ligand (Khwaja et al., 2006) show a p75^{NTR}-dependence of the regulation of progression through both the G1/S restriction point and the S phase in prostate tumor cells. Hence, it seems clear that p75^{NTR} expression selectively alters specific cell cycle regulatory molecules that retard progression through early to mid-G1, the G1/S restriction point and the S phase of the cell cycle (Figure 3). Moreover, in the absence of ligand, the ability of p75^{NTR} to retard cell cycle progression is dependent on the intracellular death domain, and that addition of NGF ligand attenuates inhibition of cell cycle progression at the level of the cyclin/cdk holoenzyme complex and related effects on Rb expression and tumor cell proliferation. Hence, p75^{NTR}-dependent regulation of specific cyclin-dependent kinases and associated cell cycle effectors to retard progression of prostate tumors through the G1 phase and entry into S phase of the cell cycle (Figure 3) provides a biochemical basis upon which p75^{NTR} inhibits prostate cell proliferation (Khwaja et al., 2006).




Fig. 3. Effects of p75^{NTR} on regulation of the cell cycle. Re-expression of p75^{NTR} induces midG1 suppression of cdk6 and up-regulation of p16^{Ink4a}. During the G1/S transition re-expression of p75^{NTR} induces suppression of cdk2 and cyclin E. Similarly, during S phase re-expression of p75^{NTR} induces suppression of cdk2 and cyclin A. The net effect of these changes is to diminish phosphorylation of Rb thereby preventing progression of the cell cycle in favor of cessation.

In addition to cell cycle effects, ectopic re-expression of p75^{NTR} has been shown to modify mitochondrial effector proteins that activate the caspase cascade leading to apoptosis (Khwaja, et al., 2006). Mitochondrial mediated apoptosis is facilitated by members of the Bcl-2 homology (Puthalakath & Strasser, 2002) family of proteins (O'Neill et al., 2004) that

include both pro- and anti-apoptotic members (Puthalakath & Strasser, 2002). In this context, a rank-order increase in p75^{NTR} levels (Khwaja et al., 2006) was associated with a decrease in the mitochondrial pro-survival effector, Bcl-xL, and concomitantly, an increase in proapoptotic effectors (Smac, Bax, Bak, Bad). In the absence of NGF ligand the ability of a death domain deleted dominant-negative antagonist of p75^{NTR} to rescue p75^{NTR}-dependent changes in Smac and BH family proteins, and the addition of NGF having a comparable effect, suggest a ligand independent p75^{NTR}-potentiated apoptosis in prostate cells that occurs via a mitochondrial stress pathway (Khwaja et al., 2006). Release of Smac from the intermembranous space of the mitochondria into the cytosol (Shiozaki & Shi, 2004) also initiates apoptosis where it competes with caspases for binding to XIAP. Indeed, to protect against inadvertent activation of apoptosis, the IAP family of proteins (Shiozaki & Shi, 2004) have been shown to bind and prevent activation of caspases. Hence, p75^{NTR}-dependent down-regulation of XIAP in prostate cells allows for activation of the caspase cascade. Significantly, a rank-order increase in p75^{NTR} levels was shown to activate both caspase-9 and caspase-7 (Khwaja et al., 2006). In the intrinsic mitochondrial dependent pathway, the assembly of the apoptosome requires the recruitment of the initiator caspase-9 (Shiozaki & Shi, 2004) which subsequently activates the effector caspase-7. Interestingly, a specific peptide inhibitor of procaspase-9 cleavage was shown to prevent cleavage/activation of procaspase-7. Since the initiator procaspase-9 is a proximate component of the caspase cascade, it is clear that procaspase-7 is downstream from procaspase-9 during mitochondrial mediated p75^{NTR} potentiation of apoptosis (Shiozaki & Shi, 2004). Moreover, in p75^{NTR} expressing prostate cells the activation of caspase-7 appears to facilitate subsequent cleavage of PARP. PARP cleavage has been shown to be a key event in the execution phase of apoptosis leading to cellular demise (Lazebnik et al., 1994). Clearly, the ability of a death domain deleted dominant-negative antagonist of p75^{NTR} and NGF ligand to rescue p75^{NTR} potentiated changes in XIAP, caspase-9, caspase-7, PARP and apoptotic nuclear fragmentation show that mitochondrial mediated apoptosis is dependent, in part, on p75^{NTR} in the prostate. Moreover, the effect of p75^{NTR}-dependent down regulation of NFκB and JNK survival pathways, regulation of cell cycle regulatory molecules of the cyclin/cdk holoenzyme complex that produce stasis in G1 and/or inhibition of progression to S phase of the cell cycle and regulation of mitochondrial effector proteins that activate the caspase cascade leading to apoptosis, in combination, all inhibit growth of prostate cells as a function of its tumor suppressor activity (Krygier & Djakiew, 2001a; Khwaja et al., 2006).

5. Constitutive p75^{NTR} expression regulates protease activity in the prostate

Ectopic re-expression of p75^{NTR} down regulates protease activities in prostate cancer cells (Nalbandian & Djakiew, 2006). The urokinase plasminogen activator (uPA) and its receptor (uPAR) are associated with tumor malignancy through an extracellular cascade of proteolysis (Andreassen et al., 2000; Reuning et al., 1998) including activation of the type IV collagen matrix metalloproteinases (MMP-2 and MMP-9) during invasion and tumor progression (Mazzier et al., 1997). Ectopic re-expression of p75^{NTR} reduces enzymatic protein levels and activity of uPA, MMP-2 and MMP-9 in prostate tumor cells (Nalbandian & Djakiew, 2006). Conversely, expression of an MMP-9 antagonist, tissue inhibitor of matrix metalloproteinase-1 (TIMP-1) exhibits an increase in protein levels with an increase in p75^{NTR} levels. Whereas, levels of TIMP-2 were not detectable (Nalbandian & Djakiew, 2006) transient transfection with an inducible death domain deleted dominant-negative antagonist of p75^{NTR} rescued uPA, MMP-

2, and MMP-9 protein levels and protease activities, and conversely suppressed TIMP-1 levels (Nalbandian & Djakiew, 2006). Since ectopic p75^{NTR} signal transduction has been shown to suppress the NF κ B and JNK pathways (Allen et al., 2005), antagonism of signaling intermediaries in these pathways, using dominant negative IKK β or dominant negative MKK-4, respectively, was shown to further decrease expression of uPA, MMP-2, and MMP-9 protein and enzymatic activity levels, and conversely up-regulate levels of TIMP-1. Hence, expression of uPA, MMP-2, MMP-9, and TIMP-1 are directly regulated by expression of p75^{NTR} and its downstream signal transduction cascade. This suggests that the metastasis suppressor activity of p75^{NTR} (Krygier & Djakiew, 2002) is mediated, in part, by down-regulation of specific proteases (uPA, type IV collagenases) implicated in cell migration and metastasis (Nalbandian & Djakiew, 2006).

6. NSAIDs selectively induce pharmacological re-expression of the p75^{NTR}

NSAIDs represent a diverse and often structurally unrelated group of compounds that exhibit a range of anti-inflammatory, anti-pyretic and analgesic activities. NSAIDs can be classified into three broad categories as carboxylic acids, enolic acids, and COX-2 inhibitors, or coxibs (Figure 4). The carboxylic acid group is further divided into salicylic acids and esters, acetic acids, and propionic acids. The enolic acids are divided into pyrazolones and oxicams. Many NSAIDs used to treat inflammation inhibit cyclooxygenase (COX) activity. Two well known isoforms of COX exist, COX-1 and COX-2. COX-1 is considered to be a house keeping gene, and is expressed constitutively and ubiquitously at low levels. COX-2 is highly inducible in response to cytokines, hormones, and growth factors. The COX enzymes catalyze the conversion of arachidonic acid to various prostaglandins, which play a role in biological processes including immune response, blood pressure regulation, angiogenesis, ovulation, pain and inflammatory responses. Nonselective NSAIDs inhibit both COX-1 and COX-2, and are frequently associated with gastrointestinal side effects. Whereas, NSAIDs that selectively inhibit COX-2 have significantly decreased gastrointestinal side effects, they often exhibit enhanced cardiovascular toxicity (Dubois et al., 2004). Overexpression of COX-2 is observed in several cancer types including colon, breast, pancreas, and lung (Sarkar et al., 2007; Mascaux et al., 2006). Increased COX-2 expression is believed to contribute to tumorigenesis through several mechanisms including stimulation of growth, promotion of angiogenesis, increased inflammation, increased invasion and migration, immune suppression, and inhibition of apoptosis (Liao et al., 2007). Hence, NSAID inhibition of COX-2 activity may suppress these mechanisms that contribute to tumorigenesis. Consistently, several reports have linked long term NSAID use to decreased cancer risk for colon (Thun et al., 1991), bladder (Castelao et al., 2000), and prostate (Nelson & Harris, 2000) and possibly other organ specific sites. Although it has been the subject of multiple reports, there is not a consensus concerning the expression and role of COX-2 in prostate cancer. Several studies show that COX-2 is overexpressed, while others found that it is low or absent in most prostate cancers (Madaan et al., 2000; Kirschenbaum et al., 2000; Yoshimura et al., 2000). In addition, expression of COX-2 is absent in the metastatic human prostate cancer cell lines LNCaP, DU145, and PC-3 (Yoshimura et al., 2000). Although reports of COX-2 expression in prostate cancer vary, long term NSAID use is associated with decreased prostate cancer risk (Nelson & Harris, 2000), and several COX

inhibitors consistently induce apoptosis in prostate cancer cells regardless of COX-2 expression (Zha et al., 2001; Jacobs et al., 2007; Roberts et al., 2002; Hsu et al., 2000; Johnson et al., 2001). Indeed, NSAIDs that do not inhibit the cyclo-oxygenases (R-flurbiprofen) or NSAIDs that inhibit COX null tumor cells demonstrates an alternate COX independent mechanism of action for selected NSAID inhibition of some tumor cells. In this context, NSAIDs exhibit selective activity to induce p75^{NTR}-dependent cell death (Khwaja et al., 2004). The propionic acids, ibuprofen and r-flurbiprofen, as well as the indolacetic acid, indomethacin, exhibit greater pharmacological activity to induce p75^{NTR}-dependent cell death than the salicylate, aspirin, or acetaminophen (Khwaja et al., 2004). Within the aryl propionic acid NSAIDs, r-flurbiprofen exhibits greater activity to induce p75^{NTR} in descending rank-order than ibuprofen, oxaprozin, fenoprofen, naproxen and least of all ketoprofen (Quann et al., 2007a). Significantly, this activity appears relatively selective for induction of p75^{NTR} (Quann et al., 2007a), since r-flurbiprofen and ibuprofen do not induce expression of Fas, p55^{TNFR}, DR3, DR4, DR5 or DR6 in prostate cancer cells (Quann et al., 2007a). Some other NSAIDs have also been reported to induce expression of certain TNF receptor super family members, particularly DR5.




Fig. 4. Classification of the Non-Steroidal Anti-Inflammatory Drugs (NSAIDs) into three categories of carboxylic acids, enolic acids and COX-2 inhibitors. Sub-classifications of NSAIDs are further delineated with specific examples.

For instance, sulindac sulfide induces DR5 associated cell death in prostate cancer cells (Huang et al., 2001) and celcoxibs induce DR5 in both prostate and colon cancer cells (He et al., 2008). Moreover, aryl propionic acid induction of p75^{NTR} appears somewhat selective for urogenital cancer cell lines such as prostate, bladder, kidney, ovary, and colon (Khwaja et al., 2004), whereas cell lines derived from the lung, or breast do not exhibit induction of the p75^{NTR} protein (Khwaja et al., 2004). Hence, induction of p75^{NTR} appears to represent a COX independent mechanism for NSAID associated reduced risk of prostate cancer.

7. NSAID induced phosphorylation of p38 MAPK is necessary for induction of p75^{NTR} expression

NSAIDs exhibit a range of efficacy to inhibit survival of prostate cancer cells (Andrews et al., 2002). Induction of p75^{NTR} dependent cell death by selected NSAIDs (R-flurbiprofen, ibuprofen) occurs through hyper-phosphorylation of p38 MAPK (Quann et al., 2007b). Although p75^{NTR} is transcribed at a high level (Quann et al., 2007b; Krygier & Djakiew, 2001b), prostate cancer cells have very little p75^{NTR} mRNA or protein due to increased mRNA instability mediated through the 3'UTR (Krygier & Djakiew, 2001b). Since, increased 3'UTR length provides increased potential for post-transcriptional regulation through the 3'UTR (Mazumder et al., 2003) R-flurbiprofen and ibuprofen appear to modulate mRNA stability as a determinant in regulating the expression level of p75^{NTR} protein (Quann et al., 2007b). Significantly, activity of the p38 MAPK pathway is an important regulator of mRNA stability (Zarubin & Han, 2005; Kennedy et al., 2007; Gaestel, 2006). Pretreatment of prostate cancer cells with a p38 MAPK selective pharmacological inhibitor or siRNA knockdown of p38 MAPK prior to R-flurbiprofen and ibuprofen treatment prevents induction of p75^{NTR} protein. In addition, the phytoestrogens biochanin A and to a lesser extent genestein, inhibit ibuprofen induced phosphorylation of p38 MAPK which in turn suppresses p75^{NTR} expression and increases cell survival (El Touny et al., 2010). R-flurbiprofen and ibuprofen cause increased p38 MAPK phosphorylation within 5 minutes of treatment (Quann et al., 2007b). Whereas, p38 MAPK can be phosphorylated by MKK6 and MKK3, R-flurbiprofen and ibuprofen induce hyperphosphorylation of only MKK6 but not MKK3 within 30 seconds of treatment (Figure 5) indicating that the target molecule of these NSAIDs is immediately proximal to MKK6. Significantly, several oxidative stress pathways converge upstream of MKK6, thereby providing a basis by which NSAID inhibition of inflammation (Masferrer et al., 1995; Tegeder et al., 2001) may regulate p38 MAPK dependent expression of the p75^{NTR} tumor suppressor to reduce the incidence of prostate cancer. Downstream of p38 MAPK R-flurbiprofen and ibuprofen induced activation of the kinase MK2 (Quann et al., 2007b). MK2 and the closely related MK3 are known to be responsible for mediating the mRNA stabilizing effects of the p38 MAPK pathway (Ronkina et al., 2007). Moreover, siRNA knockdown of both MK2 and MK3 together prevents induction of p75^{NTR} by R-flurbiprofen or ibuprofen to a greater extent than knockdown of either MK2 or MK3 separately, indicating that p38 MAPK is able to induce p75^{NTR} by acting through both MK2 and MK3 (Quann et al., 2007b).




Fig. 5. Western blot showing the time course in seconds (s), minutes (m) and hours (h) of the phosphorylation forms of MKK6, but not MKK3, in PC-3 prostate cancer cells treated with r-flurbiprofen (Flu), or ibuprofen (Ibu). The positive controls for the phosphorylated forms of both MKK6 and MKK3 are shown to the right.

The mRNA stabilizing effects of the p38 MAPK pathway (Ronkina et al., 2007) also involves the RNA binding protein HuR (Tran et al., 2003; Lin et al., 2006; Jin et al., 2007; Song et al., 2005). Its ability to stabilize target mRNAs is linked to its subcellular localization, and activation of p38 MAPK and MK2 have been shown to cause translocation of HuR from the nucleus to the cytoplasm, resulting in increased mRNA stability of a number of p38 MAPK regulated genes (Tran et al., 2003; Lin et al., 2006; Jin et al., 2007; Song et al., 2005). HuR has repeatedly been shown to stabilize transcripts containing the AUUUA sequence (Tran et al., 2003; Song et al., 2005). The human p75^{NTR} transcript contains AUUUA sites located in the 3'UTR at positions 2946 and 3124, suggesting they may be involved in regulating p75^{NTR} expression. Significantly, treatment with R-flurbiprofen or ibuprofen results in an increase in the cytoplasmic level of HuR and binding to the p75^{NTR} transcript (Quann et al., 2007b) although this is not the sole mechanism responsible for increased p75^{NTR} expression. Indeed, eIF4E is also phosphorylated by kinases downstream of p38 MAPK (Scheper & Proud, 2002). eIF4E has also been shown to control the nuclear export as well as translation of a subset of transcripts (Culjkovic et al., 2007). eIF4E has been linked to the control of mRNA stability by removal of the 5'-cap during mRNA degradation (von der Haar et al., 2004). R-flurbiprofen or ibuprofen treatment increases the level of phosphorylated eIF4E (Quann et al., 2007b) involved in translation initiation and this appears to occur at least partially through the p38 MAPK pathway since the increase in phosphorylation is substantially inhibited in the presence of p38 MAPK siRNA. Therefore, modulation of eIF4E activity is another mechanism by which the p38 MAPK pathway may control post-transcriptional events in response to R-flurbiprofen or ibuprofen treatment (Figure 6). R-flurbiprofen or ibuprofen stabilization of the p75^{NTR} transcript allows levels of the p75^{NTR} protein to accumulate (Quann et al., 1997a, Quann et al., 197b) that then exhibit biological activity to inhibit prostate cancer cell survival and induce apoptosis (Quann et al., 1997a) consistent with its tumor suppressor activity (Krygier & Djakiew, 2001a). Significantly, NSAID induction of p38 MAPK dependent p75^{NTR} expression and tumor suppressor activity may represent a new pathway to medicinal drug design. Rather than inhibit kinase dependent proliferation pathways, pharmacological induction of the p75^{NTR} tumor suppressor, as shown with NSAIDs (Figure 6), may represent an alternate approach to the targeting of cancer cell survival pathways. Presumably, target molecules that activate this pathway appear to occur immediately proximal to MKK6 and the p38 MAPK (Figure 5).

8. NSAIDs induce expression of NAG-1 down stream of p75^{NTR} via the p38 MAPK pathway

R-flurbiprofen and ibuprofen treatment of prostate cancer cells induces expression of the NSAID activated gene-1 (Nag-1) protein, a divergent member of the transforming growth factor beta (TGF- β) family (Wynne & Djakiew, 2010). Moreover, a selective pharmacological inhibitor of p38 MAPK and p38 MAPK specific siRNA, both reduce Nag-1 induction following NSAID treatment. Hence, NSAID induced Nag-1 expression is regulated by the p38 MAPK pathway (Figure 6). Interestingly, p75^{NTR} specific siRNA pretreatment abrogates Nag-1 induction by NSAIDs (Wynne & Djakiew, 2010) thereby demonstrating that Nag-1 is downstream of p75^{NTR} induction (Figure 6). Functionally, decreased survival of NSAID treated cells is rescued by p75^{NTR} specific siRNA (Quann et al., 2007a; Wynne & Djakiew, 2010) but not by Nag-1 siRNA. Transwell chamber and *in vitro* wound healing assays demonstrate decreased cell migration upon NSAID treatment (Wynne & Djakiew, 2010). Pre-treatment of prostate cancer cells with p75^{NTR} and Nag-1 specific siRNA shows that

NSAID inhibition of cell migration is mediated by Nag-1 and p75^{NTR} (Wynne & Djakiew, 2010). Additionally, prostate cancer cells stably expressing Nag-1 exhibit decreased migration relative to the parental cell line (Wynne & Djakiew, 2010), thereby independently confirming a role for this protein in reduced prostate cancer cell migration. Hence, it appears that NSAID induction of Nag-1 functions in the inhibition of cell migration, but not survival (Figure 6). Interestingly, NSAIDs have been linked to metastasis suppression in a variety of cancers including prostate cancer (Lloyd et al., 2003; Jin et al., 2010; Kamei et al., 2009). Hence, NSAID induction of Nag-1 via the p38 MAPK pathway may contribute to the metastasis suppressor activity of p75^{NTR} (Krygiel & Djakiew, 2002).




Fig. 6. Schematic diagram showing the biochemical pathway by which profen NSAIDs positively stimulate and the phytoestrogen, biochanin A, negatively inhibits phosphorylation of the p38 MAPK pathway leading to p75^{NTR} expression with consequent tumor suppressor and metastasis suppressor activity.

9. Summary and conclusions

Paracrine regulated growth of the prostate is mediated, in part, by neurotrophin dependent interactions with the Trk family of receptors and the p75^{NTR} (Djakiew, 2000). In the normal prostate neurotrophins stimulate proliferation via the Trk family of receptors and abrogate p75^{NTR} apoptotic activity. However, p75^{NTR} protein expression progressively declines in human prostate cancer (Perez et al., 1997; Pflug et al., 1995; Djakiew et al., 1996). Hence, in the absence of p75^{NTR} inhibition of growth, prostate tumor cells respond to neurotrophins with a stoichiometry that favors Trk family dependent proliferation. Even though the p75^{NTR} protein is no longer expressed in many prostate cancer cells the gene has remained intact (Krygiel & Djakiew, 2001b). Indeed, reduced p75^{NTR} protein in cancer cells occurs via loss of mRNA stability (Krygiel & Djakiew, 2001b). Ectopic re-expression of p75^{NTR} in cancer cells induces tumor suppressor (Krygiel & Djakiew, 2001a), metastasis suppressor (Krygiel & Djakiew, 2002) and differentiation phenotypes (Nalbandian et al., 2005). The tumor suppressor activity of p75^{NTR} is manifest as a suppression of the NFκB and JNK pathways, modification of cyclin-dependent holoenzyme complexes resulting in accumulations of cells in G1 and restriction of entry into the S1 phase of the cell cycle, and modification of mitochondrial effector proteins

that activate the caspase cascade leading to apoptosis. The metastasis suppressor activity of p75^{NTR} is manifest as down-regulation of urokinase plasminogen activator and type IV collagenases (MMP-2, MMP-9) as well as increased expression of the tissue inhibitor of matrix metalloproteinase-1 (Nalbandian & Djakiew, 2006). The differentiation activity of p75^{NTR} is manifest as up-regulated expression of CRABPI for localized sequestration of retinoids that are available to newly up-regulated RAR- β , RXR- α , and RXR- β proteins (Nalbandian et al., 2005). Moreover, certain aryl propionic acid type NSAIDs (profens) restore p75^{NTR} protein levels that inhibit growth via a COX independent mechanism of action. Indeed, these profens induce hyperphosphorylation of p38 MAPK that initiates a signal transduction cascade leading to mRNA stability of the p75^{NTR} transcript and increased protein levels (Quann et al., 2007a; Quann et al., 2007b). Profen NSAIDs also induce Nag-1 expression downstream of p75^{NTR} (Wynne & Djakiew, 2010). Functionally, profen dependent induction of p75^{NTR} inhibits growth and downstream Nag-1 inhibition of cell migration consistent with the tumor suppressor (Krygier & Djakiew, 2001a) and metastasis suppressor (Krygier & Djakiew, 2002) activities of p75^{NTR} expression, respectively. Hence, these profen NSAIDs induce multiple activities of the p75^{NTR} and downstream Nag-1 consistent with observations that chronic consumption of NSAIDs is associated with a reduced incidence of prostate cancer (Nelson & Harris, 2000).

10. Acknowledgements

The author thanks his former graduate students (Drs. Robert Delsite, Beth Pflug, Scott Krygier, Jeff Allen, Angele Nalbandian, Emily Quann, Shehla Wynne) and research associates (Drs. Rhaki Dalal, Ashia Tabassum, Fatima Khwaja, Lara El Touny) for their dedication and tireless efforts that contributed to much of this work. A significant amount of this research was supported by the National Institutes of Health (DK52626). Research support was also provided by the Department of Defense (PC060409) to graduate students (EQ, SW).

11. References

- Allen, J., Khwaja, F., Byers, S. & Djakiew, D. (2005). The p75^{NTR} Mediates a Bifurcated Signal Transduction Cascade through the NF κ B and JNK Pathways to Inhibit Cell Survival. *Experimental Cell Research*, Vol.304, No.1, (March 2005), pp69-80, PMID 15707575
- Allen, J., Khwaja, F. & Djakiew, D. (2004). Gene therapy of prostate xenograft tumors with a p75^{NTR} lipoplex. *Anticancer Research*, Vol.24, No.5A, (September-October 2004), pp2997-3003, PMID 15517907
- Andreasen, P., Egelund, R. & Petersen, H. (2000). The plasminogen activation system in tumor growth, invasion, and metastasis. *Cellular and Molecular Life Sciences*, Vol.57, No.1, (January 2000), pp 25-40, PMID 10949579
- Andrews, J., Djakiew, D., Krygier, S. & Andrews, P. (2002). Superior effectiveness of ibuprofen compared with other NSAIDs for reducing the survival of human prostate cancer cells. *Cancer Chemotherapy and Pharmacology*, Vol. 50, No.4, (October 2002), pp 277-284, PMID 12357301
- Angelsen, A., Sandvik, A., Syversen, U., Stridsberg, M. & Waldum, H. (1998). NGF-beta, NE-cells and prostatic cancer cell lines. A study of neuroendocrine expression in the human prostatic cancer cell lines DU-145, PC-3, LNCaP, and TSU-pr1 following stimulation of the nerve growth factor-beta. *Scandinavian Journal of Urology and Nephrology*, Vol.32, No.1 (February 1998), pp 7-13, PMID 9561567

- Berkemeier, L., Winslow, J., Kaplan, D., Nikolics, K., Goeddel, D. & Rosenthal, A. (1991). Neurotrophin-5: a novel neurotrophic factor that activates trk and trkB. *Neuron*, Vol.7, No.5, (November 1991), pp 857-866, PMID 1742028
- Berg M.M., Sternberg D.W., Parada L.F. & Chao M.V. (1992). K-252a inhibits nerve growth factor-induced trk proto-oncogene tyrosine phosphorylation and kinase activity. *Journal of Biological Chemistry*, Vol.267, No.1 (January 1992), pp13-16, PMID 1730579
- Boonstra, J. (2003). Progression through the G1-phase of the on-going cell cycle. *Journal of Cellular Biochemistry*, Vol.90, No.2, (October 2003), pp 244-252, PMID 14505341
- Bothwell, M. (1995). Functional interactions of neurotrophins and neurotrophin receptors. *Annual Reviews of Neuroscience*, Vol.18, (March 1995), pp 223-253, PMID 7605062
- Bostwick, D., Burke, H., Djakiew, D., Euling, S., Ho, S., Landolph, J., Morrison, H., Sonawane, B., Shifflett, T., Waters, D., Timms, B. (2004). Human prostate cancer risk factors. *Cancer*, Vol.101, Supplement 10, (November 2004), pp 2371-490, PMID 15495199
- Burrows, C. (2009). Surviving an Oxygen Atmosphere: DNA Damage and Repair. *ACS symposium Series American Chemical Society* (December 2009); pp.147-156, PMID 20458355
- Castelao, J., Yuan, M., Gago-Dominguez, M., Yu, M. & Ross, R. (2000). Non-steroidal anti-inflammatory drugs and bladder cancer prevention. *The British Journal of Cancer*, Vol.82, No.7, (April 2000), pp 1364-1369, PMID 10755416
- Chao, M. (1994). The p75 neurotrophin receptor. *Journal of Neurobiology*, Vol.25, No.11 (November 1994), pp1373-1385, PMID 7852992
- Chapman, B. (1995). A region of the 75kD neurotrophin receptor homologous to the death domains of TNFR-1 and Fas. *Federation of Experimental Biology Society Letters* Vol.374, No. (October 1995), pp 216-220, PMID 7589538
- Chung, L., Li, W., Gleave, M., Hsieh, J., Wu, H., Sikes, R., Zhau, H., Bandyk, M., Logothetis, J., Rubin, J. & von Eschenbach, A. (1992). Human prostate cancer model: roles of growth factors and extracellular matrices. *Journal of Cellular Biochemistry*, Supplement 16H (1992), pp 99-105, PMID 1289680
- Cobb, M. (1999). MAP kinase pathways. *Progress in Biophysics and Molecular Biology*, Vol.71, No.3-4, (April 1999), pp 479-500, PMID 10354710
- Culjkovic, B., Topisirovic, I. & Borden, K. (2007). Controlling gene expression through RNA regulons: the role of the eukaryotic translation initiation factor eIF4E. *Cell Cycle*, Vol.6, No.1, (January 2007), pp 65-69, PMID 17245113
- Dalal, R. & Djakiew, D. (1997). Molecular characterization of neurotrophin expression and the corresponding tropomyosin receptor kinases (trks) in epithelial and stromal cells of the human prostate. *Molecular and Cellular Endocrinology*, Vol.134, No.1 (October 1997), pp 15-22, PMID 9406845
- Delsite R. & Djakiew D. (1996). Anti-proliferative effect of the kinase inhibitor K252a on human prostatic carcinoma cell lines. *Journal of Andrology*, Vol.17, No.5 (September-October 1996), pp481-490, PMID 8957691
- Delsite R. & Djakiew D. (1999). Characterization of nerve growth factor precursor protein expression by human prostate stromal cells: a role in selective neurotrophin stimulation of prostate epithelial cell growth. *Prostate*, Vol.41, No.1 (September 1999), pp 39-48, PMID 10440874
- Dionne, C., Camoratto, A., Jani, J., Emerson, E., Neff, N., Vaught, J., Murakata, C., Djakiew, D., Lamb, J., Bova, S., George, D. & Isaacs, J. (1998). *Clinical Cancer Research*, Vol.4, No.8 (August 1998), pp1887-1898, PMID 9717816

- Djakiew, D. (2000). Neurotrophins, p75^{NTR} and TRK in Prostate Cancer, In: *Neurobiology of the Neurotrophins*, I. Mocchetti, (Ed.), 525-539, F. P. Graham Publishing Co., Johnson City, Tennessee, USA, ISBN 1-929675-01-1
- Djakiew, D., Delsite, R., Pflug, B., Wrathall, J., Lynch, J. & Onoda M. (1991). Regulation of growth by a nerve growth factor-like protein which modulates paracrine interactions between a neoplastic epithelial cell line and stromal cells of the human prostate. *Cancer Research*, Vol.51, No.12, (June 1991), pp 3304-3310, PMID 1710170
- Djakiew, D., Delsite, R., Dalal, R. & Pflug, B. (1996). Role of the low-affinity nerve growth factor receptor and the high-affinity Trk nerve growth factor receptor in human prostate carcinogenesis. *Radiation Oncology Investigations*, Vol.3, (1996), 333-339.
- Djakiew, D., & Pflug, B., Delsite, R., Onoda, M., Lynch, J., Arand, G. & Thompson, G. (1993). Chemotaxis and chemokinesis of human prostate tumor cell lines in response to human prostate stromal cell secretory proteins containing a nerve growth factor-like protein. *Cancer Research*, Vol.53, No.6, (March 1993), pp 1416-20, PMID 8443820
- Dubois, R., Melmed, G., Henning, J. & Laine, L. (2004). Guidelines for the appropriate use of non-steroidal anti-inflammatory drugs, cyclo-oxygenase-2-specific inhibitors and proton pump inhibitors in patients requiring chronic anti-inflammatory therapy. *Alimentary Pharmacology and Therapeutics*, Vol.19, No.2, (January 2004), pp 197-208, PMID 14723611
- El Touny, L., Henderson, F. & Djakiew, D. (2010). Biochanin A Reduces Drug-Induced p75^{NTR} expression and Enhances Cell Survival: a New *in Vitro* Assay for Screening Inhibitors of p75^{NTR} Expression. *Rejuvenation Research*, Vol.13, No.5, (October 2010), pp 527-537, PMID 20818983
- Gaestel, M. (2006). MAPKAP kinases - MKs - two's company, three's a crowd. *Nature Reviews Molecular Cell Biology*, Vol.7, (February 2007), pp 120-130, PMID 16421520
- Gee, J., Barroso, A., Ellis, I., Robertson, J. & Nicholson, R. (2000). Biological and clinical associations of c-jun activation in human breast cancer. *International Journal of Cancer*, Vol.89, No.2, (March 2000), pp 177-186, PMID 10754497
- George, D., Suzuki, H., Bova, G. & Isaacs, J. (1998). Mutational analysis of the TrkA gene in prostate cancer. *Prostate*, Vol.36, No.3 (August 1998), pp172-180, PMID 9687989
- Golsteyn, R. (2005). Cdk1 and Cdk2 complexes (cyclin dependent kinases) in apoptosis: a role beyond the cell cycle *Cancer Letters*, Vol.217, No.2, (January 2005), pp. 129-138, PMID 15617830
- Gonzalez, M., Jimenez, B., Berciano, M., Gonzalez-Sancho, J., Caelles, C., Lafarga, M. & Munoz, A. (2000). Glucocorticoids antagonize AP-1 by inhibiting the Activation/phosphorylation of JNK without affecting its subcellular distribution. *Journal of Cell Biology*, Vol.150, No.5, (September 2000), pp 1199-1208, PMID 10974006
- Graham, C., Lynch, J. & Djakiew, D. (1992). Distribution of nerve growth factor-like protein and nerve growth factor receptor in human benign prostatic hyperplasia and prostatic adenocarcinoma. *Journal of Urology*, Vol.147, No.5, (May 1992), pp. 1444-1447, PMID 1373782
- Hallböök, F., Ibáñez, C. & Persson, H. (1991). Evolutionary studies of the nerve growth factor family reveal a novel member abundantly expressed in *Xenopus* ovary. *Neuron*, Vol.6, No.5, (May 1991), pp 845-858, PMID 2025430
- Harbour, J. & Dean, D. (2000). The Rb/E2F pathway: expanding roles and emerging paradigms. *Genes and Development*, Vol.14, No.19, (October 2000), pp2393-2409, PMID 11018009
- Harper, N., Hughes, M. MacFarlane, M. & Cohen, G. (2003). Fas-associated death domain protein and caspase-8 are not recruited to the tumor necrosis factor receptor 1


- signaling complex during tumor necrosis factor-induced apoptosis. *Journal of Biological Chemistry*, Vol.278, No.28, (July 2003), pp 25534-25541, PMID 12721308
- He, Q., Luo, X., Jin, W., Huang, Y., Reddy, M., Reddy, E. & Sheikh, M. (2008). Celecoxib and a novel COX-2 inhibitor ON09310 upregulate death receptor 5 expression via GADD153/CHOP. *Oncogene*, Vol.27, No.18 (April 2008), 2656-60, PMID 17968315
- Herr, I. & Debatin, K. (2001). Cellular stress response and apoptosis in cancer therapy. *Blood*, Vol.98, No.9, (November 2001), pp 2603-2614, PMID 11675328
- Hsu, A., Ching, T., Wang, D., Song, X., Rangnekar, V. & Chen, C. (2000). The cyclooxygenase-2 inhibitor celecoxib induces apoptosis by blocking Akt activation in human prostate cancer cells independently of Bcl-2. *Journal of Biological Chemistry*, Vol.275, No.15, (April 2000), pp 11397-11403, PMID 10753955
- Huang, Y., He, Q., Hillman, M., Rong, R. & Sheikh, M. (2001). Sulindac sulfide-induced apoptosis involves death receptor 5 and the caspase 8-dependent pathway in human colon and prostate cancer cells. *Cancer Research*, Vol.61, No.18, (September 2001), pp 6918-6924, PMID 11559570
- Huang, Z., Tang, X. & Cambi, F. (2002). Down-regulation of the retinoblastoma protein (rb) is associated with rat oligodendrocyte differentiation. *Molecular and Cellular Neuroscience*, Vol.19, No.2, (February 2002), pp250-262, PMID 11860277
- Hur, G., Lewis, J., Yang, Q., Lin, Y., Nakano, H., Nedospasov, S. & Liu, Z. (2003). The death domain kinase RIP has an essential role in DNA damage-induced NF-kappa B activation. *Genes and Development*, Vol.17, No.7, (April 2003), pp 873-882, PMID 12654725
- Jacobs, E., Thun, M., Bain, E., Rodríguez, C., Henley, S. & Calle, E. (2007). A large cohort study of long-term daily use of adult-strength aspirin and cancer incidence. *Journal of the National Cancer Institute*, Vol.99, No.8, (April 2007), pp 608-15, PMID 17440162
- Jin, S., Kim, T., Yang, K. & Kim, W. (2007). Thalidomide destabilizes cyclooxygenase-2 mRNA by inhibiting p38 mitogen-activated protein kinase and cytoplasmic shuttling of HuR. *European Journal of Pharmacology*, Vol.558, No.1-3, (March 2007), pp 14-20, PMID 17208222
- Jin, H., Wang, Z., Liu, L., Gao, L., Sun, L., Li, X., Zhao, H., Pan, Y., Shi, H., Liu, N., Hong, L., Liang, J., Wu, Q., Yang Z., Wu, K. & Fan, D. (2010). R-flurbiprofen reverses multidrug resistance, proliferation and metastasis in gastric cancer cells by p75(NTR) induction. *Molecular Pharmacology*, Vol.7, No.1, (February 2010), pp 156-168, PMID 19916560
- Johnson, A., Song, X. Hsu, A. & Chen, C. (2001). Apoptosis signaling pathways mediated by cyclooxygenase-2 inhibitors in prostate cancer cells. *Advances in Enzyme Regulation*, Vol.41, pp 221-235, PMID 11384747
- Kamei, S., Sakayama, K., Tamashiro, S., Aizawa, J., Miyawaki, J., Miyazaki, T., Yamamoto, H., Norimatsu, Y., Masuno, H. (2009). Ketoprofen in topical formulation decreases the matrix metalloproteinase-2 expression and pulmonary metastatic incidence in nude mice with osteosarcoma. *Journal of Orthopaedic Research*, Vol.27, No.7, (July 2009), 909-915.
- Kang, J. & Sowers, L. (2008). Examination of hypochlorous acid-induced damage to cytosine residues in a CpG dinucleotide in DNA. *Chemical Research in Toxicology*, Vol.21, No.6, (June 2008), pp 1211-1218, PMID 18826175
- Kelliher, M., Grimm, S., Ishida, Y., Kuo, F., Stanger, B. & Leder, P. (1998). The death domain kinase RIP mediates the TNF-induced NF-kappaB signal. *Immunity*, Vol.8, No.3, (March 1998), pp 297-303, PMID 9529147

- Kennedy, N., Cellurale, C. & Davis, R. (2007). A radical role for p38 MAPK in tumor initiation. *Cancer Cell*, Vol.11, No.2, (February 2007), p101-103, PMID 17292820
- Khwaja, F., Allen, J., Lynch, J., Andrews, P. & Djakiew D. (2004). Ibuprofen inhibits survival of bladder cancer cells by induced expression of the p75^{NTR} tumor suppressor protein. *Cancer Research*, Vol.64, No.17, (September 2004), pp 6207-6213, PMID 15342406
- Khwaja, F., Tabassum, A., Allen, J. & Djakiew D. (2006). The p75^{NTR} tumor suppressor induces cell cycle arrest facilitating caspase mediated apoptosis in prostate tumor cells. *Biochemical and Biophysical Research Communications*, Vol.341, No.4, (March 2006), pp 1184-1192, PMID 16460673
- Kirschenbaum, A., Klausner, A., Lee, R., Unger, P., Yao, S., Liu, X., & Levine, A. (2000). Expression of cyclooxygenase-1 and cyclooxygenase-2 in the human prostate. *Urology*, Vol.56, No.4, (October 2000), pp 671-676, PMID 11018637
- Krygiel, S. & Djakiew, D. (2001a). The Neurotrophin Receptor p75^{NTR} is a tumor suppressor in the human prostate. *Anticancer Research* Vol.21, No.6A, (November-December 2001) pp 3749-3756, PMID 11911243
- Krygiel, S. & Djakiew, D. (2001b). Molecular characterization of p75^{NTR} loss of the expression in human prostate tumor cells. *Molecular Carcinogenesis*, Vol.31, No.1, (May 2001), pp46-55, PMID 11398197
- Krygiel, S. & Djakiew, D. (2002). Neurotrophin Receptor p75^{NTR} suppresses growth and nerve growth factor-mediated metastasis of human prostate cancer cells. *International Journal of Cancer*, Vol.98, No.1, (March 2002) pp. 1-7, PMID 11857376
- Lamballe, F., Klein, R. & Barbacid, M. (1991). trkC, a new member of the trk family of tyrosine protein kinases, is a receptor for neurotrophin-3. *Cell*, Vol.66, No.5, (September 1991), pp967-979, PMID 1653651
- Lazebnik, Y., Kaufmann, S., Desnoyers, S., Poirier, G. & Earnshaw, W. (1994). Cleavage of poly(ADP-ribose) polymerase by a proteinase with properties like ICE. *Nature*, Vol.371, No.6495, (September 1994), pp. 346-347, PMID 8090205
- Leibrock, J., Lottspeich, F., Hohn, A., Hofer, M., Hengerer, B., Masiakowski, P., Thoenen, H. & Barde, Y. (1989). Molecular cloning and expression of brain-derived neurotrophic factor. *Nature*, Vol.341, No.6238, (September 1989), pp 149-152, PMID 2779653
- Liao, Z., Mason, K. & Milas, L. (2007). Cyclo-oxygenase-2 and its inhibition in cancer: is there a role? *Drugs*, Vol.67, No.6, pp 821-45, PMID 17428102
- Lin, F., Chen, Y., Lin, Y, Tsai, J., Chen, J., Wang, H., Chen, Y., Li, C. & Lin S.. (2006). The role of human antigen R, an RNA-binding protein, in mediating the stabilization of toll-like receptor 4 mRNA induced by endotoxin: a novel mechanism involved in vascular inflammation. *Arteriosclerosis, Thrombosis, and Vascular Biology*, Vol.26, No.12, (December 2006), pp 2622-2629, PMID 16990552
- Lin, Y., Devin, A., Rodriguez, Y. & Liu, Z. (1999). Cleavage of the death domain kinase RIP by caspase-8 prompts TNF-induced apoptosis. *Genes and Development*, Vol.13, No.19, (October 1999), pp 2514-2526, PMID 10521396
- Lloyd, F., Slivova, V., Valachovicova, T. & Sliva, D. (2003). Aspirin inhibits highly invasive prostate cancer cells. *International Journal of Oncology*, Vol.23, No.5, (November 2003), pp 1277-1283, PMID 14532966
- MacEwan, D. (2002). TNF receptor subtype signaling: differences and cellular consequences. *Cell Signalling*, Vol.14, No.6, (June 2002), pp 477-492, PMID 11897488
- MacGrogan, D., Saint-Andre J. & Dicou E. (1992). Expression of nerve growth factor and nerve growth factor receptor genes in human tissues and in prostatic adenocarcinoma cell lines. *Journal of Neurochemistry*, Vol.59, No.4, (October 1992), pp 1381-1391, PMID 1383421

- Madaan, S., Abel, P. Chaudhary, K., Hewitt, R., Stott, M., Stamp, G. & Lalani, E. (2000). Cytoplasmic induction and over-expression of cyclooxygenase-2 in human prostate cancer: implications for prevention and treatment. *British Journal of Urology International*, Vol.86, No.6, (October 2000), pp 736-41, PMID 11069387
- Maisonpierre, P., Belluscio, L., Squinto, S., Ip, N., Furth, M., Lindsay, R. & Yancopoulos, G. (1990). Neurotrophin-3: a neurotrophic factor related to NGF and BDNF. *Science*, (March 1990) Vol.247, No. 4949, pp 1446-1451, PMID 2321006
- Martinon, F., Holler, N., Richard, C. & Tschopp, J. (2000). Activation of a pro-apoptotic amplification loop through inhibition of NF-kappaB-dependent survival signals by caspase-mediated inactivation of RIP. *Federation of Experimental Biology Society Letters*, Vol.468, No.2-3, (February 2000), pp 134-136, PMID 10692573
- Mascaux, C., Martin, B., Paesmans, M. Berghmans, T., Dusart, M., Haller, A., Lothaire, P., Meert, A., Lafitte, J. & Sculier, J. (2006). Has Cox-2 a prognostic role in non-small-cell lung cancer? A systematic review of the literature with meta-analysis of the survival results. *British Journal of Cancer*, Vol.95, No.2, (July 2006), pp 139-45, PMID 16786043
- Masferrer, J., Zweifel, B.S., Colburn, S.M., Ornberg, R.L., Salvemini, D., Isakson, P. & Seibert, K. (1995). The Role of Cyclooxygenase-2 in Inflammation. *American Journal of Therapeutics*, Vol.2, No.9, (September 1995) pp. 607-610, PMID11854834
- Mazumder, B., Seshadri, V. & Fox, P. (2003). Translational control by the 3'-UTR: the ends specify the means. *Trends in Biochemical Sciences*, 2003;Vol.28, No.2, (February 2003), pp 91-98, PMID 12575997
- Mazzieri, R., Masiero, L., Zanetta, L., Monea, S., Onisto, M., Garbisa, S. & Mignatti, P. (1997). Control of type IV collagenase activity by components of the urokinase-plasmin system: a regulatory mechanism with cell-bound reactants. *The EMBO Journal*, Vol.16, No.9, (May 1997), pp 2319-32, PMID 9171346
- Meyer, C., Jacobs, H. & Lehner, C. (2002). Cyclin d-cdk4 is not a master regulator of cell multiplication in Drosophila embryos. *Current Biology*, Vol.12, No.8, (April 2002), pp 661-666, PMID 11967154
- Nalbandian A. & Djakiew, D. (2006). The p75^{NTR} metastasis suppressor inhibits urokinase plasminogen activator, matrix metalloproteinase-2 and matrix metalloproteinase-9 in PC-3 prostate cancer cells. *Clinical and Experimental Metastasis*, Vol.23, No.2, (August 2006), pp. 107-116, PMID 16912916
- Nalbandian, A., Pang, A., Rennert, O., Chan, W., Ravindranath, N. & Djakiew, D. (2005). A novel function of differentiation revealed by cDNA microarray profiling of p75^{NTR}-regulated gene expression. *Differentiation*, Vol.73, No.8 (October 2005), pp385-396, PMID 16316409
- Nelson, J. & Harris, R. (2000). Inverse association of prostate cancer and non-steroidal anti-inflammatory drugs (NSAIDs): results of a case-control study. *Oncology Reports*, Vol.7, No.1, (January-February 2000), pp. 69-70, PMID 10601612
- O'Neill, J., Manion, M., Schwartz, P. & Hockenbery, D. (2004). Promises and challenges of targeting Bcl-2 anti-apoptotic proteins for cancer therapy *Biochimica Biophysica Acta*, Vol.1705, No.1, (December 2004), pp 43-51, PMID 15585172
- Paul, A., Grant, E. & Habib, F. (1992). The expression and localisation of beta-nerve growth factor (beta-NGF) in benign and malignant human prostate tissue: relationship to neuroendocrine differentiation. *British Journal of Cancer*, Vol.74, No.12, (December 1992), pp 1990-1996, PMID 8980402

- Perez, M., Regan, T., Pflug, B., Lynch, J. & Djakiew, D. (1997). Loss of low-affinity nerve growth factor receptor during malignant transformation of the human prostate. *Prostate*, Vol.30, No.4, (March 1997), pp. 274-279, PMID 9111606
- Pflug, B., Dionne, C., Kaplan, D., Lynch, J. & Djakiew, D. (1995). Expression of a Trk high affinity nerve growth factor receptor in the human prostate. *Endocrinology*, Vol.136, No.1 (January 1995), pp. 262-268, PMID 7828539
- Pflug, B. & Djakiew, D. (1998). Expression of p75^{NTR} in a human prostate epithelial tumor cell line reduces nerve growth factor-induced cell growth by activation of programmed cell death. *Molecular Carcinogenesis* Vol.23, No.2 (October 1998), pp 106-114, PMID 9808164
- Pflug, B., Onoda, M., Lynch, J. & Djakiew, D. (1992). Reduced expression of the low affinity nerve growth factor receptor in benign and malignant human prostate tissue and loss of expression in four human metastatic prostate tumor cell lines. *Cancer Research*, Vol.52, No.19 (October 1992), pp. 5403-5406, PMID 1382843
- Puthalakath, H. & Strasser, A. (2002). Keeping killers on a tight leash: transcriptional and post-translational control of the pro-apoptotic activity of BH3-only proteins *Cell Death and Differentiation*, Vol.9, No.5, (May 2002), pp505-512, PMID 11973609
- Quann, E., Khwaja, F., & Djakiew, D. (2007b). The p38 MAPK Pathway Mediates Aryl Propionic Acid-Induced messenger RNA Stability of p75^{NTR} in Prostate Cancer Cells. *Cancer Research*, Vol.67, No.23, (December 2007) pp. 11402-11410, PMID 18056468
- Quann, E., Khwaja, F., Zavitz, K. & Djakiew, D. (2007a). The Aryl Propionic Acid R-Flurbiprofen Selectively Induces p75^{NTR}-Dependent Decreased Survival of Prostate Tumor Cells. *Cancer Research* Vol.67, No.7, (April 2007) pp 3254-3262, PMID 17409433
- Reuning, U., Magdolen, V., Wilhelm, O., Fischer, K., Lutz, V., Graeff, H. & Schmitt, M. (1998). Multifunctional potential of the plasminogen activation system in tumor invasion and metastasis (review). *International Journal of Oncology*, Vol.13, No.5, (November 1998), pp. 893-906, PMID 9772277
- Roberts, R., Jacobson, D., Girman, C., Rhodes, T., Lieber, M. & Jacobsen, S. (2002). A population-based study of daily nonsteroidal anti-inflammatory drug use and prostate cancer. *Mayo Clinic Proceedings*, Vol.77, No.3 (March 2002), pp 219-25, PMID 11888024
- Ronkina, N., Kotlyarov, A., Dittrich-Breiholz, O., Kracht, M., Hitti, E., Milarski, K., Askew, R., Marusic, S., Lin, L., Gaestel, M. & Telliez, J. (2007). The mitogen-activated protein kinase (MAPK)-activated protein kinases MK2 and MK3 cooperate in stimulation of tumor necrosis factor biosynthesis and stabilization of p38 MAPK. *Molecular and Cellular Biology*, Vol.27, No.1, (January 2007), pp 170-181, PMID 17030606
- Satyanarayana, A. & Rudolph, K. (2004). p16 and ARF: activation of teenage proteins in old age. *Journal of Clinical Investigation*, Vol.114, No.9, (November 2004), pp 1237-1240, PMID 15520854
- Sarkar, F, Adsule, S., Li, Y. & Padhye, S. (2007). Back to the future: COX-2 inhibitors for chemoprevention and cancer therapy. *Mini Reviews in Medicinal Chemistry*, Vol.7, No.6, (June 2007), pp 599-608.
- Scheper, G., & Proud, C. (2002). Does phosphorylation of the cap-binding protein eIF4E play a role in translation initiation? *European Journal of Biochemistry*, Vol.269, No.22, (November 2002), pp 5350-5359, PMID 12423333
- Sellers, W. & Kaelin W. (1996). RB [corrected] as a modulator of transcription. *Biochimica Biophysica Acta*, Vol.1288, No.3, (August 1996), Pp M1-5, PMID 8764839

- Sherr, C. (1996). Cancer cell cycles. *Science*, Vol. 274, No. 5293 (December 1996), pp1672-1677, PMID 8939849
- Sherr, C. & Roberts, J. (1995). Inhibitors of mammalian G1 cyclin-dependent kinases. *Genes and Development*, Vol.9, No.10, (May 1995), pp 1149-1163, PMID 7758941
- Shintani, S., Mihara, M., Nakahara, Y., Kiyota, A., Ueyama, Y., Matsumura, T. & Wong, D. Expression of cell cycle control proteins in normal epithelium, premalignant and malignant lesions of oral cavity. *Oral Oncology*, Vol.38, No.3, (April 2002), pp235-243, PMID 11978545
- Shiozaki, E. Y. Shi (2004). Caspases, IAPs and Smac/DIABLO: mechanisms from structural biology. *Trends in Biochemical Sciences*, Vol.29, No.9, (September 2004), pp. 486-494, PMID 15337122
- Song, I., Tatebe, S., Dai, W. & Kuo, M. (2005). Delayed mechanism for induction of gamma-glutamylcysteine synthetase heavy subunit mRNA stability by oxidative stress involving p38 mitogen-activated protein kinase signaling. *Journal of Biological Chemistry*, Vol.280, No.13, (August 2005), pp 28230-28240, PMID 15946948
- Soppet, D., Escandon, E., Maragos, J., Middlemas, D., Reid, S., Blair, J., Burton, L., Stanton, B., Kaplan, D., Hunter, T., Nikolics, K. & Parada, L. (1991). The neurotrophic factors brain-derived neurotrophic factor and neurotrophin-3 are ligands for the trkB tyrosine kinase receptor. *Cell*, Vol.65, No.5, (May 1991), pp 895-903, PMID 1645620
- Tegeder, I., Pfeilschifter, J., & Geisslinger, G. (2001). Cyclooxygenase-independent actions of cyclooxygenase inhibitors. *Federation of American Society of Experimental Biology Journal*, Vol.15, No.12, (October 2001) pp. 2057-2072, PMID 11641233
- Thun, M., Namboodiri, M. & Heath, C. (1991). Aspirin use and reduced risk of fatal colon cancer. *The New England Journal of Medicine*, Vol.325, No.23, (December 1991), pp 1593-6, PMID 1669840
- Tran, H., Maurer, F. & Nagamine, Y. (2003). Stabilization of urokinase and urokinase receptor mRNAs by HuR is linked to its cytoplasmic accumulation induced by activated mitogen-activated protein kinase-activated protein kinase 2. *Molecular and Cellular Biology*, Vol.23, No.20, (October 2003), pp 7177-7188, PMID 14517288
- von der Haar, T., Gross, J., Wagner, G. & McCarthy, J. (2004). The mRNA cap-binding protein eIF4E in post-transcriptional gene expression. *Nature Structure and Molecular Biology*, Vol.11, No.6, (June 2004), pp 503-511, PMID 15164008
- Westwood, G., Dibling, B., Cuthbert-Heavens, D. & Burchill, S. (2002). Basic fibroblast growth factor (bFGF)-induced cell death is mediated through a caspase-dependent and p53-independent cell death receptor pathway. *Oncogene*, Vol.21, No.5, (January 2002), pp 809-824, PMID 11850809
- Wynne, S. & Djakiew, D. (2010). NSAID-inhibition of prostate cancer cell migration is mediated by Nag-1 induction via the p38 MAPK-p75^{NTR} pathway. *Molecular Cancer Research*, Vol.8, No.12, (December 2010), pp 1656-1664, PMID 21097678
- Yoshimura, R., Sano, H., Masuda, C., Kawamura, M., Tsubouchi, Y., Chargui, J., Yoshimura, N., Hla, T. & Wada, S. (2000). Expression of cyclooxygenase-2 in prostate carcinoma. *Cancer*, Vol.89, No.3, (August 2000), pp 589-596, PMID 10931458
- Zarubin, T. & Han, J. (2005). Activation and signaling of the p38 MAP kinase pathway. *Cell Research*, Vol.15, No.1, (January 2005), p11-18, PMID 15686620
- Zha, S., Gage, W., Sauvageot, J., Saria, E., Putzi, M., Ewing, C., Faith, D., Nelson, W., De Marzo, A. & Isaacs, W. (2001). Cyclooxygenase-2 is up-regulated in proliferative inflammatory atrophy of the prostate, but not in prostate carcinoma. *Cancer Research*, Vol.61, No.24, (December 2001), pp 8617-8623, PMID 11751373



Prostate Cancer - Original Scientific Reports and Case Studies

Edited by Dr. Philippe E. Spiess

ISBN 978-953-307-342-2

Hard cover, 238 pages

Publisher InTech

Published online 21, November, 2011

Published in print edition November, 2011

This book encompasses three sections pertaining to the topics of cancer biology, diagnostic markers, and therapeutic novelties. It represents an essential resource for healthcare professionals and scientist dedicated to the field of prostate cancer research. This book is a celebration of the significant advances made within this field over the past decade, with the hopes that this is the stepping stone for the eradication of this potentially debilitating and/or fatal malignancy.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Daniel Djakiew (2011). NSAID Induction of p75NTR in the Prostate: A Suppressor of Growth and Cell Migration Via the p38 MAPK Pathway, Prostate Cancer - Original Scientific Reports and Case Studies, Dr. Philippe E. Spiess (Ed.), ISBN: 978-953-307-342-2, InTech, Available from: <http://www.intechopen.com/books/prostate-cancer-original-scientific-reports-and-case-studies/nsaid-induction-of-p75ntr-in-the-prostate-a-suppressor-of-growth-and-cell-migration-via-the-p38-mapk>

INTECH

open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821