

Attachment and Biofilm Formation by *Salmonella* in Food Processing Environments

Efstathios Giaouris¹, Nikos Choriantopoulos²,
Panagiotis Skandamis³ and George-John Nychas³

¹Department of Food Science and Nutrition, University of the Aegean

²Veterinary Research Institute, National Agricultural Research Foundation (NAGREF)

³Department of Food Science and Technology, Agricultural University of Athens (AUA)
Greece

1. Introduction

During the last decades, it has become increasingly clear that bacteria, including foodborne pathogens such as *Salmonella enterica*, grow predominantly as biofilms in most of their natural habitats, rather than in planktonic mode. A biofilm can be broadly defined as a microbially derived sessile community characterized by cells that are irreversibly attached to a substratum or interface or to each other, are embedded in a matrix of extracellular polymeric substances (EPS) that they have produced, and exhibit an altered phenotype with respect to growth rate and gene transcription (Donlan & Costerton, 2002; Kuchma & O'Toole, 2000; Lazazzera, 2005; Shemesh et al., 2007). Interestingly, it has been observed that the resistance of biofilm cells to antimicrobials is significantly increased compared with what is normally seen with the same cells being planktonic (Gilbert et al., 2002; Mah & O'Toole, 2001). Thus, it is believed that biofilm formation enhances the capacity of pathogenic *Salmonella* bacteria to survive stresses that are commonly encountered both within food processing, as well as during host infection.

In food industry, biofilms may create a persistent source of product contamination, leading to serious hygienic problems and also economic losses due to food spoilage (Brooks & Flint, 2008; Carpentier & Cerf, 1993; Ganesh Kumar & Anand, 1998; Lindsay & von Holy, 2006; Zottola & Sasahara, 1994). Improperly cleaned surfaces promote soil build-up, and, in the presence of water, contribute to the development of bacterial biofilms which may contain pathogenic microorganisms, such as *Salmonella*. Cross contamination occurs when cells detach from biofilm structure once food passes over contaminated surfaces or through aerosols originating from contaminated equipment. Till now, there is only limited information on the presence of *Salmonella* in biofilms in real food processing environments. However, numerous studies have shown that *Salmonella* can easily attach to various food-contact surfaces (such as stainless steel, plastic and cement) and form biofilms under laboratory conditions (Chia et al., 2009; Giaouris et al., 2005; Giaouris & Nychas, 2006; Hood & Zottola, 1997a,b; Marin et al., 2009; Oliveira et al., 2006; Rodrigues et al., 2011; Vestby et al., 2009a,b).

The natural environments that most bacteria inhabit are typically complex and dynamic. Unfortunately, this complexity is not fully appreciated when growing microorganisms in monocultures under laboratory conditions. Thus, in real environments, biofilm communities

are usually inhabited by numerous different species in close proximity (Wimpenny et al., 2000). Spatial and metabolic interactions between species contribute to the organization of multispecies biofilms, and the production of a dynamic local environment (Goller & Romeo, 2008; Tolker-Nielsen & Molin, 2000). Indeed, cell-to-cell signalling and interspecies interactions have been demonstrated to play a key role in cell attachment and detachment from biofilms, as well as in the resistance of biofilm community members against antimicrobial treatments (Annous et al., 2009; Burmølle et al., 2006; Irie & Parsek, 2008; Nadell et al., 2008; Remis et al., 2010). Mixed-species biofilms are usually more stable than mono-species biofilms, while biofilm formation by *Salmonella* has also been shown to be influenced by either the natural *in situ* presence of other species, or just their metabolic by-products (Chorianopoulos et al., 2010; Girenavar et al., 2008; Habimana et al., 2010b; Jones & Bradshaw, 1997; Prouty et al., 2002; Soni et al., 2008).

In this chapter, we review up-to-date available voluminous literature on the attachment and biofilm formation by *Salmonella* strains on abiotic surfaces, simulating those encountered in food processing areas (section 4). Before this, the advantages of biofilm lifestyle for microorganisms are briefly discussed (section 2), together with the serious negative implications of biofilm formation for the food industry (section 3). Major molecular components building up *Salmonella* biofilm matrix are then reported (section 5). Finally, we review available knowledge on the influence of cell-to-cell communication (quorum sensing) on the establishment of *Salmonella* biofilms (section 6).

2. Bacterial attachment to surfaces and advantages of the biofilm lifestyle

For most of the history of microbiology, microorganisms have primarily been characterised as planktonic, freely suspended cells and described on the basis of their growth characteristics in nutritionally rich culture media. Although this traditional way of culturing bacteria in liquid media has been instrumental in the study of microbial pathogenesis and enlightening as to some of the amazing facets of microbial physiology, pure culture planktonic growth is rarely how bacteria exist in nature. On the contrary, direct observation of wide variety of natural habitats has shown that the majority of microbes persist attached to surfaces within a structured biofilm ecosystem and not as free-floating organisms (Costerton et al., 1987, 1995; Kolter & Greenberg, 2006; Verstraeten et al., 2008).

The data on which this theory is predicated came mostly from natural aquatic ecosystems, in which direct microscopic observations together with direct quantitative recovery techniques showed unequivocally that more than 99.9% of the bacteria grow as biofilms on a wide variety of surfaces. The diversity and distribution of salmonellae in fresh water biofilms has also been recently shown (Sha et al., 2011). Moreover, it is becoming clear that these natural assemblages of bacteria within the biofilm matrix function as a cooperative consortium, in a relatively complex and coordinated manner (James et al., 1995; Moons et al., 2009; Wuertz et al., 2004). Nowadays, besides natural aquatic systems, it is well established that biofilms may form on a wide variety of surfaces, including living tissues, indwelling medical devices and also industrial systems, such as pharmaceutical industries, oil drilling, paper production, waste water treatment and food processing (Hall-Stoodley et al., 2004). Thus, examples of this bacterial lifestyle are abundant in daily life: the slimy material that covers flower vases, pipelines, submerged rocks, and even the surface of teeth (Marsh, 2005; Wimpenny, 2009).

Biofilm formation occurs through sequential steps in which the initial attachment of planktonic bacteria to a solid surface is followed by their subsequent proliferation and

accumulation in multilayer cell clusters, and the final formation of the bacterial community enclosed in a self-produced polymeric matrix (Goller & Romeo, 2008; Lasa, 2006; O'Toole et al., 2000; Palmer et al., 2007; Rickard et al., 2003). The initial interaction between solid surface and bacterial cell envelope appears to be mediated by a complex array of chemical and physical interactions, with each affected by the chemical and physical environment to which the bacterial cell and the surface are currently or recently exposed (Palmer et al., 2007). Mature biofilms are highly organized ecosystems in which water channels are dispersed and can provide passages for the exchange of nutrients, metabolites and waste products (Stoodley et al., 2002). Once the biofilm structure has developed, some bacteria are released into the liquid medium, in order to colonize new surfaces, probably when surrounding conditions become less favourable (Gilbert et al., 1993; Hall-Stoodley & Stoodley, 2002, 2005; Klausen et al., 2006).

According to Darwin's theory of evolution, the only true driving force behind the course of action of any organism is reproductive fitness. Outside of the laboratory bacteria rarely, if ever, find themselves in an environment as nutrient rich as culture media, and in these conditions, there are a number of fitness advantages imparted by the biofilm mode of growth (Jefferson, 2004). The process of biofilm formation is believed to begin when bacteria sense certain environmental parameters (extracellular signals) that trigger the transition from planktonic growth to life on a surface (Lopez et al., 2010). Currently, four potential incentives behind the formation of biofilms by bacteria are considered: (i) protection from the harmful environment (as a stress response mechanism), (ii) sequestration to a nutrient rich area, (iii) utilization of cooperative benefits (through metabolic cooperativity), and (iv) acquisition of new genetic traits (Davey & O'Toole, 2000; Molin & Tolker-Nielsen, 2003).

Bacteria experience a certain degree of shelter and homeostasis when residing within a biofilm and one of the key components of this microniche is the surrounding extrapolymeric substance (EPS) matrix (Flemming & Wingender, 2010). This matrix is composed of a mixture of components, such as exopolysaccharides, proteins, nucleic acids, and other substances (Branda et al., 2005). The nature of biofilm matrix and the physiological attributes of biofilm microorganisms confer an inherent resistance to antimicrobial agents, whether these antimicrobial agents are antibiotics, disinfectants or germicides. Thus, established biofilms can tolerate antimicrobial agents at concentrations of 10-1000 times that need to kill genetically equivalent planktonic bacteria, and are also extraordinary resistant to phagocytosis, making rather difficult to eradicate biofilms from living hosts (Cos et al., 2010). Mechanisms responsible for resistance may be one or more of the following: (i) delayed penetration of the antimicrobial agent through the biofilm matrix, (ii) altered growth rate of biofilm microorganisms, and (iii) other physiological changes due to the biofilm mode of growth, e.g. existence of subpopulations of resistant phenotypes in the biofilm, which have been referred to as "persisters" (Donlan & Costerton, 2002; Gilbert et al., 2002; Lewis, 2001; Mah & O'Toole, 2001).

Scientific interest in the process of bacterial biofilm formation has erupted in recent years and studies on the molecular genetics of biofilm formation have begun to shed light on the driving forces behind the transition to the biofilm mode of existence. Evidence is mounting that up- and down-regulation of a number of genes occurs in the attaching cells upon initial interaction with the substratum (Donlan, 2002; Sauer, 2003). Thus, high-throughput DNA microarray studies have been conducted to study biofilm formation in many model microorganisms and have identified a large number of genes showing differential expression under biofilm conditions (Beloin et al., 2004; Hamilton et al., 2009; Lazazzera,

2005; Shemesh et al., 2007; Whiteley et al., 2001). In *S. Typhimurium*, 10% of its genome (i.e. 433 genes) showed a 2-fold or more change in the biofilm, using a silicone rubber tubing as a substratum for growth, compared with planktonic cells (Hamilton et al., 2009). The genes that were significantly up-regulated implicated certain cellular processes in biofilm development, including amino acid metabolism, cell motility, global regulation and tolerance to stress. Obviously, the more we learn about the genetic regulation of biofilm formation, the more we understand about the relative roles of benefits and forces that drive the switch to the biofilm mode of growth.

3. Biofilm formation in food processing environments and implications

The ability of bacteria to attach to abiotic surfaces and form biofilms is a cause of concern for many industries, including the food ones (Chmielewski & Frank, 2003). Poor sanitation of food-contact surfaces is believed to be an essential contributing factor in foodborne disease outbreaks, especially those involving *Listeria monocytogenes* and *Salmonella*. This is because the attachment of bacterial cells to such surfaces is the first step of a process which can ultimately lead to the contamination of food products. Thus, biofilms formed in food processing environments are of special importance since they may act as a persistent source of microbial contamination which may lead to food spoilage or/and transmission of diseases (Brooks & Flint, 2008; Zottola & Sasahara, 1994). While food spoilage and deterioration may result in huge economic losses, food safety is a major priority in today's globalizing market with worldwide transportation and consumption of raw, fresh and minimally processed foods (Shi & Zhu, 2009).

Besides food spoilage and safety issues, in the dairy industry, bacterial attachment in heat exchangers (a process commonly known as "biofouling") greatly reduces the heat transfer and operating efficiency of the processing equipment, while it can also causes corrosion problems (Austin & Bergeron, 1995). Additionally, in the various filtration systems, biofilm formation reduces significantly the permeability of the membranes (Tang et al., 2009). However, it should be noted that in the industry of fermented food products (sausages, cheeses etc), biofilm formation by some useful and technological bacteria (e.g. staphylococci, lactococci, lactobacilli) can be desirable, as a mean of the enhancement of food fermentation process, and more importantly as a mean of protection against the establishment of pathogenic biofilms (Chorianopoulos et al., 2008; Zhao et al., 2006).

Adhesion of *Salmonella* to food surfaces was the first published report on foodborne bacterial biofilm (Duguid et al., 1966). Since that time, many documents have described the ability of foodborne pathogens to attach to various surfaces and form biofilms, including *L. monocytogenes* (Blackman & Frank, 1996; Chorianopoulos et al., 2011; di Bonaventura et al., 2008; Poimenidou et al., 2009), *Salmonella enterica* (Chia et al., 2009; Giaouris et al., 2005; Giaouris & Nychas, 2006; Habimana et al., 2010b; Joseph et al., 2001; Kim & Wei, 2007, 2009; Oliveira et al., 2006; Marin et al., 2009; Rodrigues et al., 2011; Solomon et al., 2005; Stepanović et al., 2003, 2004), *Yersinia enterocolitica* (Kim et al., 2008), *Campylobacter jejuni* (Joshua et al., 2006) and *Escherichia coli* O157:H7 (Habimana et al., 2010a; Skandamis et al., 2009).

Modern food processing supports and selects for biofilm forming bacteria on food-contact surfaces due to mass production of products, lengthy production cycles and vast surface areas for biofilm development (Lindsay & von Holy, 2006). *In situ* biofilms have been recognised in various food processing industries, such as processors of cheese and other milk products, raw and cooked/fermented meats, raw and smoked fish etc (Austin & Bergeron, 1995; Bagge-Ravn

et al., 2003; Gounadaki et al., 2008; Gunduz & Tuncel, 2006; Sharma & Anand, 2002). Several studies were also focused on the attachment of bacterial pathogens to food surfaces such as *Escherichia coli* to beef muscle and adipose tissue (Rivas et al., 2006) and *S. Typhimurium*, *Yersinia enterocolitica* and *L. monocytogenes* to pork skin (Morild et al., 2011).

Biofilm formation depends on an interaction between three main components: the bacterial cells, the attachment surface and the surrounding medium (Van Houdt & Michiels, 2010). Adhesion of bacterial cells, the first phase of biofilm formation, is influenced by the physicochemical properties of the cells' surface, which in turn are influenced by factors such as microbial growth phase, culture conditions and strain's variability (Briandet et al., 1999; Giaouris et al., 2009). The surfaces of most bacterial cells are negatively charged, and this net negative charge of the cell surface is adverse to bacterial adhesion, due to electrostatic repulsive force. However, the bacterial cell-surface possesses hydrophobicity due to fimbriae, flagella and lipopolysaccharide (LPS) (Ukuku & Fett, 2006). Hydrophobic interactions between the cell surface and the substratum may enable the cell to overcome repulsive forces and attach irreversibly (Donlan, 2002). The properties of the attachment surface (e.g. roughness, cleanability, disinfectability, wettability, vulnerability to wear) are important factors that also affect the biofilm formation potential and thus determine the hygienic status of the material. Stainless steel type 304, commonly used in the food processing industry, is an ideal material for fabricating equipment due to its physico-chemical stability and high resistance to corrosion. Teflon and other plastics are often used for gaskets and accessories of instruments. These surfaces become rough or crevice with continuous reuse and form a harbourage to protect bacteria from shear forces in the food fluid.

Environmental factors such as pH, temperature, osmolarity, O₂ levels, nutrient composition and the presence of other bacteria play important roles in the process of biofilm formation (Giaouris et al., 2005; Hood & Zottola, 1997a; Stepanovic et al., 2003). The integration of these influences ultimately determines the pattern of behavior of a given bacterium with respect to biofilm development (Goller & Romeo, 2008). In food processing environments, bacterial attachment is additionally affected by food matrix constituents, which can be adsorbed onto a substratum and create conditioning films (Bernbom et al., 2009). For example, skim milk was found to reduce adhesion of *Staphylococcus aureus*, *L. monocytogenes*, and *Serratia marcescens* to stainless steel coupons (Barnes et al., 1999). Additionally, in real environments, the presence of mixed bacterial communities adds additional complexity to attachment and biofilm formation procedure. For instance, the presence of *Staphylococcus xylosus* and *Pseudomonas fragi* affected the numbers of *L. monocytogenes* biofilm cells on stainless steel (Norwood & Gilmour, 2001), while compounds present in *Hafnia alvei* cell-free culture supernatant inhibited the early stage of *S. Enteritidis* biofilm formation on the same material (Chorianopoulos et al., 2010).

Once biofilms have formed in the factory environment, they are difficult to be removed often resulting in persistent and endemic populations (Vestby et al., 2009b). Interestingly, persistent *L. monocytogenes* strains had the added ability of enhanced adhesion within shorter times to stainless steel surfaces compared to non-persistent strains (Lundén et al., 2000). It has been suggested that such persistence is likely due to physical adaptation of cells in biofilms, particularly resistance to cleaning and sanitizing regimes, since it is generally accepted and well documented that cells within a biofilm are more resistant to biocides than their planktonic counterparts (Carpentier & Cerf, 1993). For example, nine disinfectants commonly used in the feed industry and efficient against planktonic *Salmonella* cells, showed a bactericidal effect that varied considerably for biofilm-grown cells with products containing

70% ethanol being most effective (Møretrø et al., 2009). Other studies similarly indicated that compared to planktonic cells, biofilm cells of *Salmonella* were more resistant to trisodium phosphate (Scher et al., 2005) and to chlorine and iodine (Joseph et al., 2001). In a comparative study of different *S. Enteritidis* phage type 4 isolates it was found that those isolates that survived better on surfaces also survived better in acidic conditions and in the presence of hydrogen peroxide and showed enhanced tolerance towards heat (Humphrey et al., 1995). The cellular mechanisms underlying microbial biofilm formation and behaviour are beginning to be understood and are targets for novel specific intervention strategies to control problems caused by biofilm formation in fields ranging from industrial processes like food processing, to health-related fields, like medicine and dentistry. In food industry, various preventive and control strategies, like hygienic plant lay-out and design of equipment, choice of materials, correct selection and use of detergents and disinfectants coupled with physical methods can be suitably applied for controlling biofilm formation. Right now, bacterial biofilms have not been specifically addressed in the HACCP system that has been employed in the food processing facilities. However, surveying of biofilms in food environments and developing an effective sanitation plan should be considered in the HACCP system (Sharma & Anand, 2002). An upgraded HACCP with biofilm assessment in food plants will provide clearer information of contamination, and assist the development of biofilm-free processing systems in the food industry.

4. Attachment to food-contact surfaces and biofilm forming ability of *Salmonella*

Salmonellae represent a group of Gram-negative bacteria that are recognized worldwide as major zoonotic pathogens for both humans and animals. In the EU, salmonellosis was the second most commonly reported zoonotic infection in 2009, with 108,614 human cases confirmed and a case fatality rate of 0.08%, which approximately corresponds to 90 human deaths (EFSA-ECDC, 2011). That year, *Salmonella* was most often found in fresh broiler, turkey and pig meat where proportions of positive samples, on average 5.4%, 8.7% and 0.7%, were detected respectively. The two most common *Salmonella* serotypes, implicated in the majority of outbreaks, are Typhimurium and Enteritidis (52.3% and 23.3% respectively of all known serovars in human cases). The native habitat of salmonellae is considered to be the intestinal tract of taxonomically diverse group of vertebrates, from which salmonellae can spread to other environments through released faeces (Litrup et al., 2010).

Interestingly, salmonellae have been shown to survive for extended periods of time in non-enteric habitats, including biofilms on abiotic surfaces (White et al., 2006). Thus, several reports have demonstrated the ability of *Salmonella* to form biofilms on abiotic surfaces outside the host, such as stainless steel (Austin et al., 1998; Chorianopoulos et al., 2010; Giaouris et al., 2005; Giaouris & Nychas, 2006; Hood & Zottola, 1997a,b; Joseph et al., 2001; Kim & Wei, 2007, 2009; Møretrø et al., 2009), plastic (Asséré et al., 2008; Iibuchi et al., 2010; Jain & Chen, 2007; Joseph et al., 2001; Ngwai et al., 2006; Stepanović et al., 2003, 2004; Vestby et al., 2009a,b), rubber (Arnold & Yates, 2009), glass (Kim & Wei, 2009; Korber et al., 1997; Prouty & Gunn, 2003; Solano et al., 1998), cement (Joseph et al., 2001), marble and granite (Rodrigues et al., 2011). Taken into account, that all these surfaces are commonly encountered in farms, slaughter houses, food industries and kitchens, it is obvious that the risk for public health is quite serious.

It is strongly believed that the ability of *Salmonella* to form biofilms on inanimate surfaces contributes to its survival and persistence in non-host environments and its transmission to

new hosts. To this direction, Vestby et al. (2009b) found a correlation between the biofilm formation capacities of 111 *Salmonella* strains isolated from feed and fish meal factories and their persistence in the factory environment. Another study on colonization and persistence of *Salmonella* on egg conveyor belts indicated that the type of egg belt (i.e. vinyl, nylon, hemp or plastic) was the most important factor in colonization and persistence, while rdar morphotype, a physiological adaptation associated with aggregation and long-term survival which is conserved in *Salmonella* (White & Surette, 2006), surprisingly, was not essential for persistence (Stocki et al., 2007). Interestingly, inoculation onto fresh-cut produce surfaces, as well as onto inert surfaces, such as polyethersulfone membranes, was found to significantly increase the survival of salmonellae during otherwise lethal acid challenge (pH 3.0 for 2 hours) (Gawande & Bhagwat, 2002). Similarly, *Salmonella* strains with high biofilm productivity survived longer on polypropylene surfaces under dry conditions than strains with low productivity (Iibuchi et al., 2010).

In the food processing environments, food-contact surfaces come in contact with fluids containing various levels of food components. Under such conditions, one of the first events to occur is the adsorption of food molecules to the surface (conditioning). Both growth media and surface conditioning were found to influence the adherence of *S. Typhimurium* cells to stainless steel (Hood & Zottola, 1997b). A study of 122 *Salmonella* strains indicated that all had the ability to adhere to plastic microwell plates and that, generally, more biofilm was produced in low nutrient conditions, as those found in specific food processing environments, compared to high nutrient conditions (Stepanovic et al., 2004). A study conducted in order to identify the risk factors for *Salmonella* contamination in poultry farms, showed that the most important factors were dust, surfaces and faeces, and nearly 50% of the strains isolated from poultry risk factors were able to produce biofilm, irrespective of the origin of different serotypes (Marin et al., 2009).

There are some studies which have investigated the influence of physicochemical and surface properties (e.g. charge, hydrophobicity, surface free energy, roughness) of *Salmonella* and surface materials on the attachment process. For instance, Sinde & Carballo (2000) found that surface free energies and hydrophobicity do not affect attachment of *Salmonella* to stainless steel, rubber and polytetrafluorethylene, while Ukuku & Fett (2002) found that there was a linear correlation between bacterial cell surface hydrophobicity and charge and the strength of attachment of *Salmonella*, *E. coli* and *L. monocytogenes* strains to cantaloupe surfaces. Korber et al. (1997) found that surface roughness influences susceptibility of *S. Enteritidis* biofilms, grown in glass flow cells (with or without artificial crevices) to trisodium phosphate. Chia et al. (2009) studied the attachment of 25 *Salmonella* strains to four different materials (Teflon®, stainless steel, rubber and polyurethane) commonly found in poultry industry and found out that materials more positive in interfacial free energies had the highest number of adhering bacteria. However, in that study, authors concluded that *Salmonella* adhesion is strain-dependent, and probably influenced by surface structures, such as cell wall and membrane proteins, fimbriae, flagella and polysaccharides. This was also the conclusion of another similar study which compared the adhesion ability of four *S. Enteritidis* isolates to three different materials (polyethylene, polypropylene and granite) used in kitchens (Oliveira et al., 2006). Ngwai et al. (2006) characterized the biofilm forming ability of eleven antibiotic-resistant *S. Typhimurium* DT104 clinical isolates from human and animal sources and concluded that there was a general lack of correlation between this ability and bacterial physicochemical surface characteristics.

The persistence of *Salmonella* within the food chain has become a major health concern, as biofilms of this pathogen formed in food processing environments can serve as a reservoir for the contamination of food products. The development of materials to be used for food-contact surfaces with improved food safety profiles continues to be a challenge. One approach which has been developed to control microbial attachment is the manufacture of food-contact materials incorporating antimicrobial compounds. Triclosan-impregnated kitchen bench stones (silestone), although prone to bacterial colonization, were found to reduce *S. Enteritidis* biofilm development on them and also the viability of cells within the biofilm (Rodrigues et al., 2011).

5. Molecular components of *Salmonella* biofilms formed on abiotic surfaces

Curli fimbriae (formerly designated as thin aggregative fimbriae or Tafi) and cellulose are the two main matrix components (exopolymeric substances, EPS) in *Salmonella* biofilms (Gerstel & Römling, 2003). When co-expressed on Congo Red (CR) agar plates, curli fimbriae and the exopolysaccharide cellulose form the characteristic rdar (red, dry and rough) morphotype (also called rugose or wrinkled) (Römling, 2005). Their syntheses are co-regulated by a complex regulatory system. The LuxR type regulator CsgD protein stimulates the production of curli through transcriptional activation of the *csgBAC* (formerly *agfBAC*) operon, while the activation of cellulose production is indirect through the regulator AdrA which is a member of the GGDEF protein family regulated by *csgD* (Römling et al., 2000). García et al. (2004) demonstrated that most GGDEF proteins of *S. Typhimurium* are functionally related, probably by controlling the levels of the same final product, cyclic di-GMP, a secondary messenger that seems to regulate a variety of cellular functions including cellulose production and biofilm formation. The co-expression of curli fimbriae and cellulose leads to the formation of a highly hydrophobic network with tightly packed cells aligned in parallel in a rigid matrix and enhances biofilm formation on abiotic surfaces (Jain & Chen, 2007). Solomon et al. (2005) showed that 72% of 71 *S. enterica* strains, originating from produce, meat or clinical sources and belonging to 28 different serovars, expressed the rdar morphotype, with curli- and cellulose-deficient isolates being least effective in biofilm formation on polystyrene microtiter plates. White et al. (2006) showed that rdar morphotype significantly enhanced the resistance of *Salmonella* to desiccation and sodium hypochlorite, suggesting that this phenotype could play a role in the transmission of *Salmonella* between hosts. However, aggregation via the rdar morphotype does not seem to be a virulence adaptation in *S. Typhimurium*, since competitive infection experiments in mice showed that nonaggregative cells outcompeted rdar-positive wild-type cells in all tissues analyzed (White et al., 2008).

A variety of environmental cues such as nutrients, oxygen tension, temperature, pH, ethanol and osmolarity can influence the expression of the transcriptional regulator CsgD, which regulates the production of both cellulose and curli (Gerstel & Römling, 2003). Transcription of *csgD* is dependent upon the stationary phase-inducible sigma factor RpoS, and is maximal in the late exponential or early stationary phase of growth (Gerstel & Römling, 2001). For an extensive overview on the current understanding of the complex genetic network regulating *Salmonella* biofilm formation, reader is advised to refer to the recently published review of Steenackers et al. (2011). When *csgD* is not expressed the morphotype is a conventional smooth and white (saw) colony, which does not produce any extracellular matrix (Römling et al., 1998b). In wild type *Salmonella* strains, rdar morphotype is restricted to low temperature (below 30°C) and low osmolarity conditions, but biogenesis of curli

fimbriae occurs upon iron starvation at 37°C. Römling et al. (2003) showed that the majority (more than 90% of 800 strains) of human disease-associated *S. Typhimurium* and *S. Enteritidis* (isolated from patients, foods and animals) displayed the rdar morphotype at 28°C, but just rarely at 37°C. Interestingly, mutants in the *csgD* promoter have also been found expressing rdar morphotype independently of temperature (Römling et al., 1998b).

Curli fimbriae are amyloid cell-surface proteins, and are involved in adhesion to surfaces, cell aggregation, environmental persistence and biofilm development (Austin et al., 1998; Collinson et al., 1991; White et al., 2006). The *csg* (curli subunit genes) genes (previously called *agf* genes) involved in curli biosynthesis are organized into two adjacent divergently-transcribed operons, *csgBAC* and *csgDEFG* (Collinson et al., 1996; Römling et al., 1998a). Knocking out the gene encoding for the subunit of thin aggregative fimbriae, AgfA, results in pink colony formation, the pdar (pink, dry and rough) morphotype, which is characterised by production of cellulose without curli (Jain & Chen, 2007). Solano et al. (2002) stressed the importance of the applied biofilm system since they noticed that curli were not essential for biofilm mediated glass adherence under adherence test medium (ATM) conditions, while they were indispensable to form a tight pellicle under LB conditions.

In addition to curli, the second component of the extracellular matrix of the *Salmonella* biofilms is cellulose, a β -1 \rightarrow 4-D-glucose polymer, which is biosynthesized by the *bcsABZC-bcsEFG* genes (bacterial cellulose synthesis) (Zogaj et al., 2001). Both operons are responsible for cellulose biosynthesis in both *S. Enteritidis* and *S. Typhimurium* (Jain & Chen, 2007; Solano et al., 2002). Cellulose production impairment generates a bdar (brown, dry and rough) morphotype on congo red (CR) agar plates, characteristic of the expression of curli. Solano et al. (2002) showed that cellulose is a crucial biofilm determinant for *Salmonella*, under both LB and ATM conditions, without however affecting the virulence of the bacterium. Additionally, cellulose-deficient mutants were more sensitive to chlorine treatments, suggesting that cellulose production and biofilm formation may be an important factor for the survival of *Salmonella* in hostile environments. Prouty & Gunn (2003) identified its crucial importance for biofilm formation on glass coverslips. However, cellulose was not a major constituent of the biofilm matrix of *S. Agona* and *S. Typhimurium* strains isolated from the feed industry, but it contributed to the highly organized matrix structurization (Vestby et al., 2009a). Malcova et al. (2008) found that cellulose was not crucial for *S. Enteritidis* adherence and biofilm formation on polystyrene.

Latasa et al. (2005) also reported another matrix component, BapA, a large cell-surface protein required for biofilm formation of *S. Enteritidis*. This protein was found to be loosely associated with the cell surface, while it is secreted through the BapBCD type I protein secretion system, encoded by the *bapABCD* operon. The expression of *bapA* was demonstrated to be coordinated with the expression of curli and cellulose through the action of *csgD* (Latasa et al., 2005). Also, these authors demonstrated that a *bapA* mutant strain showed a significant lower colonization rate at the intestinal cell barrier and consequently a decreased efficiency for organ invasion compared with the wild-type strain.

Motility was found to be important for *Salmonella* biofilm development on glass (Prouty & Gunn, 2003) and polyvinyl chloride (PVC) (Mireles et al., 2001). On the contrary, Teplitski et al. (2006) noticed that the presence of the flagellum on the surface of the cell, functional or not, is inhibitory to biofilm formation on polystyrene, as mutants lacking intact flagella, showed increased biofilm formation compared to the wild-type. Flagella were not found to be important for *S. Typhimurium* rdar expression on Congo Red (CR) agar plates (Römling & Rohde, 1999). Solano et al. (2002) noticed that flagella affect *S. Enteritidis* biofilm development

only under LB but not under ATM conditions. Stafford & Hughes (2007) showed that the conserved flagellar regulon gene *flhE*, while it is not required for flagella production or swimming, appeared to play a role in flagella-dependent swarming and biofilm formation on PVC. Kim & Wei (2009) noticed that flagellar assembly was important during biofilm formation on PVC in different (meat, poultry and produce) broths and on stainless steel and glass in LB broth.

Colanic acid, a capsular extracellular polysaccharide, essential for *S. Typhimurium* biofilm development on epithelial cells was found not to be required for *Salmonella* biofilm formation on abiotic surfaces (Ledeboer & Jones, 2005; Prouty & Gunn, 2003). Solano et al. (2002) showed that colonic acid was important to form a tight pellicle under LB conditions, while it was dispensable under ATM conditions. De Rezende et al. (2005) purified another capsular polysaccharide (CP) from extracellular matrix of multiresistant *S. Typhimurium* DT104 which was found to be important for biofilm formation on polystyrene centrifuge tubes and was detected at both 25°C and 37°C. This was comprised principally of glucose and mannose, with galactose as a minor constituent. Malcova et al. (2008) confirmed the importance of this capsular polysaccharide in the biofilm formation capacity of strains unable to produce either curli fimbriae or cellulose. Due to mucoid and brown appearance on Congo Red agar plates, their morphotype was designated as sbam (smooth, brown and mucoid).

However, other capsular polysaccharides can be present in the extracellular biofilm matrix of *Salmonella* strains (de Rezende et al., 2005; Gibson et al., 2006; White et al., 2003), and the exact composition depends upon the environmental conditions in which the biofilms are formed (Prouty & Gunn, 2003). Another component of the EPS matrix of *Salmonella* bile-induced biofilms, the O-antigen (O-ag) capsule, while it was found to be crucial for *S. Typhimurium* and *S. Typhi* biofilm development on gallstones, this was not necessary for adhesion and biofilm formation on glass and plastic (Crawford et al., 2008). The formation of this O-ag capsule was also found to be important for survival during desiccation stress (Gibson et al., 2006). Anriany et al. (2006) highlighted the importance of an integral lipopolysaccharide (LPS), at both the O-antigen and core polysaccharide levels, in the modulation of curli protein and cellulose production, as well as in biofilm formation, thereby adding another potential component to the complex regulatory system which governs multicellular behavior in *S. Typhimurium*. Mireles et al. (2001) observed that for *S. Typhimurium* LT2, all of the LPS mutants examined were able to form a biofilm on polyvinyl chloride (PVC) but none were able to attach to a hydrophilic surface such as glass. Kim & Wei (2009) noticed that a *rfaA* mutant of *S. Typhimurium* DT104, showing an aberrant LPS profile, was impaired in rdar expression, pellicle formation, biofilm forming capability on PVC in meat, poultry and produce broths and biofilm formation on stainless steel and glass.

6. Cell-to-cell communication in *Salmonella* biofilms (quorum sensing)

It has been thoroughly suggested that bacterial cells communicate by releasing and sensing small diffusible signal molecules, in a process commonly known as quorum sensing (QS) (Miller & Bassler, 2001; Smith et al., 2004; Whitehead et al., 2001). Through cell-to-cell signaling mechanisms, bacteria modulate their own behaviour and also respond to signal produced by other species (Ryan & Dow, 2008). QS involves a density-dependent recognition of signaling molecules (autoinducers, AIs), resulting in modulation of gene expression (Bassler, 1999). Gram-negative bacteria primarily use a variety of *N*-acylhomoserine lactones (AHLs) as AI (autoinducer-1, AI-1), while Gram-positive bacteria

use a variety of autoinducing polypeptides (AIPs). AHLs are synthesized and recognized by QS circuits composed of LuxI and LuxR homologues, respectively (Whitehead et al., 2001). Both AHLs and AIPs are highly specific to the species that produce them. A third QS system is proposed to be universal, allowing interspecies communication, and is based on the enzyme LuxS which is in part responsible for the production of a furanone-like compound, called autoinducer-2 (AI-2) (Schauder et al., 2001).

Bacteria use QS communication circuits to regulate a diverse array of physiological activities, such as genetic competence, pathogenicity (virulence), motility, sporulation, bioluminescence and production of antimicrobial substances (Miller & Bassler, 2001). Yet, a growing body of evidence demonstrates that QS also contributes to biofilm formation by many different species (Annous et al., 2009; Davies et al., 1998; Irie & Parsek, 2008; Lazar, 2011). As biofilms typically contain high concentration of cells, autoinducer (AI) activity and QS regulation of gene expression have been proposed as essential components of biofilm physiology (Kjelleberg & Molin, 2002; Parsek & Greenberg, 2005).

To date, three QS systems have been identified in *S. enterica* and are thought to be mainly implicated in the regulation of virulence (SdiA, luxS/AI-2 and AI-3/epinephrine/norepinephrine signaling system) (Boyen et al., 2009; Walters & Sperandio, 2006). Firstly, the LuxR homologue SdiA has been characterized in *Salmonella*, but there does not appear to be a corresponding signal-generating enzyme similar to LuxI in this species (Ahmer et al., 1998). Since *Salmonella* does not possess a luxI homologue, it cannot produce its own AHLs (Ahmer, 2004). However, *Salmonella* SdiA can detect AHLs produced by a variety of bacterial species, leading to the suggestion that SdiA can be used in interspecies communication within a mixed-species community (Michael et al., 2001; Smith & Ahmer 2003). Till now, SdiA is known to activate the expression of the *rck* operon and the *srgE* gene (Ahmer et al., 1998; Smith & Ahmer, 2003). In contrast to the function of SdiA in *E. coli* adherence to HEp-2 epithelial cells and also biofilm formation on polystyrene (Lee et al., 2009; Sharma et al., 2010), no direct link between SdiA and *Salmonella* biofilms has been reported. Interestingly, Chorianopoulos et al. (2010) demonstrated that cell-free culture supernatant (CFS) of the psychrotrophic spoilage bacterium *Hafnei alvei*, containing AHLs among other unknown metabolites, negatively influenced the early stage of biofilm formation by *S. Enteritidis* on stainless steel. Similarly, Dheilly et al. (2010) reported the inhibitory activity of CFS from the marine bacterium *Pseudoalteromonas* sp. strain 3J6 against biofilm formation on glass flow cells by *S. enterica* and other Gram-negative bacteria. Taking into account that *Salmonella* possess SdiA, a receptor of AHLs which may be produced by resident flora on food-contact surfaces (Michael et al., 2001; Smith & Ahmer, 2003; Soares & Ahmer, 2011), the effect of AHLs on biofilm formation by this pathogen in multispecies real food processing environments needs to be further studied.

The second QS system of *Salmonella* uses the LuxS enzyme for the synthesis of AI-2 (Schauder et al., 2001; Soni et al., 2008). The Lsr ABC transporter is known to be involved in the detection and transport of AI-2 into the cell (Taga et al., 2001), while the *rhs* transporter has recently been suggested as an alternative AI-2 uptake system (Jesudhasan et al., 2010). A *S. Typhimurium* *luxS* deletion mutant was impaired in biofilm formation on polystyrene (De Keersmaecker et al., 2005; Jesudhasan et al., 2010). However, this phenotype could not be complemented by extracellular addition of QS signal molecules, suggesting that AI-2 is not the actual signal involved in *Salmonella* biofilm formation (De Keersmaecker et al., 2005). To this direction, Kint et al. (2010) analyzed additional *luxS* mutants for their biofilm phenotype. Interestingly, a *luxS* kanamycin insertion mutant and a partial deletion mutant,

that only lacked the 3' part of the *luxS* coding sequence, were found to be able to form mature wild-type biofilms on polystyrene, despite the fact that these strains were unable to produce AI-2. These authors concluded that a small regulatory RNA molecule, MicA, encoded in the *luxS* adjacent genomic region, rather than LuxS itself, influences *S. Typhimurium* biofilm formation phenotype. On the other hand, Prouty et al. (2002) showed that a *S. Typhimurium luxS* insertion mutant formed scattered biofilm on gallstones with little apparent EPS even after 14 days of incubation. Yoon & Sofos (2008) showed that biofilm formation by *S. Thompson* on stainless steel, under monoculture conditions (72 h at 25°C), was similar between AI-2 positive and negative strains. Altogether, these results demonstrate that the relationship between biofilm formation and the presence of an active LuxS system and AI-2 in *S. enterica* is not clear and further research is needed.

The third QS system of *Salmonella* uses the two component system PreA/B (Bearson & Bearson 2008; Merighi et al., 2006). PreA/B is similar to the *luxS*-dependent two component QseB/QseC of enterohemorrhagic *E. coli*, which has been shown to sense the QS signal AI-3, as well the eukaryotic hormones epinephrine and norepinephrine (Sperandio et al., 2002; Walters & Sperandio, 2006). In *S. Typhimurium*, the histidine sensor kinase QseC, which is able to detect norepinephrine, has been implicated in the regulation of virulence traits, such as motility and *in vivo* competitive fitness in pigs (Bearson & Bearson, 2008). Even though the role of AI-3/epinephrine/norepinephrine signaling system in the formation of biofilm by *Salmonella* is still unknown, given that motility is usually an important biofilm determinant in many bacterial species, it is quite possible that this third QS system may also affect *Salmonella* biofilm formation.

7. Conclusions

Biofilms are commonly defined as communities of microorganisms attached to a surface and producing an extracellular matrix, in which these microorganisms are embedded. Biofilms are very diverse and unique, not just to the microorganism, but to the particular environment in which they are being formed. This makes *in vitro* characterization of biofilms difficult and requires the establishment of laboratory conditions that mimic the natural setting being studied. Pathogenic biofilms have been of considerable interest in the context of food safety and have provoked interest of many research groups. In particular, biofilm formation by *Salmonella* is a serious concern in food industry, since the persistence of this bacterium in biofilms formed on food-contact surfaces may become a constant source of product contamination.

The discovery of bacterial biofilms in medical and industrial ecosystems has created an urgency to identify and characterize factors that are necessary for biofilm development, which may serve as targets for biofilm prevention and treatment. Thus, researchers in the fields of clinical, food, water, and environmental microbiology have begun to investigate microbiological processes from a biofilm perspective. As the pharmaceutical, health-care and food industries embrace this approach, novel strategies for biofilm formation and control will undoubtedly emerge. Particularly challenging is the attempt to understand the complexity of the interactions within a biofilm community, since these interactions between the different species influence the final outcome of this community. Communication between species may include extracellular compounds whose sole role is to influence gene expression, metabolic cooperativity and competition, physical contact, and the production of antimicrobial exoproducts. One or all of these interactions may be

occurring simultaneously. The challenge becomes more intriguing given that microflora on inadequately cleaned and disinfected food processing surfaces is a complex community, contrary to the laboratory studied pure-species biofilms.

Undoubtedly, a clearer understanding of the factors which influence microbial attachment to abiotic surfaces could provide the information necessary to modify processes in food processing environments in order to reduce microbial persistence and therefore reduce the contamination of food products. For instance, the understanding of bacterial attachment to solid surfaces, such as stainless steel, may help in the future development of surfaces with no or reduced attachment, or in developing an effective sanitation programme and thus reducing the potential contamination of processed products by spoilage or/and pathogenic bacteria. Undoubtedly, the ability to recognize how *Salmonella* attach to food-contact surfaces and form biofilms on them is an important area of focus, since a better understanding of this ability may provide valuable ways towards the elimination of this pathogenic bacterium from food processing environments and eventually lead to reduced *Salmonella*-associated human illness.

8. Acknowledgement

Authors would like to acknowledge European Union project ProSafeBeef (ref. Food-CT-2006-36241) within the 6th Framework Programme for the financial support of some of the studies on *Salmonella* biofilms performed on our lab.

9. References

- Ahmer, B.M.M. (2004). Cell-to-cell signalling in *Escherichia coli* and *Salmonella enterica*. *Molecular Microbiology*, Vol.52, No.4, (January 2004), pp. 933-945
- Ahmer, B.M.; van Reeuwijk, J.; Timmers, C.D.; Valentine, P.J. & Heffron, F. (1998). *Salmonella typhimurium* encodes an SdiA homolog, a putative quorum sensor of the LuxR family, that regulates genes on the virulence plasmid. *Journal of Bacteriology*, Vol.180, No.5, (March 1998), pp. 1185-1193
- Annous, B.A.; Fratamico, P.M & Smith, J.L. (2009). Quorum sensing in biofilms: why bacteria behave the way they do. *Journal of Food Science*, Vol.74, No.1., pp. R24-R37
- Anriany, Y.; Sahu, S.N.; Wessels, K.R.; McCann, L.M. & Joseph, S.W. (2006). Alteration of the rugose phenotype in *waaG* and *ddhC* mutants of *Salmonella enterica* serovar Typhimurium DT104 is associated with inverse production of curli and cellulose. *Applied and Environmental Microbiology*, Vol.72, No.7, (July 2006), pp. 5002-5012
- Arnold, J.W. & Yates, I.E. (2009). Interventions for control of *Salmonella*: clearance of microbial growth from rubber picker fingers. *Poultry Science*, Vol.88, No.6, (June 2009), pp. 1292-1298
- Asséré, A.; Oulahal, N. & Carpentier, B. (2008). Comparative evaluation of methods for counting surviving biofilm cells adhering to a polyvinyl chloride surface exposed to chlorine or drying. *Journal of Applied Microbiology*, Vol.104, No.6, (June 2008), pp. 1692-1702
- Austin, J.W. & Bergeron, G. (1995). Development of bacterial biofilms in dairy processing lines. *Journal of Dairy Research*, Vol.62, No.3, (August 1995), pp. 509-519
- Austin, J.W.; Sanders, G.; Kay, W.W. & Collinson, S.K. (1998). Thin aggregative fimbriae enhance *Salmonella enteritidis* biofilm formation. *FEMS Microbiology Letters*, Vol.162, No.2, (May 1998), pp. 295-301

- Bagge-Ravn, D.; Ng, Y.; Hjelm, M.; Christiansen, J.N.; Johansen, C. & Gram, L. (2003). The microbial ecology of processing equipment in different fish industries - analysis of the microflora during processing and following cleaning and disinfection. *International Journal of Food Microbiology*, Vol.87, No.3, (November 2003), pp. 239-250
- Barnes, L.M.; Lo, M.F.; Adams, M.R. & Chamberlain, A.H. (1999). Effect of milk proteins on adhesion of bacteria to stainless steel surfaces. *Applied and Environmental Microbiology*, Vol.65, No.10, (October 1999), pp. 4543-4548
- Bassler, B.L. (1999). How bacteria talk to each other: regulation of gene expression by quorum sensing. *Current Opinion in Microbiology*, Vol.2, No.6, (December 1999), pp. 582-587
- Bearson, B.L. & Bearson, S.M. (2008). The role of the QseC quorum-sensing sensor kinase in colonization and norepinephrine-enhanced motility of *Salmonella enterica* serovar Typhimurium. *Microbial Pathogenesis*, Vol.44, No.4, (April 2008), pp. 271-278
- Beloin, C.; Valle, J.; Latour-Lambert, P.; Faure, P.; Kzreminski, M.; Balestrino, D.; Haagenen, J.A.; Molin, S.; Prensier, G.; Arbeille, B. & Ghigo, J.M. (2004). Global impact of mature biofilm lifestyle on *Escherichia coli* K-12 gene expression. *Molecular Microbiology*, Vol.51, No.3, (February 2004), pp. 659-674
- Bernbom, N.; Ng, Y.Y.; Jørgensen, R.L.; Arpanaei, A.; Meyer, R.L.; Kingshott, P.; Vejborg, R.M.; Klemm, P. & Gram, L. (2009). Adhesion of food-borne bacteria to stainless steel is reduced by food conditioning films. *Journal of Applied Microbiology*, Vol.106, No.4, (April 2009), pp. 1268-1279
- Blackman, I.C. & Frank, J.F. (1996). Growth of *Listeria monocytogenes* as a biofilm on various food-processing surfaces. *Journal of Food Protection*, Vol.59, No.8, (August 1996), pp. 827-831
- Boyen, F.; Eeckhaut, V.; Van Immerseel, F.; Pasmans, F.; Ducatelle, R. & Haesebrouck, F. (2009). Quorum sensing in veterinary pathogens: mechanisms, clinical importance and future perspectives. *Veterinary Microbiology*, Vol.135, No.3-4, (March 2009), pp. 187-195
- Branda, S.S.; Vik, S.; Friedman, L. & Kolter, R. (2005). Biofilms: the matrix revisited. *Trends in Microbiology*, Vol.13, No.1, (January 2005), pp. 20-26
- Briandet, R.; Meylheuc, T.; Maher, C. & Bellon-Fontaine, M.N. (1999). *Listeria monocytogenes* Scott A: cell surface charge, hydrophobicity, and electron donor and acceptor characteristics under different environmental growth conditions. *Applied and Environmental Microbiology*, Vol.65, No.12, (December 1999), pp. 5328-5333
- Brooks, J.D. & Flint, S.H. (2008). Biofilms in the food industry: problems and potential solutions. *International Journal of Food Science and Technology*, Vol.43, (August 2008), pp. 2163-2176
- Burmølle, M.; Webb, J.S.; Rao, D.; Hansen, L.H.; Sørensen, S.J. & Kjelleberg, S. (2006). Enhanced biofilm formation and increased resistance to antimicrobial agents and bacterial invasion are caused by synergistic interactions in multispecies biofilms. *Applied and Environmental Microbiology*, Vol.72, No.6, (June 2006), pp. 3916-3923
- Carpentier, B. & Cerf, O. (1993). Biofilms and their consequences, with particular reference to hygiene in the food industry. *Journal of Applied Bacteriology*, Vol.75, (June 1993), pp. 499-451
- Chia, T.W.R.; Goulter, R.M.; McMeekin, T.; Dykes, G.A. & Fegan, N. (2009). Attachment of different *Salmonella* serovars to materials commonly used in a poultry processing plant. *Food Microbiology*, Vol.26, (May 2009), pp. 853-859
- Chmielewski, R.A.N. & Frank, J.F. (2003). Biofilm formation and control in food processing facilities. *Comprehensive Reviews in Food Science and Food Safety*, Vol.2, pp. 22-32

- Chorianopoulos, N.G.; Giaouris, E.D.; Kourkoutas, Y. & Nychas, G.-J. (2010). Inhibition of the early stage of *Salmonella enterica* serovar Enteritidis biofilm development on stainless steel by cell-free supernatant of a *Hafnia alvei* culture. *Applied and Environmental Microbiology*, Vol.76, No.6, (January 2010), pp. 2018-2022
- Chorianopoulos, N.G.; Giaouris, E.D.; Skandamis, P.N.; Haroutounian, S.A. & Nychas, G.-J. (2008). Disinfectant test against monoculture and mixed-culture biofilms composed of technological, spoilage and pathogenic bacteria: bactericidal effect of essential oil and hydrosol of *Satureja thymbra* and comparison with standard acid-base sanitizers. *Journal of Applied Microbiology*, Vol.104, No.6, (November 2007), pp. 1586-1596
- Chorianopoulos, N.G.; Tsoukleris, D.S.; Panagou, E.Z.; Falaras, P. & Nychas, G.J. (2011). Use of titanium dioxide (TiO₂) photocatalysts as alternative means for *Listeria monocytogenes* biofilm disinfection in food processing. *Food Microbiology*, Vol.28, No.1, (February 2011), pp. 164-170
- Collinson, S.K.; Clouthier, S.C.; Doran, J.L.; Banser, P.A. & Kay, W.W. (1996). *Salmonella enteritidis* agfBAC operon encoding thin, aggregative fimbriae. *Journal of Bacteriology*, Vol.178, No.3, (February 1996), pp. 662-667
- Collinson, S.K.; Emödy, L.; Müller, K.H.; Trust, T.J. & Kay, W.W. (1991). Purification and characterization of thin, aggregative fimbriae from *Salmonella enteritidis*. *Journal of Bacteriology*, Vol.173, No.15, (August 1991), pp. 4773-4781
- Cos, P.; Toté, K.; Horemans, T. & Maes, L. (2010). Biofilms: an extra hurdle for effective antimicrobial therapy. *Current Pharmaceutical Design*, Vol.16, No.20, pp. 2279-2295
- Costerton, J.W.; Cheng, K.J.; Geesey, G.G.; Ladd, T.I.; Nickel, J.C.; Dasgupta, M. & Marrie, T.J. (1987). Bacterial biofilms in nature and disease. *Annual Review of Microbiology*, Vol.41, pp. 435-464
- Costerton, J.W.; Lewandowski, Z.; Caldwell, D.E.; Korber, D.R. & Lappin-Scott, H.M. (1995). Microbial biofilms. *Annual Review of Microbiology*, Vol.49, pp. 711-745
- Crawford, R.W.; Gibson, D.L.; Kay, W.W. & Gunn, J.S. (2008). Identification of a bile-induced exopolysaccharide required for *Salmonella* biofilm formation on gallstone surfaces. *Infection and Immunity*, Vol.76, No.11, (November 2008), pp. 5341-5349
- Davey, M.E. & O'Toole, G.A. (2000). Microbial biofilms: from ecology to molecular genetics. *Microbiology and Molecular Biology Reviews*, Vol.64, No.4, pp. 847-867
- Davies, D.G.; Parsek, M.R.; Pearson, J.P.; Iglewski, B.H.; Costerton, J.W. & Greenberg, E.P. (1998). The involvement of cell-to-cell signals in the development of a bacterial biofilm. *Science*, Vol.280, No.5361, (April 1998), pp. 295-298
- De Keersmaecker, S.C.; Varszegi, C.; van Boxel, N.; Habel, L.W.; Metzger, K.; Daniels, R.; Marchal, K.; De Vos, D. & Vanderleyden, J. (2005). Chemical synthesis of (S)-4,5-dihydroxy-2,3-pentanedione, a bacterial signal molecule precursor, and validation of its activity in *Salmonella typhimurium*. *Journal of Biological Chemistry*, Vol.280, No.20, (May 2005), pp. 19563-19568
- De Rezende, C.E.; Anriany, Y.; Carr, L.E.; Joseph, S.W. & Weiner, R.M. (2005). Capsular polysaccharide surrounds smooth and rugose types of *Salmonella enterica* serovar Typhimurium DT104. *Applied and Environmental Microbiology*, Vol.71, No.11, (November 2005), pp. 7345-7351
- Dheilly, A.; Soum-Soutéra, E.; Klein, G.L.; Bazire, A.; Compère, C.; Haras, D. & Dufour, A. (2010). Antibiofilm activity of the marine bacterium *Pseudoalteromonas* sp. strain 3J6. *Applied and Environmental Microbiology*, Vol.76, No.11, (June 2010), pp. 3452-3461
- Di Bonaventura, G.; Piccolomini, R.; Paludi, D.; D'Orio, V.; Vergara, A.; Conter, M. & Ianieri, A. (2008). Influence of temperature on biofilm formation by *Listeria monocytogenes*

- on various food-contact surfaces: relationship with motility and cell surface hydrophobicity. *Journal of Applied Microbiology*, Vol.104, No.6, (June 2008), pp. 1552-1561
- Donlan, R.M. (2002). Biofilms: microbial life on surfaces. *Emerging and Infectious Diseases*, Vol.8, No.9, pp. 881-890
- Donlan, R.M. & Costerton, J.W. (2002). Biofilms: survival mechanisms of clinically relevant microorganisms. *Clinical Microbiology Reviews*, Vol.15, No.2, pp. 167-193
- Duguid, J.P.; Anderson, E.S. & Campbell, I. (1966). Fimbriae and adhesive properties in *Salmonellae*. *Journal of Pathology and Bacteriology*, Vol.92, No.1, (July 1966), pp. 107-138
- EFSA-ECDC. (2011). The European Union summary report on trends and sources of zoonoses, zoonotic agents and food-borne outbreaks in 2009. *The EFSA Journal*, Vol.9, No.3:2090
- Flemming, H.C. & Wingender, J. (2010). The biofilm matrix. *Nature Reviews. Microbiology*, Vol.8, No.9, (August 2010), pp. 623-633.
- Ganesh Kumar, S. & Anand, S.K. (1998). Significance of microbial biofilms in food industry: a review. *International Journal of Food Microbiology*, Vol.42, (April 1998), pp. 9-27
- García, B.; Latasa, C.; Solano, C.; García-del Portillo, F.; Gamazo, C. & Lasa, I. (2004). Role of the GGDEF protein family in *Salmonella* cellulose biosynthesis and biofilm formation. *Molecular Microbiology*, Vol.54, No.1, (October 2004), pp. 264-277
- Gawande, P.V. & Bhagwat, A.A. (2002). Inoculation onto solid surfaces protects *Salmonella* spp. during acid challenge: a model study using polyethersulfone membranes. *Applied and Environmental Microbiology*, Vol.68, No.1, (October 2001), pp. 86-92
- Gerstel, U. & Römling, U. (2001). Oxygen tension and nutrient starvation are major signals that regulate *agfD* promoter activity and expression of the multicellular morphotype in *Salmonella typhimurium*. *Environmental Microbiology*, Vol.3, No.10, (October 2001), pp. 638-648
- Gerstel, U. & Römling, U. (2003). The *csgD* promoter, a control unit for biofilm formation in *Salmonella typhimurium*. *Research in Microbiology*, Vol.154, No.10, (December 2003), pp. 659-667
- Giaouris, E.; Chapot-Chartier, M.P. & Briandet, R. (2009). Surface physicochemical analysis of natural *Lactococcus lactis* strains reveals the existence of hydrophobic and low charged strains with altered adhesive properties. *International Journal of Food Microbiology*, Vol.131, No.1, (April 2009), pp. 2-9
- Giaouris, E.; Chorianopoulos, N. & Nychas, G.-J.E. (2005). Effect of temperature, pH, and water activity on biofilm formation by *Salmonella enterica* Enteritidis PT4 on stainless steel surfaces as indicated by the bead vortexing method and conductance measurements. *Journal of Food Protection*, Vol.68, No.10, (May 2005), pp. 2149-2154
- Giaouris, E.D. & Nychas, G.-J.E. (2006). The adherence of *Salmonella* Enteritidis PT4 to stainless steel: the importance of the air-liquid interface and nutrient availability. *Food Microbiology*, Vol.23, (February 2006), pp. 747-752
- Gibson, D.L.; White, A.P.; Snyder, S.D.; Martin, S.; Heiss, C.; Azadi, P.; Surette, M. & Kay, W.W. (2006). *Salmonella* produces an O-antigen capsule regulated by AgfD and important for environmental persistence. *Journal of Bacteriology*, Vol.188, No.22, (November 2006), pp. 7722-7730
- Gilbert, P.; Allison, D.G. & McBain, A.J. (2002). Biofilms *in vitro* and *in vivo*: do singular mechanisms imply cross-resistance? *Journal of Applied Microbiology*, Vol.92, pp. 98S-110S
- Gilbert, P.; Evans, D.J. & Brown, M.R. (1993). Formation and dispersal of bacterial biofilms *in vivo* and *in situ*. *Journal of Applied Bacteriology*, Vol.74, Suppl.67S-78S

- Girenavar, B.; Cepeda, M.L.; Soni, K.A.; Vikram, A.; Jesudhasan, P.; Jayaprakasha, G.K.; Pillai, S.D. & Patil, B.S. (2008). Grapefruit juice and its furcoumarins inhibits autoinducer signaling and biofilm formation in bacteria. *International Journal of Food Microbiology*, Vol.125, (March 2008), pp. 204-208
- Goller, C.C. & Romeo, T. (2008). Environmental influences on biofilm development. *Current Topics in Microbiology and Immunology*, Vol.322, pp. 37-66
- Gounadaki, A.S.; Skandamis, P.N.; Drosinos, E.H. & Nychas, G.J. (2008). Microbial ecology of food contact surfaces and products of small-scale facilities producing traditional sausages. *Food Microbiology*, Vol.25, No.2, (April 2008), pp. 313-323
- Gunduz, G.T. & Tuncel, G. (2006). Biofilm formation in an ice cream plant. *Antonie van Leeuwenhoek*, Vol.89, pp. 329-336
- Habimana, O.; Heir, E.; Langsrud, S.; Asli, A.W. & Møretrø, T. (2010a). Enhanced surface colonization by *Escherichia coli* O157:H7 in biofilms formed by an *Acinetobacter calcoaceticus* isolate from meat-processing environments. *Applied and Environmental Microbiology*, Vol.76, No.13, (July 2010), pp. 4557-4559
- Habimana, O.; Møretrø, T.; Langsrud, S.; Vestby, L.K.; Nesse, L.L. & Heir, E. (2010b). Micro ecosystems from feed industry surfaces: a survival and biofilm study of *Salmonella* versus host resident flora strains. *BMC Veterinary Research*, Vol.6, No. 48
- Hall-Stoodley, L.; Costerton, J.W. & Stoodley, P. (2004). Bacterial biofilms: from the natural environment to infectious diseases. *Nature Reviews. Microbiology*, Vol.2, (February 2004), pp. 95-108
- Hall-Stoodley, L. & Stoodley, P. (2002). Developmental regulation of microbial biofilms. *Current Opinion in Biotechnology*, Vol.13, No.3, (June 2002), pp. 228-233
- Hall-Stoodley, L. & Stoodley, P. (2005). Biofilm formation and dispersal and the transmission of human pathogens. *Trends in Microbiology*, Vol.13, No.1, (January 2005), pp. 7-10
- Hamilton, S.; Bongaerts, R.J.; Mulholland, F.; Cochrane, B.; Porter, J.; Lucchini, S.; Lappin-Scott, H.M. & Hinton, J.C. (2009). The transcriptional programme of *Salmonella enterica* serovar Typhimurium reveals a key role for tryptophan metabolism in biofilms. *BMC Genomics*, Vol.10, (December 2009), 599
- Hood, S.K. & Zottola, E.A. (1997a). Adherence to stainless steel by foodborne microorganisms during growth in model food systems. *International Journal of Food Microbiology*, Vol.37, (June 1997), pp. 145-153
- Hood, S.K. & Zottola, E.A. (1997b). Growth media and surface conditioning influence the adherence of *Pseudomonas fragi*, *Salmonella typhimurium*, and *Listeria monocytogenes* cells to stainless steel. *Journal of Food Protection*, Vol.60, No.9, (January 1997), pp. 1034-1037
- Humphrey, T.J.; Slater, E.; McAlpine, K.; Rowbury, R.J. & Gilbert, R.J. (1995). *Salmonella* Enteritidis Phage Type 4 isolates more tolerant of heat, acid, or hydrogen peroxide also survive longer on surfaces. *Applied and Environmental Microbiology*, Vol.61, No.8, (May 1995), pp. 3161-3164
- Iibuchi, R.; Hara-Kudo, Y.; Hasegawa, A. & Kumagai, S. (2010). Survival of *Salmonella* on a polypropylene surface under dry conditions in relation to biofilm-formation capability. *Journal of Food Protection*, Vol.73, No.8, (December 2009), pp. 1506-1510
- Irie, Y. & Parsek, M.R. (2008). Quorum sensing and microbial biofilms. *Current Topics in Microbiology and Immunology*, Vol.322, pp. 67-84
- Jain, S. & Chen, J. (2007). Attachment and biofilm formation by various serotypes of *Salmonella* as influenced by cellulose production and thin aggregative fimbriae biosynthesis. *Journal of Food Protection*, Vol.70, No.11, (November 2007), pp. 2473-2479

- James, G.A.; Beaudette, L. & Costerton, J.W. (1995). Interspecies bacterial interactions in biofilms. *Journal of Industrial Microbiology*, Vol.15, pp. 257-262
- Jefferson, K.K. (2004). What drives bacteria to produce a biofilm? *FEMS Microbiology Letters*, Vol.236, (June 2004), pp. 163-173
- Jesudhasan, P.R.; Cepeda, M.L.; Widmer, K.; Dowd, S.E.; Soni, K.A.; Hume, M.E.; Zhu, J. & Pillai, S.D. (2010). Transcriptome analysis of genes controlled by *luxS*/autoinducer-2 in *Salmonella enterica* serovar Typhimurium. *Foodborne Pathogens and Disease*, Vol.7, No.4, (April 2010), pp. 399-410
- Jones, K. & Bradshaw, S.B. (1997). Synergism in biofilm formation between *Salmonella enteritidis* and a nitrogen-fixing strain of *Klebsiella pneumoniae*. *Journal of Applied Microbiology*, Vol.82, (October 1996), pp. 663-668
- Joseph, B.; Otta, S.K.; Karunasagar, I. & Karunasagar, I. (2001). Biofilm formation by *salmonella* spp. on food contact surfaces and their sensitivity to sanitizers. *International Journal of Food Microbiology*, Vol.64, No.3, (March 2001), pp. 367-372
- Joshua, G.W.; Guthrie-Irons, C.; Karlyshev, A.V. & Wren, B.W. (2006). Biofilm formation in *Campylobacter jejuni*. *Microbiology*, Vol.152(Pt2), (February 2006), pp. 387-396
- Kim, T.J.; Young, B.M. & Young, G.M. (2008). Effect of flagellar mutations on *Yersinia enterocolitica* biofilm formation. *Applied and Environmental Microbiology*, Vol.74, No.17, (September 2008), pp. 5466-5574
- Kim, S.-H. & Wei, C.-I. (2007). Biofilm formation by multidrug-resistant *Salmonella enterica* serotype Typhimurium Phage Type DT104 and other pathogens. *Journal of Food Protection*, Vol.70, No.1, (July 2006), pp. 22-29
- Kim, S.-H. & Wei, C.-I. (2009). Molecular characterization of biofilm formation and attachment of *Salmonella enterica* serovar Typhimurium DT104 on food contact surfaces. *Journal of Food Protection*, Vol.72, No.9, (September 2009), pp. 1841-1847
- Kint, G.; De Coster, D.; Marchal, K.; Vanderleyden, J. & De Keersmaecker, S.C. (2010). The small regulatory RNA molecule MicA is involved in *Salmonella enterica* serovar Typhimurium biofilm formation. *BMC Microbiology*, Vol.10, (November), 276
- Kjelleberg, S. & Molin, S. (2002). Is there a role for quorum sensing signals in bacterial biofilms? *Current Opinion in Microbiology*, Vol.5, No.3, (June 2002), pp. 254-258
- Klausen, M.; Gjermansen, M.; Kreft, J.U. & Tolker-Nielsen, T. (2006). Dynamics of development and dispersal in sessile microbial communities: examples from *Pseudomonas aeruginosa* and *Pseudomonas putida* model biofilms. *FEMS Microbiology Letters*, Vol.261, No.1, (August 2006), pp. 1-11
- Kolter, R. & Greenberg, E.P. (2006). The superficial life of microbes. *Nature*, Vol.441, No.7091, (May 2006), pp. 300-302
- Korber, D.R.; Choi, A.; Wolfaardt, G.M.; Ingham, S.C. & Caldwell, D.E. (1997). Substratum topography influences susceptibility of *Salmonella enteritidis* biofilms to trisodium phosphate. *Applied and Environmental Microbiology*, Vol.63, No.9, (September 1997), pp. 3352-3358
- Kuchma, S.L. & O'Toole, G.A. (2000). Surface-induced and biofilm-induced changes in gene expression. *Current Opinion in Biotechnology*, Vol.11, No.5, (October 2000), pp. 429-433
- Lasa, I. (2006). Towards the identification of the common features of bacterial biofilm development. *International Microbiology*, Vol.9, (January 2006), pp. 21-28
- Latasa, C.; Roux, A.; Toledo-Arana, A.; Ghigo, J.M.; Gamazo, C.; Penadés, J.R. & Lasa, I. (2005). BapA, a large secreted protein required for biofilm formation and host colonization of *Salmonella enterica* serovar Enteritidis. *Molecular Microbiology*, Vol.58, No.5, (December 2005), pp. 1322-1339

- Lazar, V. (2011). Quorum sensing in biofilms - How to destroy the bacterial citadels or their cohesion/power? *Anaerobe*, in press
- Lazazzera, B.A. (2005). Lessons from DNA microarray analysis: the gene expression profile of biofilms. *Current Opinion in Microbiology*, Vol.8, No.2, (April 2005), pp. 222-227
- Ledeboer, N.A. & Jones, B.D. (2005). Exopolysaccharide sugars contribute to biofilm formation by *Salmonella enterica* serovar Typhimurium on HEp-2 cells and chicken intestinal epithelium. *Journal of Bacteriology*, Vol.187, No.9, (May 2005), pp. 3214-3226
- Lee, J.; Maeda, T.; Hong, S.H. & Wood, T.K. (2009). Reconfiguring the quorum-sensing regulator SdiA of *Escherichia coli* to control biofilm formation via indole and N-acylhomoserine lactones. *Applied and Environmental Microbiology*, Vol.75, No.6, (March 2009), pp. 1703-1716
- Lewis, K. (2001). Riddle of biofilm resistance. *Antimicrobial Agents and Chemotherapy*, Vol.45, No.4, (April 2001), pp. 999-1007
- Lindsay, D. & von Holy, A. (2006). What food safety professionals should know about bacterial biofilms. *British Food Journal*, Vol.108, No.1, pp. 27-37
- Litrup, E.; Torpdahl, M.; Malorny, B.; Huehn, S.; Christensen, H. & Nielsen, E.M. (2010). Association between phylogeny, virulence potential and serovars of *Salmonella enterica*. *Infection, Genetics and Evolution*, Vol.10, No. 7, (July 2010), pp. 1132-1139
- López, D.; Vlamakis, H. & Kolter, R. (2010). Biofilms. *Cold Spring Harbor Perspectives in Biology*, Vol.2, No.7, (June 2010), a000398
- Lundén, J.M.; Miettinen, M.K.; Autio, T.J. & Korkeala, H.J. (2000). Persistent *Listeria monocytogenes* strains show enhanced adherence to food contact surface after short contact times. *Journal of Food Protection*, Vol.63, No.9, (September 2000), pp. 1204-1207
- Mah, T.-F.C. & O'Toole, G.A. (2001). Mechanisms of biofilm resistance to antimicrobial agents. *Trends in Microbiology*, Vol.9, No.1, (January 2001), pp. 34-39
- Malcova, M.; Hradecka, H.; Karpiskova, R. & Rychlik, I. (2008). Biofilm formation in field strains of *Salmonella enterica* serovar Typhimurium: identification of a new colony morphology type and the role of SGI1 in biofilm formation. *Veterinary Microbiology*, Vol.129, No.3-4, (June 2008), pp. 360-366
- Marin, C.; Hernandez, A. & Lainez, M. (2009). Biofilm development capacity of *Salmonella* strains isolated in poultry risk factors and their resistance against disinfectants. *Poultry Science*, Vol.88, (October 2008), pp. 424-431
- Marsh, P.D. (2005). Dental plaque: biological significance of a biofilm and community lifestyle. *Journal of Clinical Periodontology*, Vol.32, Suppl.6, pp. 7-15.
- Merighi, M.; Carroll-Portillo, A.; Septer, A.N.; Bhatiya, A. & Gunn, J.S. (2006). Role of *Salmonella enterica* serovar Typhimurium two-component system PreA/PreB in modulating PmrA-regulated gene transcription. *Journal of Bacteriology*, Vol.188, No.1, (January 2006), pp. 141-149
- Michael, B.; Smith, J.N.; Swift, S.; Heffron, F. & Ahmer, B.M. (2001). SdiA of *Salmonella enterica* is a LuxR homolog that detects mixed microbial communities. *Journal of Bacteriology*, Vol.183, No.19, (October 2001), pp. 5733-5742
- Miller, M.B. & Bassler, B.L. (2001). Quorum sensing in bacteria. *Annual Review of Microbiology*, Vol.55, pp. 165-199
- Mireles, J.R. 2nd; Toguchi, A. & Harshey, R.M. (2001). *Salmonella enterica* serovar typhimurium swarming mutants with altered biofilm-forming abilities: surfactin inhibits biofilm formation. *Journal of Bacteriology*, Vol.183, No.20, (October 2001), pp. 5848-5854


- Molin, S. & Tolker-Nielsen, T. (2003). Gene transfer occurs with enhanced efficiency in biofilms and induces enhanced stabilisation of the biofilm structure. *Current Opinion in Biotechnology*, Vol.14, pp. 255-261
- Moons, P.; Michiels, C.W. & Aertsen, A. (2009). Bacterial interactions in biofilms. *Critical Reviews in Microbiology*, Vol.35, No.3, (August 2009), pp. 157-168
- Møretrø, T.; Vestby, L.K.; Nesse, L.L.; Storheim, S.E.; Kotlarz, K. & Langsrud, S. (2009). Evaluation of efficacy of disinfectants against *Salmonella* from the feed industry. *Journal of Applied Microbiology*, Vol.106, No.3., (September 2008), pp. 1005-1012
- Morild, R.K.; Olsen, J.E. & Aabo, S. (2011). Change in attachment of *Salmonella* Typhimurium, *Yersinia enterocolitica*, and *Listeria monocytogenes* to pork skin and muscle after hot water and lactic acid decontamination. *International Journal of Food Microbiology*, Vol.145, No.1, (January 2011), pp. 353-358
- Nadell, C.D.; Xavier, J.B.; Levin, S.A. & Foster, K.R. (2008). The evolution of quorum sensing in bacterial biofilms. *PLoS Biology*, Vol.6, No.1, (January 2008), e14
- Norwood, D.E. & Gilmour, A. (2001). The differential adherence capabilities of two *Listeria monocytogenes* strains in monoculture and multispecies biofilms as a function of temperature. *Letters in Applied Microbiology*, Vol.33, No.4, (October 2001), pp. 320-324
- Ngwai, Y.B.; Adachi, Y.; Ogawa, Y. & Hara, H. (2006). Characterization of biofilm-forming abilities of antibiotic-resistant *Salmonella typhimurium* DT104 on hydrophobic abiotic surfaces. *Journal of Microbiology, Immunology and Infection*, Vol.39, (December 2005), pp. 278-291
- Oliveira, K.; Oliveira, T.; Teixeira, P.; Azeredo, J.; Henriques, M. & Oliveira, R. (2006). Comparison of the adhesion ability of different *Salmonella* Enteritidis serotypes to materials used in kitchens. *Journal of Food Protection*, Vol.69, No.10, (May 2006), pp. 2352-2356
- O'Toole, G.; Kaplan, H.B. & Kolter, R. (2000). Biofilm formation as microbial development. *Annual Review of Microbiology*, Vol.54, pp. 49-79
- Palmer, J.; Flint, S. & Brooks, J. (2007). Bacterial cell attachment, the beginning of a biofilm. *Journal of Industrial Microbiology and Biotechnology*, Vol.34, (July 2007), pp. 577-578
- Parsek, M.R. & Greenberg, E.P. (2005). Sociomicrobiology: the connections between quorum sensing and biofilms. *Trends in Microbiology*, Vol.13, No.1, (January 2005), pp. 27-33
- Poimenidou, S.; Belessi, C.A.; Giaouris, E.D.; Gounadaki, A.S.; Nychas, G.J. & Skandamis, P.N. (2009). *Listeria monocytogenes* attachment to and detachment from stainless steel surfaces in a simulated dairy processing environment. *Applied and Environmental Microbiology*, Vol.75, No.22, (November 2009), pp. 7182-7188
- Prouty, A.M. & Gunn, J.S. (2003). Comparative analysis of *Salmonella enterica* serovar Typhimurium biofilm formation on gallstones and on glass. *Infection and Immunity*, Vol.71, No.12, (December 2003), pp. 7154-7158
- Prouty, A.M.; Schwesinger, W.H. & Gunn, J.S. (2002). Biofilm formation and interaction with the surfaces of gallstones by *Salmonella* spp.. *Infection and Immunity*, Vol.70, No.5, (January 2002), pp. 2640-2649
- Remis, J.P.; Costerton, J.W. & Auer, M. (2010). Biofilms: structures that may facilitate cell-cell interactions. *ISME Journal*, Vol.4, No.9, (July 2010), pp. 1085-1087
- Rickard, A.H.; Gilbert, P.; High, N.J.; Kolenbrander, P.E. & Handley, P.S. (2003). Bacterial coaggregation: an internal process in the development of multi-species biofilms. *Trends in Microbiology*, Vol.11, No.2, (February 2003), pp. 94-100

- Rivas, L.; Dykes, G.A. & Fegan, N. (2006). Attachment of shiga toxicogenic *Escherichia coli* to beef muscle and adipose tissue. *Journal of Food Protection*, Vol.69, No.5, (May 2006), pp. 999-1006
- Rodrigues, D.; Teixeira, P.; Oliveira, R. & Azeredo, J. (2011). *Salmonella enterica* Enteritidis biofilm formation and viability on regular and triclosan-impregnated bench cover materials. *Journal of Food Protection*, Vol.74, No.1, (July 2010), pp. 32-37
- Römling, U. (2005). Characterization of the rdar morphotype, a multicellular behaviour in Enterobacteriaceae. *Cellular and Molecular Life Sciences*, Vol.62, No.11, (June 2005), pp. 1234-1246
- Römling, U.; Bian, Z.; Hammar, M.; Sierralta, W.D. & Normark, S. (1998a). Curli fibers are highly conserved between *Salmonella typhimurium* and *Escherichia coli* with respect to operon structure and regulation. *Journal of Bacteriology*, Vol.180, No.3, (February 1998), pp. 722-731
- Römling, U.; Bokranz, W.; Rabsch, W.; Zogaj, X.; Nimtz, M. & Tschäpe, H. (2003). Occurrence and regulation of the multicellular morphotype in *Salmonella* serovars important in human disease. *International Journal of Medical Microbiology*, Vol.293, No.4, (August 2003), pp. 273-285
- Römling, U. & Rohde, M. (1999). Flagella modulate the multicellular behavior of *Salmonella typhimurium* on the community level. *FEMS Microbiology Letters*, Vol.180, No.1, (November 1999), pp. 91-102
- Römling, U.; Rohde, M.; Olsén, A.; Normark, S. & Reinköster, J. (2000). AgfD, the checkpoint of multicellular and aggregative behaviour in *Salmonella typhimurium* regulates at least two independent pathways. *Molecular Microbiology*, Vol.36, No.1, (April 2000), pp. 10-23
- Römling, U.; Sierralta, W.D.; Eriksson, K. & Normark, S. (1998b). Multicellular and aggregative behaviour of *Salmonella typhimurium* strains is controlled by mutations in the *agfD* promoter. *Molecular Microbiology*, Vol.28, No.2, (April 1998), pp. 249-264
- Ryan, R.P. & Dow, J.M. (2008). Diffusible signals and interspecies communication in bacteria. *Microbiology*, Vol.154(Pt 7), (July 2008), pp. 1845-1858
- Sauer, K. (2003). The genomics and proteomics of biofilm formation. *Genome Biology*, Vol.4, No.6, (May 2003), 219
- Schauder, S.; Shokat, K.; Surette, M.G. & Bassler, B.L. (2001). The LuxS family of bacterial autoinducers: biosynthesis of a novel quorum-sensing signal molecule. *Molecular Microbiology*, Vol.41, No.2, (July 2001), pp. 463-476
- Scher, K., Romling, U. & Yaron, S. (2005). Effect of heat, acidification, and chlorination on *Salmonella enterica* serovar Typhimurium cells in a biofilm formed at the air-liquid interface. *Applied and Environmental Microbiology*, Vol.71, No.3, (September 2004), pp. 1163-1168
- Sha, Q.; Gunathilake, A.; Forstner, M.R.J. & Hahn, D. (2011). Temporal analyses of the distribution and diversity of *Salmonella* in natural biofilms. *Systematic and Applied Microbiology*, Vol.34, No.5, (July 2011), pp. 353-359
- Sharma, M. & Anand, S.K. (2002). Biofilms evaluation as an essential component of HACCP for food/dairy processing industry – a case. *Food Control*, Vol.13, (July 2001), pp. 469-477
- Sharma, V.K.; Bearson, S.M. & Bearson, B.L. (2010). Evaluation of the effects of *sdiA*, a *luxR* homologue, on adherence and motility of *Escherichia coli* O157:H7. *Microbiology*, Vol.156(Pt5), (May 2010), pp. 1303-1312

- Shemesh, M.; Tam, A. & Steinberg, D. (2007). Differential gene expression profiling of *Streptococcus mutants* cultured under biofilm and planktonic conditions. *Microbiology*, Vol.153, (May 2007), pp. 1307-1317
- Shi, X. & Zhu, X. (2009). Biofilm formation and food safety in food industries. *Trends in Food Science and Technology*, Vol.20, No.9, (September 2009), pp. 407-413
- Sinde, E. & Carballo, J. (2000). Attachment of *Salmonella* spp. and *Listeria monocytogenes* to stainless steel, rubber and polytetrafluorethylene: the influence of free energy and the effect of commercial sanitizers. *Food Microbiology*, Vol.17, pp. 439-447
- Skandamis, P.N.; Stopforth, J.D.; Ashton, L.V.; Geornaras, I.; Kendall, P.A. & Sofos, J.N. (2009). *Escherichia coli* O157:H7 survival, biofilm formation and acid tolerance under simulated slaughter plant moist and dry conditions. *Food Microbiology*, Vol.26, No.1, (February 2009), pp. 112-119
- Smith, J.N. & Ahmer, B.M. (2003). Detection of other microbial species by *Salmonella*: expression of the SdiA regulon. *Journal of Bacteriology*, Vol.185, No.4, (February 2003), pp. 1357-1366
- Smith, J.L.; Fratamico, P.M. & Novak, J.S. (2004). Quorum sensing: a primer for food microbiologists. *Journal of Food Protection*, Vol.67, No.5, (May 2004), pp. 1053-1070
- Soares, J.A. & Ahmer, B.M. (2011). Detection of acyl-homoserine lactones by *Escherichia* and *Salmonella*. *Current Opinion in Microbiology*, Vol.14, No.2, (April 2011), pp. 188-193
- Solano, C.; García, B.; Valle, J.; Berasain, C.; Ghigo, J.M.; Gamazo, C. & Lasa, I. (2002). Genetic analysis of *Salmonella enteritidis* biofilm formation: critical role of cellulose. *Molecular Microbiology*, Vol.43, No.3, (February 2002), pp. 793-808
- Solano, C.; Sesma, B.; Alvarez, M.; Humphrey, T.J.; Thorns, C.J. & Gamazo, C. (1998). Discrimination of strains of *Salmonella enteritidis* with differing levels of virulence by an in vitro glass adherence test. *Journal of Clinical Microbiology*, Vol.36, No.3, (March 1998), pp. 674-678
- Solomon, E.B.; Niemira, B.A.; Sapers, G.M. & Annous, B.A. (2005). Biofilm formation, cellulose production, and curli biosynthesis by *Salmonella* originating from produce, animal, and clinical sources. *Journal of Food Protection*, Vol.68, No.5, (May 2005), pp. 906-912
- Soni, K.A.; Jesudhasan, P.R.; Cepeda, M.; Williams, B.; Hume, M.; Russell, W.K.; Jayaraman, A. & Pillai, S.D. (2008). Autoinducer AI-2 is involved in regulating a variety of cellular processes in *Salmonella* Typhimurium. *Foodborne Pathogens and Disease*, Vol.5, No.2, pp. 147-153
- Sperandio, V.; Torres, A.G. & Kaper, J.B. (2002). Quorum sensing *Escherichia coli* regulators B and C (QseBC): a novel two-component regulatory system involved in the regulation of flagella and motility by quorum sensing in *E. coli*. *Molecular Microbiology*, Vol.43, No.3, (February 2002), pp. 809-821
- Stafford, G.P. & Hughes, C. (2007). *Salmonella typhimurium* *flhE*, a conserved flagellar regulon gene required for swarming. *Microbiology*, Vol.153(Pt2), (February 2007), pp. 541-547
- Steenackers, H.; Hermans, K.; Vanderleyden, J. & De Keersmaecker, S.C.J. (2011). *Salmonella* biofilms: an overview on occurrence, structure, regulation and eradication. *Food Research International*, in press
- Stepanović, S.; Cirković, I; Mijac, V. & Svabić-Vlahović, M. (2003). Influence of the incubation temperature, atmosphere and dynamic conditions on biofilm formation by *Salmonella* spp.. *Food Microbiology*, Vol.20, (September 2002), pp. 339-343

- Stepanović, S.; Cirković, I.; Ranin, L. & Svabić-Vlahović, M. (2004). Biofilm formation by *Salmonella* spp. and *Listeria monocytogenes* on plastic surface. *Letters in Applied Microbiology*, Vol.38, No.5, pp. 428-432
- Stocki, S.L.; Annett, C.B.; Sibley, C.D.; McLaws, M.; Checkley, S.L.; Singh, N.; Surette, M.G. & White, A.P. (2007). Persistence of *Salmonella* on egg conveyor belts is dependent on the belt type but not on the rdar morphotype. *Poultry Science*, Vol.86, (July 2007), pp. 2375-2383
- Stoodley, P.; Sauer, K.; Davies, D.G. & Costerton, J.W. (2002). Biofilms as complex differentiated communities. *Annual Review of Microbiology*, Vol.56, (January 2002), pp. 187-209
- Taga, M.E.; Semmelhack, J.L. & Bassler, B.L. (2001). The LuxS-dependent autoinducer AI-2 controls the expression of an ABC transporter that functions in AI-2 uptake in *Salmonella typhimurium*. *Molecular Microbiology*, Vol.42, No.3, (November 2001), pp. 777-793
- Tang, X.; Flint, S.H.; Bennett, R.J.; Brooks, J.D. & Morton, R.H. (2009). Biofilm growth of individual and dual strains of *Klebsiella oxytoca* from the dairy industry on ultrafiltration membranes. *Journal of Industrial Microbiology and Biotechnology*, Vol.36, No.12, (December 2009), pp. 1491-1497
- Teplitski, M.; Al-Agely, A. & Ahmer, B.M. (2006). Contribution of the SirA regulon to biofilm formation in *Salmonella enterica* serovar Typhimurium. *Microbiology*, Vol.152 (Pt 11), (November 2006), pp. 3411-3424
- Tolker-Nielsen, T. & Molin, S. (2000). Spatial organization of microbial biofilm communities. *Microbial Ecology*, Vol.40, No.2, (August 2000), pp. 75-84
- Ukuku, D.O. & Fett, W.F. (2002). Relationship of cell surface charge and hydrophobicity to strength of attachment of bacteria to cantaloupe rind. *Journal of Food Protection*, Vol.65, No.7, (July 2002), pp. 1093-1099
- Ukuku, D.O. & Fett, W.F. (2006). Effects of cell surface charge and hydrophobicity on attachment of 16 *Salmonella* serovars to cantaloupe rind and decontamination with sanitizers. *Journal of Food Protection*, Vol.69, No.8, (August 2006), pp. 1835-1843
- Van Houdt, R. & Michiels, S.W. (2010). Biofilm formation and the food industry, a focus on the bacterial outer surface. *Journal of Applied Microbiology*, Vol.109, (April 2010), pp. 1117-1131
- Verstraeten, N.; Braeken, K.; Debkumari, B.; Fauvart, M.; Franssaer, J.; Vermant, J. & Michiels, J. (2008). Living on a surface: swarming and biofilm formation. *Trends in Microbiology*, Vol.16, No.10, (October 2008), pp. 496-506
- Vestby, L.K.; Møretrø, T.; Balance, S.; Langsrud, S. & Nesse, L.L. (2009a). Survival potential of wild type cellulose deficient *Salmonella* from the feed industry. *BMC Veterinary Research*, Vol.5:43 (November 2009)
- Vestby, L.K.; Møretrø, T.; Langsrud, S.; Heir, E. & Nesse, L.L. (2009b). Biofilm forming abilities of *Salmonella* are correlated with persistence in fish meal- and feed factories. *BMC Veterinary Research*, Vol.5:20, (May 2009)
- Walters, M. & Sperandio, V. (2006). Quorum sensing in *Escherichia coli* and *Salmonella*. *International Journal of Medical Microbiology*, Vol.296, pp. 125-131
- White, A.P.; Gibson, D.L.; Collinson, S.K.; Banser, P.A. & Kay, W.W. (2003). Extracellular polysaccharides associated with thin aggregative fimbriae of *Salmonella enterica* serovar Enteritidis. *Journal of Bacteriology*, Vol.185, No.18, (September 2003), pp. 5398-5407

- White, A.P.; Gibson, D.L.; Grassl, G.A.; Kay, W.W.; Finlay, B.B.; Vallance, B.A. & Surette, M.G. (2008). Aggregation via the red, dry, and rough morphotype is not a virulence adaptation in *Salmonella enterica* serovar Typhimurium. *Infection and Immunity*, Vol.76, No.3, (March 2008), pp. 1048-1058
- White, A.P.; Gibson, D.L.; Kim, W.; Kay, W.W. & Surette, M.G. (2006). Thin aggregative fimbriae and cellulose enhance long-term survival and persistence of *Salmonella*. *Journal of Bacteriology*, Vol.188, No.9, (May 2006), pp. 3219-3227
- White, A.P. & Surette, M.G. (2006). Comparative genetics of the rdar morphotype in *Salmonella*. *Journal of Bacteriology*, Vol.188, No.24, (December 2006), pp. 8395-8406
- Whitehead, N.A.; Barnard, A.M.; Slater, H.; Simpson, N.J. & Salmond, G.P. (2001). Quorum-sensing in Gram-negative bacteria. *FEMS Microbiology Reviews*, Vol.25, No.4, (August 2001), pp. 365-404
- Whiteley, M.; Banger, M.G.; Bumgarner, R.E.; Parsek, M.R.; Teitzel, G.M.; Lory, S. & Greenberg, E.P. (2001). Gene expression in *Pseudomonas aeruginosa* biofilms. *Nature*, Vol.413, No.6858, (October 2001), pp. 860-864
- Wimpenny, J. (2009). Microbial metropolis. *Advances in Microbial Physiology*, Vol.56, (November 2009), pp. 29-84
- Wimpenny, J.; Manz, W. & Szewzyk, U. (2000). Heterogeneity in biofilms. *FEMS Microbiology Reviews*, Vol.24, No.5, (December 2000), pp. 661-671
- Wuertz, S.; Okabe, S. & Hausner, M. (2004). Microbial communities and their interactions in biofilm systems: an overview. *Water Science and Technology*, Vol.49, No.11-12, pp. 327-336
- Yoon, Y. & Sofos, J.N. (2008). Autoinducer-2 activity of Gram-negative foodborne pathogenic bacteria and its influence on biofilm formation. *Journal of Food Science*, Vol.73, No.3, (January 2008), pp. M140-M147
- Zhao, T.; Podtburg, T.C.; Zhao, P.; Schmidt, B.E.; Baker, D.A.; Cords, B. & Doyle, M.P. (2006). Control of *Listeria* spp. by competitive-exclusion bacteria in floor drains of a poultry processing plant. *Applied and Environmental Microbiology*, Vol.72, No.5, (May 2006), pp. 3314-3320
- Zogaj, X.; Nimtz, M.; Rohde, M.; Bokranz, W. & Römling, U. (2001). The multicellular morphotypes of *Salmonella typhimurium* and *Escherichia coli* produce cellulose as the second component of the extracellular matrix. *Molecular Microbiology*, Vol.39, No.6, (March 2001), pp. 1452-1463
- Zottola, E.A. & Sasahara, K.C. (1994). Microbial biofilms in the food processing industry - Should they be a concern? *International Journal of Food Microbiology*, Vol.23, (February 1994), pp. 125-148



Salmonella - A Dangerous Foodborne Pathogen

Edited by Dr. Dr. Barakat S M Mahmoud

ISBN 978-953-307-782-6

Hard cover, 450 pages

Publisher InTech

Published online 20, January, 2012

Published in print edition January, 2012

More than 2,500 serotypes of Salmonella exist. However, only some of these serotypes have been frequently associated with food-borne illnesses. Salmonella is the second most dominant bacterial cause of food-borne gastroenteritis worldwide. Often, most people who suffer from Salmonella infections have temporary gastroenteritis, which usually does not require treatment. However, when infection becomes invasive, antimicrobial treatment is mandatory. Symptoms generally occur 8 to 72 hours after ingestion of the pathogen and can last 3 to 5 days. Children, the elderly, and immunocompromised individuals are the most susceptible to salmonellosis infections. The annual economic cost due to food-borne Salmonella infections in the United States alone is estimated at \$2.4 billion, with an estimated 1.4 million cases of salmonellosis and more than 500 deaths annually. This book contains nineteen chapters which cover a range of different topics, such as the role of foods in Salmonella infections, food-borne outbreaks caused by Salmonella, biofilm formation, antimicrobial drug resistance of Salmonella isolates, methods for controlling Salmonella in food, and Salmonella isolation and identification methods.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Efstathios Giaouris, Nikos Chorianopoulos, Panagiotis Skandamis and George-John Nychas (2012). Attachment and Biofilm Formation by Salmonella in Food Processing Environments, Salmonella - A Dangerous Foodborne Pathogen, Dr. Dr. Barakat S M Mahmoud (Ed.), ISBN: 978-953-307-782-6, InTech, Available from: <http://www.intechopen.com/books/salmonella-a-dangerous-foodborne-pathogen/attachment-and-biofilm-formation-by-salmonella-in-food-processing-environments>

INTECH
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2012 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.