

# Macromolecular Synthesis in the Urinary and Reproductive Systems

Tetsuji Nagata<sup>1,2</sup>

*<sup>1</sup>Department of Anatomy and Cell Biology,  
Shinshu University School of Medicine, Matsumoto*

*<sup>2</sup>Shinshu Institute of Alternative Medicine and Welfare, Nagano  
Japan*

## 1. Introduction

This chapter deals with the third parts of the application of microscopic radioautography to the organ systems, including the urinary organs, the reproductive organs and the endocrine organs.

## 2. Macromolecular synthesis in the urinary system

The urinary system consists of the kidney and the urinary tract. We studied only the macromolecular synthesis in the kidneys of several groups of aging mice by LM and EM RAG, while the localization of an anti-allergic agent was observed in the urinary bladders of adult rats (Nagata 2005).

### 2.1 Macromolecular synthesis in the kidney

We studied only the DNA, RNA and glucides syntheses in the kidneys of several groups of aging mice by LM and EM RAG.

#### 2.1.1 The DNA synthesis in the kidney

The kidneys of mammals microscopically consist of the nephrons, which can be divided into two portions, the renal corpuscles and the uriniferous tubules. The renal corpuscles are composed of the glomeruli which are covered with the Bowman's capsules. They are localized in the outer zone of the kidney, the renal cortex, while the uriniferous tubules are composed of two portions, the proximal portions and the distal portions which can further be divided into several portions which run from the outer zone of the kidney, the renal cortex, to the inner zone, the medulla. We studied the DNA synthesis by <sup>3</sup>H-thymidine radioautography in 3 groups of ddY mouse embryos from prenatal day 13 (Fig. 16A), day 15 (Fig. 16B) to day 19 in vitro, as well as perinatal mice from embryonic day 19 to postnatal day 1, 8, 30, 60 and 365 (1 year) in vivo (Hanai 1993, Hani et al. 1993, Hanai and Nagata 1994a,b).




Fig. 16. LM RAG of the uro-genital organs. From Nagata, T.: *Special Cytochemistry in Cell Biology*, In, *Internat. Rev. Cytol.* Vol. 211, No. 1, p. 108, 2001, Academic Press, San Diego, USA, London, UK.

Fig. 16A. LM RAG of the metanephros of a prenatal day 13.5 mouse embryo labeled with  $^3\text{H}$ -thymidine, showing DNA synthesis.  $\times 1,200$ .

Fig. 16B. LM RAG of the metanephros cortex of a prenatal day 15.5 mouse embryo labeled with  $^3\text{H}$ -thymidine, showing DNA synthesis.  $\times 1,200$ .

Fig. 16C. LM RAG of the testis of a postnatal day 7 male mouse labeled with  $^3\text{H}$ -thymidine, showing DNA synthesis.  $\times 800$ .

Fig. 16D. LM RAG of the testis of a postnatal year 1 male mouse labeled with  $^3\text{H}$ -thymidine, showing DNA synthesis.  $\times 800$ .

Fig. 16E. LM RAG of the testis of a postnatal day 3 male mouse labeled with  $^3\text{H}$ -uridine in vitro, showing RNA synthesis.  $\times 1,500$ .

Fig. 16F. LM RAG of the testis of a male mouse at postnatal day 1 labeled with  $^3\text{H}$ -leucine in vitro, showing protein synthesis.  $\times 1,125$ .

Fig. 16G. LM RAG of the ovary of a postnatal day 3 female mouse labeled with  $^3\text{H}$ -thymidine in vitro, showing DNA synthesis in granulosa cells (G) and theca cells (T).  $\times 400$ .

Fig. 16H. LM RAG of the oviduct of a postnatal day 30 female mouse labeled with  $^3\text{H}$ -thymidine in vitro, showing DNA synthesis in epithelial cells.  $\times 400$ .

The labeling indices by LM RAG in glomeruli (28 to 32%) and uriniferous tubules (31 to 33%) in the superficial layer were higher than those of labeling indices (10 to 12%) and (8 to 16%) in the deeper layer from the late fetal to the suckling period, then decreased with aging from weaning to senescence (Fig. 17). EM RAG revealed the same results (Hanai and Nagata 1994a,b,c). At the same time, immunocytochemical staining of PCNA/cyclin was carried out in the same animals in several aging groups as  $^3\text{H}$ -thymidine RAG (Hanai 1993, Hanai et al. 1993). The results from the PCNA/cyclin positive indices in respective aging groups were almost the same as the labeling indices with  $^3\text{H}$ -thymidine RAG. The incorporation of  $^3\text{H}$ -thymidine was formerly observed by EM RAG in mitochondrial matrix of cultured kidney cells from chickens and mice in vitro demonstrating mitochondrial DNA synthesis (Nagata et al. 1967b).

### 2.1.2 The RNA synthesis in the kidney

The RNA synthesis by incorporation of  $^3\text{H}$ -uridine into the kidneys of aging mice was studied by LM and EM RAG (Hanai and Nagata 1994a,b, Nagata 2002). When the kidneys of several groups of aging mice from embryo to postnatal 1 year were radioautographed with  $^3\text{H}$ -uridine either in vitro (embryonic day 15, 19 and postnatal day 1) and in vivo (embryonic day 19, postnatal day 1, 7, month 1, 2, 12), RNA synthesis was observed in all the cells of the kidney at various ages. The numbers of silver grains demonstrating the incorporation of  $^3\text{H}$ -uridine in glomeruli (34.6 per cell) and uriniferous tubules (56.4 per cell) were higher in the superficial layer than those (15.6 and 18.6 per cell) in the deeper layer at embryonic day 15 and decreased gradually with aging. These results demonstrated the aging changes of RNA synthesis in the kidney.

### 2.1.3 Glucide synthesis in the kidney

The incorporations of  $^3\text{H}$ -glucosamine in the kidneys of aging mice were studied by LM RAG (Joukura 1996, Joukura and Nagata 1995) and EM RAG (Joukura et al. 1996). Silver grains were observed over all the cell type nephrons at embryonic day 19, i.e., glomerular epithelial cells, endothelial cells, mesangial cells, Bowman's capsular cells and tubule cells.

In newborn and suckling stages, from postnatal day 1, 3, 7 to 14, both the renal corpuscles and urinary tubules were well differentiated and the number of silver grains increased (Figs. 36 C, D, E F, G, H in Nagata 2002). The results from grain counting revealed that the numbers of silver grains in both the renal corpuscles and the uriniferous tubules were less in the embryonic stage, but increased postnatally and reached peaks at day 1 and 3, then decreased to senescence at 1 year. These results showed that glucide synthesis in the kidney cells also changed with aging of animals.




Fig. 17. Histogram showing aging changes of average labeling indices in respective cell types of the kidneys of aging mice labeled with  $^3\text{H}$ -thymidine. Mean  $\pm$  Standard Deviation. From Nagata, T.: Radioautography, General and Special. In, Prog. Histochem. Cytochem. Vol. 37, No. 2, p. 156, 2002, Urban & Fischer, Jena, Germany

Fig. 17A. Labeling indices of the glomeruli and the uriniferous tubules of mouse embryo from prenatal day 13 and 19.

Fig. 17B. Labeling indices of the glomeruli and the proximal and distal tubules of mouse embryo from prenatal day 19 to postnatal year 1. Abbreviations: GS=glomeruli of the superficial layer. US=uriniferous tubules of the superficial layer. GD=glomeruli of the deeper layer. UD=uriniferous tubules of the deeper layer. G=glomeruli. PT=proximal tubules. DT=distal tubules.

## 2.2 Localization of drugs in the urinary tract

The urinary tract is composed of the ureter, the urinary bladder and the urethra. We studied the urinary bladder of adult rats by LM RAG after oral administration of  $^3\text{H}$ -tranilast, an anti-allergic agent produced by Kissei Pharmaceutical Co. (Momose et al. 1989, Nishigaki et al. 1987, 1990a,b). It was found that this agent specifically localized over the transitional epithelium and the endothelium of the veins in the mucosa of normal adult rats. However, any study on the DNA synthesis in the ureter, the urinary bladder and the urethra was not carried out.

## 3. Macromolecular synthesis in the reproductive system

The reproductive system or genital organs can be divided into two parts, the male genital organs and female genital organs. We studied the DNA and RNA syntheses and protein synthesis in several groups of aging mice, both male and female, by LM and EM RAG (Nagata 2002).

### 3.1 Macromolecular synthesis in the male genital organs

The male genital organs consist of the testis and its excretory ducts such as ductuli efferentes, ductus epididymidis, ductus deferens, ejaculatory ducts, auxiliary glands and penis. We studied both DNA and RNA syntheses in these organs of several groups of ddY aging mice by LM and EMRAG using macromolecular precursors.

#### 3.1.1 The DNA synthesis in the male genital organs

Among the male genital organs, the testis was the main target of the scientific interests. Formerly, Clermont (1958, 1963) demonstrated using  $^3\text{H}$ -thymidine radioautography that several stages of development of the spermatogonia were found at different levels in the germinal epithelium of mature men and rodents, with the most primitive germ cells found at the base and the more differentiated cells located at higher levels. We studied the DNA synthesis in the testis of several groups of aging mice.

##### 3.1.1.1 The DNA synthesis in the testis

The structure of the testis of mammals is a compound tubular gland enclosed in tunica albuginea, a thick fibrous capsule. The parenchyma of the testis is composed of around 250 pyramidal compartments in men and animals, named lobules. Each lobule is made of convoluted seminiferous tubules, consisting of many spermatogenic cells differentiating to sperms among the supporting cells of Sertoli in the seminiferous epithelium, surrounded by the interstitial cells of Leydig. We first studied the macromolecular synthesis in the testis of aging male ddY mice at various ages (Gao 1993, Gao et al. 1994, 1995a,b). When testicular tissues were labeled with  $^3\text{H}$ -thymidine and observed by LM and EM RAG, many spermatogonia and myoid cells as well as Leydig cells were labeled with  $^3\text{H}$ -thymidine at various ages from embryonic day 19 to postnatal day 1, 3, 7 (Fig. 16C), 14, month 1, 2, 6, 12 (Fig. 16D) and 24 (2 years). Silver grains were localized over the nuclei and several mitochondria of the spermatogonia showing DNA synthesis. Among of the aging groups, we counted the numbers of mitochondria per cell profile area, the numbers of labeled mitochondria per cell of the spermatogonia from 4 aging groups, prenatal embryonic day 19, postnatal day 3, and adults at month 1 and 6, and the labeling indices were calculated.

The results showed that the LI of the spermatogonia increased from embryonic day 19 (17%) to postnatal day 7 (25%) and month 1 (30%), reaching the maximum, then decreased to month 6 (20%) to year 2.




Fig. 18. Transitional curves of the labeling indices of respective cell types in the testis of aging mice labeled with  $^3\text{H}$ -thymidine, showing DNA synthesis. Mean  $\pm$  Standard Deviation. From Nagata, T.: Radioautography, General and Special. In, Prog. Histochem. Cytochem. Vol. 37, No. 2, p. 160, 2002, Urban & Fischer, Jena, Germany

Fig. 18A. The spermatogonia.

Fig. 18B. The Sertoli cells.

Fig. 18C. The myoid cells.

At embryonic and neonatal stages, DNA synthesis of spermatocytes was weak and only a few labeled spermatogonia could be observed during the perinatal stages. The labeled spermatocytes were recognized at postnatal day 4 and 7 (Fig. 16C) and the number of labeled spermatogonia and spermatocytes increased from 2 and 3 weeks, keeping high level from month 1 to year 1 and 2 until senescence (Fig. 18A). However, the Sertoli's cells (Fig. 18B) and myoid cells (Fig. 18C) labeled with  $^3\text{H}$ -thymidine were frequently observed at perinatal stages from embryo to postnatal day 7, while the labeling indices of both cells decreased from young adulthood (postnatal 2 weeks) to senescence (Gao 1993, Gao et al. 1994,1995a). The interstitial cells of Leydig in the testis surrounding the seminiferous tubules shall be described in the following section of the endocrine system in detail.

### **3.1.2 The RNA synthesis in the male genital organs**

Among of the male genital organs, we studied the RNA synthesis in the testis of several groups of aging mice.

#### **3.1.2.1 The RNA synthesis in the testis**

We studied the RNA syntheses in aging mouse testis by LM and EM RAG, demonstrating the incorporations of  $^3\text{H}$ -uridine into various cells of the seminiferous tubules (Gao 1993, Gao et al. 1994, Nagata 2002). The RNA synthesis of various cells in the seminiferous tubules was studied using  $^3\text{H}$ -uridine. Silver grains due to  $^3\text{H}$ -uridine demonstrating RNA synthesis were observed over the nuclei and cytoplasm of all spermatogonia, spermatocytes, Sertoli's cells, myoid cells of immature mice at perinatal stages at day 1 and 3 (Fig. 16E), as well as in mature and senescent mice from month 1, 6 to year 1 and 2. The synthetic activities of spermatogonia, Sertoli's cells and myoid cells as shown by grain counting with  $^3\text{H}$ -uridine, as expressed by grain counting, were low (2-8 grain counts per 10 mm<sup>2</sup>) at the embryonic and neonatal stages but increased at adult stages and maintained high levels (10-20 grain counts per 10 mm<sup>2</sup>) until senescence. These results showed that DNA synthesis in myoid cells and Sertoli's cells increased at the perinatal stages and decreased from postnatal 2 weeks as described previously (Fig. 16A), while the RNA synthesis (Fig. 16E) in spermatogonia increased from postnatal 2 weeks together with DNA and protein syntheses (Fig. 16F) to senescence.

### **3.1.3 The protein synthesis in the male genital organs**

We studied the protein synthesis of the reproductive system in both the male and female reproductive organs.

#### **3.1.3.1 The protein synthesis in the testis**

We studied the protein syntheses in aging mouse testis by LM and EM RAG, demonstrating the incorporations of  $^3\text{H}$ -leucine into various cells of the seminiferous tubules (Gao 1993, Gao et al. 1994, Nagata 2002). The protein synthesis of various cells in the seminiferous tubules was first studied after administration of  $^3\text{H}$ -leucine into aging male mice at various ages from perinatal to senescence at postnatal 2 years. Silver grains due to  $^3\text{H}$ -leucine incorporation demonstrating protein synthesis were observed over the nuclei and cytoplasm of all the cells, spermatogonia, spermatocytes, Sertoli's cells, myoid cells of all male mice at respective stages from perinatal to senescence. The synthetic activities of spermatogonia, Sertoli's cells and myoid cells as shown by the number of silver grains due to  $^3\text{H}$ -leucine, as expressed by grain

counting, were low at the embryonic and neonatal stages but increased at adult stages and maintained high levels until senescence. These results showed that DNA synthesis in Sertoli's cells (Fig. 18B) and myoid cells (Fig. 18C) that increased at the perinatal stages and decreased from postnatal 2 weeks, while the DNA synthesis in spermatogonia increased from postnatal 2 weeks (Fig. 18A) together with RNA and protein syntheses to senescence.

### **3.2 Macromolecular synthesis in the female genital organs**

The female genital organs consist of the ovary, the oviduct, the uterus, the vagina and the external genitals. We studied the macromolecular synthesis in the ovary, oviduct and uterus of several litters of ddY mice in aging.

#### **3.2.1 The DNA synthesis in the female genital organ**

Among the female genital organs, we studied the DNA synthesis in the ovary, oviduct and uterus of several litters of ddY mice in aging.

##### **3.2.1.1 The DNA synthesis in the ovary**

The ovary consists of the germinal epithelium covering the surface and the stroma containing many developing ovarian follicles depending upon the age of animals.

The nucleic acids, DNA and RNA, syntheses in the developing virgin mice ovaries of 6 litters, each 3 individuals, consisting of 36 female mice at various ages in respective precursors were studied by  $^3\text{H}$ -thymidine and  $^3\text{H}$ -uridine radioautography (Li 1994, Li and Nagata 1995, Li et al. 1992). The  $^3\text{H}$ -thymidine incorporations were active in all surface epithelial cells, stromal and follicular cells of the ovaries between postnatal days 1 to 7 and decreased from day 14 (Fig. 16G) and maintained a lower level to day 60, while  $^3\text{H}$ -uridine incorporations were active in all surface epithelial cells, stromal and follicular cells of the ovaries between postnatal days 1 to 7 and maintained medium levels from day 14 on.

The labeling indices with  $^3\text{H}$ -thymidine showing DNA synthetic activity were high in all the surface epithelial cells, follicular cells and stromal cells of mice at neonatal stage from postnatal day 1 to 7, but decreased from day 40 to day 60 at mature stage (Fig. 19A). The grain counts showing RNA synthetic activity were high at neonatal stage from day 1 to day 7, and maintained medium levels from day 14 to day 60 at mature stage.

##### **3.2.1.2 The DNA synthesis in the oviduct**

The nucleic acids, DNA and RNA, syntheses in the oviducts of developing virgin mice at various ages were studied by  $^3\text{H}$ -thymidine and  $^3\text{H}$ -uridine radioautography (Li 1994, Li and Nagata 1995). The silver grains with  $^3\text{H}$ -thymidine showing the DNA synthesis were observed over many nuclei in all surface epithelial cells, stromal and smooth muscle cells at neonatal stage between postnatal day 1 to 3 and decreased from day 7 to 30 (Fig. 16H) and 60, while the silver grains showing the RNA synthesis with  $^3\text{H}$ -uridine were observed over the nuclei and cytoplasm of all the epithelial and stromal cells from postnatal day 1 to day 60. The labeling indices with  $^3\text{H}$ -thymidine were high at neonatal stage from postnatal day 1 to 3 but decreased from day 7 to day 60 (Fig. 19C). The grain counts with  $^3\text{H}$ -uridine were high at neonatal stage from postnatal day 1 to 3 and increased from day 7 to day 14 and decreased from day 30 to day 60. These results demonstrated an unparalleled alternation of DNA and RNA syntheses in the oviduct (Li and Nagata 1995).




Fig. 19. Histogram showing aging changes of average labeling indices in respective cell types of female genital organs of aging mice labeled with  $^3\text{H}$ -thymidine. Mean  $\pm$  Standard Deviation. From Nagata, T.: Radioautography, General and Special. In, Prog. Histochem. Cytochem. Vol. 37, No. 2, p. 166, 2002, Urban & Fischer, Jena, Germany

Fig. 19A. The ovary.

Fig. 19B. The uterus.

Fig. 19C. The oviduct.

### 3.2.1.3 The DNA synthesis in the uterus

The silver grains with  $^3\text{H}$ -thymidine showing DNA synthesis of the uterus was observed over some of the nuclei of all the cells in the epithelia, stroma and smooth muscles from postnatal day 1 to 60 (Li 1994, Li and Nagata 1995). The labeling indices with  $^3\text{H}$ -thymidine (Fig. 19B) were high (80-95%) at postnatal day 1 and decreased from day 3 to 60 (>10%). The silver grains showing RNA synthesis of the uterus were observed over all the nuclei and cytoplasm of all the cells in the uterine epithelia, stroma and smooth muscles from day 1 to 60. The number of silver grains in the uterine epithelium increased from postnatal day 1 to 7 and decreased from day 14 to 60, while they increased in the stroma from day 1 to 3 and decreased from day 7 to 60.

These results from the female genital organs showed that both DNA and RNA syntheses, as expressed by labeling indices and grain counting, were active in all kinds of cells, such as surface epithelial cells, stromal cells and follicular cells of the ovaries between postnatal

days 1 to 7, then they decreased from day 14 to 60. However, the DNA synthesis in the epithelial cells and the stromal cells of both the uteri and the oviducts was active at postnatal day 1 and 3 and decreased from day 7 to 60. The RNA synthesis in the uteri and oviducts was active at postnatal day 1, increased from day 1 to day 14, and decreased from day 30 to 60. The unparalleled alteration of the DNA and RNA syntheses was shown between the ovary and the uterus or oviduct (Li and Nagata 1995).

We also studied PCNA/cyclin immunostaining in the ovary, oviduct and uterus (Li 1994). It was demonstrated that PCNA/cyclin positive cells were observed in the ovarian follicular epithelium, ovarian interstitial cells, tubal epithelial cells, tubal interstitial cells, uterine epithelial cells and uterine interstitial cells. The positive cells increased from postnatal 1 day to 3 and 7 days, then decreased from 14 days to senescence. These results accorded well with the results obtained from the  $^3\text{H}$ -thymidine radioautography (Li 1994, Li and Nagata 1995). Moreover, the mucosubstance synthesis incorporating sulfuric acid was also carried out (Oliveira et al. 1991, Li et al. 1995).

#### **3.2.1.4 The DNA synthesis in gametogenesis**

The gametogenesis consists of both spermatogenesis in male germ cells and the oogenesis in female germ cells, leading to the implantation and further development of blastocysts. The macromolecular synthesis, DNA, RNA and protein synthesis, in both the testis and the ovary were already described in the sections of male and female reproductive systems (3.7.1. and 3.7.2.) previously.

#### **3.2.1.5 The DNA synthesis in implantaion**

In order to detect the changes of DNA, RNA and protein synthesis of the developing blastocysts in mouse endometrium during activation of the implantation, ovulations of female BALB/C strain mice were controlled by pregnant mare serum gonadotropin and human chorionic gonadotropin, then pregnant female mice were ovariectomized on the 4th day of pregnancy (Yamada 1993, Yamada and Nagata 1992a,b, 1993). The delay implantation state was maintained for 48 hrs and after 0 to 18 hrs of estrogen supply  $^3\text{H}$ -thymidine was injected. The three regions of the endometrium, i. e. the interinplantation site, the antimesometrial and mesometrial sides of implantation site, were taken out and processed for LM and EM RAG. It was well known that the uterus of the rodent becomes receptive to blastocyst implantation only for a restricted period. This is called the implantation window which is intercalated between refractory states of the endometrium whose cycling is regulated by ovarian hormones (Yoshinaga 1988). We studied the changes of DNA synthesis by  $^3\text{H}$ -thymidine (Yamada and Nagata 1992a,b) incorporations in the endometrial cells of pregnant-ovariectomized mice after time-lapse effect of nidatory estradiol. As the results, the endometrial cells showed topographical and chronological differences in the nucleic acid synthesis. The cells labeled with  $^3\text{H}$ -thymidine increased after nidatory estradiol effects in the stromal cells around the blastocyst, but not in the epithelial cells. The results suggested that the presence of the blastocysts in the uterine lumen induced selective changes in the behavior of endometrial cells after nidatory estradiol effect showing the changes of DNA synthesis.

As for a lower vertebrate, cell proliferation and migration of scleroblasts and their precursor cells during ethisterone-induced anal-fin process formation of the medaka, *orizias latipes*, was studied by LM RAG labeled with  $^3\text{H}$ -thymidine (Uwa and Nagata 1976). The results

showed that the labeling index in the posterior margin of the joint plate rapidly increased and the scleroblast population in the central portion increased simultaneously from the 3rd to 5th day of ethisterone treatment. These results indicated that the scleroblasts and their precursor cells migrated from the peripheral portion to the central portion along the proximal-distal axis of the joint plate.

### **3.2.2 The RNA synthesis in the female genital organs**

We studied the RNA synthesis of female reproductive organs of aging mice after the administration of  $^3\text{H}$ -uridine at various ages.

#### **3.2.2.1 The RNA synthesis in the ovary**

The ovary consists of the germinal epithelium covering the surface and the stroma containing many developing ovarian follicles depending upon the age of animals.

The nucleic acids, DNA and RNA, syntheses in the developing virgin mice ovaries of 6 litters, each 3 individuals, consisting of 36 female mice at various ages in 2 groups were studied by  $^3\text{H}$ -thymidine and  $^3\text{H}$ -uridine radioautography (Li 1994, Li and Nagata 1995, Li et al. 1992). The  $^3\text{H}$ -thymidine incorporations were active in all surface epithelial cells, stromal and follicular cells of the ovary between postnatal days 1 to 7 and decreased from day 14 (Fig. 16G) and maintained a lower level to day 60, while  $^3\text{H}$ -uridine incorporations were active in all surface epithelial cells, stromal and follicular cells of the ovary between postnatal days 1 to 7 and maintained medium levels from day 14 on.

The labeling indices with  $^3\text{H}$ -thymidine showing DNA synthetic activity were high in all the surface epithelial cells, follicular cells and stromal cells of mice at neonatal stage from postnatal day 1 to 7, but decreased from day 40 to day 60 at mature stage. The grain counts showing RNA synthetic activity were high at neonatal stage from day 1 to day 7, and maintained medium levels from day 14 to day 60 at mature stage.

#### **3.2.2.2 The RNA synthesis in the oviduct**

The nucleic acids, DNA and RNA, syntheses in the oviducts of developing virgin mice at various ages were studied by  $^3\text{H}$ -thymidine and  $^3\text{H}$ -uridine radioautography (Li 1994, Li and Nagata 1995). The silver grains with  $^3\text{H}$ -thymidine showing the DNA synthesis were observed over many nuclei in all surface epithelial cells, stromal and smooth muscle cells at neonatal stage between postnatal day 1 to 3 and decreased from day 7 to 30 (Fig. 16H) and 60, while the silver grains showing the RNA synthesis with  $^3\text{H}$ -uridine were observed over the nuclei and cytoplasm of all the epithelial and stromal cells from postnatal day 1 to day 60. The labeling indices with  $^3\text{H}$ -thymidine were high at neonatal stage from postnatal day 1 to 3 but decreased from day 7 to day 60. The grain counts with  $^3\text{H}$ -uridine were high at neonatal stage from postnatal day 1 to 3 and increased from day 7 to day 14 and decreased from day 30 to day 60. These results demonstrated an unparalleled alternation of DNA and RNA syntheses in the oviduct (Li and Nagata 1995).

#### **3.2.2.3 The RNA synthesis in the uterus**

The silver grains with  $^3\text{H}$ -uridine showing RNA synthesis of the uterus was observed over almost all the nuclei and cytoplasm of all the cells in the epithelia, stroma and smooth muscles from postnatal day 1 to 60 (Li 1994, Li and Nagata 1995). The labeling indices with

$^3\text{H}$ -thymidine were high (80-95%) at postnatal day 1 and decreased from day 3 to 60 (>10%). The silver grains showing RNA synthesis of the uterus were observed over all the nuclei and cytoplasm of all the cells in the uterine epithelia, stroma and smooth muscles from day 1 to 60. The number of silver grains in the uterine epithelium increased from postnatal day 1 to 7 and decreased from day 14 to 60, while they increased in the stroma from day 1 to 3 and decreased from day 7 to 60.

These results from the female genital organs showed that both DNA and RNA syntheses, as expressed by labeling indices and grain counting, were active in all kinds of cells, such as surface epithelial cells, stromal cells and follicular cells of the ovaries between postnatal days 1 to 7, then they decreased from day 14 to 60. However, the DNA synthesis in the epithelial cells and the stromal cells of both the uteri and the oviducts was active at postnatal day 1 and 3 and decreased from day 7 to 60. The RNA synthesis in the uteri and oviducts was active at postnatal day 1, increased from day 1 to day 14, and decreased from day 30 to 60. The unparalleled alteration of the DNA and RNA syntheses was shown between the ovary and the uterus or oviduct (Li and Nagata 1995).

We also studied PCNA/cyclin immunostaining in the ovary, oviduct and uterus (Li 1994). It was demonstrated that PCNA/cyclin positive cells were observed in the ovarian follicular epithelium, ovarian interstitial cells, tubal epithelial cells, tubal interstitial cells, uterine epithelial cells and uterine interstitial cells. The positive cells increased from postnatal 1 day to 3 and 7 days, then decreased from 14 days to senescence. These results accorded well with the results obtained from the  $^3\text{H}$ -thymidine radioautography (Li 1994, Li and Nagata 1995). Moreover, the mucosubstance synthesis incorporating sulfuric acid was also carried out (Oliveira et al. 1991, 1995, Li et al. 1992).

#### **3.2.2.4 The RNA synthesis in gametogenesis**

The gametogenesis consists of both spermatogenesis in male germ cells and the oogenesis in female germ cells, leading to the implantation and further development of blastocysts. The macromolecular synthesis, DNA, RNA and protein synthesis, in both the testis and the ovary were already described in the sections of male and female reproductive systems (8.1.1. and 8.1.2.) previously.

#### **3.2.2.5 The RNA synthesis in implantation**

In order to detect the changes of DNA, RNA and protein synthesis of the developing blastocysts in mouse endometrium during activation of the implantation, ovulations of female BALB/C strain mice were controlled by pregnant mare serum gonadotropin and human chorionic gonadotropin, then pregnant female mice were ovariectomized on the 4th day of pregnancy (Yamada 1993, Yamada and Nagata 1992a,b, 1993). The delay implantation state was maintained for 48 hrs and after 0 to 18 hrs of estrogen supply  $^3\text{H}$ -thymidine was injected. The three regions of the endometrium, i. e. the interimplantation site, the antimesometrial and mesometrial sides of implantation site, were taken out and processed for LM and EM RAG. It was well known that the uterus of the rodent becomes receptive to blastocyst implantation only for a restricted period. This is called the implantation window which is intercalated between refractory states of the endometrium whose cycling is regulated by ovarian hormones (Yoshinaga 1988). We studied the changes of DNA synthesis by  $^3\text{H}$ -thymidine (Yamada and Nagata 1992a,b) incorporations in the endometrial cells of pregnant-ovariectomized mice after time-lapse effect of nidatory

estradiol. As the results, the endometrial cells showed topographical and chronological differences in the nucleic acid synthesis. The cells labeled with  $^3\text{H}$ -thymidine increased after nidatory estradiol effects in the stromal cells around the blastocyst, but not in the epithelial cells. The results suggested that the presence of the blastocysts in the uterine lumen induced selective changes in the behavior of endometrial cells after nidatory estradiol effect showing the changes of DNA synthesis.

As for a lower vertebrate, cell proliferation and migration of scleroblasts and their precursor cells during ethisterone-induced anal-fin process formation of the medaka, *orizias latipes*, was studied by LM RAG labeled with  $^3\text{H}$ -thymidine (Uwa and Nagata 1976). The results showed that the labeling index in the posterior margin of the joint plate rapidly increased and the scleroblast population in the central portion increased simultaneously from the 3rd to 5th day of ethisterone treatment. These results indicated that the scleroblasts and their precursor cells migrated from the peripheral portion to the central portion along the proximal-distal axis of the joint plate.

### **3.2.3 The protein synthesis of the female genital organs**

We studied the protein synthesis of female reproductive organs of aging mice after the administration of  $^3\text{H}$ -leucine at various ages.

#### **3.2.3.1 The protein synthesis in the uterus**

We studied the protein synthesis of the developing blastocysts in female mouse endometrium during activation of the implantation. The ovulations of female BALB/C strain adult mice were controlled by pregnant mare serum gonadotropin and human chorionic gonadotropin, then pregnant female mice were ovariectomized on the 4th day of pregnancy (Yamada 1993, Yamada and Nagata 1992a,b, 1993). The delay implantation state was maintained for 48 hrs and after 0 to 18 hrs of estrogen supply. After the mice were injected with  $^3\text{H}$ -leucine, they were sacrificed and the uteri were processed for LM and EMRAG. We studied the changes of protein synthesis by  $^3\text{H}$ -leucine incorporations (Yamada 1993, Yamada and Nagata 1992a). As the results, the endometrial cells showed topographical and chronological differences in the protein synthesis. The cells labeled with  $^3\text{H}$ -leucine were observed in both epithelial cells and stromal cells. Quantitative analysis revealed that the number of silver grains increased from 0 hr to 3 and 6 hr, reaching the peak at 6 hr and decreased from 12 to 18 hr. The protein synthesis in the decidual cells of pregnant mice uteri was compared to the endometrial cells of virgin mice uteri using  $^3\text{H}$ -proline and  $^3\text{H}$ -tryptophane incorporations. The results demonstrated that silver grains were localized over the endoplasmic reticulum and the Golgi apparatus of fibroblasts and accumulated over collagen fibrils in the extracellular matrix suggesting that the decidual cells produced collagen in the matrix. The collagen synthesis in the mouse decidual cells by  $^3\text{H}$ -proline showed that silver grains were localized over the endoplasmic reticulum and Golgi apparatus of fibroblasts and accumulated over collagen fibrils in the extracellular matrix (Oliveira et al. 1991, 1995). However, analytical studies on protein synthesis in aging mice at various ages were not yet carried out.

#### **3.2.3.2 The protein synthesis in the implantation**

In order to detect the changes of DNA, RNA and protein synthesis of the developing blastocysts in female mouse endometrium during activation of the implantation, ovulations

of female BALB/C strain mice were controlled by pregnant mare serum gonadotropin and human chorionic gonadotropin, then pregnant female mice were ovariectomized on the 4th day of pregnancy (Yamada 1993, Yamada and Nagata 1992a,b, 1993). The delay implantation state was maintained for 48 hrs and after 0 to 18 hrs of estrogen supply and  $^3\text{H}$ -leucine was injected. The three regions of the endometrium, i. e. the interimplantation site, the antimesometrial and mesometrial sides of implantation site, were taken out and processed for LM and EM RAG. It was well known that the uterus of the rodent becomes receptive to blastocyst implantation only for a restricted period. This is called the implantation window which is intercalated between refractory states of the endometrium whose cycling is regulated by ovarian hormones (Yoshinaga 1988). We studied the changes of protein synthesis by  $^3\text{H}$ -leucine (Yamada 1993, Yamada and Nagata 1992a) incorporations in the endometrial cells of pregnant-ovariectomized mice after time-lapse effect of nidatory estradiol. As the results, the endometrial cells showed topographical and chronological differences in the nucleic acid and protein synthesis. The cells labeled with  $^3\text{H}$ -leucine were observed in both epithelial cells and stromal cells. Quantitative analysis revealed that the number of silver grains as expressed by grain counting per  $\text{mm}^2$  in both the stromal and epithelial cells on the antimesometrial side with  $^3\text{H}$ -leucine increased from 0 hr to 3 and 6 hr, reaching the peak at 6 hr and decreased from 12 to 18 hr. These results suggested that the presence of the blastocysts in the uterine lumen induced selective changes in the behavior of endometrial cells after nidatory estradiol effect showing the changes of DNA, RNA and protein synthesis. The time coincident peak of RNA and protein synthesis detected in the endometrial cells at the anti-mesometrial side of the implantation site, probably reflected the activation moment of the implantation window. The protein synthesis in the decidual cells of pregnant mice uteri was compared to the endometrial cells of virgin mice uteri using  $^3\text{H}$ -proline and  $^3\text{H}$ -tryptophane incorporations. The results demonstrated that silver grains were localized over the endoplasmic reticulum and the Golgi apparatus of fibroblasts and accumulated over collagen fibrils in the extracellular matrix suggesting that the decidual cells produced collagen in the matrix. On the other hand, collagen synthesis in the mouse decidual cells was studied by LM and EM RAG using  $^3\text{H}$ -proline (Oliveira et al. 1991, 1995). Silver grains were localized over the endoplasmic reticulum and Golgi apparatus of fibroblasts and accumulated over collagen fibrils in the extracellular matrix. The results suggested that the decidual cells produced collagen into the matrix. The quantitative analysis showed that both incorporations in the decidual cells and the matrix increased in the pregnant mice than the endometrial cells in virgin mice.

### **3.2.4 The glucide synthesis in the reproductive system**

Among the reproductive organs, only the mucosubstance synthesis with radiosulfate,  $^{35}\text{SO}_4$ , was studied in the ovaries of mice during the estrus cycle.

#### **3.2.4.1 The glucide synthesis in the ovary**

Litter mate groups of female ddY mice, aged 8-10 weeks, were divided into 4 groups, diestrus, proestrus, estrus and metestrus according to the vaginal smears. The ovaries were taken out, labeled with  $^{35}\text{SO}_4$  in vitro and radioautographed. In all the animals, silver grains were localized over the granulosa and theca cells. Almost all compartments of the ovaries were labeled. The grain counts per cell changed according to cell cycle. From the results, it was concluded that all the cells of the ovary incorporated mucosubstances throughout the estrus cycle (Li et al. 1992).

#### 4. References

- Chen, S., Gao, F., Kotani, A., Nagata, T.: Age-related changes of male mouse submandibular gland: A morphometric and radioautographic study. *Cell. Mol. Biol.* 41, 117-124, 1995.
- Clermont Y.: The contractile elements in the limiting membrane of the seminiferous tubules of rats. *Exp. Cell Res.* 15, 438-342, 1958.
- Clermont, Y.: Renewal of spermatogonia in man. *Amer. J. Anat.* 112, 35-51, 1963.
- Cui, H.: Light microscopic radioautographic study on DNA synthesis of nerve cells in the cerebella of aging mice. *Cell. Mol. Biol.* 41, 1139-1154, 1995.
- Cui, H., Gao, F., Nagata, T.: Light microscopic radioautographic study on protein synthesis in perinatal mice corneas. *Acta Histochem. Cytochem.* 33, 31-37, 2000.
- Duan, H., Gao, F., Li, S., Hayashi, K., Nagata, T.: Aging changes and fine structure and DNA synthesis of esophageal epithelium in neonatal, adult and old mice. *J. Clin. Electron Microsc.* 25, 452-453, 1992.
- Duan, H., Gao, F., Li, S., Nagata, T.: Postnatal development of esophageal epithelium in mouse: a light and electron microscopic radioautographic study. *Cell. Mol. Biol.* 39, 309-316, 1993.
- Duan, H., Gao, F., Oguchi, K., Nagata, T.: Light and electron microscopic radioautographic study on the incorporation of  $^3\text{H}$ -thymidine into the lung by means of a new nebulizer. *Drug Res.* 44, 880-883, 1994.
- Feulgen, R., Rossenbeck, H.: Mikroskopische-chemischer Nachweis einer Nukeinsaeure von Thymus der Thymonukeinsaeure *Z. Physik. Chem.* 135, 203-248, 1924.
- Fujii, Y., Ohno, S., Yamabayashi, S., Usuda, N., Saito, H., Furuta, S., Nagata, T.: Electron microscopic radioautography of DNA synthesis in primary cultured cells from an IgG myeloma patient. *J. Clin. Electr. Microsc.* 13, 582-583, 1980.
- Gao, F.: Study on the macromolecular synthesis in aging mouse seminiferous tubules by light and electron microscopic radioautography. *Cell. Mol. Biol.* 39, 659-672, 1993.
- Gao, F., Toriyama, K., Nagata, T.: Light microscopic radioautographic study on the DNA synthesis of prenatal and postnatal aging mouse retina after labeled thymidine injection. *Cell. Mol. Biol.* 38, 661-668, 1992a.
- Gao, F., Li, S., Duan, H., Ma, H., Nagata, T.: Electron microscopic radioautography on the DNA synthesis of prenatal and postnatal mice retina after labeled thymidine injection. *J. Clin. Electron Microsc.* 25, 721-722, 1992b.
- Gao, F., Toriyama, K., Ma, H., Nagata, T.: Light microscopic radioautographic study on DNA synthesis in aging mice corneas. *Cell. Mol. Biol.* 39, 435-441, 1993.
- Gao, F., Ma, H., Sun, L., Jin, C., Nagata, T.: Electron microscopic radioautographic study on the nucleic acids and protein synthesis in the aging mouse testis. *Med. Electron Microsc.* 27, 360-362, 1994.
- Gao, F., Chen, S., Sun, L., Kang, W., Wang, Z., Nagata, T.: Radioautographic study of the macromolecular synthesis of Leydig cells in aging mice testis. *Cell. Mol. Biol.* 41, 145-150, 1995a.
- Gao, F., Jin, C., Ma, H., Sun, L., Nagata, T.: Ultrastructural and radioautographic studies on DNA synthesis in Leydig cells of aging mouse testis. *Cell. Mol. Biol.* 41, 151-160, 1995b.
- Gunarso, W.: Radioautographic studies on the nucleic acid synthesis of the retina of chick embryo. I. Light microscopic radioautography. *Shinshu Med. J.* 32, 231-240, 1984a.

- Gunarso, W.: Radioautographic studies on the nucleic acid synthesis of the retina of chick embryo. II. Electron microscopic radioautography. *Shinshu Med. J.* 32, 241-248, 1984b.
- Gunarso, W., Gao, F., Cui, H., Ma, H., Nagata, T.: A light and electron microscopic radioautographic study on RNA synthesis in the retina of chick embryo. *Acta Histochem.* 98, 309-32, 1996.
- Gunarso, W., Gao, F., Nagata, T.: Development and DNA synthesis in the retina of chick embryo observed by light and electron microscopic radioautography. *Cell. Mol. Biol.* 43, 189-201, 1997.
- Hanai, T.: Light microscopic radioautographic study of DNA synthesis in the kidneys of aging mice. *Cell. Mol. Biol.* 39, 81-91, 1993.
- Hanai, T., Nagata, T.: Electron microscopic radioautographic study on DNA and RNA synthesis in perinatal mouse kidney. In, *Radioautography in Medicine*, Nagata, T., Ed., pp. 127-131, Shinshu University Press, Matsumoto, 1994a.
- Hanai, T., Nagata, T.: Study on the nucleic acid synthesis in the aging mouse kidney by light and electron microscopic radioautography. In, *Radioautography in Medicine*, Nagata, T., Ed., pp. 209-214, Shinshu University Press, Matsumoto, 1994b.
- Hanai, T., Nagata, T.: Electron microscopic study on nucleic acid synthesis in perinatal mouse kidney tissue. *Med. Electron Microsc.* 27, 355-357, 1994c.
- Hanai, T., Usuda, N., Morita, T., Shimizu, T., Nagata, T.: Proliferative activity in the kidneys of aging mice evaluated by PCNA/cyclin immunohistochemistry. *Cell. Mol. Biol.* 39, 181-191, 1993.
- Hayashi, K., Gao, F., Nagata, T.: Radioautographic study on  $^3\text{H}$ -thymidine incorporation at different stages of muscle development in aging mice. *Cell. Mol. Biol.* 39, 553-560, 1993.
- Ito, M.: The radioautographic studies on aging change of DNA synthesis and the ultrastructural development of mouse adrenal gland. *Cell. Mol. Biol.* 42, 279-292, 1996.
- Ito, M., Nagata, T.: Electron microscopic radioautographic studies on DNA synthesis and ultrastructure of aging mouse adrenal gland. *Med. Electron Microsc.* 29, 145-152, 1996.
- Izumiyama, K., Kogure, K., Kataoka, S., Nagata, T.: Quantitative analysis of glucose after transient ischemia in the gerbil hippocampus by light and electron microscope radioautography. *Brain Res.* 416, 175-179, 1987.
- Jamieson, J. D., Palade, G. E.: Intracellular transport of secretory proteins in the pancreatic exocrine cells. *J. Cell Biol.* 34, 577-615, 1967.
- Jin, C.: Study on DNA synthesis of aging mouse colon by light and electron microscopic radioautography. *Cell. Mol. Biol.* 42, 255-268, 1996.
- Jin, C., Nagata, T.: Light microscopic radioautographic study on DNA synthesis in cecal epithelial cells of aging mice. *J. Histochem. Cytochem.* 43, 1223-1228, 1995a.
- Jin, C., Nagata, T.: Electron microscopic radioautographic study on DNA synthesis in cecal epithelial cells of aging mice. *Med. Electron Microsc.* 28, 71-75, 1995b.
- Joukura, K.: The aging changes of glycoconjugate synthesis in mouse kidney studied by  $^3\text{H}$ -glucosamine radioautography. *Acta Histochem. Cytochem.* 29, 57-63, 1996.
- Joukura, K., Nagata, T.: Aging changes of  $^3\text{H}$ -glucosamine incorporation into mouse kidney observed by radioautography. *Acta Histochem. Cytochem.* 28, 494-494, 1995.

- Joukura, K., Usuda, N., Nagata, T.: Quantitative study on the aging change of glycoconjugates synthesis in aging mouse kidney. Proc. Xth Internat. Cong. Histochem. Cytochem., Acta Histochem. Cytochem. 29, Suppl. 507-508, 1996.
- Kobayashi, K., Nagata, T.: Light microscopic radioautographic studies on DNA, RNA and protein syntheses in human synovial membranes of rheumatoid arthritis patients. J. Histochem. Cytochem. 42, 982-982, 1994.
- Komiyama, K., Iida, F., Furihara, R., Murata, F., Nagata, T.: Electron microscopic radioautographic study on  $^{125}\text{I}$ -albumin in rat gastric mucosal epithelia. J. Clin. Electron Microsc. 11, 428-429, 1978.
- Kong, Y.: Electron microscopic radioautographic study on DNA synthesis in perinatal mouse retina. Cell. Mol. Biol. 39, 55-64, 1993.
- Kong, Y., Nagata, T.: Electron microscopic radioautographic study on nucleic acid synthesis of perinatal mouse retina. Med. Electron Microsc. 27, 366-368, 1994.
- Kong, Y., Usuda, N., Nagata, T.: Radioautographic study on DNA synthesis of the retina and retinal pigment epithelium of developing mouse embryo. Cell. Mol. Biol. 38, 263-272, 1992a.
- Kong, Y., Usuda, N., Morita, T., Hanai, T., Nagata, T.: Study on RNA synthesis in the retina and retinal pigment epithelium of mice by light microscopic radioautography. Cell. Mol. Biol. 38, 669-678, 1992b.
- Leblond, C. P.: Localization of newly administered iodine in the thyroid gland as indicated by radioiodine. J. Anat. 77, 149-152, 1943.
- Leblond, C. P.: The life history of cells in renewing systems. Am. J. Anat. 160, 113-158, 1981.
- Leblond, C. P., Messier, B.: Renewal of chief cells and goblet cells in the small intestine as shown by radioautography after injection of thymidine- $^3\text{H}$  into mice. Anat. Rec. 132: 247-259. 1958.
- Li, S.: Relationship between cellular DNA synthesis, PCNA expression and sex steroid hormone receptor status in the developing mouse ovary, uterus and oviduct. Histochemistry 102, 405-413, 1994.
- Li, S., Nagata, T.: Nucleic acid synthesis in the developing mouse ovary, uterus and oviduct studied by light and electron microscopic radioautography. Cell. Mol. Biol. 41, 185-195, 1995.
- Li, S., Gao, F., Duan, H., Nagata, T.: Radioautographic study on the uptake of  $^{35}\text{SO}_4$  in mouse ovary during the estrus cycle. J. Clin. Electron Microsc. 25, 709-710, 1992.
- Liang, Y.: Light microscopic radioautographic study on RNA synthesis in the adrenal glands of aging mice. Acta Histochem. Cytochem. 31, 203-210, 1998.
- Liang, Y., Ito, M., Nagata, T.: Light and electron microscopic radioautographic studies on RNA synthesis in aging mouse adrenal gland. Acta Anat. Nippon. 74, 291-300, 1999.
- Ma, H.: Light microscopic radioautographic study on DNA synthesis of the livers in aging mice. Acta Anat. Nippon. 63, 137-147, 1988.
- Ma, H., Nagata, T.: Electron microscopic radioautographic study on DNA synthesis of the livers in aging mice. J. Clin. Electron Microsc. 21, 335-343, 1988a.
- Ma, H., Nagata, T.: Studies on DNA synthesis of aging mice by means of electron microscopic radioautography. J. Clin. Electron Microsc. 21, 715-716, 1988b.
- Ma, H., Nagata, T.: Electron microscopic radioautographic studies on DNA synthesis in the hepatocytes of aging mice as observed by image analysis. Cell. Mol. Biol. 36, 73-84, 1990a.

- Ma, H., Nagata, T.: Study on RNA synthesis in the livers of aging mice by means of electron microscopic radioautography. *Cell. Mol. Biol.* 36, 589-600, 1990b.
- Ma, H., Nagata, T.: Collagen and protein synthesis in the livers of aging mice as studied by electron microscopic radioautography. *Ann. Microsc.* 1, 13-22, 2000.
- Ma, H., Gao, F., Olea, M. T., Nagata, T.: Protein synthesis in the livers of aging mice studied by electron microscopic radioautography. *Cell. Mol. Biol.* 37, 607-615, 1991.
- Matsumura, H., Kobayashi, Y., Kobayashi, K., Nagata, T.: Light microscopic radioautographic study of DNA synthesis in the lung of aging salamander, *Hynobius nebulosus*. *J. Histochem. Cytochem.* 42, 1004-1004, 1994.
- Momose, Y., Nagata, T.: Radioautographic study on the intracellular localization of a hypolipidemic agent, bezafibrate, a peroxisome proliferator, in cultured rat hepatocytes. *Cell. Mol. Biol.* 39, 773-781, 1993a.
- Momose, Y., Naito, J., Nagata, T.: Radioautographic study on the localization of an anti-allergic agent, tranilast, in the rat liver. *Cell. Mol. Biol.* 35, 347-355, 1989.
- Momose, Y., Shibata, N., Kiyosawa, I., Naito, J., Watanabe, T., Horie, S., Yamada, J., Suga, T., Nagata, T.: Morphometric evaluation of species differences in the effects of bezafibrate, a hypolipidemic agent, on hepatic peroxisomes and mitochondria. *J. Toxicol. Pathol.* 6, 33-45, 1993b.
- Momose, Y., Naito, J., Suzawa, H., Kanzawa, M., Nagata, T.: Radioautographic study on the intracellular localization of bezafibrate in cultured rat hepatocytes. *Acta Histochem. Cytochem.* 28, 61-66, 1995.
- Morita, T.: Radioautographic study on the aging change of  $^3\text{H}$ -glucosamine uptake in mouse ileum. *Cell. Mol. Biol.* 39, 875-884, 1993.
- Morita, T., Usuda, N., Hanai, T., Nagata, T.: Changes of colon epithelium proliferation due to individual aging with PCNA/cyclin immunostaining comparing with  $^3\text{H}$ -thymidine radioautography. *Histochemistry* 101, 13-20, 1994.
- Murata, F., Momose, Y., Yoshida, K., Nagata, T.: Incorporation of  $^3\text{H}$ -thymidine into the nucleus of mast cells in adult rat peritoneum. *Shinshu Med. J.* 25, 72-77, 1977a.
- Murata, F., Momose, Y., Yoshida, K., Ohno, S., Nagata, T.: Nucleic acid and mucosubstance metabolism of mastocytoma cells by means of electron microscopic radioautography. *Acta Pharmacol. Toxicol.* 41, 58-59, 1977b.
- Murata, F., Yoshida, K., Ohno, S., Nagata, T.: Ultrastructural and electron microscopic radioautographic studies on the mastocytoma cells and mast cells. *J. Clin. Electron Microsc.* 11, 561-562, 1978.
- Murata, F., Yoshida, K., Ohno, S., Nagata, T.: Mucosubstances of rabbit granulocytes studied by means of electron microscopic radioautography and X-ray microanalysis. *Histochemistry* 61, 139-150, 1979.
- Nagata, T.: On the relationship between cell division and cytochrome oxidase in the Yoshida sarcoma cells. *Shinshu Med. J.* 5: 383-386, 1956.
- Nagata, T.: Studies on the amitosis in the Yoshida sarcoma cells. I. Observation on the smear preparation under normal conditions. *Med. J. Shinshu Univ.* 2: 187-198, 1957a.
- Nagata, T.: Studies on the amitosis in the Yoshida sarcoma cells. II. Phase-contrast microscopic observations under normal conditions. *Med. J. Shinshu Univ.* 2: 199-207, 1957b.
- Nagata, T.: Cell divisions in the liver of the fetal and newborn dogs. *Med. J. Shinshu Univ.* 4: 65-73, 1959.

- Nagata, T.: A radioautographic study of the DNA synthesis in rat liver, with special reference to binucleate cells. *Med. J. Shinshu Univ.* 7, 17-25, 1962.
- Nagata, T.: A quantitative study on the ganglion cells in the small intestine of the dog. *Med. J. Shinshu Univ.* 10, 1-11, 1965.
- Nagata, T.: A radioautographic study on the RNA synthesis in the hepatic and the intestinal epithelial cells of mice after feeding with special reference to binuclearity. *Med. J. Shinshu Univ.* 11, 49-61, 1966.
- Nagata, T.: On the increase of binucleate cells in the ganglion cells of dog small intestine due to experimental ischemia. *Med. J. Shinshu Univ.* 12, 93-113, 1967a.
- Nagata, T.: A radioautographic study on the protein synthesis in the hepatic and the intestinal epithelial cells of mice, with special reference to binucleate cells. *Med. J. Shinshu Univ.* 12, 247-257, 1967b.
- Nagata, T.: Chapter 3. Application of microspectrophotometry to various substances. In , *Introduction to Microspectrophotometry.* Isaka, S., Nagata, T., Inui, N., Eds., Olympus Co., Tokyo, pp. 49-155, 1972a.
- Nagata, T.: Electron microscopic dry-mounting autoradiography. *Proc. 4th Internat. Cong. Histochem. Cytochem.* Kyoto, pp. 43-44, 1972b.
- Nagata, T.: Electron microscopic radioautography of intramitochondrial RNA synthesis of HeLa cells in culture. *Histochemie* 32, 163-170, 1972c.
- Nagata, T.: Quantitative electron microscope radioautography of intramitochondrial nucleic acid synthesis. *Acta Histochem. Cytochem.* 5, 201-203, 1972d.
- Nagata, T.: Electron microscopic observation of target cells previously observed by phase-contrast microscopy: Electron microscopic radioautography of laser beam irradiated cultured cells. *J. Clin. Electron Microsc.* 17, 589-590, 1984.
- Nagata, T.: Principles and techniques of radioautography. In, *Histo- and Cyto-chemistry 1985*, Japan Society of Histochemistry and Cytochemistry, Ed., Gakusai Kikaku Co., Tokyo, pp. 207-226, 1985.
- Nagata, T.: Electron microscopic radioautography and analytical electron microscopy. *J. Clin. Electron Microsc.* 24, 441-442, 1991.
- Nagata, T.: Radiolabeling of soluble and insoluble compounds as demonstrated by light and electron microscopy. *Recent Advances in Cellular and Molecular Biology*, Wegmann, R. J., Wegmann, M. A., Eds. Peters Press, Leuven, Vol. 6, pp. 9-21, 1992.
- Nagata, T.: Quantitative analysis of histochemical reactions: Image analysis of light and electron microscopic radioautograms. *Acta Histochem. Cytochem.* 26, 281-291, 1993a.
- Nagata, T.: Quantitative light and electron microscopic radioautographic studies on macromolecular synthesis in several organs of prenatal and postnatal aging mice. *Chinese J. Histochem. Cytochem.* 2: 106-108, 1993b.
- Nagata, T.: Electron microscopic radioautography with cryo-fixation and dry-mounting procedure. *Acta Histochem. Cytochem.* 27: 471-489, 1994a.
- Nagata, T.: Application of electron microscopic radioautography to clinical electron microscopy. *Med. Electron Microsc.* 27; 191-212, 1994b.
- Nagata, T.: *Radioautography in Medicine.* Shinshu University Press, 268pp, Matsumoto, 1994c.
- Nagata, T.: Radioautography, general and special. In, *Histo- and Cyto-chemistry 1994*, Japan Society of Histochemistry and Cytochemistry, ed, pp. 219-231, Gakusai Kikaku Co., Tokyo, 1994d.

- Nagata, T., Application of electron microscopic radioautography to clinical electron microscopy. *Med. Electron Microsc.* 27, 191-212, 1994e.
- Nagata, T.: Light and electron microscopic radioautographic study on macromolecular synthesis in digestive organs of aging mice. *Cell. Mol. Biol.* 41, 21-38, 1995a.
- Nagata, T.: Histochemistry of the organs: Application of histochemistry to anatomy. *Acta Anat. Nippon.* 70, 448-471, 1995b.
- Nagata, T.: Three-dimensional observation of whole mount cultured cells stained with histochemical reactions by ultrahigh voltage electron microscopy. *Cell. Mol. Biol.* 41, 783-792, 1995c.
- Nagata, T.: Morphometry in anatomy: image analysis on fine structure and histochemical reactions with special reference to radioautography. *Ital. J. Anat.* 100 (Suppl. 1), 591-605, 1995d.
- Nagata, T.: Technique and application of electron microscopic radioautography. *J. Electron Microsc.* 45, 258-274, 1996a.
- Nagata, T.: Techniques of light and electron microscopic radioautography. In, *Histochemistry and Cytochemistry 1996. Proc. Xth Internat. Congr. Histochem. Cytochem. Acta Histochem. Cytochem.* 29 (Suppl.), 343-344, 1996b.
- Nagata, T.: Remarks: Radioautographology, general and special. *Cell. Mol. Biol.* 42 (Suppl.), 11-12, 1996c.
- Nagata, T.: On the terminology of radioautography vs. autoradiography. *J. Histochem. Cytochem.* 44, 1209-1209, 1996d.
- Nagata, T.: Techniques and applications of microscopic radioautography. *Histol. Histopathol.* 12, 1091-1124, 1997a.
- Nagata T.: Three-dimensional observation on whole mount cultured cells and thick sections stained with histochemical reactions by high voltage electron microscopy. In, *Recent Advances in Microscopy of Cells, Tissues and Organs*, Motta, P., Ed., Antonio Delfino Editore, Roma, pp. 37-44, 1997b.
- Nagata, T.: Radioautographic study on collagen synthesis in the ocular tissues. *J. Kaken Eye Res.* 15, 1-9, 1997c.
- Nagata, T.: Techniques of radioautography for medical and biological research. *Braz. J. Biol. Med. Res.* 31, 185-195, 1998a.
- Nagata, T.: Radioautographology, the advocacy of a new concept. *Braz. J. Biol. Med. Res.* 31, 201-241, 1998b.
- Nagata, T.: Radioautographic studies on DNA synthesis of the bone and skin of aging salamander. *Bull. Nagano Women's Jr. College* 6, 1-14, 1998c.
- Nagata, T.: 3D observation of cell organelles by high voltage electron microscopy. *Microscopy and Analysis, Asia Pacific Edition*, 9, 29-32, 1999a.
- Nagata, T.: Application of histochemistry to anatomy: Histochemistry of the organs, a novel concept. *Proc. XV Congress of the International Federation of Associations of Anatomists, Ital. J. Anat. Embryol.* 104 (Suppl. 1), 486-486, 1999b.
- Nagata, T.: Aging changes of macromolecular synthesis in various organ systems as observed by microscopic radioautography after incorporation of radiolabeled precursors. *Methods Find. Exp. Clin. Pharmacol.* 21, 683-706, 1999c.
- Nagata, T.: Radioautographic study on protein synthesis in mouse cornea. *J. Kaken Eye Res.* 8, 8-14, 1999d.
- Nagata, T.: Radioautographology, general and special: a novel concept. *Ital. J. Anat. Embryol.* 104 (Suppl. 1), 487-487, 1999e.

- Nagata, T.: Three-dimensional observations on thick biological specimens by high voltage electron microscopy. *Image Analysis Stereolog.* 19, 51-56, 2000a.
- Nagata, T.: Biological microanalysis of radiolabeled and unlabeled compounds by radioautography and X-ray microanalysis. *Scanning Microscopy International*, 14, on line, 2000b.
- Nagata, T.: Electron microscopic radioautographic study on protein synthesis in pancreatic cells of perinatal and aging mice. *Bull. Nagano Women's Jr. College* 8, 1-22, 2000c.
- Nagata, T.: Light microscopic radioautographic study on radiosulfate incorporation into the tracheal cartilage in aging mice. *Acta Histochem. Cytochem.* 32, 377-383, 2000d.
- Nagata, T.: Introductory remarks: Special radioautography. *Cell. Mol. Biol.* 46 (Congress Suppl.), 161-161, 2000e.
- Nagata, T.: Special radioautography: the eye. *J. Kaken Eye Res.* 18, 1-13, 2000f.
- Nagata, T.: Three-dimensional high voltage electron microscopy of thick biological specimens. *Micron* 32, 387-404, 2001a.
- Nagata, T.: Three-dimensional and four-dimensional observation of histochemical and cytochemical specimens by high voltage electron microscopy. *Acta Histochem. Cytochem.* 34, 153-169, 2001b.
- Nagata, T.: Special cytochemistry in cell biology. In, *Internat. Rev. Cytol.* Jeon, K.W., ed., Vol. 211, Chapter 2, Academic Press, New York, pp. 33-154, 2001c.
- Nagata, T.: Radioautography General and Special, In, *Prog. Histochem. Cytochem.*, Graumann, W., Ed., Vol. 37 No. 2, Urban & Fischer, Jena, pp. 57-226, 2002.
- Nagata T.: Light and electron microscopic study on macromolecular synthesis in amitotic hepatocyte mitochondria of aging mice. *Cell. Mol. Biol.* 49, 591-611, 2003.
- Nagata, T.: X-ray microanalysis of biological specimens by high voltage electron microscopy. In, *Prog. Histochem. Cytochem.*, Graumann, W., Ed., Vol. 39, No. 4, Urban & Fischer Verlag, Jena, pp. 185-320, 2004.
- Nagata T.: Aging changes of macromolecular synthesis in the uro-genital organs as revealed by electron microscopic radioautography. *Ann. Rev. Biomed. Sci.* 6, 13-78, 2005.
- Nagata T.: Electron microscopic radioautographic study on protein synthesis in hepatocyte mitochondria of developing mice. *Ann. Microsc.* 6, 43-54, 2006a.
- Nagata T.: Electron microscopic radioautographic study on nucleic acids synthesis in hepatocyte mitochondria of developing mice. *The Sci. World J.* 6: 1583-1598, 2006b.
- Nagata T.: Macromolecular synthesis in hepatocyte mitochondria of aging mice as revealed by electron microscopic radioautography. I: Nucleic acid synthesis. In, *Modern Research and Educational Topics in Microscopy.* Mendez-Vilas, A. and Diaz, J. Eds., Formatex Microscopy Series No. 3, Vol. 1, Formatex, Badajoz, Spain, pp. 245-258, 2007a.
- Nagata T.: Macromolecular synthesis in hepatocyte mitochondria of aging mice as revealed by electron microscopic radioautography. II: Protein synthesis. In, *Modern Research and Educational Topics in Microscopy.* Mendez-Vilas, A. and Diaz, J. eds., Formatex Microscopy Series No. 3, Vol. 1, Formatex, Badajoz, Spain, pp. 259-271, 2007b.
- Nagata, T.: Electron microscopic radioautographic study on macromolecular synthesis in hepatocyte mitochondria of aging mouse. *J. Cell Tissue Res.* 7, 1019-1029, 2007c.
- Nagata, T.: Electron microscopic radioautographic study on nucleic acids synthesis in hepatocyte mitochondria of developing mice. *Trends Cell Molec. Biol.* 2, 19-33, 2007d.

- Nagata, T.: Aging changes of macromolecular synthesis in the mitochondria of mouse hepatocytes as revealed by microscopic radioautography. *Ann. Rev. Biomed. Sci.* 9, 30-36, 2007e.
- Nagata, T.: Radioautographology, *Bull. Shinshu Institute Alternat. Med.* 2, 3-32, 2007f.
- Nagata, T.: Electron microscopic radioautographic study on mitochondrial DNA synthesis in adrenal cortical cells of developing mice. *J. Cell. Tis. Res.* 8, 1303-1312, 2008a.
- Nagata T.: Electron microscopic radioautographic study on mitochondrial DNA synthesis in adrenal cortical cells of developing and aging mice. *The Sci. World J.* 8, 683-697. 2008b.
- Nagata, T.: Sexual difference between the macromolecular synthesis of hepatocyte mitochondria in male and female mice in aging as revealed by electron microscopic radioautography. Chapter 22. In, *Women and Aging: New Research*, H. T. Bennninghouse, A. D. Rosset, Eds. Nova Biomed. Books, New York, pp. 461-487, 2009a
- Nagata, T.: Protein synthesis in hepatocytes of mice as revealed by electron microscopic radioautography. In, *Protein Biosynthesis*. Esterhouse, T. E. and Petrinis, L. B., Eds., Nova Biomed. Books, New York, pp. 133-161, 2009b.
- Nagata, T.: Electron microscopic radioautographic studies on macromolecular synthesis in mitochondria of various cells. 18EMSM Conference Proc. 9th Asia-Pacific Microscopy Conference (APMC9), Kuala Lumpur, Malaysia, pp. 48-50, 2009c.
- Nagata, T.: Recent studies on macromolecular synthesis labeled with  $^3\text{H}$ -thymidine in various organs as revealed by electron microscopic radioautography. *Current Radiopharmaceutics* 2, 118-128, 2009d.
- Nagata, T.: Electron microscopic radioautographic study on mitochondrial DNA synthesis in adrenal medullary cells of developing and aging mice. *J. Cell Tissue Res.* 9, 1793-1802, 2009e.
- Nagata, T.: Applications of high voltage electron microscopy to thick biological specimens. *Ann. Microsc.* 9, 4-40, 2009f.
- Nagata, T.: Electron microscopic radioautographic study on DNA synthesis of mitochondria in adrenal medullary cells of aging mice. *Open Anat. J.* 1, 14-24, 2009g.
- Nagata, T.: Electron microscopic radioautographic studies on macromolecular synthesis in mitochondria of animal cells in aging. *Ann. Rev. Biomed. Sci.* 11, 1-17, 2009h.
- Nagata, T.: Electron microscopic radioautographic studies on macromoleclular synthesis in mitochondria of some organs in aging animals. *Bull. Shinshu Inst. Alternat. Med. Welfare* 4, 15-38, 2009i.
- Nagata, T.: Electron microscopic radioautographic study on mitochondrial DNA synthesis in adreno-cortical cells of aging ddY mice. *Bull. Shinshu Inst. Alternat. Med. Welfare* 4, 51-66, 2009j.
- Nagata T.: Electron microscopic radioautographic study on mitochondrial RNA synthesis in adrenocortical cells of aging mice. *Open Anat J.* 2, 91-97, 2010a.
- Nagata T. Electron microscopic radioautographic study on mitochondrial RNA synthesis in adrenal medullary cells of aging and senescent mice. *J Cell Tissue Res.* 10, 2213-2222, 2010b.
- Nagata, T.: Macromolecular synthesis in the livers of aging mice as revealed by electron microscopic radioautography. In, *Prog. Histochem. Cytochem.*, Sasse, D., Ed., Elsevier, Amsterdam, Boston, London, New York, Oxford, Paris, Philadelphia, San Diego, St. Louis, Vol. 45, No. 1, pp. 1-80, 2010c.

- Nagata, T.: Electron microscopic radioautographic study on protein synthesis of mitochondria in adrenal medullary cells of aging mice. *Bulletin Shinshu Inst Alternat Med Welfare* 5, 25-37, 2010d.
- Nagata, T.: Electron microscopic radioautographic study on mitochondrial RNA synthesis in adrenal cortical and medullary cells of aging mice. *J. Biomed. Sci. Engineer.* 4, 219-232, 2010e.
- Nagata, T.: Electron microscopic radioautographic study on protein synthesis of mitochondria in adrenal cortical cells of aging mice. *Bulletin Shinshu Inst. Alternat. Med. Welfare* 5, 38-52, 2010f.
- Nagata T.: Electron microscopic radioautographic study on mitochondrial DNA, RNA and protein synthesis in adrenal cells of aging mice. *Formatex Microscopy Series No. 3, Vol. 3, Formatex, Badajoz, Spain, in press, 2010g.*
- Nagata, T.: Electron microscopic radioautographic studies on macromolecular synthesis in mitochondria of animal cells in aging. *Ann. Rev. Biomed. Sci.* 12, 1-29, 2010h.
- Nagata, T., Cui, H., Gao, F.: Radioautographic study on glycoprotein synthesis in the ocular tissues. *J. Kaken Eye Res.* 13, 11-18, 1995.
- Nagata, T., Cui, H., Kong, Y.: The localization of TGF- $\beta$ 1 and its mRNA in the spinal cords of prenatal and postnatal aging mice demonstrated with immunohistochemical and in situ hybridization techniques. *Bull. Nagano Women's Jr. College*, 7, 75-88, 1999a.
- Nagata, T., Cui, H., Liang, Y.: Light microscopic radioautographic study on the protein synthesis in the cerebellum of aging mouse. *Bull. Nagano Women's Jr. College*, 9, 41-60 (2001).
- Nagata, T., Fujii, Y., Usuda, N.: Demonstration of extranuclear nucleic acid synthesis in mammalian cells under experimental conditions by electron microscopic radioautography. *Proc. 10th Internat. Congr. Electr. Microsc.* 2, 305-306, 1982b.
- Nagata, T., Hirano, I., Shibata, O., Nagata, T.: A radioautographic study on the DNA synthesis in the hepatic and the pancreatic acinar cells of mice during the postnatal growth, with special reference to binuclearity. *Med. J. Shinshu Univ.* 11, 35-42, 1966.
- Nagata, T., Ito, M., Chen, S.: Aging changes of DNA synthesis in the submandibular glands of mice as observed by light and electron microscopic radioautography. *Ann. Microsc.* 1, 4-12, 2000a.
- Nagata, T. Ito, M., Liang, Y.: Study of the effects of aging on macromolecular synthesis in mouse steroid secreting cells using microscopic radioautography. *Methods Find. Exp. Clin. Pharmacol.* 22, 5-18, 2000b.
- Nagata, T., Iwadare, I., Murata, F.: Electron microscopic radioautography of nucleic acid synthesis in cultured cells treated with several carcinogens. *Acta Pharmacol. Toxicol.* 41, 64-65, 1977c.
- Nagata, T., Kawahara, I.: Radioautographic study of the synthesis of sulfomucin in digestive organs of mice. *J. Trace Microprobe Analysis* 17, 339-355, 1999.
- Nagata, T., Kawahara, I., Usuda, N., Maruyama, M., Ma, H.: Radioautographic studies on the glycoconjugate synthesis in the gastrointestinal mucosa of the mouse. In, *Glycoconjugate in Medicine*, Ohyama, M., Muramatsu, T., Eds, pp. 251-256, Professional Postgrad. Service, Tokyo, 1988a.
- Nagata, T., Kong, Y.: Distribution and localization of TGF $\beta$ 1 and bFGF, and their mRNAs in aging mice. *Bull. Nagano Women's Jr. College* 6, 87-105, 1998.

- Nagata, T., Ma, H., Electron microscopic radioautographic study on mitochondrial DNA synthesis in hepatocytes of aging mouse. *Ann. Microsc.* 5, 4-18, 2005a.
- Nagata, T., Ma, H., Electron microscopic radioautographic study on RNA synthesis in hepatocyte mitochondria of aging mouse. *Microsc. Res. Tech.* 67, 55-64, 2005b.
- Nagata, T., Momoze, S.: Aging changes of the amitotic and binucleate cells in dog livers. *Acta Anat. Nipponica* 34, 187-190, 1959.
- Nagata, T., Morita, T., I. Kawahara, I.: Radioautographic studies on radi sulfate incorporation in the digestive organs of mice. *Histol. Histopathol.* 14, 1-8, 1999b.
- Nagata, T., Murata, F.: Electron microscopic dry-mounting radioautography for diffusible compounds by means of ultracryotomy. *Histochemistry* 54, 75-82, 1977.
- Nagata, T., Murata, F., Yoshida, K., Ohno, S., Iwadare, N.: Whole mount radioautography of cultured cells as observed by high voltage electron microscopy. *Proc. Fifth Internat. Conf. High Voltage Electron Microsc.* 347-350, 1977d.
- Nagata, T., Nawa, T.: A modification of dry-mounting technique for radioautography of water-soluble compounds. *Histochemie* 7, 370-371, 1966a.
- Nagata, T., Nawa, T.: A radioautographic study on the nucleic acids synthesis of binucleate cells in cultivated fibroblasts of chick embryos. *Med. J. Shinshu Univ.* 11, 1-5, 1966b.
- Nagata, T., Nawa, T., Yokota, S.: A new technique for electron microscopic dry-mounting radioautography of soluble compounds. *Histochemie* 18, 241-249, 1969.
- Nagata, T., Nishigaki, T., Momose, Y.: Localization of anti-allergic agent in rat mast cells demonstrated by light and electron microscopic radioautography. *Acta Histochem. Cytochem.* 19, 669-683, 1986b.
- Nagata, T., Ohno, S., Kawahara, I., Yamabayashi, S., Fujii, Y., Murata, F.: Light and electron microscopic radioautography of nucleic acid synthesis in mitochondria and peroxisomes of rat hepatic cells during and after DEHP administration. *Acta Histochem. Cytochem.* 16, 610-611, 1979.
- Nagata, T., Ohno, S., Murata, F.: Electron microscopic dry-mounting radioautography for soluble compounds. *Acta Pharmacol. Toxicol.* 41, 62-63, 1977a.
- Nagata, T., Ohno, S., Yoshida, K., Murata, F.: Nucleic acid synthesis in proliferating peroxisomes of rat liver as revealed by electron microscopical radioautography. *Histochem. J.* 14, 197-204, 1982a.
- Nagata, T., Olea, M. T.: Electron microscopic radioautographic study on the protein synthesis in aging mouse spleen. *Bull. Nagano Women's Jr. College* 7, 1-9, 1999.
- Nagata, T., Shibata, O., Omochi, S.: A new method for radioautographic observation on isolated cells. *Histochemie* 2, 255-259, 1961
- Nagata, T., Shibata, O., Nawa, T.: Simplified methods for mass production of radioautograms. *-Acta Anat. Nippon.* 42, 162-166, 1967a.
- Nagata, T., Shibata, O., Nawa, T.: Incorporation of tritiated thymidine into mitochondrial DNA of the liver and kidney cells of chickens and mice in tissue culture. *Histochemie* 10, 305-308, 1967b.
- Nagata, T., Shimamura, K., Onozawa, M., Kondo, T., Ohkubo, K., Momoze, S.: Relationship of binuclearity to cell function in some organs. I. Frequencies of binucleate cells in some organs of toads in summer and winter. *Med. J. Shinshu Univ.* 5, 147-152, 1960a.
- Nagata, T., Shimamura, K., Kondo, T., Onozawa, M., Momoze, S., Okubo, M.: Relationship of binuclearity to cell function in some organs. II. Variation of frequencies of

- binucleate cells in some organs of dogs owing to aging. *Med. J. Shinshu Univ.* 5, 153-158, 1960b.
- Nagata, T., Steggerda, F. R.: Histological study on the deganglionated small intestine of the dog. *Physiologist* 6, 242-242, 1963.
- Nagata, T., Steggerda, F. R.: Observations on the increase of binucleate cells in the ganglion cells of the dog's intestine due to experimental ischemia. *Anat. Rec.* 148, 315-315, 1964.
- Nagata, T., Toriyama, K., Kong, Y., Jin, C., Gao, F.: Radioautographic study on DNA synthesis in the ciliary bodies of aging mice. *J. Kaken Eye Res.* 12, 1-11, 1994.
- Nagata, T., Usuda, N.: Image processing of electron microscopic radioautograms in clinical electron microscopy. *J. Clin. Electron. Microsc.* 18, 451-452, 1985.
- Nagata, T., Usuda, N.: Studies on the nucleic acid synthesis in pancreatic acinar cells of aging mice by means of electron microscopic radioautography. *J. Clin. Electron Microsc.* 19, 486-487, 1986.
- Nagata, T., Usuda, N.: Electron microscopic radioautography of protein synthesis in pancreatic acinar cells of aging mice. *Acta Histochem. Cytochem.* 26, 481-481, 1993a.
- Nagata, T., Usuda, N.: In situ hybridization by electron microscopy using radioactive probes. *J. Histochem. Cytochem.* 41, 1119-1119, 1993b.
- Nagata, T., Usuda, N., Ma, H.: Electron microscopic radioautography of nucleic acid synthesis in pancreatic acinar cells of prenatal and postnatal aging mice. *Proc. XIth Intern. Cong. Electr. Microsc.* 3, 2281-2282, 1984.
- Nagata, T., Usuda, N., Ma, H.: Electron microscopic radioautography of lipid synthesis in pancreatic cells of aging mice. *J. Clin. Electr. Microsc.* 23, 841-842, 1990.
- Nagata, T., Usuda, N., Maruyama, M., Ma, H.: Electron microscopic radioautographic study on lipid synthesis in perinatal mouse pancreas. *J. Clin. Electr. Microsc.* 21, 756-757, 1988b.
- Nagata, T., Usuda, N., Suzawa, H., Kanzawa, M.: Incorporation of  $^3\text{H}$ -glucosamine into the pancreatic cells of aging mice as demonstrated by electron microscopic radioautography. *J. Clin. Electron Microsc.* 25, 646-647, 1992.
- Nagata, T., Yamabayashi, S.: Intracellular localization of  $^3\text{H}$ -bepunolol by means of electron microscopic radioautography of cryo-fixed ultrathin sections. *J. Clin. Electron Microsc.* 16, 737-738, 1983.
- Nagata, T., Yoshida, K., Murata, F.: Demonstration of hot and cold mercury in the human thyroid tissues by means of radioautography and chemography. *Acta Pharmacol. Toxicol.* 41, 60-61, 1977b.
- Nagata, T., Yoshida, K., Ohno, S., Murata, F.: Ultrastructural localization of soluble and insoluble  $^3\text{H}$ -methyl prednisolone as revealed by electron microscopic dry-mounting radioautography. *Proc. 9th Internat. Congr. Electr. Microsc.* 2, 40-41, 1978b.
- Nishigaki, T., Momose, Y., Nagata, T.: Light microscopic radioautographic study of the localization of anti-allergic agent, tranilast, in rat mast cells. *Histochem. J.* 19, 533-536, 1987.
- Nishigaki, T., Momose, Y., Nagata, T.: Electron microscopic radioautographic study of the localization of an anti-allergic agent, tranilast, in rat mast cells. *Cell. Mol. Biol.* 36, 65-71, 1990a.

- Nishigaki, T., Momose, Y., Nagata, T.: Localization of the anti-allergic agent tranilast in the urinary bladder of rat as demonstrated by light microscopic radioautography. *Drug Res.* 40, 272-275, 1990b.
- Oguchi, K., Nagata, T.: A radioautographic study of activated satellite cells in dystrophic chicken muscle. In, *Current Research in Muscular Dystrophy Japan. The Proc. Ann. Meet. Muscular Dystrophy Res.* 1980, pp. 16-17, Ministry of Welfare of Japan, Tokyo, 1980.
- Oguchi, K., Nagata, T.: Electron microscopic radioautographic observation on activated satellite cells in dystrophy chickens. In, *Clinical Studies on the Etiology of Muscular Dystrophy. Annual Report on Neurological Diseases* 1981, pp. 30-33, Ministry of Welfare of Japan, Tokyo, 1981.
- Ohno, S., Fujii, Y., Usuda, N., Endo, T., Hidaka, H., Nagata, T.: Demonstration of intracellular localization of calmodulin antagonist by wet-mounting radioautography. *J. Electron Microsc.* 32, 1-12, 1983.
- Ohno, S., Fujii, Y., Usuda, N., Nagata, T., Endo, T., Tanaka, T., Hidaka, H.: Intracellular localization of calmodulin antagonists (W-7). In, *Calmodulin and intracellular Ca<sup>2+</sup> receptors.* Kakiuchi, S., Hidaka, H, Means, A. R., Eds., pp. 39-48, Plenum Publishing Co., New York, 1982.
- Olea, M. T.: An ultrastructural localization of lysosomal acid phosphatase activity in aging mouse spleen: a quantitative X-ray microanalytical study. *Acta Histochem. Cytochem.* 24, 201-208, 1991.
- Olea, M. T., Nagata, T.: X-ray microanalysis of cerium in mouse spleen cells demonstrating acid phosphatase activity using high voltage electron microscopy, *Cell. Mol. Biol.* 37, 155-163, 1991.
- Olea, M. T., Nagata, T. : Simultaneous localization of <sup>3</sup>H-thymidine incorporation and acid phosphatase activity in mouse spleen: EM radioautography and cytochemistry. *Cell. Mol. Biol.* 38, 115-122, 1992a.
- Olea, M. T., Nagata, T.: A radioautographic study on RNA synthesis in aging mouse spleen after <sup>3</sup>H-uridine labeling in vitro. *Cell. Mol. Biol.* 38, 399-405, 1992b.
- Oliveira, S. F., Nagata, T., Abrahamsohn, P. A., Zorn, T. M. T.: Electron microscopic radioautographic study on the incorporation of <sup>3</sup>H-proline by mouse decidual cells. *Cell. Mol. Biol.* 37, 315-323, 1991.
- Oliveira, S. F., Abrahamsohn, P. A., Nagata, T., Zorn, T. M. T.: Incorporation of <sup>3</sup>H-amino acids by endometrial stromal cells during decidualization in the mouse. A radioautographical study. *Cell. Mol. Biol.* 41, 107-116, 1995.
- Pearse, A. G. E.: *Histochemistry, Theoretical and Applied.* 4th Ed. Vol. 1. 439 pp., 1980, Vol. 2. 1055 pp., 1985, Vol. 3. Ed. with P. Stoward, 728 pp. Churchill Livingstone, Edinburgh, London and New York, 1991.
- Sakai, Y., Ikado, S., Nagata, T.: Electron microscopic radioautography of satellite cells in regenerating muscles. *J. Clin. Electr. Microsc.* 10, 508-509, 1977.
- Sato, A.: Quantitative electron microscopic studies on the kinetics of secretory granules in G-cells. *Cell Tissue Res.* 187, 45-59, 1978.
- Sato, A., Iida, F., Furihara, R., Nagata, T.: Electron microscopic radioautography of rat stomach G-cells by means of <sup>3</sup>H-amino acids. *J. Clin. Electron Microsc.* 10, 358-359, 1977.

- Shimizu, T., Usuda, N., Yamanda, T., Sugeno, A., Iida, F.: Proliferative activity of human thyroid tumors evaluated by proliferating cell nuclear antigen/cyclin immunohistochemical studies. *Cancer* 71, 2807-2812, 1993.
- Sun, L.: Age related changes of RNA synthesis in the lungs of aging mice by light and electron microscopic radioautography. *Cell. Mol. Biol.* 41, 1061-1072, 1995.
- Sun, L., Gao, F., Duan, H., Nagata, T.: Light microscopic radioautography of DNA synthesis in pulmonary cells in aging mice. In, *Radioautography in Medicine*, Nagata, T. Ed., pp. 201-205, Shinshu University Press, Matsumoto, 1994.
- Sun, L., Gao, F., Nagata, T.: Study on the DNA synthesis of pulmonary cells in aging mice by light microscopic radioautography. *Cell. Mol. Biol.* 41, 851-859, 1995a.
- Sun, L., Gao, F., Jin, C., Duan, H., Nagata, T.: An electron microscopic radioautographic study on the DNA synthesis of pulmonary tissue cells in aging mice. *Med. Electron. Microsc.* 28, 129-131, 1995b.
- Sun, L., Gao, F., Jin, C., Nagata, T.: DNA synthesis in the tracheae of aging mice by means of light and electron microscopic radioautography. *Acta Histochem. Cytochem.* 30, 211-220, 1997a.
- Sun, L., Gao, F., Nagata, T.: A Light Microscopic radioautographic study on protein synthesis in pulmonary cells of aging mice. *Acta Histochem. Cytochem.* 30, 463-470, 1997b.
- Suzuki, K., Imada, T., Gao, F., Ma, H., Nagata, T.: Radioautographic study of benidipine hydrochloride: localization in the mesenteric artery of spontaneously hypertensive rat. *Drug Res.* 44, 129-133, 1994.
- Terauchi, A., Mori, T., Kanda, H., Tsukada, M., Nagata, T.: Radioautographic study of  $^3\text{H}$ -taurine uptake in mouse skeletal muscle cells. *J. Clin. Electron Microsc.* 21, 627-628, 1988.
- Terauchi, A., Nagata, T.: Observation on incorporation of  $^3\text{H}$ -taurine in mouse skeletal muscle cells by light and electron microscopic radioautography. *Cell. Mol. Biol.* 39, 397-404, 1993.
- Terauchi, A., Nagata, T.: In corporation of  $^3\text{H}$ -taurine into the blood capillary cells of mouse skeletal muscle. *Radioautography in Medicine*, Nagata, T. ed., Shinshu University Press, Matsumoto, 1994.
- Toriyama, K.: Study on the aging changes of DNA and protein synthesis of bipolar and photo-receptor cells of mouse retina by light and electron microscopic radioautography. *Cell. Mol. Biol.* 41, 593-601, 1995.
- Tsukahara, S., Yoshida, K., Nagata, T.: A radioautographic study on the incorporation of  $^{14}\text{C}$ -bupranolol (beta-blocking agent) into the rabbit eye. *Histochemistry* 68, 237-244, 1980.
- Usuda, N., Nagata, T.: Electron microscopic radioautography of acyl-CoA mRNA by in situ hybridization. *J. Clin. Electron Microsc.* 25, 332-333, 1992.
- Usuda, N., Nagata, T.: The immunohistochemical and in situ hybridization studies on hepatic peroxisomes. *Acta Histochem. Cytochem.* 28, 169-172, 1995.
- Usuda, N., Hanai, T., Morita, T., Nagata, T.: Radioautographic demonstration of peroxisomal acyl-CoA oxidase mRNA by in situ hybridization. In, *Recent advances in cellular and molecular biology*, Vol. 6. Molecular biology of nucleus, peroxisomes, organelles and cell movement. Wegmann, R. J., Wegmann, M., Eds, pp.181-184, Peeters Press, Leuven, 1992.

- Uwa, H., Nagata, T.: Cell population kinetics of the scleroblast during ethisterone-induced anal-fin process formation in adult females of the Medaka. *Dev. Growth Different.* 9, 693-694, 1976.
- Watanabe, I., Makiyama, M. C. K., Nagata, T.: Electron microscopic radioautographic observation of the submandibular salivary gland of aging mouse. *Acta Microscopica* 6. 130-131, 1997.
- Yamabayashi, S., Gunarso, W., Tsukahara, S., Nagata, T.: Incorporation of  $^3\text{H}$ -bepunolol (beta blocking agent) into melanin granules of ocular tissues in the pigmented rabbits. I. Light microscopic radioautography. *Histochemistry* 73, 371-375, 1981.
- Yamada, A. T.: Timely and topologically defined protein synthesis in the periimplanting mouse endometrium revealed by light and electron microscopic radioautography. *Cell. Mol. Biol.* 39, 1-12, 1993.
- Yamada, A., Nagata, T.: Ribonucleic acid and protein synthesis in the uterus of pregnant mouse during activation of implantation window. *Med. Electron Microsc.* 27, 363-365, 1992a.
- Yamada, A., Nagata, T.: Light and electron microscopic radioautography of DNA synthesis in the endometria of pregnant-ovariectomized mice during activation of implantation window. *Cell. Mol. Biol.* 38, 763-774, 1992b.
- Yamada, A., Nagata, T.: Light and electron microscopic radioautography of RNA synthesis of peri-implanting pregnant mouse during activation of receptivity for blastocyst implantation. *Cell. Mol. Biol.* 38, 211-233, 1993.
- Yoshinaga, K.: Uterine receptivity for blastocyst implantation. *Ann. N. Y. Acad. Sci. USA*, 541, 424-431, 1988.
- Yoshizawa, S., Nagata, A., Honma, T., Oda, M., Murata, F., Nagata, T.: Study of ethionine pancreatitis by means of electron microscopic radioautography. *J. Clin. Electron Microsc.* 7, 349-350, 1974.
- Yoshizawa, S., Nagata, A., Honma, T., Oda, M., Murata, F., Nagata, T.: Radioautographic study of protein synthesis in pancreatic exocrine cells of alcoholic rats. *J. Clin. Electron. Microsc.* 10, 372-373, 1977.



## **Senescence**

Edited by Dr. Tetsuji Nagata

ISBN 978-953-51-0144-4

Hard cover, 850 pages

**Publisher** InTech

**Published online** 29, February, 2012

**Published in print edition** February, 2012

The book "Senescence" is aimed to describe all the phenomena related to aging and senescence of all forms of life on Earth, i.e. plants, animals and the human beings. The book contains 36 carefully reviewed chapters written by different authors, aiming to describe the aging and senescent changes of living creatures, i.e. plants and animals.

### **How to reference**

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Tetsuji Nagata (2012). Macromolecular Synthesis in the Urinary and Reproductive Systems, *Senescence*, Dr. Tetsuji Nagata (Ed.), ISBN: 978-953-51-0144-4, InTech, Available from:  
<http://www.intechopen.com/books/senescence/macromolecular-synthesis-in-the-urinary-and-reproductive-systems>

**INTECH**  
open science | open minds

### **InTech Europe**

University Campus STeP Ri  
Slavka Krautzeka 83/A  
51000 Rijeka, Croatia  
Phone: +385 (51) 770 447  
Fax: +385 (51) 686 166  
[www.intechopen.com](http://www.intechopen.com)

### **InTech China**

Unit 405, Office Block, Hotel Equatorial Shanghai  
No.65, Yan An Road (West), Shanghai, 200040, China  
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元  
Phone: +86-21-62489820  
Fax: +86-21-62489821

© 2012 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.