

---

# Effect of Mucilage Extraction on the Functional Properties of Chia Meals

---

Marianela I. Capitani, Susana M. Nolasco and  
Mabel C. Tomás

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/53171>

---

## 1. Introduction

Chia (*Salvia hispanica* L.) is an annual herbaceous plant that belongs to the *Lamiaceae* family, which is native to southern Mexico and northern Guatemala. The *Salvia hispanica* fruit consists of four nutlets, similar to an indehiscent achene, which contain a single seed. These nutlets are commonly called “seeds” [1]. Chia seed, together with corn, beans, and amaranth were important crops for pre-Columbian civilizations in America, including the Mayan and Aztec populations [2, 3]. With time its use was abandoned, but by at the end of the last century there was a resurgence of interest in chia due to its nutritional value [4]. Chia is considered an alternative crop to diversify and stabilize the economy of Northwestern Argentina [5]. The plant produces numerous small white and dark seeds that mature in autumn [6]. These seeds contain about 30% oil, and they mainly consist of unsaturated fatty acids [4, 7]. Chia seeds are a natural source of omega-3 fatty acids, antioxidants, proteins, vitamins, minerals and dietary fiber [5, 7, 8].

Chia meal (residue of the seeds after oil extraction) is a good source of proteins (19-23%) [9], dietary fiber (33.9-39.9%) [10], and compounds with antioxidant activity [7]. It also exhibits some interesting functional properties for its use in the food industry [11]. Functional properties are generally associated with the presence of proteins [12] and also of dietary fiber [13-16].

Dietary fiber (DF) consists of a heterogeneous mixture of compounds that are classified according to their physical properties and effects of their intake into: soluble dietary fiber (SDF) and insoluble dietary fiber (IDF) [17], referring to the solubility of fibers in water. Plant secretions such as pectins and gums, components such as mucilage, and chelating

agents such as phytates are sources of SDF; whereas cellulose, lignin, some fractions of hemicellulose, phenolic compounds such as tannins and lipid structures such as waxes, suberins and cutins constitute IDF [18].

The functional properties of food components can be defined as any physicochemical property that affects and/or modifies some of its characteristics and that contributes to the quality of the final product. Knowledge about of functional properties such as color, particle size, water holding, absorption and adsorption capacity, as well as those linked to the affinity for lipid components is very useful for the food industry, because during the processing some modifications can occur that must be taken into account according to the usage of the final product and its marketing conditions [19]. For example, water-holding capacity (WHC) is related to the freshness and softness effect present in bakery products, and the oil-holding capacity (OHC) is related to the un-fatty effect in fried food when it is low and to the juiciness and texture in meat products when it is high [18, 20, 21].

In addition to the characteristics mentioned above, it is important to consider the physiological effects of the DF intake. Given the capacity of SDF to form gels, it increases the viscosity of the bolus in the gastrointestinal tract, slowing the intestinal transit, making digestion and the absorption of nutrients more efficient, providing more of a feeling of satiety. Soluble fiber are fermentable fibers that can be microbiologically decomposed in the colon, producing gases such as carbon dioxide, hydrogen and methane, and short-chain fatty acids (acetic, propionic and butyric) which are absorbed and used as energy sources. Some of the most important beneficial effects of SDF is that it regulates blood sugar and lower cholesterol levels. On the other hand, IDF is responsible for adding bulk to the stool, speeding the passage of stool through the intestine by promoting peristalsis, alleviating constipation and other gastrointestinal disorders [22, 23]. Both types of fiber may also reduce the risk of obesity, hypertension, appendicitis, and other disorders [24]. The beneficial effects noted above show the important role that DF play in human intake, and that is why a daily intake of 25-30 g is recommended, with a good SDF/IDF balance (a minimum of 30% SDF and 70% IDF, optimum 50/50 ratio) in order to benefit from both fractions of fiber [25, 26].

Chia mucilage (SDF), a complex carbohydrate of high molecular weight, is an important component of the seed due to its physiological role. The mucilage is secreted when the seed comes into contact with water, generating high-viscosity solutions [27, 28]. Many studies have examined the functional properties of different types of gums (*Linum usitatissimum*, *Opuntia Picus indica*, *Alyssum homolocarpum*, *Psyllium plantago*) [29-32]. However, little information has been reported on the functionality of chia seed mucilage as a stabilizing or thickening agent of food products.

The objective of the present work was to perform a comparative evaluation of the functional properties of chia meals (*Salvia hispanica* L.) obtained from seeds with and without mucilage.

## 2. Materials and methods

### 2.1. Seeds

Chia seeds were obtained from commercial sources in Salta, Argentina (25° S and 65.5° W). They were cleaned manually by removing the foreign matter such as stones, dirt and broken seeds. They were packed in hermetic plastic vessels and stored at 5°C until further use.

### 2.2. Meal without mucilage (Msm)

Meal without mucilage refers to the residue obtained after the oil extraction process using a Soxhlet apparatus following the IUPAC Standard Method [33] (*n*-hexane under reflux, 8 h, 90 °C,) of seeds that had previously had the mucilage extracted.

#### 2.2.1. Mucilage extraction

The mucilage was extracted of chia seeds previously soaked in water (1:4) for 4 hours at room temperature. This mixture was distributed into plastic trays and covered with aluminum foil and frozen at -80°C, lyophilized, and the mucilage was removed by a sieving process (20 sieve mesh ASTM, 840 µm) (3 sections of 15 min each).

### 2.3. Meal with mucilage (Ms)

The data corresponding to the meal with mucilage that was considered in the analysis of the results corresponds to that reported by Capitani *et al.* [11]. The meal was obtained after oil solvent extraction (*n*-hexane) in a Soxhlet apparatus (Buenos Aires, Argentina) by thermal cycles at 80°C for 8 h, following the IUPAC Standard Method [33], of chia seeds previously ground in a laboratory grinder (Moulinex, horizontal blade grinder, Buenos Aires, Argentina).

Meals (Msm and Ms) were homogenized and stored in plastic vessels at 5°C until further use.

### 2.4. Scanning Electron Microscopy (SEM)

The whole seeds and seeds after mucilage removal were adhered to a cover slip, coated with a thin gold film (600 Å) in a sputter coater (Pelco 91000) and observed in a scanning electron microscope (LEO model EVO 40) at 5 kV. Longitudinal sections were sliced with a razor blade, after being plunged into liquid nitrogen to ensure the maintenance of their internal structure, and analyzed by microscopy using the same procedure and magnification ranges between x136 and x5000.

### 2.5. Characterization of meals

#### 2.5.1. Proximate composition

Moisture, crude fiber and ash content were determined according to AOCS recommended practices Ba 2<sup>a</sup>-38, Ba 6-84 and Ba 5<sup>a</sup>-49, respectively [34]. Oil and nitrogen content (N) were

determined following IUPAC Standard Method [33] and AOAC Method [35], respectively. Protein content was calculated as nitrogen  $\times$  6.25. Carbohydrate content was estimated by calculating the nitrogen-free extract (NFE) by difference using Eq. (1).

$$\text{NFE:100} - (\text{oil} + \text{protein} + \text{crude fiber} + \text{ash}) \quad (1)$$

### 2.5.2. Total, soluble and insoluble dietary fiber

Total dietary fiber (TDF), soluble dietary fiber (SDF) and insoluble dietary fiber (IDF) were determined according to the enzymatic gravimetric method [36].

### 2.5.3. Neutral Detergent fiber (NDF), Acid Detergent fiber (ADF), lignin, cellulose and hemicellulose

The vegetable cell was separated into two parts (Van Soest method): cell content (highly digestible) and cell wall (partially digestible). The cell wall was analyzed and its components (cellulose, hemicellulose and lignin) were determined. The technique makes use of acidic and neutral detergent [35, 37].

### 2.5.4. Antioxidant activity

The extraction of phenolic compounds was carried out according to the method of Re *et al.* [38]. Ten mL ethanol were added to 1 g sample, then it was homogenized in Vortex for 2 min, decanted and filtered (0.45  $\mu\text{m}$  nylon paper). The supernatant was transferred into a flask and evaporated using a rotavapor apparatus (BUCHI R124, Germany) to concentrate the sample. It was then redissolved in 1000  $\mu\text{L}$  ethanol.

A spectrophotometric method was used to determine the antioxidant activity using a Hitachi U-1900 UVeVIS spectrophotometer (Japan). The antioxidant activity was quantified by a dyeing assay of the radical cation ABTS<sup>+</sup> measuring ABTS<sup>+</sup> reduction as the percentage of absorption inhibition at 734 nm, just 6 min later. The radical cation ABTS and potassium persulfate were obtained from Sigma Aldrich. Chlorogenic acid was used as standard antioxidant. Results were expressed as  $\mu\text{mol/L}$  Trolox g/sample, considering that chlorogenic acid diminishes twice the amount of absorption than Trolox [39].

### 2.5.5. Functional properties

#### 2.5.5.1. Water-Holding (WHC) and Oil-Holding Capacity (OHC)

Water and oil holding capacities were determined according to the method of Chau *et al.* [40]. Briefly, 1 g (dry base (d.b.)) sample was weighed and then stirred into 10 mL distilled water or corn oil (density 0.92 g/mL, Arcor). These suspensions were centrifuged at 2200  $\times$  g for 30 min (Rolco Centrifuge Refrigerate, Model CR-5850, 22 cm radius, Buenos Aires, Argentina) and the supernatant volumes were measured. Water-holding capacity was expressed as gram water held per gram sample, and oil-holding capacity as gram oil held per gram sample.

#### 2.5.5.2. Water Absorption Capacity ( $WA_bC$ )

This property was determined according to the AACC method 88-04 [41]. Approximate water absorption capacity was first determined by weighing out 2 g (d.b.) sample, adding water until saturation (approx. 35 mL) and centrifuging at  $2000 \times g$  for 10 min in a Rolco Model CR-5850, 22-cm radius centrifuge (Buenos Aires, Argentina). Approximate water absorption capacity was calculated by dividing the increase in sample weight (g), by initial weight, quantifying the water needed to complete the original sample weight (2 g d.b.) to 15 g. Water absorption capacity ( $WA_bC$ ) was then determined by placing samples in four tubes, adding different quantities of water (1.5 and 0.5 mL water above original weight, and 1.5 and 0.5 mL water below; one in each tube), agitating vigorously, and centrifuging the samples at  $2000 \times g$  for 10 min in a Rolco Model CR 5850. The supernatant was discarded and the residue weighed. Average water absorbed was calculated, and  $WA_bC$  was determined and expressed as gram water absorbed per gram sample.

#### 2.5.5.3. Organic Molecule Absorption Capacity (OMAC)

This capacity was determined according to the method of Zambrano *et al.* [19]. A three gram (d b.) sample was placed in excess quantity corn oil (approx. 25 mL) for 24 h at room temperature, and then centrifuged at  $2000 \times g$  for 15 min in a Rolco Model CR-5850. OMAC was expressed as the absorbed hydrophobic component and calculated in terms of sample weight gain (g oil/sample g).

#### 2.5.5.4. Emulsifying Activity (EA) and Emulsion Stability (ES)

These properties were evaluated according to the method [36] of Chau *et al.* [40]. Briefly, 100 mL 2 g/100 mL suspension was homogenized using an Ultra-Turrax T25 disperser (Janke & Kunkel, IKA-Labortechnik, Germany) at 7800 rpm for 2 min. Then, 100 mL corn oil (density 0.92 g/mL, Arcor) were added and homogenized at 15,000 rpm for 2 min. Emulsions were centrifuged in a 15 mL graduated centrifuge tube at  $455 \times g$  for 10 min, and then emulsion volume was measured. The EA was expressed as the remaining volume of the centrifuged emulsion corresponding to 100 mL of initial emulsion. The emulsion stability was determined by heating the emulsions to  $80^\circ\text{C}$  for 30 min, cooling them to room temperature and then centrifuging the samples at  $455 \times g$  for 10 min. ES was expressed as the remaining volume of the centrifuged emulsion corresponding to 100 mL of initial emulsion. On the other hand, all emulsions were evaluated by optical characterization using a Vertical Scan Analyzer (QuickSCAN, Beckman Coulter, Fullerton, USA). The QuickSCAN head scans the entire length of the sample (approximately 65 mm), collecting backscattering (BS) data every  $40\mu\text{m}$ . Thus, it is possible to obtain curves showing the percentage of backscattering light flux, relative to external standards, as a function of the sample height in mm [42]. Coalescence kinetics were determined by measuring the mean values of BS as a function of time in the 25-30 mm zone (Backscattering % 25-30 mm).

## 2.6. Statistical analysis

The results obtained were analyzed using ANOVA and Tukey's test ( $p \leq 0.05$ ), using Infostat software [43].

## 3. Results and discussion


The nutlet of *Salvia hispanica* consists of the seed and a pericarp surrounding the seed. The true seed, in turn, consists of a coat (testa), the endosperm and the embryo, consisting mainly of two cotyledons [1]. Basically, the pericarp of the chia seed is similar to that of other Nepeptideae because it shows cuticle, exocarp, mesocarp, layers of sclereids and endocarp. The cells of the mesocarp and exocarp are parenchymatic. Figure 1 shows a scanning electron microscopy of a *Salvia hispanica* nutlet. In the exocarp there often are cells that produce mucilage when the nutlets get wet (Figure 1D).

Figure 2 shows SEM microscopy of *Salvia hispanica* L. seeds after mucilage extraction. In these images it can be observed that the mixocarp phenomenon occurs in the outer layers (cuticle and exocarp). After removing the mucilage, the nutlet surface is characterized by small hill-like eminences, spaced, that cover the entire surface, corresponding to the mesocarp cells. Chia seeds presented a similar structure to that of two mucilaginous species (*Carrichtera annua* and *Anastatica hierochuntica*), which could be associated with the presence of concentric aggregates of glucuronic acid [44]. The retention of the mucilage close to the seed can be due to the association of the mucilage with the columella (a secondary wall cell produced after mucilage secretion) and portions of the cell wall [45].


The proximate composition of chia meals with and without mucilage is presented in Table 1. Both meals were characterized by a high protein content, higher than that reported for sunflower meals of different origin (20.6-23.1%) [46] and canola meals (36.1-40.0%) [47, 48], and within the range of the values reported for linseed meals (38.9-43.3%) [48, 49].

On the other hand, both types of meals presented a high crude fiber content, with values higher than those reported for sesame, soybean, linseed and canola meals, 5.8%, 3.5%, 5.27% and 11.54%, respectively [50, 48].

In Table 2 it is possible to observe that both types of meals presented a high content of TDF, consisting mainly of IDF. Even though the value of SDF was relatively low, this could be attributed to the fact that, during the determination of this fiber, some components were not quantified because they cannot precipitate during the treatment with ethanol, and thus SDF was underestimated [26]. It is worth noting that the meal obtained from seeds that previously had their mucilage extracted (Msm) exhibited a statistically higher content of IDF ( $p < 0.05$ ) than that for Ms, at the expense of a significant decrease in its SDF content. These results are consistent with the data obtained from the analysis of NDF, consisting of cellulose, hemicellulose and lignin (structural polysaccharides that contribute to the IDF fraction), which was statistically higher ( $p < 0.05$ ) in Msm (Table 3). Regarding Ms, it presented a better IDF/SDF balance, with a 89/11 ratio.



**Figure 1.** SEM microscopy of *Salvia hispanica* nutlets. (A) Whole nutlet, lateral view (x150), (B) pericarp surface (x5000), (C) broken nutlet, longitudinal section (x149), (D) broken nutlet (x1500).



**Figure 2.** SEM microscopy of *Salvia hispanica* L. nutlets after mucilage extraction. (A) Whole nutlet (x136), (B) nutlet surface (x5000), (C) broken nutlet (x150), (D) broken nutlet (x3500).

Component	Msm	Ms #
Moisture	10.66 ± 0.04 <sup>a</sup>	10.47 ± 0.16 <sup>a</sup>
Protein*	42.43 ± 0.71 <sup>a</sup>	41.36 ± 0.28 <sup>a</sup>
Crude fiber	27.75 ± 0.97 <sup>a</sup>	27.57 ± 0.07 <sup>a</sup>
Ash	7.82 ± 0.13 <sup>a</sup>	7.24 ± 0.15 <sup>a</sup>
Oil	0.22 ± 0.25 <sup>a</sup>	0.21 ± 0.08 <sup>a</sup>
NFE	24.17 ± 0.76 <sup>a</sup>	23.62 ± 0.94 <sup>a</sup>

# Capitani *et al.* [11]

Mean value (n = 3)

Values followed by different letters differ significantly ( $p \leq 0.05$ ), according to Tukey's test.

\*Factor: 6.25; NFE: nitrogen-free extract.

**Table 1.** Proximate composition of chia (*Salvia hispanica* L.) meals (% d.b.)

Component	Msm	Ms #
TDF	47.13 ± 0.17 <sup>a</sup>	46.06 ± 0.86 <sup>a</sup>
IDF	45.62 ± 0.37 <sup>b</sup>	41.13 ± 0.47 <sup>a</sup>
SDF	1.51 ± 0.24 <sup>a</sup>	4.93 ± 0.65 <sup>b</sup>

# Capitani *et al.* [11]

Mean value (n = 3)

Values followed by different letters differ significantly ( $p \leq 0.05$ ), according to Tukey's test.

**Table 2.** Total (TDF), soluble (SDF) and insoluble (IDF) dietary fiber of chia (*Salvia hispanica* L.) meals (% d.b.)

Component	Msm	Ms #
NDF	63.6 ± 2.1 <sup>b</sup>	53.9 ± 0.3 <sup>a</sup>
ADF	30.8 ± 1.2 <sup>a</sup>	38.1 ± 1.2 <sup>b</sup>
Lignin	6.9 ± 0.5 <sup>a</sup>	4.5 ± 0.7 <sup>a</sup>
Cellulose	23.1 ± 0.9 <sup>a</sup>	34.6 ± 1.3 <sup>b</sup>
Hemicellulose	33.6 ± 1.0 <sup>b</sup>	14.8 ± 1.2 <sup>a</sup>

# Capitani *et al.* [11]

Mean value (n = 3)

Values followed by different letters differ significantly ( $p \leq 0.05$ ), according to Tukey's test.

**Table 3.** Fiber composition of chia meals analyzed according to the method of Van Soest (% d.b.)

The antioxidant activity of the two chia meals compared with other types of meals is shown in Table 4. Both for Ms and Msm, the activity was high, without a significant difference between them ( $p > 0.05$ ). These values were higher than those found for wheat bran and sorghum and barley whole grain meals. But they were significantly lower than those found for chia meal obtained as a byproduct of cold-pressing oil extraction. The latter could be attributed to the fact that the meal obtained by pressing shows a higher percentage of residual oil (11.39% d.b.), which contains tocopherols, a class of compound with natural antioxidant activity [11].

Sample	Trolox equivalent antioxidant coefficient (TEAC, $\mu\text{mol/g}$ )
Msm	$187.4 \pm 33.21^a$
Ms <sup>1</sup>	$226.6 \pm 4.13^a$
Chia meal from oil pressing extraction <sup>1</sup>	$557.2 \pm 28.18^b$
Wheat bran <sup>2</sup>	48.5
Sorghum meal <sup>3</sup>	51.7
Barley meal <sup>3</sup>	14.9

<sup>1</sup> Capitani *et al.*, [11]

<sup>2</sup> Iqbal *et al.* [57]

<sup>3</sup> Ragaee *et al.* [58]

Mean value (n = 3)

Values followed by different letters differ significantly ( $p \leq 0.05$ ), according to Tukey's test.

**Table 4.** Antioxidant activity of chia (*Salvia hispanica* L.) meals compared with other meals

Regarding the functional properties, the meal with mucilage (Ms) exhibited a statistically higher absorption and water holding capacity ( $p < 0.05$ ) than that of the meal without mucilage (Msm) (Table 5). This behavior can be associated with the presence of mucilage in Ms, which acts as soluble dietary fiber, capable of holding water inside its matrix [51]. The WAbC of both meals was higher than that observed for canola and soybean meals (3.90 g/g and 3.28 g/g, respectively) and similar to that of linseed meal (6.03 g/g) [48].

Both types of chia meals presented a low absorption of organic molecules and oil-holding capacity, being significantly higher in Msm. These differences could be explained in terms of the particle size and the cellulose content of the meal [52, 53]. The determination of OHC is

important because it is related to the capacity that food components have to hold oil, affecting their flavor and mouthfeel [54]. The OMAC is associated with the interaction of the fiber with fats, bile acids, cholesterol, drugs, and toxic and carcinogenic compounds at the intestinal level. Due to their low OHC levels, both types of chia meals could be considered an important ingredient in the manufacture of fried products due to their low fatty mouthfeel contribution. It is noteworthy that the OHC values were higher than those reported for the *Jessenia polycarpa* fruit meal [55] and similar to those of the fibrous residue of *Canavalia ensiformis* and barley [53, 56] and those reported by Khattab and Arntfield [48] for linseed and canola meals (2.01 g/g and 2.09 g/g, respectively).


Property	Msm	Ms #
WHC (g/g)	5.25 ± 0.39 <sup>a</sup>	10.64 ± 0.60 <sup>b</sup>
WA <sub>6</sub> C (g/g)	4.79 ± 0.49 <sup>a</sup>	6.45 ± 0.41 <sup>b</sup>
OHC (g/g)	2.94 ± 0.14 <sup>b</sup>	2.03 ± 0.08 <sup>a</sup>
OMAC (g/g)	2.22 ± 0.01 <sup>b</sup>	1.64 ± 0.02 <sup>a</sup>

# Capitani *et al.* [11]  
 Mean value (n = 3)  
 Values followed by different letters differ significantly ( $p \leq 0.05$ ), according to Tukey's test.


**Table 5.** Functional properties of chia (*Salvia hispanica* L.) meals

In Figure 3 it can be noted that the differences observed in the emulsifying activity were not significant, although the stability of the emulsion prepared with Ms was statistically higher than that with Msm ( $p < 0.05$ ). This effect can be associated with the capacity of mucilage to act as a thickening agent due to its ability to increase the viscosity of the aqueous phase in an O/W emulsion, thus hindering movement of the oil droplets of the dispersed phase [59]. This property is similar in the linseed mucilage, which has a strong thickening capacity, favorably affecting the water-holding capacity and the emulsifying properties of defatted linseed flour [60].

The behavior of the meals studied with respect to emulsion stability, examined by their optical characterization with a vertical scan analyzer (QuickScan), is shown in Figure 4. Both meals presented a high initial emulsifying capacity (66.3 and 62.2 %BS. for Ms and Msm, respectively), which remained approximately constant for all the time span studied (60 min) for Ms. However, the emulsion stability of the meal without mucilage decreased markedly by the end of the 60 min (50.4 %BS).



**Figure 3.** Activity and stability of O/W emulsions (50:50 p/p) with chia meal with and without mucilage. Values followed by different letters differ significantly (Tukey's test,  $p \leq 0,05$ )



**Figure 4.** Destabilization kinetics of O/W emulsions (50:50 p/p) with chia meal

## 4. Conclusions

The results obtained show that both types of chia meals (with and without mucilage) present interesting functional properties for the food industry, which could be applied for example in the manufacture of bakery products, powdered beverages, yogurts, ice-creams, sauces and creams. They also suggest the potential use of two chia byproducts: the residual meal obtained after the oil extraction of whole chia seeds, and the use of a byproduct of the mucilage extraction. As regards the formulation of stable emulsions, the meal with mucilage is recommended for use given the role of mucilage as a thickening agent. From a physiological point of view, the presence of mucilage becomes a potentially interesting food ingredient due to its health benefits, since it has the capacity to form high-viscosity gels, slowing the intestinal transit, providing more of a feeling of satiety, and helping to prevent diseases such as obesity, colon cancer, hypercholesterolemia and diabetes.

## Acknowledgments

This work was supported by grants from Agencia Nacional de Promoción Científica y Tecnológica (ANPCyT), Argentina (PICT 2002-10768), PICT 2007, 1085, PIP 6097 CONICET, Universidad Nacional de La Plata (UNLP) (11/X390), Universidad Nacional del Centro de la Provincia de Buenos Aires (UNCPBA), Argentina.

Author M.I. Capitani is a recipient of a fellowship from the Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), Argentina.

Author S.M. Nolasco is a Scientific and Technological Researcher and Professor at the Facultad de Ingeniería de la Universidad Nacional del Centro de la Provincia de Buenos Aires (UNCPBA), Argentina.

Author M.C. Tomás is member of the career of Scientific and Technological Researcher of the Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), Argentina.

## Author details

Marianela I. Capitani<sup>1,2</sup>, Susana M. Nolasco<sup>2</sup> and Mabel C. Tomás<sup>1</sup>

1 Centro de Investigación y Desarrollo en Criotecología de Alimentos (CIDCA), (CCT- La Plata – CONICET) Facultad de Ciencias Exactas, UNLP, La Plata, Buenos Aires, Argentina

2 Grupo de Investigaciones TECSE, Departamento de Ingeniería Química, Facultad de Ingeniería, UNCPBA, Olavarría, Buenos Aires, Argentina

## References

- [1] Ixtaina VY, Nolasco SM, Tomás MC. Characterization of chia (*Salvia hispanica* L.) white and dark seeds and oils. Ed/ MC. Tomás/, *Advances in Fats and Oils Research*, Transworld Research Network Kerala, India, pp. 135-147. 2010.
- [2] Ayerza R, Coates W. Ground chia seed and chia oil effects on plasma lipids and fatty acids in the rat. *Nutrition Research* 2005; 25 995-1003.
- [3] Álvarez-Chávez LM, Valdivia-López MA, Alberto-Juárez ML, Tecante A. Chemical characterization of the lipid fraction of mexican chia seed (*Salvia hispanica* L.). *International Journal of Food Properties* 2008; 11 687-697.
- [4] Bushway A, Belya. Chia seed as a source of oil, polysaccharide and protein. *Journal of Food Science* 1981; 46 1349-1356.
- [5] Coates W, Ayerza R. Production potencial of chia in northwestern Argentina. *Industrial Crops and Products* 1996; 5 229-233.
- [6] Sahagun B. 1950-1982, Florentine codex: general history of the things of New Spain. In: *Monographs of the School of America Research*. Anderson AJO and Dibble CE, eds. Salt Lake City: University of Utah Press (Originally written 1575-1577 or 1578-1580).
- [7] Taga MS, Miller EE, Pratt DE. Chia seeds as a source of natural lipid antioxidants. *Journal of the American Oil Chemists Society* 1984; 61 928-931.
- [8] Ixtaina VY, Martínez ML, Spotorno V, Mateo CM, Maestri DM, Diehl BW, Nolasco SM, Tomás MC. Characterization of chia seed oils obtained by pressing and solvent extraction. *Journal of Food Composition and Analysis* 2011; 24(2) 166-174.
- [9] Craig R, Sons M. Application for approval of whole chia (*Salvia hispanica* L.) seed and ground whole chia as novel food ingredients. Advisory committee for novel food and processes. Ireland, Company David Armstrong, 2004. pp. 1-29.
- [10] Reyes-Caudillo E, Tecante A, Valdivia-López MA. Dietary fiber content and antioxidant activity of phenolic compounds present in Mexican chia (*Salvia hispanica* L.) seeds. *Journal of Food Chemistry* 2008; 107 656-663.
- [11] Capitani MI, Spotorno V, Nolasco SM, Tomás MC. Physicochemical and functional characterization of by-products from chia (*Salvia hispanica* L.) seeds of Argentina. *Lebensmittel Wissenschaft and Technology - Food Science Technology* 2012; 45 94-102.
- [12] Badui S. *Química de los Alimentos*. Longman / Alhambra; 1999.
- [13] Chau C, Huang Y. Comparison of the chemical composition and physicochemical properties of different fibers prepared from the peel of *Citrus sinensis* L. cv. Liucheng. *Journal of Agriculture and Food Chemistry* 2003; 51 2615-2618.

- [14] Wong K, Cheung P. Dietary fibers from mushroom sclerotia: 1. Preparation and physicochemical and functional properties. *Journal and Agriculture Food Chemistry* 2005; 53 9395-9400.
- [15] Yoshimoto M, Yamakawa O, Tanoë H. Potencial chemopreventive properties and varietal difference of dietary fiber from sweetpotato (*Ipomea batatas* L.) root. *Japan Agricultural Research Quarterly* 2005; 39 37-43.
- [16] Raghavendra S, Ramachandra S, Rastogi N, Raghavarao K, Kumar S, Tharanathan R. Grinding characteristics and hydration properties of coconut residue: A source of dietary fiber. *Journal of Food Engineering* 2006; 72 281-286.
- [17] Gorinstein S, Zachwieja Z, Folta M, Barton H, Piotrowicz J, Sembré M, Weisz M, Trakhtenberg S, Martín-Belloso O. Comparative content of dietary fiber, total phenolics, and minerals in persimmons and apples. *Journal of Agriculture Food Chemistry* 2001; 49 952-957.
- [18] Sánchez GBS. Caracterización Físicoquímica y Funcional de la Fibra dietética del fruto del Nispero (*Eriobotrya japonica*) y de la cáscara de mango Obo (*Mangifera indica* L.). Tesis de Grado, Universidad Tecnológica de la Mixteca, Oaxaca, México. 2005.
- [19] Zambrano-Zaragoza ML, Meléndez R, Gallardo Y. Propiedades funcionales y metodología para su evaluación en fibra dietética. Ed/ FM. Lajolo, F Saura-Calixto, E Witting and Wenzel de Menezes/, *Fibra dietética en Iberoamérica: Tecnología y Salud. Obtención, caracterización, efecto fisiológico y aplicación en alimentos*. Brasil: Livraria LTDA. pp. 195-209. 2001.
- [20] Peraza GM. Caracterización de los residuos fibrosos de *Canavalia ensiformis* L. y *Phaseolus lunatus* L. y su incorporación a un product alimenticio. Tesis de Maestría. Facultad de Ingeniería Química, Universidad Autónoma de Yucatán, México. 2000.
- [21] Ator CJ. Caracterización química de la fibra dietetic de subproductos obtenidos de la leguminosa *Vigna Unguiculata* L. Walps. Tesis de Maestría, Facultad de Ingeniería Química, Universidad Autónoma de Yucatán, Mérida, Yucatán, México, 2000.
- [22] Larrauri JA, Goñi I, Martín-Carrón N, Rupérez P, Saura-Calixto F. Measurement of health-promoting properties in fruti dietary fibres: antioxidant capacity, fermentability and glucosa retardation index. *Journal of Agriculture and Food Chemistry* 1996; 71 515-519.
- [23] McDougall GJ, Morrison IM, Stewart D, Hillman RJ. Plant cell walls as dietary fibre: range, structure, processing and function. *Journal of Science and Food Agriculture* 1996; 70 133-150.
- [24] Rehinan Z, Rashid M, Shah WH. Insoluble dietary fibre components of food legumes as affected by soaking and cooking processes, *Journal of Food Chemistry* 2004; 85 245-249.
- [25] Martín-Belloso O, Gorinstein S, Grijelmo-Miguel N. Characterization of peach dietary fibre concentrate as a food ingredient. *Food Chemistry* 1999; 65 175-181.

- [26] Saura-Calixto F, García-Alonso A. Metodología para el análisis de fibra y carbohidratos. Ed/ FM. Lajolo, F Saura-Calixto, E Witting and Wenzel de Menezes/, *Fibra dietética en Iberoamérica: Tecnología y Salud. Obtención, caracterización, efecto fisiológico y aplicación en alimentos.* Brasil: Livraria LTDA. pp. 17-25. 2001.
- [27] Lin KY, Daniel JR. Structure of chia seed polysaccharide exudates. *Carbohydrate Polymer* 1994; 23 13-18.
- [28] Marin Flores FM, Acevedo MJ, Tamez RM, Nevero MJ, Garay AL WO/2008/0044908 Method for obtaining mucilage from *Salvia hispánica* L. Word Internacional Property Organization. 2008.
- [29] Mazza G, Biliaderis CG. Functional properties of flax seed mucilage. *Journal of Food Science* 1989; 54 1302-1305.
- [30] Sáenz C, Sepúlveda E, Matsushiro B. Opuntia spp mucilage's: a functional component with industrial perspectives. *Journal of Arid Environments* 2004; 57 275-290.
- [31] Koocheki A, Mortazavi SA, Shahidi F, Razavi SMA, Taherian AR. Rheological properties of mucilage extracted from Alyssum homolocarpum seed as a new source of thickening agent. *Journal of Food Engineering* 2009; 91, 490-496.
- [32] Cheng Z, Blackford J, Wang Q, Yu L. Acid treatment to improve psyllium functionality. *Journal of Functional Foods* 2009; 1 44-49.
- [33] IUPAC (1992), International Union of Pure and Applied Chemistry – Standard Methods for the Analysis of Oils, Fats and Derivates. 7<sup>th</sup> edn. Paquot, C. and Hautfene, A., Blackwell Scientific Publications, Inc., Oxford.
- [34] AOCS (1998), Official and Recommended Practices of the American Oil Chemists' Society, 5th edn., edited by D. Firestone, AOCS Press, Champaign Illinois.
- [35] AOAC (1997), In W. Horwitz (Ed.), Official methods of analysis (17<sup>th</sup> ed.), Washington, D.C., USA, Association of Official Analytical Chemists.
- [36] Prosky L, Asp N, Schweizer T, Devries S, Furda I. Determination of insoluble, soluble and total dietary fiber in food and food products: interlaboratory study. *Journal of the Association of Official Analytical Chemists* 1988; 71 1017-1023.
- [37] Guiragossian VY, Van Scoyoc SW, Auxtell JD. Chemical and Biological Methods for Grain and Forage Sorghum, Departament of Agronomy Internacional Programs in Agricultura, Purdue University, West Lafayette, Indiana, USA. 1979.
- [38] Re R, Pellegrini N, Proteggente A, Pannala A, Yang M, Rice-Evans C. Antioxidant activity applying an improved ABTS radical cation decolorization assay. *Free Radical Biological Medicine* 1999; 26 (9/10) 1231-1237.
- [39] Walker RB, Everette JD Comparative reaction rates of various antioxidants with ABTS radical cation. *Journal of Agriculture and Food Chemists* 2009; 57 1156-1161.

- [40] Chau C, Cheung K, Wong Y. Functional properties of protein concentrate from three Chinese indigenous legume seeds, *Journal of Agricultural and Food Chemistry* 1997; 45 2500-2503.
- [41] AACC (1984), *Official methods of análisis* (12th ed.). St. Paul, MN, USA: American Association of Cereal Chemists.
- [42] Pan LG, Tomás MC, Añón MC. Effect of sunflower lecithins on the stability of water in oil (W/O) and oil in water (O/W) emulsions. *Journal of Surfactant and Detergents* 2002; 5(2) 135-143.
- [43] Infostat Group, Infostat version 2004, Facultad de Ciencias Agrarias, Universidad Nacional de Córdoba, Argentina 2004.
- [44] Gutterman Y, Shem-Tov S, Mucilaginous seed coat structure of *Carrichtera annua* and *Anastatica hierochuntica* from the Negev Desert highlands of Israel, and its adhesion to the soil crust. *Journal of the Arid Environment* 1997; 35 695-705.
- [45] Western T, Skinner D, Haughn G. Differentiation of mucilage secretory cells of the *Arabidopsis* seed coat. *Plant Physiological* 2000; 122 345-355.
- [46] Pérez EE, Carelli AA, Crapiste GH. Chemical characterization of oils and meals from wild sunflower (*Helianthus petiolaris* Nutt). *Journal of the American Oil Chemists` Society* 2004; 81 1-5.
- [47] Sun M, Xu L, Saldaña DA. Comparison of canola meals obtained with conventional methods and supercritical CO<sub>2</sub> with and without ethanol. *Journal of the American Oil Chemists` Society* 2008; 85 667-675.
- [48] Khattab RY, Arntfield SD. Functional properties of raw and processed canola meal, *LWT – Food Science and Technology* 2009; 42 1119-1124.
- [49] Mueller K, Eisner P, Yoshie-Stark Y, Nakada R, Kirchhoff E. Functional properties and chemical composition of fractionated brown and yellow linseed meal (*Linum usitatissimum* L.). *Journal of Food Engineering* 2010; 98 453-460.
- [50] Egbekun MK, Ehieze MU. Proximate composition and functional properties of fullfat and defatted beniseed (*Sesamum indicum* L.) flour. *Plant Foods for Human Nutrition* 1997; 51 35-41.
- [51] Figuerola F, Hurtado ML, Estévez AM, Chiffelle I, Asenjo F. Fibre concentrates from apple pomace and citrus peel as potential fibre sources for food enrichment. *Food Chemistry* 2005; 91 395-401.
- [52] López G, Ros G, Rincón F, Periago M, Martínez M, Ortuño J. Relationship between physical and hydration properties of soluble and insoluble fiber of artichoke. *Journal of the Agricultural and Food Chemistry* 1996; 44 277-2778.
- [53] Betancur-Ancona D, Peraza-Mercado G, Moguer-Ordoñez Y, Fuertes-Blanco S. Physicochemical characterization of lima vean (*Phaseolus lunatus*) and jack bean (*Canavalia ensiformis*) fibrous residues, *Food Chemistry* 2004; 84 287-295.

- [54] Kinsella JE. Functional properties of protein foods, *Critical Review of Food Science and Nutrition* 1976; 1 219-229.
- [55] Belén D, Alemán R, Alvarez F, Álvarez M. Evaluación de algunas propiedades funcionales y reológicas de harinas de coroba (*Jessenia polycarpa Karst*). *Revista de la Facultad de Agronomía (LUZ, Venezuela)* 2004; 21 161-170.
- [56] Mongeau R, Brassard R. Insoluble dietary fibre from breakfast cereals and brands, bile salt binding and water holding capacity in relation to particle size. *Cereal Chemistry* 1982; 59(5) 413-415.
- [57] Iqbal S, Bhanger M, Anwar F. Antioxidant properties and components of some commercially available varieties of rice bran in Pakistan. *Food Chemistry* 2005; 93 265-272.
- [58] Ragaee S, Abdel-Aal M, Norman M. Antioxidant activity and nutrient composition of selected cereals for food use. *Food Chemistry* 2006; 98 32-38.
- [59] McClements DJ. *Food emulsions: Principles, practice, and techniques*. Boca Raton, FL: CRC Press. 1999.
- [60] Dev DK, Quensel E. Functional and microestructural characteristics of linseed (*Linum usitatissimum* L.) flour and a protein isolate, *Lebensmittel Wissenschaft and Technology - Food Science Technology* 1986; 19 331-337.

