
Anesthetic Management for Laparoscopic Cholecystectomy

Somchai Amornyotin

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/52742>

1. Introduction

Laparoscopic surgery aims to minimize trauma of the interventional process but still achieve a satisfactory therapeutic result. It is commonly performed because of various advantages such as reduced postoperative pain, faster recovery and more rapid return to normal activities, shorter hospital stay, and reduced postoperative pulmonary complications. The operative technique requires inflating gas into the abdominal cavity to provide a surgical procedure. An intra-abdominal pressure (IAP) of 10-15 mmHg is used. Carbon dioxide (CO₂) is commonly used because it does not support combustion, is cleared more rapidly than other gases, and is highly soluble in blood. However, the disadvantage of CO₂ is that the absorption of CO₂ can cause hypercapnia and respiratory acidosis [1].

Laparoscopic cholecystectomy (LC) procedure offers several advantages such as a reduction in stress response, postoperative pain, postoperative wound infection rate, intraoperative bleeding, impairment of respiratory function and pulmonary complications, short recovery time, and cosmetic appearance [1,2]. LC reduces hospital stay but has no overall effect on postoperative mortality [3]. The risk factors for perioperative complications in patients undergoing LC can be estimated based on patient characteristics, clinical findings and the surgeon's experience [4]. The advantages should to be balanced with potential adverse effects caused by CO₂ pneumoperitoneum.

The physiological effects of intra-abdominal CO₂ insufflation combined with the variations in patient positioning can have a major impact on cardiorespiratory function. In addition, the sequential effects of anesthesia combine to produce a characteristic hemodynamic response. A thorough understanding of these physiological changes is fundamental for optimal anesthetic care. Several anesthetic techniques can be performed for LC. General anesthesia using balanced anesthetic technique including intravenous drugs, inhalation

agents and muscle relaxants is usually used. Short acting drugs such as propofol, atracurium, vecuronium, sevoflurane or desflurane represent the maintenance drugs of choice. Pre-procedure assessment and preparation, appropriate monitoring and a high index of suspicion can result in early diagnosis and treatment of complications.

2. Pathophysiological effects during laparoscopic cholecystectomy

2.1. Physiological effects of pneumoperitoneum

Carbon dioxide was shown to be affected by raising the intra-abdominal pressure (IAP) above the venous pressure which prevents CO₂ resorption leading to hypercapnia. Hypercapnia activates the sympathetic nervous system leading to an increase in blood pressure, heart rate, arrhythmias and myocardial contractility as well as it also sensitizes myocardium to catecholamines [5]. Increased IAP may compress venous vessels causing an initial increase in preload, followed by a sustained decrease in preload.

2.2. Respiratory effects

The changes in pulmonary function during LC include reduction in lung volumes, decrease in pulmonary compliance, and increase in peak airway pressure [6]. Increased IAP shifts the diaphragm cephalad and reduces diaphragmatic excursion, resulting in early closure of smaller airways leading to intraoperative atelectasis with a decrease in functional residual capacity. Additionally, the upward displacement of diaphragm leads to preferential ventilation of nondependent parts of lung, which results in ventilation-perfusion (V/Q) mismatch with a higher degree of intrapulmonary shunting. Oxygenation is minimally affected with no significant change in alveolar arterial oxygen gradient [7]. Higher IAP reduces the thoracic compliance and may cause pneumothorax and pneumomediastinum due to the increased in alveolar pressures [6].

2.3. Cardiovascular effects

Hemodynamic changes include the alterations in arterial blood pressure, arrhythmias and cardiac arrest. These cardiovascular changes depend on the interaction of several factors including patient positioning, neurohumoral response and the patient factors such as cardiorespiratory status and intravascular volume. The principal responses are an increase in systemic vascular resistance, mean arterial blood pressure and myocardial filling pressures, with little change in heart rate [2]. CO₂ pneumoperitoneum is associated with increased preload and afterload in patients undergoing LC. It also decreased heart performance (fractional shortening), but does not affect cardiac output [8]. The patients with normal cardiovascular function are able to well tolerate these hemodynamic changes. At IAP levels greater than 15 mmHg, venous return decreases leading to decreased cardiac output and hypotension [9]. However, these changes are short lived and have no statistical significance at 10 minutes from the time that the patient undergoes pneumoperitoneum [10].

Bradycardias are attributed to vagal stimulation caused by insertion of the needle or the trocar, peritoneal stretch, stimulation of the fallopian tube during bipolar electrocauterization, or carbon dioxide embolization [11]. These may induce cardiovascular collapse during laparoscopy even in the healthy patients. Increased concentrations of CO₂ and catecholamines can create tachycardias. Paroxysmal tachycardia and hypertension, followed by ventricular fibrillation, have been reported [12].

2.4. Effects of other systems

Increases in IAP, cardiovascular responses to peritoneal insufflations, changes in patient position and alterations in CO₂ concentration can alter intracranial pressure (ICP) and cerebral perfusion. ICP shows a significant further increase. Cerebral blood flow has been shown to increase significantly during CO₂ insufflation.

Pneumoperitoneum reduces renal cortical and medullary blood flow with an associated reduction in glomerular filtration rate (GFR), urinary output and creatinine clearance [2]. The reduction of renal blood flow may be due to a direct pressure effect on renal cortical blood flow and renal vascular compression as well as an increase in antidiuretic hormone (ADH), aldosterone and renin. Pretreatment with an ADH antagonist improves urine output and urea excretion despite an unaltered GFR.

Increased in IAP reduces femoral venous blood flow. This is due to increased pressure on the inferior vena cava and iliac veins, which reduces venous blood flow in the lower extremities. It also has been shown to reduce the portal blood flow, which may lead to transient elevation of liver enzymes.

The C-reactive protein and interleukin-6 levels are less elevated after laparoscopy compared to the open surgery, suggesting an attenuation of the surgical inflammatory response [13].

Patient positions can further compromise cardiac and respiratory functions, can increase the risk of regurgitation and can result in peripheral nerve injuries. Head-up position reduces venous return, cardiac output, cardiac index and mean arterial blood pressure as well as an increase in peripheral and pulmonary vascular resistance [5,14]. Head-down position increases volume and cardiac output back towards normal. Respiratory function is impaired because of the cephalad shifting of diaphragm is exaggerated. Intracranial pressure is increased.

3. Anesthetic management

3.1. Preoperative assessment

The general health status of each patient must be evaluated. History and physical examinations are generally sufficient techniques. The patients with cardiorespiratory diseases require additional investigation. To aid in assessment risk, the American Society of Anesthesiologists (ASA) has developed a classification system for patients, which categoriz-

es individuals on a general health basis. In this preoperative assessment, there are no differences in a routine practice between the laparoscopy and the open surgery.

3.2. Patient monitoring

Appropriate patient selection with proper monitoring to detect and reduce complications must be used to ensure optimal anesthesia care during LC. Standard intraoperative monitoring including noninvasive blood pressure, electrocardiogram, pulse oximeter, airway pressure, end tidal carbon dioxide (ETCO₂), body temperature and peripheral nerve stimulation is routinely used. Invasive hemodynamic monitoring may be appropriate in the patients with hemodynamic unstable or those with compromised cardiopulmonary function [1].

ETCO₂ is most commonly used as a noninvasive indicator of PaCO₂ in evaluating the adequacy of ventilation. Careful consideration should be taken for the gradient between PaCO₂ and the tension of CO₂ in expired gas (PECO₂) because of V/Q mismatch. However, in the patients with compromised cardiopulmonary function, the gradient between PaCO₂ and PECO₂ increases to become unpredictable. Direct arterial blood gas analysis may be considered to detect hypercarbia. Generally, the airway pressure monitor is routinely used during intermittent positive pressure ventilation. The high airway pressure can help detection of excessive elevation in IAP.

3.3. Anesthetic techniques

Various anesthetic techniques can be performed for LC. However, general anesthesia with endotracheal intubation for controlled ventilation is the most common anesthetic technique. In short procedures and in certain patients, ventilation using supraglottic airway device can be used as an alternative. General anesthesia without endotracheal intubation can be used safely and effectively with a ProSeal laryngeal mask airway in non-obese patients [15]. The use of laryngeal mask airway results in less sore throat and provide smoother emergence with less post-extubation coughing compared with endotracheal intubation [16].

3.3.1. General anesthesia

General anesthesia using balanced anesthesia technique including inhalation agents, intravenous drugs and muscle relaxant drugs is usually used. The uses of rapid and short acting volatile anesthetics such as sevoflurane and desflurane as well as rapid and short acting intravenous drugs such as propofol, etomidate, remifentanyl, fentanyl, atracurium, vecuronium and rocuronium are commonly used and have allowed anesthesiologists to more consistently achieve a recovery profile. Propofol is effective and safe even in children and elderly patients [17-21].

Ventilation should be adjusted to keep ETCO₂ of around 35 mmHg by adjusting the minute ventilation [1]. In patients with chronic obstructive pulmonary disease and in patients with a history of spontaneous pneumothorax or bullous emphysema, an increase in respiratory rate rather than tidal volume is preferable to avoid increased alveolar inflation and reduce the risk of pneumothorax [22].

Furthermore, the use of an auditory evoked potential or Bispectral index monitor to titrate the volatile anesthetics leads to a significant reduction in the anesthetic requirement, resulting in a shorter postanesthesia care stay and an improved quality of recovery from the patient's perspective [23].

Combination of local anesthetic wound infiltration, intraperitoneum spray of local anesthetic, paracetamol and non-steroidal anti-inflammatory drugs or cyclooxygenase 2 inhibitors provides the most effective pain relief, which can be supplemented with small doses of opioids.

3.3.2. Regional anesthesia

Several advantages of regional anesthesia technique are quicker recovery, decreased postoperative nausea and vomiting, fewer hemodynamic changes, less postoperative pain, shorter hospital stay, early diagnosis of complications, improved patient satisfaction and cost effectiveness [24]. This anesthetic technique requires a cooperative patient, low IAP to reduce pain and ventilation disturbances, gentle surgical technique and a supportive operating room staff. However, regional anesthesia technique is not commonly used for LC. This technique should be performed in combination with other anesthetic techniques. Local anesthetic infiltration at the trocar site combined with general anesthesia significantly reduces postoperative pain and decreases medication usage costs [25]. Additionally, subcostal transversusabdominis block provides superior postoperative analgesia, improves theater efficiency by reducing time to discharge from the recovery unit and reduces opioid requirement following LC [26]. Bilateral paravertebral blockade at T5-6 level combined with general anesthesia can be used for LC [27].

Mehta and college had been conducted a prospective, randomized, controlled trial to compare spinal anesthesia with the gold standard general anesthesia for elective LC in the healthy patients. Their study demonstrated that spinal anesthesia was adequate and safe for LC in otherwise healthy patients and offered better postoperative pain control than general anesthesia without limiting the recovery [28]. The interim analysis of a controlled randomized trial is also confirmed [29]. Thoracic epidural anesthesia with 0.75% ropivacaine and fentanyl for elective LC is also efficacious and has preserved ventilation and hemodynamic changes within physiological limits during pneumoperitoneum with minimal treatable side effects [30]. In addition, epidural anesthesia might be applicable for LC. However, the incidence rate of intraoperative referred pain is high, and so careful patient recruitment and management of shoulder pain should be considered [31].

4. Intraoperative complications

Misplacement of the needle can lead to intravascular, subcutaneous tissue, preperitoneal space, bowel, and omentum. Inadvertent insufflation of gas into intravascular vessels, tear of abdominal wall or peritoneal vessels, can produce to gas embolism. Although, it is rare but it is a potentially lethal complication and can result in severe hypotension, cyanosis, ar-

rhythmias and asystole. Subcutaneous emphysema may occur after direct subcutaneous gas insufflations. The majority of subcutaneous emphysema has no specific intervention. It can resolve soon after the abdomen is deflated and nitrous oxide is discontinued to avoid expansion of closed space.

Pneumothorax can occur when the airway pressure is high. The gas traverses into the thorax through the tear of visceral peritoneum, parietal pleura during dissection, or spontaneous rupture of pre-existing emphysematous bulla [1]. Pneumothorax can be asymptomatic or can increase the peak airway pressure, decrease oxygen saturation, hypotension, and even cardiac arrest in severe cases. The treatment is according to the severity of cardiopulmonary compromise [32].

Extension of subcutaneous emphysema into thorax and mediastinum can lead to pneumomediastinum. Pneumopericardium can occur when the gas is forced through the inferior vena cava into the mediastinum and pericardium. Their managements depend on the severity of the cardiovascular dysfunction.

The other complications can be presented. Accidental insertion of the trocar or needle into the major or minor vessels, gastrointestinal tract injuries and urinary tract injuries can occur [32].

5. Postoperative period

The efficacy of post-anesthesia care units is therefore important to facilitate return to normal functions. In the early postoperative period, respiratory rate and ETCO_2 of laparoscopic patients breathing spontaneously are higher as compared with open surgery. So, the ventilation requirement is increased. The patients with respiratory dysfunction can have problems excreting excessive CO_2 load, which results in more hypercapnia. Additionally, the patients with cardiovascular diseases are more prone to hemodynamic changes and instabilities.

Although LC results in less discomfort compared with the open surgery, postoperative pain still can be considerable. Several medications used intraoperatively for prevention and treatment of postoperative pain are the uses of local anesthesia, opioids, nonsteroidal anti-inflammatory drugs, and multimodal analgesia techniques. Additionally, preprocedure administration of parecoxib is clinically effective [33].

Postoperative nausea and vomiting (PONV) is a common and distressing symptom following LC. The use of multimodal analgesia regimens and the reduction of opioid doses are likely to reduce the incidence of PONV. Propofol-based anesthesia has been associated with reduced PONV [34]. Ondansetron has been found to provide effective prophylaxis against PONV [35]. Administration of ondansetron at the end of surgery produces a significantly greater anti-emetic effect compared to pre-induction dosing. Reduced preoperative anxiety by providing more information should also relieve postoperative adverse effects in order to promote faster and better postoperative recovery period.

6. Summary

Laparoscopic cholecystectomy has proven to be a major advance in the treatment of patients with symptomatic gall bladder diseases. Several advantages from this procedure are minimal tissue trauma, reduction of postoperative pain, quicker recovery, shortening the hospital stay. Pneumoperitoneum induces intraoperative cardiorespiratory changes. Arterial CO₂ increases because of CO₂ absorption from the pneumoperitoneum. Improved knowledge of pathophysiological changes in the patients allows for successful anesthetic management. Proper patient selection and preparation as well as adequate monitoring should be performed. General anesthesia and controlled ventilation comprise the accepted anesthetic technique. Balanced anesthesia technique including inhalation agent, intravenous drug and muscle relaxant is commonly used. Intraoperative complications may arise due to physiologic changes associated with patient positioning and pneumoperitoneum. Multimodal analgesic regimen combining opioids, non-steroidal anti-inflammatory drugs, and local anesthetic infiltration is the most effective regimen for postoperative pain management.

Author details

Somchai Amorniyotin

Department of Anesthesiology and Siriraj GI Endoscopy Center, Faculty of Medicine, Siriraj Hospital, Mahidol University, Bangkok, Thailand

References

- [1] Gerges FJ, Kanazi GE, Jabbour-Khoury SI. (2006). Anesthesia for laparoscopy: a review. *Journal of Clinical Anesthesia* 2001; 18(1): 67-78.
- [2] Leonard IE, Cunningham AJ. Anesthetic consideration for laparoscopic cholecystectomy. *Best Practice & Research Clinical Anesthesiology* 2002; 16(1): 1-20.
- [3] McMahon AJ, Fischbacher CM, Frame SH, MacLeod MC. Impact of laparoscopic cholecystectomy: a population-based study. *Lancet* 2000; 356(11): 1632-1637.
- [4] Giger UF, Michel JM, Opitz I, et al. Risk factors for perioperative complications in patients undergoing laparoscopic cholecystectomy: analysis of 22,953 consecutive cases from the Swiss Association of laparoscopic and thoracoscopic surgery database. *Journal of American College of Surgeons* 2006; 203(5): 723-728.
- [5] Gutt CN, Oniu T, Mehrabi A, et al. Circulatory and respiratory complications of carbon dioxide insufflations. *Digestive Surgery* 2004; 21(2): 95-105.

- [6] Rauh R, Hemmerling TM, Rist M, Jacobi KE. Influence of pneumoperitoneum and patient positioning on respiratory system compliance. *Journal of Clinical Anesthesia* 2001; 13(5): 361-365.
- [7] Sharma KC, Brandstetter RD, Brensilver JM, Jung LD. Cardiopulmonary physiology and pathophysiology as a consequence of laparoscopic surgery. *Chest* 1996; 110(3): 810-815.
- [8] Larsen JF, Svendsen FM, Pedersen V. Randomized clinical trial of the effect of pneumoperitoneum on cardiac function and hemodynamics during laparoscopic cholecystectomy. *British Journal of Surgery* 2004; 91(7): 848-854.
- [9] Odeberg S, Ljungqvist O, Svenberg T, et al. Hemodynamic effects of pneumoperitoneum and the influence of posture during anesthesia for laparoscopic surgery. *Acta Anaesthesiologica Scandinavica* 1994; 38(3): 276-283.
- [10] Zuckerman RS, Heneghan S. The duration of hemodynamic depression during laparoscopic cholecystectomy. *Surgical Endoscopy* 2002; 16(8): 1233-1236.
- [11] Sprung J, Abdelmalak B, Schoenwald PK. Recurrent complete heart block in a healthy patient during laparoscopic electrocauterization of the fallopian tube. *Anesthesiology* 1998; 88(5): 1401-1403.
- [12] Cheong MA, Kim YC, Park HK, et al. Paroxysmal tachycardia and hypertension with or without ventricular fibrillation during laparoscopic adrenalectomy: two case reports in patients with noncatecholamine-secreting adrenocortical adenomas. *Journal of Laparoendoscopic & Advanced Surgical Techniques A* 1999; 9(3): 277-281.
- [13] Grabowski JE, Talamini MA. Physiological effects of pneumoperitoneum. *Journal of Gastrointestinal Surgery* 2009; 13(5): 1009-1016.
- [14] Hirvonen EA, Poikolainen EO, Paakkonen ME, Nuutinen LS. The adverse hemodynamic effects of anesthesia, head-up tilt, and carbon dioxide pneumoperitoneum during laparoscopic cholecystectomy. *Surgical Endoscopy* 2000; 14(3): 272-277.
- [15] Maltby JR, Beriault MT, Watson NC, Liepert D, Fick GH. The LMA-ProSeal is an effective alternative to tracheal intubation for laparoscopic cholecystectomy. *Canadian Journal of Anesthesia*, 2002; 49(8): 857-862
- [16] Cook TM, Lee G, Nolan JP. The ProSeal laryngeal mask airway: a review of the literature. *Canadian Journal of Anesthesia* 2005; 52(7): 739-760.
- [17] Amornyotin S, Chalayonnavin W, Kongphlay S. Assisted sedation for percutaneous endoscopic gastrostomy in sick patients in a developing country. *Gastroenterology Insights* 2010; 2(e5): 17-20.
- [18] Amornyotin S, Prakanrattana U, Chalayonnavin W, Kongphlay S, Kachintorn U. (2010). Propofol based sedation does not increase perforation rate during colonoscopic procedure. *Gastroenterology Insights* 2010; 2(e4): 13-16.

- [19] Amornyotin S, Chalayonnawin W, Kongphlay S. Propofol-based sedation does not increase rate of complication during percutaneous endoscopic gastrostomy procedure. *Gastroenterology Research and Practice* 2011 Article ID 134819; 6 pages, doi: 10.1155/2011/134819.
- [20] Amornyotin S, Srikureja W, Pausawasdi N, Prakanrattana U, Kachintorn U. Intravenous sedation for gastrointestinal endoscopy in very elderly patients of Thailand. *Asian Biomedicine* 2011; 5(4): 485-491.
- [21] Amornyotin S, Kachintorn U, Chalayonnawin W, Kongphlay S. Propofol-based deep sedation for endoscopic retrograde cholangiopancreatography procedure in sick elderly patients in a developing country. *Therapeutics and Clinical Risk Management* 2011; 7: 251-255.
- [22] Salihoglu Z, Demiroglu S, Dikmen Y. Respiratory mechanics in morbid obese patients with chronic obstructive pulmonary disease and hypertension during pneumoperitoneum. *European Journal of Anaesthesiology* 2003; 20(8): 658-661.
- [23] Recart A, Gasanova I, White PF, et al. (2003). The effect of cerebral monitoring on recovery after general anesthesia: a comparison of the auditory evoked potential and Bispectral index devices with standard clinical practice. *Anesthesia Analgesia* 2003; 97(6): 1667-1674.
- [24] Collins LM, Vaghadia H. Regional anesthesia for laparoscopy. *Anesthesiology Clinic of North America* 2001; 19(1): 43-55.
- [25] Hasaniya NW, Zayed FF, Faiz H, Severino R. Preinsertion local anesthesia at the trocar site improves perioperative pain and decreases costs of laparoscopic cholecystectomy. *Surgical Endoscopy* 2001; 15(9): 962-964.
- [26] Tolchard S, Davies R, Martindale S. Efficacy of the subcostal transversusabdominis plane block in laparoscopic cholecystectomy: comparison with conventional port-site infiltration. *Journal of Anaesthesiology Clinical Pharmacology* 2012; 28(3): 339-343.
- [27] Naja MZ, Ziade MF, Lonnqvist PA. General anesthesia combined with bilateral paravertebral blockade (T5-6) vs. general anesthesia for laparoscopic cholecystectomy: a prospective, randomized clinical trial. *European Journal of Anaesthesiology* 2004; 21(6): 489-495.
- [28] Mehta PJ, Chavda HR, Wadhvana AP, Porecha MM. Comparative analysis of spinal versus general anesthesia for laparoscopic cholecystectomy: a controlled, prospective, randomized trial. *Anesthesia: Essays and Researches* 2010; 4(2): 91-95.
- [29] Tzovaras G, Fafoulakis F, Pratsas K, et al. Spinal vs general anesthesia for laparoscopic cholecystectomy: an interim analysis of a controlled randomized trial. *Archives of Surgery* 2008; 143(5): 497-501.
- [30] Gupta A, Gupta K, Gupta PK, Agarwal N, Rastogi B. Efficacy of thoracic epidural anesthesia for laparoscopic cholecystectomy. *Anesthesia: Essays and Researches* 2011; 5(2): 138-141.

- [31] Lee JH, Huh J, Kim DK, et al. Laparoscopic cholecystectomy under epidural anesthesia: a clinical feasibility study. *Korean Journal of Anesthesiology* 2010; 59(6): 383-388.
- [32] Joshi GP. Complications of laparoscopy. *Anesthesiology Clinic of North America* 2001; 19(1): 89-105.[33]
- [33] Amornytin S, Chalayonnawin W, Kongphlay S. A randomized controlled trial of preprocedure administration of parecoxib for therapeutic endoscopic retrograde cholangiopancreatography. *Journal of Pain Research* 2012; 5: 251-256.
- [34] Fujii Y. Management of postoperative nausea and vomiting in patients undergoing laparoscopic cholecystectomy. *Surgical Endoscopy* 2011; 25(3): 691-695.
- [35] Wu SJ, Xiong XZ, Cheng TY, Lin YX, Cheng NS. Efficacy of ondansetronvs metoclopramide in prophylaxis of postoperative nausea and vomiting after laparoscopic cholecystectomy: a systematic review and meta-analysis. *Hepatogastroenterology* 2012; 59(119), Doi: 10.5754/hge11811 [Epub ahead of print]