

PUBLISHED BY

# INTECH

open science | open minds

World's largest Science,  
Technology & Medicine  
Open Access book publisher



**3,350+**  
OPEN ACCESS BOOKS



**108,000+**  
INTERNATIONAL  
AUTHORS AND EDITORS



**114+ MILLION**  
DOWNLOADS



**BOOKS**  
DELIVERED TO  
151 COUNTRIES

AUTHORS AMONG  
**TOP 1%**  
MOST CITED SCIENTIST



**12.2%**  
AUTHORS AND EDITORS  
FROM TOP 500 UNIVERSITIES



Selection of our books indexed in the  
Book Citation Index in Web of Science™  
Core Collection (BKCI)

**WEB OF SCIENCE™**

Chapter from the book *Lipoproteins - From Bench to Bedside*

Downloaded from: <http://www.intechopen.com/books/lipoproteins-from-bench-to-bedside>

Interested in publishing with IntechOpen?  
Contact us at [book.department@intechopen.com](mailto:book.department@intechopen.com)

---

# Lipoprotein-Associated Phospholipase A<sub>2</sub> – Pathophysiological Role and Clinical Significance as a Cardiovascular Biomarker

---

Sanja Stankovic and Milika Asanin

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/60608>

---

## Abstract

Within the last decade, a broad range of biomarkers associated with an increased risk for death and cardiovascular/cerebrovascular endpoints have been identified. Epidemiological studies clearly indicate that lipoprotein-associated phospholipase A<sub>2</sub> (Lp-PLA<sub>2</sub>) has the potential to become clinically useful emerging biomarker in the true sense, linking plaque biology with cardiovascular/cerebrovascular event rate. Lipoprotein-associated phospholipase A<sub>2</sub> is a specific vascular inflammatory marker, a risk factor, a prognostic biomarker, and also a therapeutic target. This chapter will summarize our current knowledge on Lp-PLA<sub>2</sub> with emphasis on its potential pathophysiological mechanisms of action and on clinical relevance as cardiovascular/cerebrovascular biomarker. This chapter gives comprehensive, systematic review of studies assessing the significance of Lp-PLA<sub>2</sub> in cardiovascular/cerebrovascular diseases with emphasis on clinical benefit of pharmacologic inhibition of Lp-PLA<sub>2</sub>.

**Keywords:** biomarker, cardiovascular disease, lipoprotein-associated phospholipase A<sub>2</sub>

---

## 1. Introduction

Although an atherogenic lipoprotein phenotype was reported as a predictor of cardiovascular/cerebrovascular disease, numerous recent studies have recognized additional lipid-related markers as emerging biomarkers to identify patients with cardiovascular/cerebrovascular disease risk. Among them the most promising biomarkers for cardiovascular/cerebrovascular risk assessment is lipoprotein-associated phospholipase A<sub>2</sub> (Lp-PLA<sub>2</sub>).

---

## 2. Lipoprotein-associated phospholipase A<sub>2</sub>: structure and biology

Lipoprotein-associated phospholipase A<sub>2</sub> (Lp-PLA<sub>2</sub>), also known as platelet-activating factor acetylhydrolase (PAF-AH), belongs to the phospholipase A<sub>2</sub> superfamily [1,2]. This Ca<sup>2+</sup>-independent phospholipase is encoded by *PLA2G7* gene that consists of 12 exons and 11 introns located on chromosome 6p21.2 to 12 [3,4]. Lp-PLA<sub>2</sub> is protein of 45,4 kDa that consists of 441 amino acid residues [5]. The major sources of Lp-PLA<sub>2</sub> in plasma are T lymphocytes, monocytes/macrophages, activated bone marrow-derived mast cells, and liver cells [6-8]. The secreted Lp-PLA<sub>2</sub> circulates in plasma in active form. It predominantly binds to LDLs, and in a much smaller extent to HDLs, Lp(a), lipoprotein remnants, and platelet-borne microparticles [6,9-12]. Indeed, Lp-PLA<sub>2</sub> is highly associated with the smallest LDL and HDL subclasses [13] and with electronegative LDL, which overlaps with small dense LDL [14]. Lp-PLA<sub>2</sub> bound to HDL has a much lower specific activity compared to when bound to LDL [10]. Different distribution of Lp-PLA<sub>2</sub> in various plasma lipoproteins affects its functions.

Lp-PLA<sub>2</sub> catalyzes hydrolysis of the acetyl group at sn-2 position of PAF to generate lyso-PAF and acetate [15]. On the other side, Lp-PLA<sub>2</sub> cleaves oxidatively modified lipoproteins from the sn-2 position of the apoB100-containing lipoproteins into oxidized nonesterified fatty acids (oxFFAs) and lysophosphatidylcholine.

## 3. Actions of lipoprotein-associated phospholipase A<sub>2</sub>

We can think about Lp-PLA<sub>2</sub> as a friend and foe at the same time. On the one hand, it functions as PAF-AH, which hydrolyzes inflammatory mediator PAF, inhibits foam cell formation, enhances cholesterol efflux in macrophages, and exerts its atheroprotective role [16].

On the other hand, Lp-PLA<sub>2</sub> has a proatherogenic role. Lp-PLA<sub>2</sub> relates to a number of different proatherogenic biological processes. Lp-PLA<sub>2</sub> hydrolyzes oxidized phospholipids on modified LDL particles within the arterial intima, and thus contributes to the initiation and progression of atheroma. Lp-PLA<sub>2</sub> is produced by macrophages and foam cells within atherosclerotic plaque. Its expression is mainly confined to plaque areas with massive lipid accumulation and leukocyte infiltration, cellular necrosis, and calcification, suggesting that Lp-PLA<sub>2</sub> is a marker for rupture-prone plaque [16]. The amount of Lp-PLA<sub>2</sub> and its by-product, lysophosphatidylcholine in the coronary circulation is proportional to the extent of the atheroma and indirectly affects local endothelial function [17]. Proatherogenic activities of lysophosphatidylcholine are: expression of adhesion molecules; upregulation of cytokines and CD40 ligand by T-cells, cytotoxic at concentrations higher than 30–50 μM; stimulation of macrophage proliferation; release of arachidonic acid from endothelial cells; induction of MCP-1 and genes for growth factors; release of myeloperoxidase; migration of vascular smooth muscle cells; chemoattractant for monocytes, macrophages, and T-cells; induction of apoptosis in smooth muscle cells and macrophages; involvement in the antigenicity of oxidized LDL; inhibition of endothelium-derived nitric oxide [18-20]. Furthermore, oxFFAs promote atherosclerosis by direct and

indirect increasing of oxidative stress and the presence of oxidized LDL and other lipoproteins in the plasma and arterial walls, thereby initiating fatty streak formation [21].

#### 4. Assays for Lp-PLA<sub>2</sub> mass and activity determination

It is well known that circulating Lp-PLA<sub>2</sub> can be measured by different assays ascertaining either its mass or activity. There is no definite decision about the potential superiority of the tests.

According to the literature data, Lp-PLA<sub>2</sub> levels were determined preferentially using PLAC® test. Lp-PLA<sub>2</sub> mass was determined by diaDexus PLAC Test ELISA, sandwich enzyme immunoassay, followed by diaDexus PLAC® test based for Lp-PLA<sub>2</sub> mass measurement on turbidimetric immunoassay technology. More recently, enzyme assay PLAC® Test for measurement of Lp-PLA<sub>2</sub> activity has been developed and commercially available. The preferred sample type is EDTA plasma, and serum is also acceptable. It should be noted that methodological issues associated with Lp-PLA<sub>2</sub> measurement make comparisons between studies difficult.

Recently published meta-analysis of 32 prospective studies in persons with stable vascular disease or recent acute ischemic event revealed a moderate correlation between mass and activity of Lp-PLA<sub>2</sub> [24]. The PROVEIT-TIMI 22 study [25] has also found a moderate correlation between Lp-PLA<sub>2</sub> activity and mass measured at baseline and at 30-day after ACS. There is still controversy about the method of estimating Lp-PLA<sub>2</sub> level. While Koenig et al. [26] reported that Lp-PLA<sub>2</sub> mass was the better risk predictor of future cardiovascular events than Lp-PLA<sub>2</sub> activity, Persson et al. [27] has reached the opposite conclusion. Jenny et al. [28] showed no difference in Lp-PLA<sub>2</sub> activity and mass with respect to risk prediction.

Currently, there is no consensus on the best method to estimate Lp-PLA<sub>2</sub> level. A consensus panel recommendation for incorporating Lp-PLA<sub>2</sub> testing into the cardiovascular disease risk assessment guidelines used Lp-PLA<sub>2</sub> mass for stratifying patients [23].

Blood samples should be refrigerated after processing and should be kept frozen for long-term storage. There are no restrictions to the time of day that the sample should be drawn and no dietary restrictions. In contrast to other emerging risk markers, a very minimal biological variation in Lp-PLA<sub>2</sub> concentrations has been demonstrated among individuals monitored serially over several weeks [29].

In addition, Lp-PLA<sub>2</sub> levels are typically unaffected by conditions of systemic inflammation, such as osteoarthritis and chronic obstructive pulmonary disease, whereas markers of inflammatory response are often elevated by these conditions. The normal population medians for men and women are in the range of 230–250 ng/mL, and a value of >300 ng/mL may be considered elevated [30].

Gender differences in Lp-PLA<sub>2</sub> levels were found; men had significantly higher levels than women. Also, a significant association between Lp-PLA<sub>2</sub> levels and smoking was noticed.

These finding has been observed in previous studies [31,32,33,34]. Lower Lp-PLA<sub>2</sub> levels in women could be explained by estrogen-mediated down-regulation of Lp-PLA<sub>2</sub> expression, due to lower concentrations of LDL cholesterol in women or estrogen-related decrease in platelet activating factor acetyl hydrolase activity [35,36]. Estrogen-replacement therapy can significantly reduce Lp-PLA<sub>2</sub> activity in healthy postmenopausal women [37], while administration of steroids with progesterone-like activity increases Lp-PLA<sub>2</sub> activity [38]. Smoking may increase the carrier (LDL) and the substrate (oxidized LDL) for Lp-PLA<sub>2</sub> [39].

## 5. Association of Lp-PLA<sub>2</sub> with cardiovascular disease risk

Lp-PLA<sub>2</sub> is an emerging inflammatory biomarker, characterized by high vascular specificity and low biovariability. High Lp-PLA<sub>2</sub> levels are indicative of rupture-prone plaque [22]. Numerous epidemiological and clinical studies examined association between Lp-PLA<sub>2</sub> concentration/enzyme activity and cardiovascular disease risk in apparently healthy individuals, in subjects with stable cardiovascular disease, acute coronary syndrome (ACS), heart failure (HF), stroke, transient ischemic attack (TIA) support, trying to answer in which patients would determination of Lp-PLA<sub>2</sub> be the most valuable. There are novel data about the prognostic significance of Lp-PLA<sub>2</sub> as a predictor of short-term or long-term outcome in patients with cardiovascular disease. To evaluate this, systematic literature review concerning the association of Lp-PLA<sub>2</sub> with cardiovascular disease risk and prognostic implications was done. The studies on this issue were extracted from relevant electronic databases (Medline (<http://www.ncbi.nlm.nih.gov/pubmed/>), Embase (<http://www.embase.com/>), Google Scholar (<http://scholar.google.com/>), Yahoo (<http://www.yahoo.com/>), Kobson (<http://www.kobson.nb.rs/>), ClinicalTrials.gov.) and the obtained results were included in the text.

In 2008, the panel's recommendation [23] incorporated Lp-PLA<sub>2</sub> testing as an adjunct to traditional risk factors in assessing future cardiovascular risks. It endorses Lp-PLA<sub>2</sub> testing in moderate-risk persons determined as having simply two risk factors. An Lp-PLA<sub>2</sub> >200 ng/mL warrants reclassification of the moderate-risk patient as high cardiovascular risk and should prompt reduction of the LDL-C target from 130 mg/dL to 100 mg/dL. The panel also recommends Lp-PLA<sub>2</sub> testing for patients with coronary artery disease (CAD) or CAD risk equivalents (diabetes, ischemic stroke, etc.) be considered at very high risk when Lp-PLA<sub>2</sub> is elevated, warranting reduction in the LDL-C target from 100 mg/dL to 70 mg/dL. Today, Lp-PLA<sub>2</sub> measurement for cardiovascular disease risk stratification of patients is included in four international clinical guidelines: 2012 European Guidelines on CVD Prevention in Clinical Practice European Society of Cardiology – Lp-PLA<sub>2</sub> may be measured as a part of a refined risk assessment in patients at high risk of a recurrent acute atherothrombotic event. Class IIb; 2010 ACCF/AHA Guideline for Assessment of Cardiovascular Risk in Asymptomatic Adults – Lp-PLA<sub>2</sub> testing may be considered in intermediate-risk asymptomatic adults; 2011 AHA/ASA Guidelines for the Primary Prevention of Stroke Measurement of inflammatory markers such as high sensitive C-reactive protein (hs-CRP) or Lp-PLA<sub>2</sub> in patients without CVD may be considered to identify patients who may be at increased risk of stroke; 2012 AACE Guidelines for Management of Dyslipidemia and Prevention of Atherosclerosis Test for Lp-

PLA<sub>2</sub>, which in some studies has demonstrated more specificity than highly sensitive CRP, when it is necessary to further stratify a patient's CVD risk [40-43].

Some studies suggested that Lp-PLA<sub>2</sub> predicts risk complementary to hs-CRP, although when compared with hs-CRP, Lp-PLA<sub>2</sub> in some studies is suggested to be a more promising marker of risk prediction. In multiple previous studies, correlation analysis revealed no association of Lp-PLA<sub>2</sub> with CRP [28,31,44-47]. It can be expected because CRP is an acute-phase reactant and its elevation can be caused by a wide range of inflammatory conditions. CRP shows intraindividual variability of ~40%. On the other hand, Lp-PLA<sub>2</sub> is not affected by systemic inflammation; it is a specific marker of vascular inflammation. Also, it shows significantly lower biologic variability than CRP, and higher stability in states of myocardial ischemia [48].

Lp-PLA<sub>2</sub> is suggested to be a more promising marker of risk prediction than CRP [49]. Winkler et al. [50] showed that increased Lp-PLA<sub>2</sub> levels in moderate-risk patients with hs-CRP < 3 mg/L doubled the risk for cardiac death. The WOSCOPS study [44] found that Lp-PLA<sub>2</sub> was significantly associated with cardiovascular risk, compared with hs-CRP. Stankovic et al.'s [32] results confirmed that Lp-PLA<sub>2</sub> have better risk prediction than CRP.

Although, the majority of published studies showed a significant relationship between Lp-PLA<sub>2</sub> levels and cardiovascular events, there are several important differences across ethnic groups; for example, African-Americans and Caucasians with respect to Lp-PLA<sub>2</sub>. Activity of Lp-PLA<sub>2</sub> was higher among African-Americans with CAD. The difference in Lp-PLA<sub>2</sub> activity levels between CAD and non-CAD patients was higher among African-Americans. Also, the Lp-PLA<sub>2</sub> index was independently associated with the extent of CAD among African-Americans [51-53].

From the WOSCOPS publication (West of Scotland Coronary Prevention Study, WOSCOPS) [44], which revealed a positive association between elevated circulating concentrations of Lp-PLA<sub>2</sub> and the risk of coronary heart disease (CHD), the interest in Lp-PLA<sub>2</sub> as a biomarker for cardiovascular disease rapidly increased. The vast body of evidence derived from prospective epidemiologic studies, two meta-analyses (79,036 participants in 32 prospective studies, and 52,995 subjects participated in 33 studies) and review revealed the positive association of elevated Lp-PLA<sub>2</sub> with cardiovascular risk [24,54-56]. Furthermore, Lp-PLA<sub>2</sub> has been confirmed to predict the presence of CAD, even among patients undergoing coronary angiography. Uniquely, Lp-PLA<sub>2</sub> predicted the risk of CAD death, but not all cause death [57].

Lp-PLA<sub>2</sub> has been considered as a prognostic marker in patients with CAD. Li et al. [58] identified prognostic value of Lp-PLA<sub>2</sub> baseline measurement for major adverse cardiac event (MACE) (cardiovascular death, nonfatal myocardial infarction, and target vessel revascularization) in ACS patients during follow-up for a median of 6 months. Mockel et al. [59] demonstrated that Lp-PLA<sub>2</sub> levels in the first 7 hours after onset of symptoms may be an independent predictor of MACE within 42 days in ACS patients. PROVEIT-TIMI 22 study [25] has shown that Lp-PLA<sub>2</sub> in patients randomized to atorvastatin or pravastatin measured 30 days after ACS are associated with an increased risk of cardiovascular events (death, myocardial infarction, unstable angina, revascularization, or stroke) over 24 months of follow-up. Gerber et al. MACE [48] found that Lp-PLA<sub>2</sub> level measured early after myocardial infarction is

strongly and independently associated with 1-year mortality. Stankovic et al. [32] demonstrated that the Lp-PLA<sub>2</sub> may have short-term predictive value in pure STEMI patients treated by primary percutaneous coronary intervention (PCI). They concluded that pre-interventional plasma Lp-PLA<sub>2</sub> level is an independent predictor of 30-day MACE in patients with first anterior STEMI treated by primary PCI, and suggested that Lp-PLA<sub>2</sub> level could help in very early risk stratification of STEMI patients treated by PCI.

Previously published studies examined and suggested the association between Lp-PLA<sub>2</sub> and heart failure (HF) incidence in a population-based cohort of healthy individuals, and in people older than 65 years [60]. Baseline Lp-PLA<sub>2</sub> levels are associated with a high risk of developing heart failure in 3,991 adults older than 65 years, independent of coronary risk factors [61]. Lp-PLA<sub>2</sub> activity is significantly associated with congestive heart failure in 5,531 persons older than 65 years [62]. Gerber et al. [63] evaluated the association of Lp-PLA<sub>2</sub> with mortality in subjects with diagnosed HF. Lp-PLA<sub>2</sub> was strongly and independently associated with mortality in patients under 80 years of age. Moldoveanu et al.'s study [64] in 208 patients with HF found significantly increased Lp-PLA<sub>2</sub> activity in HF patients with preserved ejection fraction (EF) than in HF with reduced EF. The literature data about the association of Lp-PLA<sub>2</sub> and heart failure on admission in patients with acute myocardial infarction are missing. Stankovic et al. [32] suggested that patients with the first anterior STEMI who had higher levels of Lp-PLA<sub>2</sub> had a worse prognosis, but not with a greater probability of developing HF. In Raichlin et al.'s [65] study in heart transplant patients, Lp-PLA<sub>2</sub> correlated with the progression of cardiac allograft vasculopathy and increased risk of cardiovascular events/death suggesting that it could be therapeutic target in heart transplant patients.

In 2005, the US Food and Drug Administration (FDA) approved Lp-PLA<sub>2</sub> blood test for assessing patients at risk for ischemic stroke. The Rotterdam Study was the first population-based study that determined the impact of elevated Lp-PLA<sub>2</sub> on stroke. It identified that Lp-PLA<sub>2</sub> activity was an independent predictor of ischemic stroke in the middle-aged healthy men and women population [31]. In the ARIC study, healthy middle-aged adults with increased levels of both Lp-PLA<sub>2</sub> and hs-CRP had an 11-fold higher incidence of stroke than individuals with low Lp-PLA<sub>2</sub> and hs-CRP levels [45]. This association between Lp-PLA<sub>2</sub> mass/activity and first ischemic stroke was confirmed in the Malmo Diet and Cancer Study [66], Bruneck Study [67], and Cardiovascular Health Study [28]. The Tsekepis et al. study [68] showed that Lp-PLA<sub>2</sub> correlated with the intima-media thickness in patients with beta-thalassemia, suggesting that Lp-PLA<sub>2</sub> may be implicated in premature carotid atherosclerosis.

Although some reports inconclusively found this positive association of Lp-PLA<sub>2</sub> and first-ever and recurrent stroke, little is known about its influence on stroke outcome. Elkind et al. [69] measured Lp-PLA<sub>2</sub> mass and activity in relation to outcome in first ischemic stroke patients, determined as recurrent stroke, recurrence of vascular events, and mortality. Lp-PLA<sub>2</sub> was a good predictor of recurrent stroke risk. Delgado et al. [70] investigated the temporal profile of Lp-PLA<sub>2</sub> mass and activity within the first 24 hours after stroke and found significant changes in Lp-PLA<sub>2</sub> concentrations early after stroke onset. Patients with higher Lp-PLA<sub>2</sub> mass were more likely to be resistant to intravenous t-PA administration with very low early recanalization rates.

Lp-PLA<sub>2</sub>, an inflammatory biomarker, has been described as able to predict risk of first-ever or recurrent stroke and myocardial infarction [71]. Moreover, Lp-PLA<sub>2</sub> may also have a role in the pathophysiology of cerebrovascular disease, particularly in strokes of atherosclerotic etiology, since its expression is enhanced in atherosclerotic carotid lesions together with markers of oxidative damage, inflammation, and instability [72].

Sarlon-Bartoli et al. [73] reported that Lp-PLA<sub>2</sub> mass is increased in patients with high-grade carotid stenosis and unstable plaque and suggest that Lp-PLA<sub>2</sub> could be an important biomarker for classifying carotid plaque as vulnerable and predict neurological risk of a carotid stenosis in asymptomatic subjects. Although these findings have the potential to improve cerebrovascular disease stratification, correlation with ultrasonic or MRI markers of plaque instability or the presence of infarction on brain imaging must be performed [74].

Shoamanesh et al. [75] investigated the association between circulating biomarkers of inflammation including Lp-PLA<sub>2</sub> and MRI markers of cerebral small vessel disease in 1,763 stroke-free Framingham offspring. They observed higher levels of lipoprotein-associated phospholipase A<sub>2</sub> mass in patients with greater white matter hyperintensity volumes and silent cerebral infarcts. These results could improve stroke risk prognosis.

The accumulating results of numerous studies demonstrate a significant positive association between Lp-PLA<sub>2</sub> levels and incident cardiovascular disease and heart failure. A dissociation could be noted between mass and activity in terms of risk prediction. Several of the epidemiology studies have measured only Lp-PLA<sub>2</sub> mass (26,30,33,44,47,57,63,65,69), whereas others measured only enzyme activity (31,50,62,64,67) and a few have measured both in the same study population (25,27,28, 76). Recently published meta-analysis assessed the 32 prospective studies of Lp-PLA<sub>2</sub> and cardiovascular outcomes. For each standard deviation Lp-PLA<sub>2</sub> increase, the relative risks for the primary endpoint of coronary heart disease were 1.10 (1.05–1.16) and 1.11 (1.07–1.16) for Lp-PLA<sub>2</sub> activity and mass, respectively. The relative adjusted risks for ischemic stroke were 1.08 (0.97–1.20) and 1.14 (1.02–1.27); vascular mortality 1.16 (1.09–1.24) and 1.13 (1.05–1.22); and nonvascular mortality 1.10 (1.04–1.17); and 1.10 (1.03–1.18) for Lp-PLA<sub>2</sub> activity and mass, respectively. The final decision whether to measure the Lp-PLA<sub>2</sub> mass or its enzyme activity may help transition of this biomarker from research to routine clinical practice.

## 6. Lp-PLA<sub>2</sub> and genetic influences

Several Lp-PLA<sub>2</sub> gene polymorphisms and their role in affecting the regulation or production of LpPLA<sub>2</sub> assessed as activity and mass were described; many in small studies and some in recently published genome-wide association studies. It is known that genetic factors account for 62% of the variation in Lp-PLA<sub>2</sub> activity [76,77].

Although familial factors explain about one-half and one-quarter of the variance in Lp-PLA<sub>2</sub> activity and mass, respectively [78,79], few genetic determinants of Lp-PLA<sub>2</sub> have been identified. The first genome-wide association study using data from 6,668 Caucasian subjects

in population-based Framingham Heart Study identified one locus associated with Lp-PLA<sub>2</sub> mass, and four loci associated with Lp-PLA<sub>2</sub> activity [80]. Twelve SNPs in the region of chromosome 6p12.3 near the gene for PLA2G7 were associated with Lp-PLA<sub>2</sub> mass at a genome-wide level of significance. The top hit SNP rs1805017, is a nonsynonymous change (H92R) within the PLA2G7 gene. It was found that T allele that corresponds to the amino acid histidine was associated with higher Lp-PLA<sub>2</sub> mass. On the other side, four loci achieving genome-wide significance for association with Lp-PLA<sub>2</sub> activity was identified: a) first within the APOE/APOC1 gene cluster on chromosome 19q13.32 (rs41377151); b) second locus on chromosome 1p13.3, which includes the genes PSRC1 (rs599839), CELSR2 (rs4970834); c) third locus within an intron of SCARB1 on chromosome 12q24.31 (rs10846744); d) fourth locus in ZNF259 gene and BUD13 gene on chromosome 11q23.3 near the apolipoprotein gene cluster APOA5/APOA4/APOC3/APOA1 (rs12286037, rs11820589). Investigation from Suchindran's study was extended by Grallert et al. who made meta-analysis with additional four cohorts [81]. They performed genome-wide association study as part of the Cohorts for Heart and Aging Research in Genomic Epidemiology (CHARGE) Consortium in five population-based studies with the aim to identify genetic loci related to Lp-PLA<sub>2</sub> mass/activity [82]. This study, in the population of 13,664 subjects, revealed the association of PLA2G7 loci variants with Lp-PLA<sub>2</sub> mass/activity, and genetic variants (APOC1, CELSR2, LDL, ZNF259, SCARB1) related to low-density lipoprotein cholesterol levels with Lp-PLA<sub>2</sub> activity.

Also, Lp-PLA<sub>2</sub> mass was associated with SNP rs247616 on chromosome 16 within the cholesteryl ester transfer protein gene. The T allele of rs247616 was associated with higher HDL-C concentration and higher mean Lp-PLA<sub>2</sub> mass [83]. The additional value of this study was the revealed significant association of four polymorphisms (in APOC1, CELSR2, SCARB1, ZNF259), but not PLA2G7 with coronary heart disease. Large-scale analysis focusing on variants in PLA2G7 did not identify any SNP that achieved experiment-wide statistical significance [79].

The most frequently studied SNPs within PLA2G7 are: variants in exon 9 (Val279Phe; rs45619133), exon 11 (Val379Ala; rs1051931), exon 7 (Ile198Thr, Iso195Thr; rs1805018), and exon 4 (Arg92His; rs1805017) [84-86].

The V279F SNP was found in subjects of Asian ancestry. The Val279Phe substitution is located within the catalytical domain of Lp-PLA<sub>2</sub> and leads to reduction of enzyme activity in V/F heterozygous individuals or complete loss of enzymatic activity in homozygous F/F individuals. The 279F null allele is relatively frequent in Japan, with approximately 25% and 2% of the population carrying one or two copies. Its prevalence declines toward the West, has intermediate frequencies in China and Korea, is rare in the Middle East, and almost completely absent in European populations. The results from the association studies on this V279F variant and coronary artery disease have thus far been inconclusive. Li et al. [87] in Chinese Han population, found significant association of V279F and coronary artery disease, indicating that carrier of F allele increases the risk of coronary artery disease. This result was consistent with the Yamada et al. [88] study (850 cases/1,684 controls), which found that subjects carrying the mutant allele are at higher risk of arterial events (MI or stroke), and the Shimokata study (3,085 subjects with coronary artery disease/2,163 controls) [89]. The Jang et al. [90] study in two

different patient sets (2,890 men diagnosed with CAD before age 60/3,128 male controls and 877 CAD cases/1,230 controls) confirmed that deficiency in Lp-PLA<sub>2</sub> activity due to carriage of PLA2G7 279F allele protects from CAD in Korean men. Nevertheless, in the South Korean population (532 cases/670 controls) and Chinese study (827 cases/947 controls), V279F variant results in an unexpectedly opposite outcome [91,92]. Meta-analysis of 14 association studies focusing on R92H polymorphism in PLA2G7 gene and risk of CHD in 8,280 cases/5,656 controls indicate 92H allele had probably increased the risk of CHD [92]. The missense polymorphisms I198T and A379V are identified mainly in Caucasians. A379V variant in which alanine is substituted by valine is functional. Some studies explored the contribution of A379V to Lp-PLA<sub>2</sub> activity with contradictory results; some found the association of A379V variant with increased [86,93] and some with decreased Lp-PLA<sub>2</sub> activity [94].

There are also different reports about the association between the A379V variant and cardiovascular disease [94-97]. A recent Taiwanese population study reported that subjects carrying the 379 V allele had increased severity of coronary atherosclerosis [94]; two studies in Caucasian subjects reported that the 379 V allele was associated with decreased atherosclerosis risk [93,96]; a large meta-analysis showed that the 379 V allele was associated with Lp-PLA<sub>2</sub> activity, but not with cardiovascular risk markers [97]; and another study showed no association [95].

Intriguingly, in the study conducted by Liu and colleagues [94], the outcome was quite contradictory. They found that in the Chinese Taiwan Han population, A379V variant is significantly associated with Lp-PLA<sub>2</sub> activity and the severity of coronary atherosclerosis. Recently, a meta-analysis including a total of 12 studies shows that in the populations from European ancestry, among the 7 SNPs, A379V variant shows the strongest association with Lp-PLA<sub>2</sub> activity; however, no significant correlation is found between PLA2G7 variants and cardiovascular risk markers, coronary atheroma, or CHD [97].

## 7. Lp-PLA<sub>2</sub> as a therapeutic target

It is well known that lipid-altering medications, including statins, fenofibrate, prescription of omega-3 fatty acids, for weight loss, have been shown to reduce Lp-PLA<sub>2</sub> levels. The degree of its reduction correlates with the extent of lipid lowering.

Lp-PLA<sub>2</sub> was identified as a potential novel target of therapy. The most used therapy targeting Lp-PLA<sub>2</sub> in plasma in advanced stages of clinical investigation is darapladib. Darapladib is a selective, potent, reversible, oral inhibitor of lipoprotein-associated phospholipase A<sub>2</sub>. The basic idea of applying darapladib is to improve patient outcomes in addition to evidence-based treatments and potentially reduce cardiovascular and cerebrovascular events by decreasing cytokines concentrations, stabilizing atherosclerotic plaque, inhibiting macrophage infiltration, and thickening of the connective tissue cap.

The first study that showed as a secondary endpoint a change in coronary artery plaque necrotic core with darapladib after 12 months of treatment was Integrated Biomarkers and

Imaging Study (IBIS)-2 [98]. It was a multicenter, randomized, double-blind placebo-controlled study that included 330 patients with angiographically confirmed coronary artery disease. Inhibition of Lp-PLA<sub>2</sub> with darapladib also prevented necrotic core expansion of coronary plaque as measured on intravascular ultrasound.

One multicenter, randomized, double-blind placebo-controlled study examined the effects of darapladib on biomarkers of cardiovascular risk in 959 CAD and CAD-risk equivalent patients who were previously randomized to atorvastatin 20 mg or 80 mg and then randomized to oral darapladib 40, 80, 160 mg, or placebo for 12 weeks. Overall dose-dependent inhibition of Lp-PLA<sub>2</sub> activity was sustained over the study period and was present in both atorvastatin dose groups, at different baseline LDL cholesterol < or ≥70 mg/dl, and high-density lipoprotein cholesterol HDL-C < or ≥40 mg/dl [99].

The first study that examined the effects of darapladib on Lp-PLA<sub>2</sub> activity in Japanese dyslipidemic patients with/without the Val279Phe single-nucleotide polymorphism (SNP) of the *PLA2G7* gene showed that darapladib produced sustained inhibition of Lp-PLA<sub>2</sub> activity [100].

Two large-scale studies with hard clinical endpoints were completed: the Stabilization of Atherosclerotic Plaque by Initiation of Darapladib Therapy (STABILITY) trial, with 15,828 randomized patients, and the Stabilization of Plaques using Darapladib-Thrombolysis In Myocardial Infarction 52 Trial (SOLID-TIMI 52), with an estimated recruitment of 13,026 patients [101-103].

The SOLID-TIMI 52 was a randomized, double-blind, placebo-controlled, multicenter, and event-driven trial that determined the clinical benefit of direct inhibition of Lp-PLA<sub>2</sub> activity with darapladib in patients after an acute coronary syndrome (non-ST-elevation or ST-elevation myocardial infarction). Subjects were randomized to darapladib (160 mg enteric-coated tablet daily) or matching placebo within 30 days after acute coronary syndrome. The primary endpoint was the composite of coronary heart disease death, nonfatal myocardial infarction, or nonfatal stroke, and secondary endpoints were major and total coronary events, individual components of the primary endpoint, and all-cause mortality. Patients were followed up for a median of 2.5 years. In patients who experienced an ACS event, direct inhibition of Lp-PLA<sub>2</sub> with darapladib added to optimal medical therapy and initiated within 30 days of hospitalization did not reduce the risk of major coronary events.

The STAbilization of Atherosclerotic plaque By Initiation of darapLadIb TherapY (STABILITY) trial was a double-blind trial in patients randomized more than one month after a myocardial infarction to double-blind darapladib (160 mg daily) or placebo (daily). The primary endpoint was a composite of cardiovascular death, myocardial infarction, or stroke. Secondary endpoints included the components of the primary endpoint as well as major coronary events (death from coronary heart disease, myocardial infarction, or urgent coronary revascularization for myocardial ischemia) and total coronary events (death from coronary heart disease, myocardial infarction, hospitalization for unstable angina, or any coronary revascularization). Patients were followed up for a median of 3.7 years. In patients with stable coronary heart disease, darapladib did not significantly reduce the risk of the primary composite endpoint of cardiovascular death, myocardial infarction, or stroke.

Darapladib is a selective Lp-PLA<sub>2</sub> inhibitor that is under investigation for its potential to stabilize high-risk atherosclerotic plaques and potentially reduce cardiovascular events [104,105]. A phase 2 clinical trial in 959 patients with CHD or CHD-risk equivalents demonstrated that darapladib was effective at producing sustained inhibition of plasma Lp-PLA<sub>2</sub> activation in patients on atorvastatin therapy. This study was a post hoc analysis of this phase 2 trial studying high-risk patients with a diagnosis of peripheral arterial disease (PAD). Despite a more aggressive baseline risk factor profile, darapladib was equally effective at reducing Lp-PLA<sub>2</sub> in patients with and without PAD [106]. Johnson et al. [107] assessed the effects of darapladib on both plasma and plaque Lp-PLA<sub>2</sub> activity in patients undergoing elective carotid endarterectomy randomized to darapladib 40 mg (n = 34), 80 mg (n = 34), or placebo (n = 34) for 14 days. Patients were followed by carotid endarterectomy 24 hours after the last dose of study medication. Darapladib reduced plasma and plaque Lp-PLA<sub>2</sub> activity compared with placebo.

## 8. Conclusion

Within the last decade, a broad range of biomarkers associated with an increased risk for death and cardiovascular/cerebrovascular endpoints have been identified. Epidemiological studies clearly indicated that Lp-PLA<sub>2</sub> has the potential to become a clinically useful biomarker because it promotes independent information in the diagnosis, and especially cardiovascular/cerebrovascular risk stratification. In the future, we can expect new drugs (new Lp-PLA<sub>2</sub> inhibitors) that will affect patient's management, and assessing the effect of Lp-PLA<sub>2</sub> inhibition on cardiovascular endpoints can provide definitive answers.

However, further clinical validation in well-designed observational and interventional studies is needed before these recommendations can be properly evaluated in order to include them in the clinical diagnostic algorithms.

Currently, Lp-PLA<sub>2</sub> measurement has only been reserved to patients with moderate and high cardiovascular risk, rather than healthy population or low-risk patients, since the values of Lp-PLA<sub>2</sub> in these population groups are insignificant. Also, the evaluation of Lp-PLA<sub>2</sub> in combination with noninvasive imaging could be expected. The formation of best-case model from Lp-PLA<sub>2</sub> and other biomarkers can yield the best patient stratification algorithm. The next step could be cost-effectiveness analysis of more accurate risk stratification with biomarker testing. Future studies need to focus on exploring the potential of this biomarker and evaluating the effects of Lp-PLA<sub>2</sub> inhibition on human populations.

The final decision on which test to use for Lp-PLA<sub>2</sub> determination, test based on mass or activity of Lp-PLA<sub>2</sub>, together with the development of commercially available automated, robust, valid, high throughput, cost-effective test capable of increasing agility and reducing the analytical imprecision can be adopted routinely in clinical practice for better risk stratification and therapeutic choice in patients with cardiovascular/cerebrovascular disease.

## Author details

Sanja Stankovic\* and Milika Asanin

\*Address all correspondence to:

Clinical Center of Serbia, Belgrade, Serbia

## References

- [1] Schaloske RH, Dennis EA. The phospholipase A2 superfamily and its group numbering system. *Biochim Biophys Acta* 2006;1761(11):1246-1259.
- [2] Burke JE, Dennis EA. Phospholipase A2 biochemistry. *Cardiovasc Drugs Ther* 2009; 23:49-59.
- [3] Tew DG, Southan C, Rice SQ, Lawrence MP, Li H, Boyd HF, Moores K, Gloger IS, Macphee CH. Purification, properties, sequencing, and cloning of a lipoprotein-associated, serine-dependent phospholipase involved in the oxidative modification of low-density lipoproteins. *Arterioscler Thromb Vasc Biol* 1996;16(4):591-599.
- [4] Zalewski A, Macphee C. Role of lipoprotein-associated phospholipase A2 in atherosclerosis: biology, epidemiology, and possible therapeutic target. *Arterioscler Thromb Vasc Biol* 2005;25(5):923-931.
- [5] Tjoelker LW, Wilder C, Eberhardt C, Stafforini DM, Dietsch G, Schimpf B, Hooper S, Le Trong H, Cousens LS, Zimmerman GA, Yamada Y, McIntyre TM, Prescott SM, Gray PW. Anti-inflammatory properties of a platelet-activating factor acetylhydrolase. *Nature* 1995;374(6522):549-553.
- [6] Stafforini DM, Elstad MR, McIntyre TM, Zimmerman GA, Prescott SM. Human macrophages secrete platelet-activating factor acetylhydrolase. *J Biol Chem* 1990;265(17): 9682-9687.
- [7] Asano K, Okamoto S, Fukunaga K, Shiomi T, Mori T, Iwata M, Ikeda Y, Yamaguchi K. Cellular source(s) of platelet-activating-factor acetylhydrolase activity in plasma. *Biochem Biophys Res Commun* 1999;261(2):511-514.
- [8] Tarbet EB, Stafforini DM, Elstad MR, Zimmerman GA, McIntyre TM, Prescott SM. Liver cells secrete the plasma form of platelet-activating factor acetylhydrolase. *J Biol Chem* 1991;266(25):16667-16673.
- [9] Mitsios JV, Vini MP, Stengel D, Ninio E, Tselepis AD. Human platelets secrete the plasma type of platelet-activating factor acetylhydrolase primarily associated with microparticles. *Arterioscler Thromb Vasc Biol* 2006;26(8):1907-1913.

- [10] Tellis CC, Tselepis AD. The role of lipoprotein-associated phospholipase A<sub>2</sub> in atherosclerosis may depend on its lipoprotein carrier in plasma. *Biochim Biophys Acta* 2009;1791(5):327-338.
- [11] Ostermann G, Kostner GM, Gries A, Malle E, Till U. The contribution of individual lipoproteins to the degradation of platelet-activating factor in human serum. *Haemostasis* 1989;19(3):160-168.
- [12] Blencowe C, Hermetter A, Kostner GM, Deigner HP. Enhanced association of platelet-activating factor acetylhydrolase with lipoprotein (a) in comparison with low density lipoprotein. *J Biol Chem* 1995;270(52): 31151-31157.
- [13] Tselepis AD, Dentan C, Karabina SA, Chapman MJ, Ninio E. PAF degrading acetylhydrolase is preferentially associated with dense LDL and VHDL-1 in human plasma. Catalytic characteristics and relation to the monocyte-derived enzyme. *Arterioscler Thromb Vasc Biol* 1995;15:1764-1773.
- [14] Benítez S, Sánchez-Quesada JL, Ribas V, Jorba O, Blanco-Vaca F, González-Sastre F, Ordóñez-Llanos J. Platelet-activating factor acetylhydrolase is mainly associated with electronegative low-density lipoprotein subfraction. *Circulation* 2003;108(1):92-96.
- [15] Stafforini DM, McIntyre TM, Carter ME, Prescott SM. Human plasma platelet-activating factor acetylhydrolase. Association with lipoprotein particles and role in the degradation of platelet activating factor. *J Biol Chem* 1987;262:4215-4222.
- [16] Bhatti S, Hakeem A, Cilingiroglu M. Lp-PLA(2) as a marker of cardiovascular diseases. *Curr Atheroscler Rep* 2010; 12(2):140-144.
- [17] Lavi S, McConnell JP, Rihal CS, Prasad A, Mathew V, Lerman LO, Lerman A. Local production of lipoprotein-associated phospholipase A<sub>2</sub> and lysophosphatidylcholine in the coronary circulation: association with early coronary atherosclerosis and endothelial dysfunction in humans. *Circulation* 2007;115:2715-2721.
- [18] Macphee CH. Lipoprotein-associated phospholipase A<sub>2</sub>: a potential new risk factor for coronary artery disease and a therapeutic target. *Curr Opin Pharmacol* 2001;1(2): 121-125.
- [19] Kume N, Cybulsky MI, Gimbrone MA Jr. Lysophosphatidylcholine, a component of atherogenic lipoproteins, induces mononuclear leukocyte adhesion molecules in cultured human and rabbit arterial endothelial cells. *J Clin Invest* 1992;90:1138-1144.
- [20] Quinn MT, Parthasarathy S, Steinberg D. Lysophosphatidylcholine: a chemotactic factor for human monocytes and its potential role in atherogenesis. *Proc Natl Acad Sci USA* 1988;85:2805-2809.
- [21] Vittos Oana, Toana B, Vittos A, Moldoveanu E. Lipoprotein-associated phospholipase A<sub>2</sub> (Lp-PLA<sub>2</sub>): a review of its role and significance as a cardiovascular biomarker. *Biomarkers* 2012;17(4): 289-302.

- [22] Zalewski A, Nelson JJ, Hegg L, Macphee C. Lp-PLA2: a new kid on the block. *Clin Chem* 2006;52(9):1645-1650.
- [23] Davidson MH, Corson MA, Alberts MJ, Anderson JL, Gorelick PB, Jones PH, Lerman A, McConnell JP, Weintraub HS. Consensus panel recommendation for incorporating lipoprotein-associated phospholipase A2 testing into cardiovascular disease risk assessment guidelines. *Am J Cardiol* 2008;101:51F-57F.
- [24] Thompson A, Gao P, Orfei L, Watson S, Di Angelantonio E, Kaptoge S, Ballantyne C, Cannon CP, Criqui M, Cushman M, Hofman A, Packard C, Thompson SG, Collins R, Danesh J, for the Lp-PLA(2) Studies Collaboration. Lipoprotein-associated phospholipase A(2) and risk of coronary disease, stroke, and mortality: collaborative analysis of 32 prospective studies. *Lancet* 2010;375:1536-1544.
- [25] O'Donoghue M, Morrow DA, Sabatine MS, Murphy SA, McCabe CH, Cannon CP, Braunwald E. Lipoprotein-associated phospholipase A2 and its association with cardiovascular outcomes in patients with acute coronary syndromes in the PROVE IT-TIMI 22 (PRavastatin Or atorVastatin Evaluation and Infection Therapy – Thrombolysis In Myocardial Infarction) trial. *Circulation* 2006;113(14):1745-1752.
- [26] Koenig W, Khuseynova N, Lowel H, Trischler G, Meisinger C. Lipoprotein-associated phospholipase A2 adds to risk prediction of incident coronary events by C-reactive protein in apparently healthy middle-aged men from the general population: results from the 14-year follow-up of a large cohort from southern Germany. *Circulation* 2004;110:1903-1908.
- [27] Persson M, Hedblad B, Nelson JJ, Berglund G. Elevated Lp-PLA2 levels add prognostic information to the metabolic syndrome on incidence of cardiovascular events among middle-aged nondiabetic subjects. *Arterioscler Thromb Vasc Biol* 2007; 27(6): 1411-1416.
- [28] Jenny NS, Solomon C, Cushman M, Tracy RP, Nelson JJ, Psaty BM, Furberg CD. Lipoprotein-associated phospholipase A(2) (Lp-PLA(2)) and risk of cardiovascular disease in older adults: results from the Cardiovascular Health Study. *Atherosclerosis* 2010; 209(2):528-532.
- [29] Wolfert RL, Kim NW, Selby RG. Biological variability and specificity of lipoprotein associated phospholipase A2, a novel marker of cardiovascular risk. *Circulation* 2004;110(Suppl III):III-309. (Abstract).
- [30] Ballantyne CM, Hoogeveen RC, Bang H, Coresh J, Folsom AR, Heiss G, Sharrett AR. Lipoprotein-associated phospholipase A2, high-sensitivity C-reactive protein, and risk for incident coronary heart disease in middle-aged men and women in the Atherosclerosis Risk in Communities (ARIC) study. *Circulation* 2004;109:837-842.
- [31] Oei HH, van der Meer IM, Hofman A, Koudstaal PJ, Stijnen T, Breteler MM, Witteman JC. Lipoprotein-associated phospholipase A2 activity is associated with risk of

- coronary heart disease and ischemic stroke: the Rotterdam Study. *Circulation* 2005;111(5):570-575.
- [32] Stankovic S, Asanin M, Trifunovic D, Majkic-Singh N, Miljković A, Ignjatovic S, Mrdovic I, Matic D, Savic L, Ostojic M, Vasiljevic Z. Utility of lipoprotein-associated phospholipase A2 for prediction of 30-day major adverse coronary event in patients with the first anterior ST-segment elevation myocardial infarction treated by primary percutaneous coronary intervention. *Clin Lab* 2012;58(11-12):1135-1144.
- [33] Sabatine MS, Morrow DA, O'Donoghue M, Jablonksi KA, Rice MM, Solomon S, Rosenberg Y, Domanski MJ, Hsia J; PEACE Investigators. Prognostic utility of lipoprotein-associated phospholipase A2 for cardiovascular outcomes in patients with stable coronary artery disease. *Arterioscler Thromb Vasc Biol* 2007;27(11):2463-2469.
- [34] Hatoum IJ, Nelson JJ, Cook NR, Hu FB, Rimm EB. Dietary, lifestyle, and clinical predictors of lipoprotein-associated phospholipase A2 activity in individuals without coronary artery disease. *Am J Clin Nutr* 2010;91(3):786-793.
- [35] Miyaura S, Maki N, Byrd W, Johnston JM. The hormonal regulation of platelet-activating factor acetylhydrolase activity in plasma. *Lipids* 1991;26:1015-1020.
- [36] Mendelsohn ME, Karas RH. The protective effects of estrogen on the cardiovascular system. *N Engl J Med* 1999; 340:1801-1811.
- [37] Yoshimura T, Ohshige A, Maeda T, Ito M, Okamura H. Estrogen replacement therapy decreases platelet-activating factor acetylhydrolase activity in post-menopausal women. *Maturitas* 1999;31:249-253.
- [38] Ohshige A, Ito M, Koyama H, Maeda T, Yoshimura T, Okamura H. Effects of estrogen and progesterone on platelet activating factor acetylhydrolase activity in ovariectomised rats. *Artery* 1994;21:234-242.
- [39] Yanbaeva DG, Dentener MA, Creutzberg EC, Wesseling G, Wouters EF. Systemic effects of smoking. *Chest* 2007;131:1557-1566.
- [40] Perk J, De Backer G, Gohlke H, Graham I, Reiner Z, Verschuren M, Albus C, Benlian P, Boysen G, Cifkova R, Deaton C, Ebrahim S, Fisher M, Germano G, Hobbs R, Hoes A, Karadeniz S, Mezzani A, Prescott E, Ryden L, Scherer M, Syväne M, Scholte op Reimer WJ, Vrints C, Wood D, Zamorano JL, Zannad F; European Association for Cardiovascular Prevention & Rehabilitation (EACPR); ESC Committee for Practice Guidelines (CPG). European Guidelines on cardiovascular disease prevention in clinical practice (version 2012). The Fifth Joint Task Force of the European Society of Cardiology and Other Societies on Cardiovascular Disease Prevention in Clinical Practice (constituted by representatives of nine societies and by invited experts). *Eur Heart J* 2012;33(13):1635-1701.
- [41] Greenland P, Alpert JS, Beller GA, Benjamin EJ, Budoff MJ, Fayad ZA, Foster E, Hlatky MA, Hodgson JM, Kushner FG, Lauer MS, Shaw LJ, Smith SC Jr, Taylor AJ, Weintraub WS, Wenger NK, Jacobs AK, Smith SC Jr, Anderson JL, Albert N, Buller CE,

- Creager MA, Ettinger SM, Guyton RA, Halperin JL, Hochman JS, Kushner FG, Nishimura R, Ohman EM, Page RL, Stevenson WG, Tarkington LG, Yancy CW; American College of Cardiology Foundation; American Heart Association. 2010 ACCF/AHA guideline for assessment of cardiovascular risk in asymptomatic adults: a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines. *J Am Coll Cardiol* 2010;56(25):e50-e103.
- [42] Goldstein LB, Bushnell CD, Adams RJ, Appel LJ, Braun LT, Chaturvedi S, Creager MA, Culebras A, Eckel RH, Hart RG, Hinchey JA, Howard VJ, Jauch EC, Levine SR, Meschia JF, Moore WS, Nixon JV, Pearson TA; American Heart Association Stroke Council; Council on Cardiovascular Nursing; Council on Epidemiology and Prevention; Council for High Blood Pressure Research; Council on Peripheral Vascular Disease; and Interdisciplinary Council on Quality of Care and Outcomes Research. Guidelines for the primary prevention of stroke: a guideline for healthcare professionals from the American Heart Association/American Stroke Association. *Stroke* 2011;42(2):517-584.
- [43] Jellinger PS, Smith DA, Mehta AE, Ganda O, Handelsman Y, Rodbard HW, Shepherd MD, Seibel JA; AACE Task Force for Management of Dyslipidemia and Prevention of Atherosclerosis. American Association of Clinical Endocrinologists' Guidelines for Management of Dyslipidemia and Prevention of Atherosclerosis. *Endocr Pract* 2012;18 (Suppl 1):1-78.
- [44] Packard CJ, O'Reilly DS, Caslake MJ, McMahon AD, Ford I, Cooney J, Macphee CH, Suckling KE, Krishna M, Wilkinson FE, Rumley A, Lowe GD. Lipoprotein-associated phospholipase A2 as an independent predictor of coronary heart disease. West of Scotland Coronary Prevention Study Group. *N Engl J Med* 2000;343(16):1148-1155.
- [45] Ballantyne CM, Hoogeveen RC, Bang H, Coresh J, Folsom AR, Chambless LE, Myerson M, Wu KK, Sharrett AR, Boerwinkle E. Lipoprotein-associated phospholipase A2, high-sensitivity C-reactive protein, and risk for incident ischemic stroke in middle-aged men and women in the Atherosclerosis Risk in Communities (ARIC) study. *Arch Intern Med* 2005;165(21):2479-2484.
- [46] Corsetti JP, Rainwater DL, Moss AJ, Zareba W, Sparks CE. High lipoprotein-associated phospholipase A2 is a risk factor for recurrent coronary events in postinfarction patients. *Clin Chem* 2006;52:1331-1338.
- [47] Brilakis ES, McConnell JP, Lennon RJ, Elesber AA, Meyer JG, Berger PB. Association of lipoprotein-associated phospholipase A2 levels with coronary artery disease risk factors, angiographic coronary artery disease, and major adverse events at follow-up. *Eur Heart J* 2005;26:137-144.
- [48] Gerber Y, McConnell JP, Jaffe AS, Weston SA, Killian JM, Roger VL. Lipoprotein-associated phospholipase A2 and prognosis after myocardial infarction in the community. *Arterioscler Thromb Vasc Biol* 2006;26:2517-2522.

- [49] Anderson JL. Lipoprotein-associated phospholipase A2: an independent predictor of coronary artery disease events in primary and secondary prevention. *Am J Cardiol* 2008;101:23F-33F.
- [50] Winkler K, Hoffmann MM, Winkelmann BR, Friedrich I, Schäfer G, Seelhorst U, Wellnitz B, Wieland H, Boehm BO, März W. Lipoprotein-associated phospholipase A2 predicts 5-year cardiac mortality independently of established risk factors and adds prognostic information in patients with low and medium high-sensitivity C-reactive protein (the Ludwigshafen risk and cardiovascular health study). *Clin Chem* 2007;53(8):1440-1447.
- [51] Yusuf S, Hawken S, Ounpuu S, Dans T, Avezum A, Lanas F, McQueen M, Budaj A, Pais P, Varigos J, Lisheng L. Effect of potentially modifiable risk factors associated with myocardial infarction in 52 countries (the INTERHEART study): case-control study. *Lancet* 2004; 364:937-952.
- [52] 2002 Third Report of the National Cholesterol Education Program (NCEP) Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III) final report. *Circulation* 2002; 106:3143-3421.
- [53] Anuurad E, Ozturk Z, Enkhmaa B, Pearson TA, Berglund L. Association of lipoprotein-associated phospholipase A2 with coronary artery disease in African-Americans and Caucasians. *J Clin Endocrinol Metab* 2010;95(5):2376-2383.
- [54] Corson MA, Jones PH, Davidson MH. Review of the evidence for the clinical utility of lipoprotein-associated phospholipase a2 as a cardiovascular risk marker. *Am J Cardiol* 2008;101[suppl]:41F-50F.
- [55] Madjid M, Ali M, Willerson JT. Lipoprotein-associated phospholipase A2 as a novel risk marker for cardiovascular disease: a systematic review of the literature. *Tex Heart Inst J* 2010;37:25-39.
- [56] Mallat Z, Lambeau G, Tedgui A. Lipoprotein-associated and secreted phospholipases A2 in cardiovascular disease: roles as biological effectors and biomarkers. *Circulation* 2010;122:2183-2200.
- [57] May HT, Horne BD, Anderson JL, Wolfert RL, Muhlestein JB, Renlund DG, Clarke JL, Kolek MJ, Bair TL, Pearson RR, Sudhir K, Carlquist JF. Lipoprotein-associated phospholipase A2 independently predicts the angiographic diagnosis of coronary artery disease and coronary death. *Am Heart J* 2006;152 (5):997-1003.
- [58] Li N, LiS, Yu C, Gu S. Plasma Lp-PLA2 in acute coronary syndrome: association with major adverse cardiac events in a community-based cohort. *Postgrad Med* 2010;122(4): 200-205.
- [59] Möckel M, Müller R, Vollert J, Müller C, Danne O, Gareis R, Störk T, Dietz R, Koenig W. Lipoprotein-associated phospholipase A2 for early risk stratification in patients with suspected acute coronary syndrome: a multi-marker approach: the North

- Wuerttemberg and Berlin Infarction Study-II (NOBIS-II). *Clin Res Cardiol* 2007;96(9):604-612.
- [60] van Vark LC, Kardys I, Bleumink GS, Knetsch AM, Deckers JW, Hofman A, Stricker BH, Witteman JC. Lipoprotein-associated phospholipase A2 activity and risk of heart failure: the Rotterdam Study. *Eur Heart J* 2006;27:2346-2352.
- [61] Suzuki T, Solomon C, Jenny NS, Tracy R, Nelson JJ, Psaty BM, Furberg C, Cushman M. Lipoprotein-associated phospholipase A(2) and risk of congestive heart failure in older adults: the Cardiovascular Health Study. *Circ Heart Fail* 2009;2:429-436.
- [62] Furberg CD, Nelson JJ, Solomon C, Cushman M, Jenny NS, Psaty BM. Distribution and correlates of lipoprotein-associated phospholipase A2 in an elderly cohort: the Cardiovascular Health Study. *J Am Geriatr Soc* 2008;56:792-799.
- [63] Gerber Y, Dunlay SM, Jaffe AS, McConnell JP, Weston SA, Killian JM, Roger VL. Plasma lipoprotein-associated phospholipase A2 levels in heart failure: association with mortality in the community. *Atherosclerosis* 2009;203(2):593-598.
- [64] Moldoveanu E, Serban M, Marta D, Serban I, Huica R. Lipoprotein-associated phospholipase A2 in patients with preserved left ventricular ejection fraction. *Biomarkers* 2011;16(7):587-589.
- [65] Raichlin E, McConnell JP, Bae JH, Kremers WK, Lerman A, Frantz RP. Lipoprotein-associated phospholipase A2 predicts progression of cardiac allograft vasculopathy and increased risk of cardiovascular events in heart transplant patients. *Transplantation* 2008;85:963-968.
- [66] Persson M, Berglund G, Nelson JJ, Hedblad B. Lp-PLA2 activity and mass are associated with increased incidence of ischemic stroke: a population-based cohort study from Malmö, Sweden. *Atherosclerosis* 2008;200(1):191-198.
- [67] Tsimikas S, Willeit J, Knoflach M, Mayr M, Egger G, Notdurfter M, Witztum JL, Wiedermann CJ, Xu Q, Kiechl S. Lipoprotein-associated phospholipase A2 activity, ferritin levels, metabolic syndrome, and 10-year cardiovascular and non-cardiovascular mortality: results from the Bruneck study. *Eur Heart J* 2009;30:107-115.
- [68] Tselepis AD, Hahalis G, Tellis CC, Papavasiliou EC, Mylona PT, Kourakli A, Alexopoulos DC. Plasma levels of lipoprotein-associated phospholipase A(2) are increased in patients with  $\beta$ -thalassemia. *J Lipid Res* 2010;51(11):3331-3341.
- [69] Elkind MS, Tai W, Coates K, Paik MC, Sacco RL. High-sensitivity C-reactive protein, lipoprotein-associated phospholipase A2, and outcome after ischemic stroke. *Arch Intern Med* 2006;166:2073-2080.
- [70] Delgado P, Chacón P, Penalba A, Pelegrí D, Merino C, Ribó M, Rubiera M, Álvarez-Sabin J, Montaner J. Temporal profile and prognostic value of Lp-PLA2 mass and activity in the acute stroke setting. *Atherosclerosis* 2012;220(2):532-536.

- [71] Gorelick PB. Lipoprotein-associated phospholipase A2 and risk of stroke. *Am J Cardiol* 2008;101:34F-40F.
- [72] Mannheim D, Herrmann J, Versari D, Gössl M, Meyer FB, McConnell JP, Lerman LO, Lerman A. Enhanced expression of Lp-PLA2 and lysophosphatidylcholine in symptomatic carotid atherosclerotic plaques. *Stroke* 2008;39:1448-1455.
- [73] Sarlon-Bartoli G, Boudes A, Buffat C, Bartoli MA, Piercecchi-Marti MD, Sarlon E, Arnaud L, Bennis Y, Thevenin B, Squarcioni C, Nicoli F, Dignat-George F, Sabatier F, Magnan PE; RISC Study Group. Circulating lipoprotein-associated phospholipase A2 in high-grade carotid stenosis: a new biomarker for predicting unstable plaque. *Eur J Vasc Endovasc Surg* 2012;43(2):154-159.
- [74] Kakkos SK, Tsolakis IA. The emerging role of lipoprotein-associated phospholipase A2 in cerebrovascular disease. *Eur J Vasc Endovasc Surg* 2012; 43:160.
- [75] Shoamanesh A, Preis SR, Beiser AS, Vasan RS, Benjamin EJ, Kase CS, Wolf PA, DeCarli C, Romero JR, Seshadri S. Inflammatory biomarkers, cerebral microbleeds, and small vessel disease: Framingham Heart Study Neurology 2015;84(8):825-32.
- [76] Caslake MJ, Packard CJ, Suckling KE, Holmes SD, Chamberlain P, Macphhee CH. Lipoprotein-associated phospholipaseA(2), platelet-activating factor acetylhydrolase: a potential new risk factor for coronary artery disease. *Atherosclerosis* 2000;150:413-419.
- [77] Ishihara M, Iwasaki T, Nagano M, Ishii J, Takano M, Kujiraoka T, Tsuji M, Hattori H, Emi M. Functional impairment of two novel mutations detected in lipoprotein-associated phospholipase A2 (Lp-PLA2) deficiency patients. *J Hum Genet* 2004;49:302-307.
- [78] Guerra R, Zhao B, Mooser V, Stafforini D, Johnston JM, Cohen JC. Determinants of plasma platelet-activating factor acetylhydrolase: heritability and relationship to plasma lipoproteins. *J Lipid Res* 1997;38: 2281-2288.
- [79] Schnabel R, Dupuis J, Larson MG, Lunetta KL, Robins SJ, Zhu Y, Rong J, Yin X, Stirnadel HA, Nelson JJ, Wilson PW, Keaney JF, Vasan RS, Benjamin EJ. Clinical and genetic factors associated with lipoprotein-associated phospholipase A2 in the Framingham Heart Study. *Atherosclerosis* 2009;204:601-607.
- [80] Suchindran S, Rivedal D, Guyton JR, Milledge T, Gao X, Benjamin A, Rowell J, Ginsburg GS, McCarthy JJ. Genome-wide association study of Lp-PLA2 activity and mass in the Framingham Heart Study. *PLoS Genet* 2010;6:e1000928.
- [81] Grallert H, Dupuis J, Bis JC, Dehghan A, Barbalic M, Baumert J, Lu C, Smith NL, Uitterlinden AG, Roberts R, Khuseyinova N, Schnabel RB, Rice KM, Rivadeneira F, Hoogeveen RC, Fontes JD, Meisinger C, Keaney JF Jr, Lemaitre R, Aulchenko YS, Vasan RS, Ellis S, Hazen SL, van Duijn CM, Nelson JJ, März W, Schunkert H, McPherson RM, Stirnadel-Farrant HA, Psaty BM, Gieger C, Siscovick D, Hofman A, Illig T, Cushman M, Yamamoto JF, Rotter JI, Larson MG, Stewart AF, Boerwinkle E, Witteman JC, Tracy RP, Koenig W, Benjamin EJ, Ballantyne CM. Eight genetic loci associated with variation in lipoprotein-associated phospholipase A2 mass and activity

- and coronary heart disease: meta-analysis of genome-wide association studies from five community-based studies. *Eur Heart J* 2012; 33(2):238-251.
- [82] Psaty BM, O'Donnell CJ, Gudnason V, Lunetta KL, Folsom AR, Rotter JI, Uitterlinden AG, Harris TB, Witteman JC, Boerwinkle E, on behalf of the CHARGE Consortium. Cohorts for Heart and Aging Research in Genomic Epidemiology (CHARGE) Consortium: design of prospective meta-analyses of genomewide association studies from 5 cohorts. *Circ Cardiovasc Genet* 2009;2:73-80.
- [83] Willer CJ, Sanna S, Jackson AU, Scuteri A, Bonnycastle LL, Clarke R, Heath SC, Timpson NJ, Najjar SS, Stringham HM, Strait J, Duren WL, Maschio A, Busonero F, Mulas A, Albai G, Swift AJ, Morken MA, Narisu N, Bennett D, Parish S, Shen H, Galan P, Meneton P, Hercberg S, Zelenika D, Chen WM, Li Y, Scott LJ, Scheet PA, Sundvall J, Watanabe RM, Nagaraja R, Ebrahim S, Lawlor DA, Ben-Shlomo Y, Davey-Smith G, Shuldiner AR, Collins R, Bergman RN, Uda M, Tuomilehto J, Cao A, Collins FS, Lakatta E, Lathrop GM, Boehnke M, Schlessinger D, Mohlke KL, Abecasis GR. Newly identified loci that influence lipid concentrations and risk of coronary artery disease. *Nat Genet* 2008;40:161-169.
- [84] Bell R, Collier DA, Rice SQ, Roberts GW, MacPhee CH, Kerwin RW, Price J, Gloger IS. Systematic screening of the LDL-PLA2 gene for polymorphic variants and case control analysis in schizophrenia. *Biochem Biophys Res Commun* 1997;241:630-635.
- [85] Kruse S, Mao XQ, Heinzmann A, Blattmann S, Roberts MH, Braun S, Gao PS, Forster J, Kuehr J, Hopkin JM, Shirakawa T, Deichmann KA. The Ile198Thr and Ala379Val variants of plasmatic PAF-acetylhydrolase impair catalytical activities and are associated with atopy and asthma. *Am J Hum Genet* 2000;66:1522-1530.
- [86] Hoffmann MM, Winkler K, Renner W, Winkelmann BR, Seelhorst U, Wellnitz B, Boehm BO, Marz W. Genetic variants and haplotypes of lipoprotein associated phospholipase A2 and their influence on cardiovascular disease (The Ludwigshafen Risk and Cardiovascular Health Study). *J Thromb Haemost* 2009;7:41-48.
- [87] Li L, Qi L, Lv N, Gao Q, Cheng Y, Wei Y, Ye J, Yan X, Dang A. Association between lipoprotein associated phospholipase A2 gene polymorphism and coronary artery disease in the Chinese Han population. *Ann Hum Genet* 2011;75(5):605-611.
- [88] Yamada Y, Ichihara S, Fujimura T, Yokota M. Identification of the G994→ T missense in exon 9 of the plasma platelet-activating factor acetylhydrolase gene as an independent risk factor for coronary artery disease in Japanese men. *Metabolism* 1998;47(2):177-181.
- [89] Shimokata K, Yamada Y, Kondo T, Ichihara S, Izawa H, Nagata K, Murohara T, Ohno M, Yokota M. Association of gene polymorphisms with coronary artery disease in individuals with or without nonfamilial hypercholesterolemia. *Atherosclerosis* 2004;172(1):167-173.
- [90] Jang Y, Waterworth D, Lee JE, Song K, Kim S, Kim HS, Park KW, Cho HJ, Oh IY, Park JE, Lee BS, Ku HJ, Shin DJ, Lee JH, Jee SH, Han BG, Jang HY, Cho EY, Vallance

- P, Whittaker J, Cardon L, Mooser V. Carriage of the V279F null allele within the gene encoding Lp-PLA2 is protective from coronary artery disease in South Korean males. *PLoS One* 2011;6(4): e18208.
- [91] Hou L, Chen S, Yu H, Lu X, Chen J, Wang L, Huang J, Fan Z, Gu D. Associations of PLA2G7 gene polymorphisms with plasma lipoprotein-associated phospholipase A2 activity and coronary heart disease in a Chinese Han population: the Beijing Atherosclerosis Study. *Hum Genet* 2009;125:11-20.
- [92] Wang Q, Hao Y, Mo X, Wang L, Lu X, Huang J, Cao J, Li H, Gu D. PLA2G7 gene polymorphisms and coronary heart disease risk: a meta-analysis. *Thromb Res* 2010;126(6):498-503.
- [93] Ninio E, Tregouet D, Carrier JL, Stengel D, Bickel C, Perret C, Rupprecht HJ, Cambien F, Blankenberg S, Tiret L. Platelet-activating factor-acetylhydrolase and PAF-receptor gene haplotypes in relation to future cardiovascular event in patients with coronary artery disease. *Hum Mol Genet* 2004;13:1341-1351.
- [94] Liu PY, Li YH, Wu HL, Chao TH, Tsai LM, Lin LJ, Shi GY, Chen JH. Platelet-activating factor acetylhydrolase A379V (exon11) gene polymorphism is an independent and functional risk factor for premature myocardial infarction. *J Thromb Haemost* 2006;4:1023-1028.
- [95] Wootton PT, Stephens JW, Hurel SJ, Durand H, Cooper J, Ninio E, Humphries SE, Talmud PJ. Lp-PLA2 activity and PLA2G7 A379V genotype in patients with diabetes mellitus. *Atherosclerosis* 2006;189:149-156.
- [96] Abuzeid AM, Hawe E, Humphries SE, Talmud PJ. Association between the Ala379Val variant of the lipoprotein associated phospholipase A2 and risk of myocardial infarction in the north and south of Europe. *Atherosclerosis* 2003;168:283-288.
- [97] Casas JP, Ninio E, Panayiotou A, Palmén J, Cooper JA, Ricketts SL, Sofat R, Nicolaidis AN, Corsetti JP, Fowkes FG, Tzoulaki I, Kumari M, Brunner EJ, Kivimaki M, Marmot MG, Hoffmann MM, Winkler K, März W, Ye S, Stirnadel HA, Boekholdt SM, Khaw KT, Humphries SE, Sandhu MS, Hingorani AD, Talmud PJ. PLA2G7 genotype, lipoprotein-associated phospholipase A2 activity, and coronary heart disease risk in 10 494 cases and 15 624 controls of European Ancestry. *Circulation* 2010;121(21):2284-2293.
- [98] Serruys PW, García-García HM, Buszman P, Erne P, Verheye S, Aschermann M, Duckers H, Bleie O, Dudek D, Bøtker HE, von Birgelen C, D'Amico D, Hutchinson T, Zambanini A, Mastik F, van Es GA, van der Steen AF, Vince DG, Ganz P, Hamm CW, Wijns W, Zalewski A; Integrated Biomarker and Imaging Study-2 Investigators. Effects of the direct lipoprotein-associated phospholipase A(2) inhibitor darapladib on human coronary atherosclerotic plaque. *Circulation* 2008; 118:1172-1182.
- [99] Mohler ER 3rd, Ballantyne CM, Davidson MH, Hanefeld M, Ruilope LM, Johnson JL, Zalewski A; Darapladib Investigators. The effect of darapladib on plasma lipopro-

tein-associated phospholipase A2 activity and cardiovascular biomarkers in patients with stable coronary heart disease or coronary heart disease risk equivalent: The results of a multicenter, randomized, double-blind, placebo-controlled study. *J Am Coll Cardiol* 2008; 51:1632-1641.

- [100] Daida H, Iwase T, Yagi S, Ando H, Nakajima H. Effect of darapladib on plasma lipoprotein-associated phospholipase A2 activity in Japanese dyslipidemic patients, with exploratory analysis of a PLA2G7 gene polymorphism of Val279Phe. *Circ J* 2013;77(6):1518-1525.
- [101] O'Donoghue ML, Braunwald E, White HD, Serruys P, Steg PG, Hochman J, Maggioni AP, Bode C, Weaver D, Johnson JL, Cicconetti G, Lukas MA, Tarka E, Cannon CP. Study design and rationale for the Stabilization of pLaques using Darapladib-Thrombolysis in Myocardial Infarction (SOLID-TIMI 52) trial in patients after an acute coronary syndrome. *Am Heart J* 2011;162(4):613-619.
- [102] O'Donoghue ML, Braunwald E, White HD, Lukas MA, Tarka E, Steg PG, Hochman JS, Bode C, Maggioni AP, Im K, Shannon JB, Davies RY, Murphy SA, Crugnale SE, Wiviott SD, Bonaca MP, Watson DF, Weaver WD, Serruys PW, Cannon CP; SOLID-TIMI 52 Investigators, Steen DL. Effect of darapladib on major coronary events after an acute coronary syndrome: the SOLID-TIMI 52 randomized clinical trial. *JAMA* 2014;312(10):1006-1015.
- [103] STABILITY Investigators, White HD, Held C, Stewart R, Tarka E, Brown R, Davies RY, Budaj A, Harrington RA, Steg PG, Ardissino D, Armstrong PW, Avezum A, Aylward PE, Bryce A, Chen H, Chen MF, Corbalan R, Dalby AJ, Danchin N, De Winter RJ, Denchev S, Diaz R, Elisaf M, Flather MD, Goudev AR, Granger CB, Grinfeld L, Hochman JS, Husted S, Kim HS, Koenig W, Linhart A, Lonn E, López-Sendón J, Manolis AJ, Mohler ER 3rd, Nicolau JC, Pais P, Parkhomenko A, Pedersen TR, Pella D, Ramos-Corrales MA, Ruda M, Sereg M, Siddique S, Sinnaeve P, Smith P, Sritara P, Swart HP, Sy RG, Teramoto T, Tse HF, Watson D, Weaver WD, Weiss R, Viigimaa M, Vinereanu D, Zhu J, Cannon CP, Wallentin L. Darapladib for preventing ischemic events in stable coronary heart disease. *N Engl J Med* 2014;370(18):1702-1711.
- [104] White H. Editorial: why inhibition of lipoprotein-associated phospholipase A2 has the potential to improve patient outcomes. *Curr Opin Cardiol* 2010;25:299-301.
- [105] Wilensky RL, Shi Y, Mohler ER 3rd, Hamamdzic D, Burgert ME, Li J, Postle A, Fenning RS, Bollinger JG, Hoffman BE, Pelchovitz DJ, Yang J, Mirabile RC, Webb CL, Zhang L, Zhang P, Gelb MH, Walker MC, Zalewski A, Macphee CH. Inhibition of lipoprotein-associated phospholipase A2 reduces complex coronary atherosclerotic plaque development. *Nat Med* 2008;14:1059-1066.
- [106] Berger JS, Ballantyne CM, Davidson MH, Johnson JL, Tarka EA, Lawrence D, Trivedi T, Zalewski A, Mohler ER 3rd. Peripheral artery disease, biomarkers, and darapladib. *Am Heart J* 2011;161(5):972-978.

- [107] Johnson JL, Shi Y, Snipes R, Janmohamed S, Rolfe TE, Davis B, Postle A, Macphee CH. Effect of darapladib treatment on endarterectomy carotid plaque lipoprotein-associated phospholipase A2 activity: a randomized, controlled trial. *PLoS One* 2014;9(2):e89034.

INTECH

INTECH

