

Key Enzymes and Proteins in Amyloid-Beta Production and Clearance

Cecília R A Santos¹, Isabel Cardoso^{2,3} and Isabel Gonçalves¹

¹*University of Beira Interior- Health Sciences Research Centre, Covilhã*

²*Escola Superior de Tecnologia da Saúde do Porto, Instituto Politécnico do Porto, Vila Nova de Gaia,*

³*Molecular Neurobiology Unit, IBMC- Instituto de Biologia Molecular e Celular, Porto, Portugal*

1. Introduction

Last century, in 1907, Alois Alzheimer first described the disease that now bears his name, relating a 54-year-old female case with presenile dementia. Alzheimer's disease (AD), an irreversible progressive neurodegenerative disorder, is actually the most common form of dementia. AD affects more than 24 million people all over the world, and is predicted to double every 20 years, becoming one of the medical burdens of our days. More than 90% of AD cases are sporadic late-onset, and have a complex idiopathic aetiology. The small number of early onset AD cases, is related to hereditary monogenic defects, and has provided important clues for understanding the AD pathology (Bertram, et al. 2007). Although the available drugs are able to delay the symptoms and progression of the disease and to positively influence the quality of life of the patients, at present there is still no cure for AD.

AD is characterized clinically by progressive decline in cognitive function and neuropathologically by the presence of neuropil threads and neuron loss, in addition to the molecular hallmarks of neurofibrillar tangles and neuritic (or senile) plaques in the brain. Neuritic plaques are extracellular amyloid deposits found abundantly in the hippocampus, in the neocortex and in the amygdala of AD brains. These pathological brain changes may occur 20 to 30 years prior to the onset of the clinical symptoms and the symptomatic phase of AD can last from 5 to 12 years (DeKosky and Marek 2003). The extracellular neuritic plaque deposits of amyloid were first investigated by (Glennner and Wong 1984), when they purified microvascular amyloid deposits from AD brains, and provided a partial sequence of a 4kDa subunit protein, that they named amyloid-beta ($A\beta$) peptide. Around the same time, the hyperphosphorylated tau (p-tau), a microtubule assembly protein, was identified as the main constituent of the neurofibrillar tangles (NFTs) that accumulate inside many neurons in AD brains (Grundke-Iqbal, et al. 1986). Amyloid deposition and neurofibrillar tangles, occur with some frequency in brains of young adults with Down's syndrome (Schochet, et al. 1973). The discovery that amyloid deposits in the brain from Down syndrome were composed of $A\beta$ peptide (Glennner and Wong 1984), as well as the cloning of the beta amyloid precursor protein (APP), with its localization to the chromosome 21

(Korenberg, et al. 1989), led the scientists to search AD-causing mutations in the APP gene. Since that, several mutations associated with familial early onset forms of AD have been described, either in the APP gene (Kowalska 2003) or in preselin 1 (PS1) or preselin 2 (PS2) genes (Bertram et al. 2007). Either PS1 or PS2 can be the catalytic subunit of γ -secretase, which is the final endoprotease in the pathways that generate the A β peptide (see section 2). All these findings led to the amyloid cascade hypothesis, articulated by John Hardy and others (Hardy and Higgins 1992), in which the accumulation of A β peptide, generated from the proteolytic cleavage of APP in the brain, could trigger a complex downstream cascade that results in the symptoms of AD. This hypothesis states that gradual accumulation and aggregation of the hydrophobic A β peptide initiates a cascade that leads to synaptic alterations, astrocytic and microglial activation, the modification of the soluble tau protein into insoluble paired helical filaments, and progressive neuronal loss associated with multiple neurotransmitter deficiencies and cognitive failure (Hardy and Selkoe 2002). The cascade hypothesis suggests that stopping or slowing formation of the A β plaques would delay the onset of the disease symptoms. A β is found in the extracellular fluids of the brain, including cerebrospinal fluid (CSF), and in the interstitial fluid surrounding neurons and glial cells in brain lobes (Seubert, et al. 1992; Vigo-Pelfrey, et al. 1993). Over the last years, several key proteins have been described as being implicated in A β production and clearance, but further elucidation of the mechanisms involved in the process will be important for identifying new potential therapies to reduce A β accumulation and combat AD. This book chapter reviews the production of A β from APP and the proteins involved in its degradation and clearance .

2. Generation of amyloid beta peptides

The β amyloid precursor protein, APP, takes a central position in AD pathogenesis, as it is processed by the sequential action of β - and γ -secretase, generating the A β peptide, which is deposited as amyloid plaques in brains of AD individuals. APP is an integral membrane protein, with a large N-terminal extracellular domain and a short C-terminal cytoplasmatic domain, which is expressed ubiquitously in neuronal and non-neuronal cells.

The human APP gene is located on chromosome 21 (Korenberg et al. 1989) and alternative splicing results in protein isoforms of various lengths: two isoforms predominant in non-neuronal tissues (751- and 770-), and the 695-amino acid form, that is the predominant isoform in neurons (Kang and Muller-Hill 1990). APP belongs to a protein family that includes APP-like protein 1 (APLP1) and 2 (APLP2) (Eggert, et al. 2004), a group of type-I transmembrane proteins that are processed in the same fashion. APP is hydrolyzed into different fragments (Figure 1) during its intracellular trafficking, and these metabolites mediate various functions (Haass 2004; Haass and Selkoe 1993). APP is first cleaved by either α - or β -secretase at the α - or β -sites, respectively, which lie in the extracellular domain of the APP. These proteases compete for APP, originating: soluble APP α (sAPP α , for α -secretase) or soluble APP β (sAPP β , for β -secretase), which are released to the extracellular space, and a membrane anchored C-terminal end (C83 for α -secretase or C99 for β -secretase). Subsequently, in the lipid bilayer, γ -secretase acts in the C-terminal end, C83 or C99. The γ -cleavage of C83 generates the APP intracellular domain (AICD), with 6kDa, and the N-terminal peptide with 3kDa (p3) into the extracellular space. γ -cleavage of C99, in a specific sequence (A β domain) generates A β peptide and the AICD. This pathway of APP processing by β -secretase followed by γ -secretase leading to A β peptide is called the

amyloidogenic pathway. Aberrant and/or cumulative A β production, have been postulated to be the main etiological basis of AD. The alternative pathway of APP processing by α -secretase followed by γ -secretase, in which no A β is formed, is termed nonamyloidogenic pathway.




Fig. 1. Schematic diagram of APP processing (not drawn in scale). TM-transmembrane; EC- extracellular; IC- intracellular

A β is a ~4kDa peptide with 38 to 43 amino acids, depending on the site of γ -secretase cleavage. A β peptide is normally produced by cells and under physiological conditions there are two major species: A β 40 and A β 42. The major species produced is A β 40 and corresponds to 90% of the total A β peptide. The minor species produced, A β 42, is more prone to aggregation due to two additional hydrophobic amino acids, and therefore it is the predominant species accumulated in AD brain plaques.

A number of proteins influence the subcellular trafficking itinerary of APP and β -secretase between the cell surface, endosomes and the Trans-Golgi-Network (TGN). APP is synthesized in the endoplasmic reticulum (ER) and is transported through the Golgi apparatus to the TGN, where the highest concentration is found in steady state neurons (Greenfield, et al. 1999). From TGN, the APP can be transported, in TGN-derived vesicles, to cell surface where it is either cleaved by α -secretase to produce sAPP α , or internalized via the endosomal-lysosomal pathway (Caporaso, et al. 1994). The A β peptide is generated either in the ER and TGN (Greenfield et al. 1999), as in the endosomal/lysosomal system (Haass and Selkoe 1993). Available evidence suggests that co-residence of APP and β -secretase in the endosomal compartments promotes amyloidogenesis. Retrograde transport of APP, out of the endosome to the TGN, reduces A β production, while APP routed to, and kept at the cell surface, enhances its non-amyloidogenic processing (Tang 2009).

Very little is known about the physiological function of APP and its proteolytic products. APP knockout mice (KO) are viable and fertile, showing a slight abnormal phenotype

(Dawson, et al. 1999). APLP1 and APLP2 KO mice are also viable and fertile, but APP/APLP2 and APLP1/APLP2 double null mice and APP/APLP1/APLP2 triple null mice show early postnatal lethality (Heber, et al. 2000; von Koch, et al. 1997). APP/APLP1 double null mice are viable, suggesting redundant functions of amyloid precursor protein family members (Heber et al. 2000). Putative suggested roles for APP include trafficking, neurotrophic signalling, control of cell adhesion, neuritic outgrowth and synaptogenesis, apoptosis and transcription regulation (Zheng and Koo 2006). As APP is proteolysed in the cell, the net effect of full-length APP on cellular activity may be a combination of the function of its proteolytic products, depending on the proportion levels of each of them.

2.1 Alpha-secretase

APP is cleaved by α -secretase in the center of A β domain, precluding A β peptide generation, and a soluble domain of APP is released: sAPP α . Three related proteases, all from the ADAM family, had been suggested to exert the α -secretase activity: ADAM-9, ADAM-10 and ADAM-17 (Asai, et al. 2003). Like full length APP, members of the ADAM family (a desintegrin and metalloprotease family) are type-I transmembrane proteins, possessing both potential adhesion and protease domains. Several studies suggested α -secretase activity for ADAM-9. However, as RNAi of ADAM-9 has no effect in sAPP α generation (Kuhn, et al. 2010), ADAM-9 seems to be involved only in regulated α -cleavage and not in constitutive α -secretase activity. Different studies strongly suggest that ADAM-10 is the main constitutive α -secretase that is active at the cell surface (Lammich, et al. 1999) (Jorissen, et al. 2010), but there may exist some functional redundancy in α -cleavage by the ADAM protease family. ADAM-17 is an 824 amino acid polypeptide containing a secretory signal sequence, a desintegrin domain and a metalloprotease domain, that also seems to be involved in regulated α -secretase activity (Merlos-Suarez, et al. 2001).

sAPP α has important roles in neuronal plasticity/survival, stem cell proliferation in CNS, and is able to rescue the abnormalities of APP deficient mice, indicating that most of APP's physiological function is mediated by sAPP α (Zhang, et al. 2011).

2.2 Beta-secretases: BACEs

β -secretase cleavage is the first critical step in the APP amyloidogenic pathway, and increased β -secretase activity levels have been correlated with brain A β deposition in late onset AD patients (Li, et al. 2004). A β peptide is generated from APP by a sequential two-step proteolytic process involving β - and γ -secretases (Haass 2004), being BACE 1 (β -site APP cleaving enzyme 1) the major β -secretase in the cell (Vassar, et al. 1999). BACE1 is a member of the pepsin family of aspartyl proteases, and its activity on APP generates the membrane bound C-terminal fragment (CTF β or C99). BACE1 requires acidic environment for optimal activity and cleaves APP at the known β -site locations, Asp1 and Glu11. Overexpression of BACE1 induces cleavage of APP at β -sites and is mainly found in the early Golgi, late Golgi/early endosomes and endosomes with acidic environment. BACE1 is also found at the cell surface (Huse, et al. 2002; Vassar et al. 1999).

BACE2 is an additional β -secretase, mapped in 21q22.3 region (Solans, et al. 2000), that also cleaves β -secretase substrates. However, BACE2 expression in neurons is lower than BACE1 (Bennett, et al. 2000). BACE1 null mice died in the first weeks and those that survived, were smaller, presented hyperactive behaviour, were affected by hypomyelination of peripheral nerves and had altered neurological behaviours such as elevated pain sensitivity

(Dominguez, et al. 2005). BACE2 KO mice are healthy overall, while a deficiency of both BACE1 and BACE2 enhanced the BACE1 KO lethality phenotype, suggesting functional redundancy (Dominguez et al. 2005).

Cathepsin B has been proposed as another β -secretase. Although its inhibition has been found to reduce $A\beta$ production, its physiological activity is not well established (Hook, et al. 2009).

2.3 The gamma secretase complex

γ -secretase is a protein complex, of high molecular weight, responsible for the membrane cleavage of the APP C-terminal remnants after cleavage by either α - or β -secretase (C83 and C99, respectively). The cleavage of C83 and C99 by γ -secretase generates p3 and the $A\beta$ peptide, respectively. In addition to C83 and C99 peptides, several non-APP substrates are cleaved by γ -secretase and all of them are type-I transmembrane proteins that require ectodomain shedding as a prerequisite to γ -secretase cleavage (Haapasalo and Kovacs 2011). γ -secretase complex comprises four core components which include: presenilin (PS1 or PS2), nicastrin, anterior pharynx-defective-1 (APH-1), and presenilin enhancer-2 (PEN2) (Kimberly and Wolfe 2003; Takasugi, et al. 2003). All of these four core components of γ -secretase are necessary for the enzymatic activity of the complex. PS1 and PS2 are two presenilin homologs, and several mutations in the corresponding genes have been described as the major cause of familial AD cases (Bertram et al. 2007). Different kinds of experimental evidences suggest that presenilins are transmembrane proteins with crucial catalytic roles in γ -secretase activity (Zhang et al. 2011). Nicastrin, other component of γ -secretase, is a type-I transmembrane glycoprotein considered the scaffolding protein of the complex (Vassar and Citron 2000). Finally, the other two components of γ -secretase are: APH-1, that interacts with nicastrin to form a stable intermediate in the early assembly stages of the complex and PEN-2, that regulates PS endoproteolysis (Vassar and Citron 2000). Besides the four components of γ -secretase complex, some factors, playing a modulatory role, have been described. Example of that is CD147 that when down regulated increases $A\beta$ production and TMP21/p23 that regulates γ -cleavage (Zhang et al. 2011).

Experimental evidences support the idea that γ -secretase resides in ER, Golgi-TGN, endosomes and intermediate compartments, most of which (except TGN) do not correspond to the main localizations of APP (Cupers, et al. 2001).

3. Enzymes and peptides involved in amyloid-beta degradation and clearance

The clearance of $A\beta$ in the human central nervous system (CNS), is roughly 8% of total production per hour. It is controlled by $A\beta$ degradation in the brain and by its efflux from the CNS to the peripheral circulation through the blood-brain barrier or the blood-cerebrospinal fluid barrier (Zlokovic 2004). The high clearance of $A\beta$ is in part due to the presence of cryptidases in several cellular compartments and reduction of $A\beta$ degradation by these enzymes may be implicated in the progression of AD cases. $A\beta$ accumulation, and the concomitant formation of amyloid plaques observed in AD brains occurs in the extracellular space. It is also in this compartment, that $A\beta$ can be degraded by cell surface and/or secreted cryptidases, such as insulin degrading enzyme and neprylisin. The subcellular distribution of $A\beta$ degrading cryptidases is a strong indicator that $A\beta$ degradation can be controlled at multiple subcellular compartments, such as the mitochondria, ER, Golgi, endosomes and lysosomes (Malito, et al. 2008).

In the brain, A β metabolism is mainly regulated by the activity of neprilysin (NEP) and insulin-degrading enzyme (IDE), but presequence peptidase, endothelin converting enzyme (ECE), angiotensin-converting enzyme (ACE), the uPA/tPA- plasmin system and matrix metalloproteinases are also involved in the process (Table 1). In addition, some proteins as transthyretin (TTR), gelsolin, alpha-2-macroglobulin and apolipoprotein E do also play an important role in A β clearance and degradation. Some of them do actually have a direct function in A β catabolism such as TTR, plasmin, and gelsolin, which have the capacity to cleave the peptide, others because their interaction with A β enable its degradation or prevent its neurotoxicity as it is the case of alpha-2-macroglobulin and apolipoprotein E (Table 2).

Enzymes	Function	Brain distribution	References
<i>IDE</i>	Hydrolises peptide bonds of A β 40 and A β 42	Neurons, microglia, endothelial cells, choroid plexus epithelial cells	(Bora et al. 2010; Bora and Prabhakar 2010; Malito et al. 2008)
<i>NEP</i>	Hydrolises peptide bonds of A β 42	Neurons, microglia, endothelial cells, choroid plexus epithelial cells	(Malito et al. 2008; Meilandt et al. 2009)
<i>PreP</i>	Hydrolises peptide bonds of A β 40 and A β 42 in mitochondria		(Falkevall et al. 2006)
<i>ECE</i>	Hydrolises peptide bonds of A β 40 and A β 42	Neurons, astrocytes, endothelial cells	(Miners et al. 2008b)
<i>ACE</i>	Converts A β 42 to A β 40 Cleaves A β 40	Neurons, endothelial cells, choroid plexus epithelial cells	(Miners et al. 2008a; Zou et al. 2007)
<i>Plasmin</i>	Degrades aggregated and non-aggregated A β 40 and A β 42 Inhibits A β fibrillogenesis	Neurons, microglia	(Tucker et al. 2002; Tucker et al. 2000b)
<i>MMP-2</i>	Hydrolises peptide bonds of A β 40 and A β 42 Beta-secretase activity	Microglia, astrocytes, Schwann cells	(Miners et al. 2008a)
<i>MMP-9</i>	Hydrolises peptide bonds of A β 40 and A β 42 and aggregated A β fibrils	Neurons, endothelial cells	(Miners et al. 2008a; Yan et al. 2006)

Table 1. Functions and brain distribution of key enzymes in A β clearance.

Key players	Function	Brain distribution	References
<i>TTR</i>	Hydrolises peptide bonds of A β 40 and A β 42, and aggregated A β oligomers and fibrils	choroid plexus epithelial cells meninges	(Costa et al. 2008; Schwarzman et al. 1994)
<i>Gelsolin</i>	Inhibits fibrillization of A β and defibrillates preformed fibrils	Oligodendrocytes, microglia, choroid plexus epithelial cells	(Chauhan et al. 2008; Ray et al. 2000)
<i>Alpha-2 macroglobulin</i>	A β carrier, binds A β protecting it from proteolysis, reduces A β aggregation and fibril formation	astroglia	(Du et al. 1998; Du et al. 1997)
<i>Apo E</i>	Binds A β enhancing the proteolytic activity of NEP, and IDE	Astrocytes, neurons, microglia	(Jiang et al. 2008)
<i>MT-2</i>	Diminishes A β binding to TTR Prevents copper mediated aggregation of A β	Cortical neurons	(Chung et al. 2010; Martinho et al. 2010)
<i>MT-3</i>	Increases A β binding to TTR Inhibits formation of A β aggregates	Cortical neurons	(Irie and Keung, 2001, 2003; Martinho et al. 2010)

Table 2. Functions and brain distribution of key proteins in A β clearance.

3.1 IDE – Insulin degrading enzyme

Insulin degrading enzyme (IDE) is a ~ 110 kDa zinc-containing metalloendopeptidase that degrades monomeric forms of A β peptides and insulin, which has also high nanomolar affinity for other substrates with different sequences and structures. The common features of IDE substrates are that they are all amyloidogenic in nature. So, besides A β and insulin, IDE also cleaves, insulin-like growth factor 2, atrial natriuretic peptide, bradykinin, endorphin, and glucagon. However, *in vivo* relevance of this degrading activity has only been demonstrated for A β and insulin (Malito et al. 2008). Interestingly, patients with type 2 diabetes are under an increased risk of AD. In addition, A β is a direct competitive inhibitor of insulin binding and action, and this is a likely justification for the increased levels of A β observed in insulin resistant AD patients (Xie, et al. 2002). This dual effect of IDE stresses the importance of the development of inhibitors and activators of its activity for the treatment of diabetes and AD, respectively.

The first evidences of the capacity of IDE to degrade A β were reported by Kurochkin and Goto (1994) who identified a protein of 110,000 Da present in cytosol fractions from rat brain and liver that cross-linked to ¹²⁵I-labeled synthetic A β . A few years later IDE was actually identified as the main soluble A β degrading enzyme in human brain and in neuronal cell cultures, where its action takes places in the extracellular milieu (McDermott and Gibson 1997). Besides its ability to degrade A β , IDE activity is also associated with oligomerization of synthetic A β at physiological levels in the conditioned media of cultured cells (Qiu, et al. 1998). In addition, evidence that membrane-associated and secreted IDE isoforms carry out the degradation and clearance of A β secreted by neurons and microglia was provided a couple of years later (Bertram, et al. 2000).

IDE is composed of four homologous domains that share 15–24% sequence similarity. These domains form two functional N- and C-terminal domains that are joined by an extended 28 amino acid residue loop, creating a large catalytic chamber which can accommodate substrates of the order of 6 kDa. Substrate binding is assisted by the C-terminal domain and, its hydrolysis occurs at the N-terminal domain (Li, et al. 2006). The active site of IDE encompasses the His-Glu-aa-aa-His sequence and requires zinc. IDE enzyme hydrolyses several peptide bonds of both A β 40 and A β 42, but is particularly efficient at hydrolysing the Lys28-Gly29 peptide bond followed by the Phe19-Phe20 and His14-Gln15 bonds of these substrates (Bora, et al. 2010; Bora and Prabhakar 2010).

IDE is expressed by cortical and subcortical neurons, and has been detected in the cytoplasm of the three major components of the vascular wall: endothelial cells, pericytes and smooth muscle cells (Dorfman, et al. 2010; Gao, et al. 2004). IDE is also expressed towards the apical surface of the choroid plexus tissue where its inhibition leads to disrupted metabolism of A β and its concurrent accumulation at the blood-CSF barrier (Behl, et al. 2009). Besides the cytoplasm, IDE is found in endosomes, on the cell surface, and in the extracellular milieu. The type of cell in the nervous system expressing IDE establishes whether it is secreted or associated with the cell surface. Primary mouse microglia and the BV-2 cell line are found to secrete IDE, but in primary hippocampal neurons and differentiated PC12 cells only membrane associated IDE has been found. IDE is also present in mitochondria, and in the dendrites of neurons (Malito et al. 2008).

Evidence that IDE activity in AD brains is reduced compared to age-matched controls has been giving support to the hypothesis that reduced IDE activity may contribute to A β accumulation in the brain (Perez, et al. 2000). For example, membrane-bound IDE protein concentrations and activity decrease during the conversion from mild cognitive impairment (MCI) to mild-severe AD in the hippocampus, which correlates negatively with brain A β 42 content in MCI and in AD brain (Zhao, et al. 2007). Still, other studies indicate that IDE may be less important in the process of A β clearance (Wang, et al. 2010). There is also evidence that in transgenic mice brain, A β plaques induce cortical mRNA and protein levels of IDE in parallel with increased A β 40 and A β 42 production, suggesting a positive feedback regulatory mechanism of A β regulation (Vepsalainen, et al. 2008).

3.2 NEP – Neprilysin

Neprilysin (NEP) is a 90–110 kDa plasma membrane glycoprotein of the neutral zinc metalloendopeptidase family. It consists of a short N-terminal cytoplasmic tail, a single transmembrane helix and a large C-terminal extracellular ectodomain. The ectodomain of neprilysin is largely made of α -helices with six disulfide bridges and at least three N-

glycosylation sites. The ectodomain encompasses the catalytic site, with the conserved HEXxH motif necessary for zinc coordination and a proteolytic chamber (Malito et al. 2008). NEP is widely expressed in several tissues such as the brush-border of intestinal and kidney epithelial cells, neutrophils, thymocytes, lung, prostate, testes, and brain. In the brain, it is expressed on the plasma membranes of neurons, pre- and post-synaptically, and is most abundant in the nigrostriatal pathway, and in brain areas vulnerable to amyloid plaque deposition, such as the hippocampus (Wang, et al. 2006). It is also expressed in the tunica media and endothelium of cortical and leptomeningeal blood vessels where it participates in the regulation of the vascular tone, and in pyramidal neurons. NEP is also involved in the regulation of neuropeptide signalling (Dorfman et al. 2010; Miners, et al. 2008b).

NEP has been implicated in the degradation of A β . It degrades A β 42 *in vivo*, and when NEP is inhibited, A β degradation in rat hippocampus is blocked with a concomitant increase of A β 42 plaques in this brain region, and in the cortical region outside the hippocampus. In addition, the capacity to degrade A β 42 of exogenous origin is severely compromised in NEP knockout homozygote mice in which the levels of A β 40 and A β 42 remain very high. These studies are corroborated by correlative studies in patients with sporadic AD and healthy age matched controls. Overexpression of NEP in brains of human amyloid precursor protein (hAPP) transgenic mice decreases overall A β levels and amyloid plaque burdens by 50% and effectively prevented early A β deposition in the neocortex and hippocampus. However, it did not reduce levels of A β oligomers or improved deficits in spatial learning and memory. The differential effect of NEP on plaques and oligomers suggests that NEP-dependent degradation of A β affects plaques more than oligomers and that these structures may form through distinct assembly mechanisms (Meilandt, et al. 2009). Moreover, the expression of NEP on the surface of leukocytes, through lentivirus transplantation of bone marrow cells, reduced soluble brain A β levels by ~30% and lowered the accumulation of A β peptides by 50–60% when transplantation was performed at both young and early adult age. This peripheral NEP expression reduced amyloid dependent performance deficits of these animals in the Morris Water Maze (Guan, et al. 2009). *Ex-vivo* gene delivery of a soluble form of NEP, via fibroblasts, into transgenic APP mice also demonstrated increased clearance of plaques. Interestingly, mRNA and protein levels of NEP can be induced by intracranial injections of A β 42, which also reduced the accumulation of amyloid plaques (Malito et al. 2008). Microglia from old PS1-APP mice, but not from younger mice, have a twofold to fivefold decrease in expression of NEP, compared with their littermate controls (Hickman, et al. 2008). Despite all the evidence sustaining the proteolytic action of NEP on A β , the molecular basis of the interaction between NEP and A β remains largely unexplained because the volume of the proteolytic chamber of NEP is about half the size of A β (Malito et al. 2008).

3.3 Presequence peptidase

Presequence peptidase (PreP) is a 110 kDa metalloprotease, ubiquitously expressed, but with higher abundance in heart and skeletal muscle. It is responsible for degradation of targeting peptides, which have been cleaved off inside the mitochondrial matrix after protein import, but it also cleaves other unstructured peptides up to 65 amino acids (Stahl, et al. 2002). More recently, PreP was shown to completely degrade A β 40 and A β 42, which are present in mitochondria. More relevance is added to this action of PreP as it has been shown that hPreP is actually the only protease responsible for degradation of A β in the

mitochondria (Falkevall, et al. 2006) and A β -induced mitochondrial toxicity has been associated with AD (Tillemont, et al. 2011).

Human PreP consists of 4 domains, creating two halves connected by a hinge region. The two halves can create a large catalytic chamber of 10 000 Å where A β fits. The inverted zinc-binding motif is located in the N-terminal region, but also includes residues located in the C-terminal half, about 800 amino acids distant from the zinc-binding motif (Johnson, et al. 2006). Unlike IDE, PreP cannot degrade insulin, making PreP a better therapeutic agent candidate than IDE as its use would preclude deleterious side effects associated with the degradation of insulin. Another interesting feature of PreP is that its proteolytic activity against A β is abolished under oxidizing conditions, probably due to the formation of a disulphide bridge between Cys90 and Cys527 that inhibits the substrate from entering the catalytic chamber. These findings indicate a possible inhibition of hPreP under elevated ROS production in mitochondria implicated in AD, and might therefore be relevant in this disease (Alikhani, et al. 2009).

3.4 ECE – Endothelin converting enzyme

Endothelin-converting enzymes 1 and 2 (ECE-1 and ECE-2) were originally implicated in the processing of pro-hormone forms of endothelin. They are type II integral membrane zinc metalloendopeptidases that are primarily localized to the endothelium throughout the human vasculature. They share common catalytic substrates and are responsible for cleaving big endothelins to produce potent vasoconstrictor endothelins (Miners, et al. 2008a; Miners et al. 2008b).

ECE sequences and domain organization are similar to NEP's and are also capable of degrading A β *in vitro* and *in vivo* (Eckman, et al. 2001). Homozygous knockouts for ECE-1 are lethal and heterozygous animals show an increased amount of A β 40 and A β 42 in the brain. Homozygous ECE-2 knockout mice show a gene dose dependent increase of both forms of A β in the brain as well (Eckman, et al. 2006; Eckman, et al. 2003).

ECE-1 is present in neurons, specifically in pyramidal neurons of the hippocampus and layer V of the neocortex, and to a less extent in astrocytes (Sluck, et al. 1999). In mice, ECE-2 is largely confined to the brain; in the rat brain, ECE-1 occurs in the cerebrovascular endothelium, whereas ECE-2 is predominantly neuronal, with special incidence on hippocampal pyramidal neurons (Miners et al. 2008a; Miners et al. 2008b).

There is evidence of significant decrease in ECE-2 gene expression in AD patients (Weeraratna, et al. 2007). The contribution of ECE-1 to the accumulation of A β or reduction in local microvascular blood flow in AD seems to be detrimental, with abnormal production of ET-1 being more likely to reflect A β -mediated upregulation of ECE-2 (Palmer, et al. 2010).

3.5 ACE – Angiotensin-converting enzyme

Angiotensin-converting enzyme (ACE) is a membrane-bound zinc metalloprotease expressed in blood vessels throughout the body. ACE is extremely important for the regulation of fluid homeostasis and blood pressure. It converts angiotensin I to the potent vasoconstrictor angiotensin II. Within the brain, ACE has been detected in cortical pyramidal neurons and in the cerebral vasculature. Its highest levels occur in circumventricular organs, such as the subfornical organ, area postrema, and the median eminence but was detected in other areas such as the caudate nucleus, putamen, substantia

nigra pars reticularis, nucleus of the solitary tract, dorsal motor nucleus, median preoptic nucleus, and choroid plexus in rat, human, rabbit, sheep and monkey (Miners et al. 2008a).

ACE has two homologous domains, each having a functional active site. The N-domain of ACE is responsible for converting A β 42 to A β 40, whereas the angiotensin-converting activity is found predominantly in the C-domain of ACE. N-linked glycosylation is essential for both A β 42 to A β 40 conversion and angiotensin-converting activities and protects ACE from proteolysis (Zou, et al. 2009).

The first correlations between ACE and AD were found in 1985 by Zubenko and colleagues (Zubenko, et al. 1985) who showed that the CSF of patients with moderate degrees of senile dementia of the AD type exhibited about half the ACE activity of age and sex-matched control individuals, raising the possibility that ACE activity in the CSF could be an index of AD. In a posterior study, however, no differences in ACE activity in the CSF were found between AD patients and age matched controls raising some controversy on this issue (Konings, et al. 1993), but other studies reported reduced activity of ACE in the parietal cortex of AD brains (Ichai, et al. 1994).

ACE was found to significantly inhibit A β aggregation in 2001 (Hu, et al. 2001). The inhibition of aggregation was specifically blocked by preincubation of ACE with an ACE inhibitor, lisinopril. ACE degraded A β by cleaving A β 40 at the site Asp(7)-Ser(8). Compared with A β 40, aggregation and cytotoxic effects of the degradation products A β -(1-7) and A β -(8-40) peptides were reduced or virtually absent. These findings led to the hypothesis that previous associations of ACE genotype with AD susceptibility (Farrer, et al. 2000; Hu, et al. 1999; Isbir, et al. 2001; Yang, et al. 2000) could rely on its capacity to degrade A β and prevent the accumulation of amyloid plaques *in vivo*. Besides A β 40 degradation, cellular expression of ACE also promotes degradation of naturally secreted A β 42, and leads to the clearance of both species (Hemming and Selkoe 2005). In addition, ACE also converts A β 42 to A β 40 which is less neurotoxic (Zou, et al. 2007).

Pharmacological inhibition of ACE with a widely prescribed drug, captopril, promotes the accumulation of cell-derived A β in the media of APP expressing cells and questions if prescribed ACE inhibitors against hypertension could elevate cerebral A β levels in humans (Hemming and Selkoe 2005). To address this question the ACE inhibitor captopril was administered to two lines of APP transgenic mice presenting low or high levels of A β with associated plaque deposition. In both models, captopril did not affect cerebral A β levels nor plaque deposition or peripheral A β levels (Hemming, et al. 2007).

Epidemiological data obtained from a male population, however, suggests that angiotensin receptor blockers are more efficient in reducing the incidence and progression of AD and dementia compared with ACE inhibitors or other cardiovascular drugs (Li, et al. 2010). However, there are strong indications that ACE inhibitors may reduce the risk for and progression of dementias (Shah, et al. 2009). Still, these effects remain rather controversial, and prescription of ACE inhibitors against hypertension should be regarded with caution (Hajjar, et al. 2008; Kehoe and Wilcock 2007; Shah et al. 2009). Alternatively, ACE inhibitors could be designed to specifically target the angiotensin-converting C-domain, without inhibiting the A β 42 to A β 40-converting activity of ACE or increasing neurotoxic A β 42 (Zou et al. 2009).

3.6 uPA/tPA-plasmin system

Plasmin is a serine protease generated from the proteolytic cleavage of inactive plasminogen, by tissue-type (tPA) or by urokinase-type plasminogen activator (uPA). In the

brain, plasminogen is synthesized in neurons, whereas uPA and tPA expression can be induced in neurons as well as in microglia. This system is involved in many normal neural functions, such as neuronal plasticity, learning, and memory (Wang et al. 2006; Zhao and Pei 2008).

uPA and tPA are induced by A β treatment in primary rat embryonic cortical cultures *in vitro* as well as in a murine model of A β accumulation *in vivo* (Tucker, et al. 2000b). In addition, it plasmin degrades both non-aggregated and aggregated A β 40 (Tucker, et al. 2000a), and also degrades A β 42, preventing A β 42 aggregation into beta-pleated sheet structures (Exley and Korchazhkina 2001). Plasmin also inhibits A β toxicity, reduces A β deposition *in vitro*, and inhibits A β fibrillogenesis (Tucker, et al. 2002). The anti-amyloidogenic effect of plasmin is further enhanced by its capacity to increase the processing of human APP, and efficiently degrading secreted amyloidogenic and non-amyloidogenic APP fragments. Consistent with these observations, brains from AD patients contain reduced levels of plasmin (Ledesma, et al. 2000). Moreover, A β injected into the hippocampus of mice lacking either tPA or plasminogen persists causing activation of microglial cells and neuronal damage. Conversely, A β injected into wild-type mice is rapidly cleared and does not cause neuronal degeneration (Melchor, et al. 2003). Assembly of A β 42, on the other hand, seems to promote the up-regulation of the tPA/plasminogen proteolytic system, which can modulate the deposition of amyloid plaques *in vivo*, in a negative feedback mechanism (Lee, et al. 2007).

3.7 Matrix metalloproteinases

Matrix metalloproteinases (MMPs) are zinc- and calcium dependent endopeptidases, produced by neurons and glial cells. Metalloproteinase 2 (MMP-2) can be detected in the walls of some blood vessels and scattered white matter glia. Metalloproteinase 3 (MMP-3) is present in and around some neurons and within occasional amyloid plaques, and metalloproteinase 9 (MMP-9) is present in many neurons.

MMP-2 and MMP-9 all have A β -degrading activity *in vitro*, although the A β degrading activity of these MMPs has not received so much attention *in vivo*. However MMP-2 and MMP-9 activity are elevated in homogenates of hippocampal tissue from AD brains, and several cell types (glial, neuronal and vascular) up-regulate endogenous MMP-2 -3 and -9 expression in response to A β stimulation (Deb and Gottschall 1996; Miners et al. 2008b; Gottschall 1996).

MMP-2 also known as gelatinase A/type IV collagenase/matrix hydrolyses A β 40 and A β 42 peptides at Lys 16-Leu 17, at Leu 34-Met 35, and Met 35-Val 36 peptide bonds (Roher, et al. 1994). Besides its A β degrading activity, MMP-2 has also beta-secretase activity (LePage, et al. 1995). In AD brains, MMP-2 is present in the reactive microglia located in the center of senile plaques, and in Schwann cells (Yamada, et al. 1995). A β stimulates the expression and activation of MMP-2 to a large extent due to the increased expression of membrane type-1 (MT1)-MMP expression, the primary MMP-2 activator (Jung, et al. 2003).

There seems to be a complex regulation of MMP2 expression by oligomeric A β in astrocytes. Oligomeric A β directly down-regulates MMP2 expression and activation in astrocytes. However, it also induces the production of pro-inflammatory cytokines which stimulate production of MMP2 (Li, et al. 2011).

Increased MMP-9 immunolabelling has been detected in neurons of AD patients, as well as in neurofibrillary tangles, plaques and blood vessel walls, but not in granular neurons or glial cells. MMP-9 is also detected in the vicinity of extracellular amyloid plaques (Backstrom, et al. 1996). MMP-9, unlike ECE-1, NEP and IDE, is capable of cleaving aggregated A β fibrils (Miners et al. 2008b). It cleaves A β at several sites, predominantly at Leu34-Met35 within the membrane-spanning domain (Backstrom et al. 1996; Yan, et al. 2006). The presence of the apoE4 significantly dampens A β -induced MMP-9 in primary cultures of rat astrocytes. This effect may affect A β clearance and promote A β deposition in AD brains (Guo, et al. 2006). There is also evidence that reduction of A β levels through the action of MMP-9 may also result from the direct processing of cell surface APP with an alpha-secretase like activity, substantially reducing the levels of secreted A β peptide (Talamagas, et al. 2007).

Aged APP/presenilin 1 mouse astrocytes surrounding amyloid plaques, showed enhanced expression of MMP-2 and MMP-9. Astrocyte-conditioned medium obtained from these animals degraded A β , producing several small fragments. In the brains of MMP-2 and MMP-9 KO mice, significant increases in the A β levels were found in comparison to wild-type controls. This study reinforces previous *in vitro* evidences that MMP-2 and -9 may contribute to extracellular brain A β clearance through the degradation of A β (Yin, et al. 2006).

Several MMPs polymorphisms have been examined in relation to the risk of developing dementia with controversial results (Baig, et al. 2008; Helbecque, et al. 2007; Reitz, et al. 2008; Reitz, et al. 2010). MMPs may also be considered potential plasmatic and cerebrospinal fluid markers of AD as the levels of MMP-3 in the plasma and CSF of AD patients was found higher than in controls, whereas MMP-2 was significantly decreased in CSF but unchanged in plasma (Horstmann, et al. 2010).

3.8 Transthyretin

Transthyretin (TTR) is a 55 kDa homotetrameric protein secreted mainly by the liver and choroid plexus into the plasma and CSF, respectively (Soprano, et al. 1985). The name "transthyretin" discloses its dual physiological role as a carrier for thyroid hormones (Woeber and Ingbar 1968) and retinol, the latter through the binding to retinol-binding protein (Goodman 1985). TTR plasma concentration is age dependent and in healthy newborns it is about half of that in adults (Stabilini, et al. 1968; Vahlquist, et al. 1975), varying from 20 to 40 mg/ dL. In spite of the low TTR levels in CSF (~2 mg/ dL), the choroid plexus is presented as the major site of TTR expression, expressed as a ratio of tissue/mass, corresponding to a 30-fold higher than that found in plasma and represents 20% of the total CSF proteins (Weisner and Kauerz 1983).

Over a hundred TTR mutations have been associated with Familial Amyloid Polyneuropathy (FAP), some of which are very common in Portuguese patients, such as the valine at position 30 substituted by a methionine (TTR V30M) (Saraiva, et al. 1984), others provide a very aggressive phenotype, such as the TTR L55P (Jacobson, et al. 1992). Curiously, TTR T119M was described as an example of a non-aggressive mutation that inclusively has a protective role against the disease (Almeida, et al. 2000).

The first report that associates TTR to A β and AD as a protective molecule is from Schwarzman et al. who describes the capacity of normal CSF to inhibit amyloid formation and concluded that TTR was the major A β binding protein in the CSF, that could also

decrease the aggregation state of the peptide and its toxicity (Schwarzman, et al. 1994). Prior to this finding, TTR was found associated to senile plaques, NFTs and microangiopathic lesions (Shirahama, et al. 1982). The sequestration hypothesis was raised suggesting that normally produced A β is sequestered by certain extracellular proteins, thereby preventing amyloid formation and A β cytotoxicity; when sequestration fails amyloid formation occurs (Schwarzman and Goldgaber 1996). The observation that TTR is reduced in the CSF of AD patients further supported the idea of a TTR protective role in this pathology (Serot, et al. 1997). Authors also observed a decrease in TTR levels with age which could be related to the epithelial atrophy in the choroid plexus. Along time several reports described TTR decrease in CSF of AD patients (Davidsson, et al. 2002; Serot et al. 1997; Gloeckner, et al. 2008; Hansson, et al. 2004) although it remains unclear whether this reduction is restricted to AD, or on the contrary is common to other neurodegenerative disorders (Chiang, et al. 2009). It is also uncertain if the TTR decrease in the CSF happens early in disease development (or even before symptoms appears) or if it is a latter event, and thus studies involving patients with mild cognitive impairment (MCI) and early-staged probable AD patients are needed.

Other studies using a transgenic model of *Caenorhabditis elegans* expressing A β 42 supported a TTR role in AD as administration of TTR rescued the neurodegeneration observed in this model (Link 1995). Mammalian models used to recapitulate AD features were never completely successful as AD transgenic mice did not show NFTs and demonstrate little or no neuronal cell loss (Holcomb, et al. 1998; Hsiao, et al. 1996; Irizarry, et al. 1997a; Irizarry, et al. 1997b; Stein and Johnson 2002). However, in some of the models, animals showed increased TTR expression in the hippocampus (Stein and Johnson 2002); TTR was then described to be a survival gene (Stein and Johnson 2002) and although this work is controversial because TTR expression is thought to be confined to the choroid plexus and meninges (in the case of the brain), authors further showed that when a chronic infusion of an antibody against TTR was applied into the hippocampus of mice expressing human APP, an increase of A β , tau phosphorylation, neuronal loss and apoptosis were observed (Stein, et al. 2004). Underlying these observations is, according to authors, sAPP α that leads to increased expression of protective genes, such as TTR, to confer neuroprotection (Stein et al. 2004). Other studies, using transgenic APP mice hemizygous for endogenous TTR showed accelerated A β deposition (Choi, et al. 2007), while double transgenic mice for APP and TTR presented lower deposition (Buxbaum, et al. 2008). However, in other models, TTR was described to have the opposite effect and was associated with increased vascular A β deposition (Wati, et al. 2009).

Regarding the nature of TTR/A β interaction, different researchers confirmed TTR binding to A β (Carro, et al. 2002; Liu and Murphy 2006; Costa, et al. 2008) not only to the monomer but also to A β oligomers and fibrils, raising the hypothesis that TTR may be involved in the formation of senile plaques (Costa et al. 2008); TTR was also able to inhibit and to disrupt A β fibrils. However, which TTR conformation binds A β peptide is still controversial. Du Murphy et al. showed that TTR tetramers interact preferably with A β aggregates rather than A β monomers enhancing A β aggregation, whereas TTR monomers arrest A β aggregate growth (Du and Murphy 2010). On the other hand, studies by Costa et al. (Costa et al. 2008) showed that soluble A β binds to different TTR variants correlating negatively with the amyloidogenic potential of the TTR mutant. Thus, TTR119M presented the highest affinity to A β , contrarily to what was observed by Du and Murphy (Du and Murphy 2010). Other

studies also indicated a different relation between TTR variant/amyloidogenic potential and binding to the peptide, with amyloidogenic mutants binding less to A β peptide and inhibiting less its aggregation (Schwarzman, et al. 2004). In this work it was also shown that TTRs were functional tetramers. Hence, other studies are necessary to completely understand the nature of the TTR/A β interaction, concerning both A β and TTR species involved. Structural analysis, obtained from computer-assisted modeling (Schwarzman et al. 2004) predicted the existence of an A β binding domain on the surface of each TTR monomer; residues 30-60, especially the 38-42 region of TTR seemed to be the key structure of the A β binding domain (Schwarzman and Goldgaber 1996; Schwarzman, et al. 2005). Du and Murphy, identified the A strand, in the inner β -sheet of TTR, as well as the EF helix, as regions of TTR that are involved with A β association (Du and Murphy 2010).

The discussion on the TTR interaction with A β and consequent inhibition of aggregation and toxicity reduction raised the hypothesis that mutations in the TTR gene or conformational changes in the protein induced by aging, could affect the sequestration properties. A study was conducted with the aim of identifying mutations in the TTR gene in the AD population but no correlation was found (Palha, et al. 1996). Finally, and concerning the mechanism underlying TTR protective role in AD, Costa and colleagues found that TTR is able to proteolytically process A β peptide (Costa et al. 2008). Regarding TTR ability to degrade A β , several cleavage sites were identified and the newly generated A β peptides shown to have decreased amyloidogenic potential, when compared to the full length counterpart (Costa et al. 2008); TTR is also able to degrade aggregated forms of the peptide and inhibition of the TTR activity resulted in increased A β fibril formation (Costa et al. 2008).

3.9 Gelsolin

Gelsolin can be found both as an intrinsic cytoplasmic protein and as a secreted protein in plasma and CSF (Kwiatkowski, et al. 1985). Intracellular gelsolin regulates actin polymerization by binding to actin, and it also caps and breaks the actin filaments (Janmey, et al. 1985; Matsumoto, et al. 2003). Secreted (plasma/CSF) and intracellular gelsolin originate from the alternative splicing of a single gene, but their disulphide structure is different with recognised functional implications. All of the five cysteine (Cys) residues present in human cytoplasmic gelsolin are free thiols, whereas in plasma/CSF gelsolin, three Cys residues are free thiols and the other two Cys residues are disulfide-linked (Wen, et al. 1996). The five free thiol groups are likely to confer anti-oxidant properties to the molecule. Secreted gelsolin has an extension of 25 amino acids at its N-terminal, which is absent in the cytoplasmic form (Chauhan, et al. 2008).

Gelsolin found in the CSF may originate from oligodendrocytes or microglia (Chauhan et al. 2008). It was also suggested that choroid plexus may be responsible for the presence of gelsolin in CSF (Matsumoto et al. 2003), where it may have also an important function counteracting the neurotoxicity of A β (Antequera, et al. 2009; Vargas, et al. 2010).

Plasma and cytosolic gelsolin both bind A β and the A β -gelsolin complex exists in the plasma and in the cytosol (Chauhan, et al. 1999; Ji, et al. 2008). Both inhibit the fibrillation of A β , and defibrillate preformed fibrils (Ray, et al. 2000). In addition it has been demonstrated that when administered to transgenic mouse models of AD reduces the A β 40/A β 42 and the amyloid load (Hirko, et al. 2007; Matsuoka, et al. 2003). Therefore, gelsolin may also be looked as a potential therapeutic agent against AD.

3.10 Alpha2-macroglobulin

Alpha2-macroglobulin (a2M) is a 720 kDa soluble glycoprotein composed of four identical 180 kDa subunits, each encoded by a single-copy gene on human chromosome 12. Each subunit contains at least five binding sites: the bait region, the internal thiol ester, the receptor binding site, the A β binding site, and the zinc binding site. The bait region, the internal thiol ester and the receptor binding site are crucial for the activation and internalization of a2M. The bait region binds any known protease. The four bait regions in the tetramer are in close contact and get cleaved by the bound proteases, triggering activation of a2M. This conformational change results in the exposure of the four thiol esters, and of the four receptor binding sites, to the extracellular environment. The receptor binding site of each monomer, a 27 residue consensus sequence located at their C-terminal tail, will then bind to the receptor and mediate the internalization of the complex. The A β binding site is located between the bait and the receptor binding regions of each monomer (Borth 1992; Du, et al. 1997; Hughes, et al. 1998; Kovacs 2000).

Binding to A β 42 occurs with high affinity ($K_d = 3.8 \times 10^{-10}$ M) and protects the peptide from proteolysis by exogenous trypsin, suggesting that a2M may function as a carrier protein for A β and may serve to regulate clearance of A β from such tissues as the brain (Du et al. 1997; Hughes et al. 1998). a2M co-incubated with A β significantly reduces aggregation and fibril formation *in vitro*, and cultured fetal rat cortical neurons are less vulnerable to the toxic actions of A β following pretreatment with a2M, being likely that a2M has the capacity to keep A β in a soluble state, preventing fibril formation and associated neurotoxicity (Du, et al. 1998). a2M also inhibits both A β fibril formation and A β -induced cytotoxicity in PC12 cells (Monji, et al. 2000). The inhibition of the formation of amyloid fibrils are probably due to the interaction of a2M with prefibrillar species to maintain the solubility of A β (Yerbury, et al. 2009), and therefore their aggregation state in the extracellular milieu. In addition, Qiu et al described a serine protease that binds to a2M to form a stable high molecular weight complex capable of efficiently cleaving A β (Qiu, et al. 1996). This serine protease-a2M complex was capable of proteolytically digesting both A β 40 and A β 42, resulting in disruption of the central region of the peptide (residues 10–35), which is believed to mediate the conformational change that underlies A β self-aggregation (Qiu et al. 1996).

a2M is also a physiological ligand for the low-density lipoprotein receptor-related protein (LRP) abundantly expressed in the CNS. The a2M/A β complexes can be degraded by glioblastoma cells and fibroblasts via LRP, but the degradation of free A β must be mediated via an LRP-independent pathway. These results suggest that LRP can function as a clearance receptor for A β via a2M (Narita, et al. 1997). The effect of self-aggregation and LRP-1 ligands on the elimination of human A β 40 from the rat brain across the blood-brain barrier has been investigated recently. In the first instance it was demonstrated that the elimination rate of 125 I hA β 40 dimer was 92.7% decreased compared to the 125 I hA β 40 monomer. When pre-incubated with LRP-1 ligands, such as activated a2M, apolipoprotein E2 (apoE2), apoE3, apoE4, and lactoferrin, the elimination of 125 I hA β 40 was reduced. There seems that dimerization and LRP-1-ligand complex formation prevents the elimination of A β 40 from the brain across the blood-brain barrier (Ito, et al. 2007).

There is also an over-representation of a common a2M polymorphism, Val1000 (GTC)/Ile1000 (ATC) in AD patients, which correlates with an increase in A β burden (Kovacs 2000; Liao, et al. 1998), further sustaining the importance of a2M in A β clearance.

3.11 Apolipoprotein E

From all the proteins and peptides associated with AD and amyloid beta metabolism, apolipoprotein E (ApoE) is probably the most thoroughly studied, and several recent reviews give excellent and comprehensive overviews about its structure and function (Zhong and Weisgraber 2009a, b), and the pathways in which it is involved in AD (Kim, et al. 2009), either A β dependent or A β independent (Huang 2010; Mahley, et al. 2006). Therefore, this section will just give a brief overview about ApoE and will focus only on its association with AD via A β dependent pathways.

ApoE is a 34,2 kDa glycoprotein containing 299 amino acids. The protein contains two structural domains that are responsible for different functions of apoE. The amino-terminal domain (residues 1-191) contains the lipoprotein receptor binding region (residues 136-150), and the carboxyl-terminal domain (residues 216-299) contains the major lipoprotein (lipid) binding domain.

There are three common isoforms of ApoE, ApoE2, ApoE3 and ApoE4, which are encoded by three alleles (e2, e3 and e4, respectively) of a single gene. Sequence differences among the ApoE isoforms reside only on amino acids 122 or 158 which may be cysteine or arginine, but are sufficient to determine significant functional consequences (Huang 2010; Mahley et al. 2006).

ApoE4 is the major known genetic risk factor for AD and as much as 65-80% of all AD patients are ApoE4 carriers. This allele is over-represented in late-onset familial AD in several populations and in late-onset sporadic AD (Farrer, et al. 1997). In addition, as the number of Apo E alleles increases, the risk of onset of AD also increases while the age of onset decreases; as the number of ApoE4 alleles increases from 0 to 2, the risk of developing late-onset AD increases from 20% to 90%, and the mean age of onset decreases from 84 to 68 years (Corder, et al. 1993; Frangione, et al. 1996). The N and C terminal domains of ApoE interact in ApoE4 (called domain interaction), which is mediated by the formation of a salt bridge between Arg-61 in the N- terminus and Glu-255 in the C- terminus of ApoE4. This interaction might be the molecular basis for the detrimental effects of ApoE4 in AD pathogenesis (Huang 2010).

ApoE is expressed in several organs but is mainly expressed in the liver, followed by the brain. Non-neuronal cells, mainly astrocytes and to some extent microglia, are the major cell types that express apoE in the brain (Boyles, et al. 1985; Pitas, et al. 1987). Neurons also produce ApoE under certain conditions, particularly in response to brain injury (Boschert, et al. 1999).

ApoE (E2 and E3) are important for the distribution of lipids among cells throughout the body and within the CNS, where the principal apolipoproteins are, and where they transport lipids and cholesterol for cell repair and neurite outgrowth. These properties are not shared by Apo E4 which seems to counteract the features of Apo E2 and ApoE3 with detrimental effects (Mahley et al. 2006).

ApoE is also essential for astrocytes to bind, internalize and degrade A β deposits (Koistinaho, et al. 2004). ApoE binds A β with high affinity. It was demonstrated that lipid-free ApoE4 and ApoE3 form stable complexes with A β , with ApoE4 being more effective in the formation of the complex, but inducing a pathological β -sheet conformational change in A β ((Wisniewski and Frangione 1996). This interaction requires the N-terminal and the C-terminal domains, and there are strong indications that the interaction is affected by the lipid content (Weisgraber and Mahley 1996).

ApoE, but not ApoE4 plays a role in facilitating the proteolytic clearance of soluble A β from the brain. The endolytic degradation of A β peptides within microglia by NEP and related enzymes is dramatically enhanced by ApoE. Similarly, A β degradation extracellularly by IDE is facilitated by ApoE. The capacity of ApoE to promote A β degradation is dependent upon the ApoE isoform and its lipidation status. The enhanced expression of lipidated ApoE, through the activation of liver X receptors, stimulates A β degradation (Jiang, et al. 2008).

In contrast, Apo E4 inhibits A β clearance and stimulates A β deposition (Huang, et al. 2004). It also enhances A β production and potentiates A β induced chromosomal leakage and apoptosis (Mahley and Huang 2006). In addition, ApoE4 enhances A β 42 oligomer induced toxicity (Manelli, et al. 2007). Taken together these observations sustain the effects of ApoE4 in the enhancement of the overall A β burden via interaction with the peptide.

3.12 Metallothioneins 2 and 3

Metallothioneins (MTs) were discovered as cadmium binding proteins in equine kidney cortex. MTs is a generic name for a superfamily of low molecular weight cysteine- and metal-rich proteins with high affinity for divalent metals, such as zinc, cadmium and copper with four major isoforms, MT-1 to MT-4, identified in humans. They occur in all living organisms from the simplest forms of life, such as prokaryotes to the most complex, such as higher plants and vertebrate animals. It is clear that MTs are multipurpose proteins with unquestionable metal binding and anti-oxidant properties. In addition, there is increasing evidence that MT-1 and MT-2 (MT-1/2), and MT-3 display such diverse physiological actions as inhibition of pro-apoptotic mechanisms, enhancement of cell survival, tissue regeneration, and have anti-inflammatory properties. Concurrent with this wide array of functions, MT-1/2 have been implicated in neuroprotection and neurodegeneration, and particularly in AD (Hidalgo, et al. 2009; Penkowa, et al. 2006).

It is interesting that a considerable body of work has related an increase of MT-1/2 brain levels with aging and AD. In AD patients elevated levels of cytokines and IL-1 may induce MT-1/2 production in astrocytes. Studies in animal models of AD, showed that the MT-1/2 levels were higher, when compared to WT mice, while MT-3 levels were unaltered or reduced, suggesting that these proteins may have a relevant role in providing long term protection against inflammation (Hidalgo et al. 2009). This up-regulation of MT-1/2 in animal models of AD may have detrimental consequences in A β clearance as MT-2 diminishes the binding of TTR to A β (Martinho, et al. 2010). Considering TTR an A β scavenger as explained above, a less efficient removal of A β would be expected when MT-2 levels are increased, and this appears to be the case in AD. Furthermore, inhibition of homeodomain interacting protein kinase 2 activity in AD patients, might be involved in p53 misfolding, most likely through MT-2A upregulation (Lanni, et al. 2010). In addition, it was also shown that MT-2A may also prevent copper-mediated A β aggregation and neurotoxicity, by a mechanism which primarily involves a specific metal exchange interaction between Zn7.MT-2A and Cu(2).A β , and subsequent inhibition of H₂O₂ generation (Chung, et al. 2010). Furthermore, a metal swap between Zn7.MT-3 and soluble and aggregated A β 1-40-Cu(II) abolishes ROS production and related cellular toxicity (Meloni, et al. 2008).

These studies relating metal exchange and A β aggregation, conducted to a recent interest in the use of metal-chelation drugs as a potential therapy for AD. For example, the

administration of the copper and zinc chelating drug, clioquinol, prevents plaque formation in transgenic AD mice (Hegde, et al. 2009).

Since the discovery of MT-3 as a growth inhibitory factor, with reduced expression levels in AD brains compared to age-matched controls, several studies provide evidence that MT-3 is related to the aetiology of AD. Some studies indicate that MT-3 may potentially promote the clearance of A β plaques, while others show an opposite trend or even no differences (Howells, et al. 2010). MT-3 alone antagonizes the toxic effect of A β because it inhibits the formation of SDS-resistant A β aggregates, thereby protecting cortical neurons from its toxic effects. Both the full-length and the N-terminal domain of MT-3 promote neuron survival at low concentrations but inhibited it at high concentrations. These observations suggest that the anti- A β activity of MT-3 is different from its neuronal growth inhibitory activity. Other possible mechanisms underlying the protection of MT-3 from A β toxicity may be related to its interaction with TTR by improving its A β degrading capacity (Martinho et al. 2010).

4. Conclusion

The extracellular neuritic plaque deposits of amyloid found in AD brains contain A β , and according to the amyloid cascade hypothesis, the accumulation of A β peptide, can trigger gradual synaptic alterations, astrocytic and microglial activation, and modification of the soluble tau protein into insoluble paired helical filaments, with progressive neuronal loss and cognitive failure (Hardy and Selkoe 2002). The cascade hypothesis suggests that stopping or slowing formation of the A β plaques would delay the onset of the disease symptoms. Therefore, it is crucial to thoroughly elucidate the regulation of A β production and clearance to design new and effective therapies against AD.

APP, takes a central position in AD pathogenesis, as A β arises from its proteolytic cleavage through the sequential action of β - and γ -secretase. APP is first cleaved by either α - or β -secretase at the α - or β -sites, respectively, which lie in the extracellular domain of APP. These proteases originate: soluble APP α (sAPP α , for α -secretase) or soluble APP β (sAPP β , for β -secretase), which are released to the extracellular space, and a membrane anchored C-terminal end (C83 for α -secretase or C99 for β -secretase). γ -cleavage of C99, generates the A β peptide. This pathway is known as the amyloidogenic pathway. Misregulation of all the intervenients in this process may lead to A β accumulation. If APP cleavage by β -secretase or γ -cleavage of C99 is more effective than α -secretase, then the all process shifts into the accumulation of A β . Also if APP synthesis is enhanced, as it is the case of several mutated forms of APP associated with early onset AD, the production of A β is also increased with similar results (Ertekin-Taner 2007). So, alterations in APP synthesis and processing by both secretases, or downstream in the γ -secretase complex action, may account for more severe forms of disease. Another good example of that are the mutations associated with presenilins 1 and 2, which are part of the γ -secretase complex, known to be responsible for early onset inherited forms of AD (Ertekin-Taner 2007).

Once A β has been produced, other important players come into action (Figure 2). These are all the enzymes and proteins that, either by directly cleaving A β into less harmful peptides or making them more prone to proteolysis or less susceptible to aggregation through protein-protein interactions, will reduce the A β load in vulnerable areas of the brain.

Reduced degradation of A β by proteases is generally accepted to enhance plaque pathology in AD brains, and may depend on several factors such as decreased mRNA expression or decreased protein levels or activity, either in the brain cortex, hippocampus or in brain

microvessels; and post-translational modifications, such as oxidation and deposition of the enzymes in the diseased brain with consequent loss of its native structure and functionality (Dorfman et al, 2010).

IDE and NEP are well accepted as A β degrading enzymes, but although few studies compare the relative contribution of each of these enzymes to the overall A β load, NEP seems to be the major protease involved in A β degradation. In transgenic mice models of familial AD, over-expression of IDE or NEP prevents amyloid plaque pathology and consequent early death (Leissring, et al. 2003) indicating that degradation of A β by these metalloproteinases may be of high therapeutic interest for AD patients. A recent study carried out in AD patients showed that, NEP mRNA, protein levels, and activity are decreased compared to normal controls without any cognitive impairment. In these patients IDE activity was unchanged, though mRNA levels increased. In the same study ECE-1 expression or activity in AD brains was not different from age-matched controls. Correlation analyses suggested that NEP expression was correlated with A β accumulation and clinical diagnosis, being lower in AD than in controls, whereas no correlations of IDE and ECE-1 with A β levels or clinical diagnosis has been found. These findings provide additional support for NEP as the major protease involved in A β degradation (Wang et al. 2010).

Age is the major risk factor for AD. A β -synthesizing enzyme activities increase with age, coinciding with declining soluble A β and increasing insoluble A β (Miners, et al, 2008b). In addition, there is an overall ageing-related down regulation of A β degrading proteases which is particularly relevant for NEP but not for IDE in transgenic Tg2576 mice brains (Dorfman et al., 2010). Nevertheless further studies assessing the relative contribution of each of these enzymes, IDE, NEP, Prep, ECE, ACE and plasmin are necessary to understand why the system redundancy is not always effective, and which of these enzymes may be more adequate as candidate therapeutic targets.

Regarding proteins with the capacity to degrade A β or of interfering in the process, TTR is probably the most promising, with potential as a biomarker of the disease and even as a putative therapeutic agent. The literature shows that TTR plays an important role in the modulation of A β aggregation and toxicity. The use of TTR as a sera biomarker for diagnosis purposes in AD patients should also be explored, with obvious advantages, for both patients and researchers, and there are already indications that sera TTR is decreased in AD patients when compared to age-matched controls (Han et al., 2011). Moreover, it is important to investigate the factors that affect TTR/A-Beta binding and/or TTR alterations that lead to its decreased in AD, such as protein oxidation (Biroccio et al., 2006). Further studies are necessary to unravel the mechanism underlying TTR protective role in AD, to establish if TTR decline is a cause or consequence of disease, and the cellular pathways involved.

Other proteins as gelsolin, a2M and Apo E counteract A β deposition through interactions with the peptide, but apparently none of them seem to have the capacity to cleave the peptide. In general these proteins bind A β and the effects of these interactions translate into preventing A β fibrillation, and eventually on disaggregation of pre-existent fibrils (gelsolin and a2M); binding of enzymes conferring them the capacity to degrade A β (a2M) or may enhance the capacity of A β proteases, as NEP and IDE to cleave A β (ApoE). MT 1/2 and MT3 bind TTR affecting its capacity to cleave A β and prevent metal associated A β aggregation or inhibit their formation.

The scientific achievements acquired over the past decade on all the pathways and key players involved in the amyloid cascade will hopefully contribute to the development of more adequate therapies against AD in a closer future.




Fig. 2. Schematic diagram depicting APP processing: key proteins in amyloid-beta production and clearance (not drawn in scale). IDE- Insulin degrading enzyme; NEP- Nephylisin; PreP- Presequence peptidase; ECE -Endothelin-converting enzyme; ACE- Angiotensin-converting enzyme; MMP- Matrix metalloproteinase; TTR-Transthyretin; a2M- Alpha2-macroglobulin; ApoE- apolipoprotein E.

5. Acknowledgments

Portuguese Foundation for Science and Technology (FCT) research grants PTDC/SAU-NMC/114800/2009 and POCI/SAU-NEU/64593/2006, by COMPETE (Programa operacional Temático Factores de Competitividade) and POCI 2010 (Programa Operacional Ciência e Inovação 2010), respectively, with the participation of the European Communitarian Fund FEDER

6. References

- Alikhani N, Ankarcona M & Glaser E 2009 Mitochondria and Alzheimer's disease: amyloid-beta peptide uptake and degradation by the presequence protease, hPreP. *J Bioenerg Biomembr* 41 447-451.
- Almeida MR, Alves IL, Terazaki H, Ando Y & Saraiva MJ 2000 Comparative studies of two transthyretin variants with protective effects on familial amyloidotic

- polyneuropathy: TTR R104H and TTR T119M. *Biochem Biophys Res Commun* 270 1024-1028.
- Antequera D, Vargas T, Ugalde C, Spuch C, Molina JA, Ferrer I, Bermejo-Pareja F & Carro E 2009 Cytoplasmic gelsolin increases mitochondrial activity and reduces Abeta burden in a mouse model of Alzheimer's disease. *Neurobiol Dis* 36 42-50.
- Asai M, Hattori C, Szabo B, Sasagawa N, Maruyama K, Tanuma S & Ishiura S 2003 Putative function of ADAM9, ADAM10, and ADAM17 as APP alpha-secretase. *Biochem Biophys Res Commun* 301 231-235.
- Backstrom JR, Lim GP, Cullen MJ & Tokes ZA 1996 Matrix metalloproteinase-9 (MMP-9) is synthesized in neurons of the human hippocampus and is capable of degrading the amyloid-beta peptide (1-40). *J Neurosci* 16 7910-7919.
- Baig S, Kehoe PG & Love S 2008 MMP-2, -3 and -9 levels and activity are not related to Abeta load in the frontal cortex in Alzheimer's disease. *Neuropathol Appl Neurobiol* 34 205-215.
- Behl M, Zhang Y & Zheng W 2009 Involvement of insulin-degrading enzyme in the clearance of beta-amyloid at the blood-CSF barrier: Consequences of lead exposure. *Cerebrospinal Fluid Res* 6 11.
- Bennett BD, Babu-Khan S, Loeloff R, Louis JC, Curran E, Citron M & Vassar R 2000 Expression analysis of BACE2 in brain and peripheral tissues. *J Biol Chem* 275 20647-20651.
- Bertram L, Blacker D, Mullin K, Keeney D, Jones J, Basu S, Yhu S, McInnis MG, Go RC, Vekrellis K, et al. 2000 Evidence for genetic linkage of Alzheimer's disease to chromosome 10q. *Science* 290 2302-2303.
- Bertram L, McQueen MB, Mullin K, Blacker D & Tanzi RE 2007 Systematic meta-analyses of Alzheimer disease genetic association studies: the AlzGene database. *Nat Genet* 39 17-23.
- Biroccio A, Del Boccio P, Panella M, Bernardini S, Di Ilio C, Gambi D, Stanzione P, Sacchetta P, Bernardi G, Martorana A, Federici G, Stefani A, Urbani A. Differential post-translational modifications of transthyretin in Alzheimer's disease: a study of the cerebral spinal fluid. *Proteomics*. 2006 Apr;6(7):2305-13.
- Biroccio A, Del Boccio P, Panella M, Bernardini S, Di Ilio C, Gambi D, Stanzione P, Sacchetta P, Bernardi G, Martorana A, Federici G, Stefani A, Urbani A. Differential post-translational modifications of transthyretin in Alzheimer's disease: a study of the cerebral spinal fluid. *Proteomics*. 2006 Apr;6(7):2305-13. Bora RP, Ozbil M & Prabhakar R 2010 Elucidation of insulin degrading enzyme catalyzed site specific hydrolytic cleavage of amyloid beta peptide: a comparative density functional theory study. *J Biol Inorg Chem* 15 485-495.
- Bora RP & Prabhakar R 2010 Elucidation of interactions of Alzheimer amyloid beta peptides (Abeta40 and Abeta42) with insulin degrading enzyme: a molecular dynamics study. *Biochemistry* 49 3947-3956.
- Borth W 1992 Alpha 2-macroglobulin, a multifunctional binding protein with targeting characteristics. *FASEB J* 6 3345-3353.
- Boschert U, Merlo-Pich E, Higgins G, Roses AD & Catsicas S 1999 Apolipoprotein E expression by neurons surviving excitotoxic stress. *Neurobiol Dis* 6 508-514.
- Boyles JK, Pitas RE, Wilson E, Mahley RW & Taylor JM 1985 Apolipoprotein E associated with astrocytic glia of the central nervous system and with nonmyelinating glia of the peripheral nervous system. *J Clin Invest* 76 1501-1513.

- Buxbaum JN, Ye Z, Reixach N, Friske L, Levy C, Das P, Golde T, Masliah E, Roberts AR & Bartfai T 2008 Transthyretin protects Alzheimer's mice from the behavioral and biochemical effects of Abeta toxicity. *Proc Natl Acad Sci U S A* 105 2681-2686.
- Caporaso GL, Takei K, Gandy SE, Matteoli M, Mundigl O, Greengard P & De Camilli P 1994 Morphologic and biochemical analysis of the intracellular trafficking of the Alzheimer beta/A4 amyloid precursor protein. *J Neurosci* 14 3122-3138.
- Carro E, Trejo JL, Gomez-Isla T, LeRoith D & Torres-Aleman I 2002 Serum insulin-like growth factor I regulates brain amyloid-beta levels. *Nat Med* 8 1390-1397.
- Chauhan V, Ji L & Chauhan A 2008 Anti-amyloidogenic, anti-oxidant and anti-apoptotic role of gelsolin in Alzheimer's disease. *Biogerontology* 9 381-389.
- Chauhan VP, Ray I, Chauhan A & Wisniewski HM 1999 Binding of gelsolin, a secretory protein, to amyloid beta-protein. *Biochem Biophys Res Commun* 258 241-246.
- Chiang HL, Lyu RK, Tseng MY, Chang KH, Chang HS, Hsu WC, Kuo HC, Chu CC, Wu YR, Ro LS, et al. 2009 Analyses of transthyretin concentration in the cerebrospinal fluid of patients with Guillain-Barre syndrome and other neurological disorders. *Clin Chim Acta* 405 143-147.
- Choi SH, Leight SN, Lee VM, Li T, Wong PC, Johnson JA, Saraiva MJ & Sisodia SS 2007 Accelerated Abeta deposition in APP^{swe}/PS1^{deltaE9} mice with hemizygous deletions of TTR (transthyretin). *J Neurosci* 27 7006-7010.
- Chung RS, Howells C, Eaton ED, Shabala L, Zovo K, Palumaa P, Sillard R, Woodhouse A, Bennett WR, Ray S, et al. 2010 The native copper- and zinc-binding protein metallothionein blocks copper-mediated Abeta aggregation and toxicity in rat cortical neurons. *PLoS One* 5 e12030.
- Corder EH, Saunders AM, Strittmatter WJ, Schmechel DE, Gaskell PC, Small GW, Roses AD, Haines JL & Pericak-Vance MA 1993 Gene dose of apolipoprotein E type 4 allele and the risk of Alzheimer's disease in late onset families. *Science* 261 921-923.
- Costa R, Ferreira-da-Silva F, Saraiva MJ & Cardoso I 2008 Transthyretin protects against A-beta peptide toxicity by proteolytic cleavage of the peptide: a mechanism sensitive to the Kunitz protease inhibitor. *PLoS One* 3 e2899.
- Cupers P, Bentahir M, Craessaerts K, Orlans I, Vanderstichele H, Saftig P, De Strooper B & Annaert W 2001 The discrepancy between presenilin subcellular localization and gamma-secretase processing of amyloid precursor protein. *J Cell Biol* 154 731-740.
- Davidsson P, Westman-Brinkmalm A, Nilsson CL, Lindbjerg M, Paulson L, Andreasen N, Sjogren M & Blennow K 2002 Proteome analysis of cerebrospinal fluid proteins in Alzheimer patients. *Neuroreport* 13 611-615.
- Dawson GR, Seabrook GR, Zheng H, Smith DW, Graham S, O'Dowd G, Bowery BJ, Boyce S, Trumbauer ME, Chen HY, et al. 1999 Age-related cognitive deficits, impaired long-term potentiation and reduction in synaptic marker density in mice lacking the beta-amyloid precursor protein. *Neuroscience* 90 1-13.
- Deb S & Gottschall PE 1996 Increased production of matrix metalloproteinases in enriched astrocyte and mixed hippocampal cultures treated with beta-amyloid peptides. *J Neurochem* 66 1641-1647.
- DeKosky ST & Marek K 2003 Looking backward to move forward: early detection of neurodegenerative disorders. *Science* 302 830-834.
- Dominguez D, Tournoy J, Hartmann D, Huth T, Cryns K, Deforce S, Serneels L, Camacho IE, Marjaux E, Craessaerts K, et al. 2005 Phenotypic and biochemical analyses of BACE1- and BACE2-deficient mice. *J Biol Chem* 280 30797-30806.

- Dorfman VB, Pasquini L, Riudavets M, Lopez-Costa JJ, Villegas A, Troncoso JC, Lopera F, Castano EM & Morelli L 2010 Differential cerebral deposition of IDE and NEP in sporadic and familial Alzheimer's disease. *Neurobiol Aging* 31 1743-1757.
- Du J & Murphy RM 2010 Characterization of the interaction of beta-amyloid with transthyretin monomers and tetramers. *Biochemistry* 49 8276-8289.
- Du Y, Bales KR, Dodel RC, Liu X, Glinn MA, Horn JW, Little SP & Paul SM 1998 Alpha2-macroglobulin attenuates beta-amyloid peptide 1-40 fibril formation and associated neurotoxicity of cultured fetal rat cortical neurons. *J Neurochem* 70 1182-1188.
- Du Y, Ni B, Glinn M, Dodel RC, Bales KR, Zhang Z, Hyslop PA & Paul SM 1997 alpha2-Macroglobulin as a beta-amyloid peptide-binding plasma protein. *J Neurochem* 69 299-305.
- Eckman EA, Adams SK, Troendle FJ, Stodola BA, Kahn MA, Fauq AH, Xiao HD, Bernstein KE & Eckman CB 2006 Regulation of steady-state beta-amyloid levels in the brain by neprilysin and endothelin-converting enzyme but not angiotensin-converting enzyme. *J Biol Chem* 281 30471-30478.
- Eckman EA, Reed DK & Eckman CB 2001 Degradation of the Alzheimer's amyloid beta peptide by endothelin-converting enzyme. *J Biol Chem* 276 24540-24548.
- Eckman EA, Watson M, Marlow L, Sambamurti K & Eckman CB 2003 Alzheimer's disease beta-amyloid peptide is increased in mice deficient in endothelin-converting enzyme. *J Biol Chem* 278 2081-2084.
- Eggert S, Paliga K, Soba P, Evin G, Masters CL, Weidemann A & Beyreuther K 2004 The proteolytic processing of the amyloid precursor protein gene family members APLP-1 and APLP-2 involves alpha-, beta-, gamma-, and epsilon-like cleavages: modulation of APLP-1 processing by n-glycosylation. *J Biol Chem* 279 18146-18156.
- Ertekin-Taner N 2007 Genetics of Alzheimer's disease: a centennial review. *Neurol Clin* 25 611-667, v.
- Exley C & Korchazhkina OV 2001 Plasmin cleaves Abeta42 in vitro and prevents its aggregation into beta-pleated sheet structures. *Neuroreport* 12 2967-2970.
- Falkevall A, Alikhani N, Bhushan S, Pavlov PF, Busch K, Johnson KA, Eneqvist T, Tjernberg L, Ankarcrona M & Glaser E 2006 Degradation of the amyloid beta-protein by the novel mitochondrial peptidase, PreP. *J Biol Chem* 281 29096-29104.
- Farrer LA, Cupples LA, Haines JL, Hyman B, Kukull WA, Mayeux R, Myers RH, Pericak-Vance MA, Risch N & van Duijn CM 1997 Effects of age, sex, and ethnicity on the association between apolipoprotein E genotype and Alzheimer disease. A meta-analysis. APOE and Alzheimer Disease Meta Analysis Consortium. *Jama* 278 1349-1356.
- Farrer LA, Sherbatich T, Keryanov SA, Korovaitseva GI, Rogaeva EA, Petruk S, Premkumar S, Moliaka Y, Song YQ, Pei Y, et al. 2000 Association between angiotensin-converting enzyme and Alzheimer disease. *Arch Neurol* 57 210-214.
- Frangione B, Castano EM, Wisniewski T, Ghiso J, Prelli F & Vidal R 1996 Apolipoprotein E and amyloidogenesis. *Ciba Found Symp* 199 132-141; discussion 141-135.
- Gao W, Eisenhauer PB, Conn K, Lynch JA, Wells JM, Ullman MD, McKee A, Thatté HS & Fine RE 2004 Insulin degrading enzyme is expressed in the human cerebrovascular endothelium and in cultured human cerebrovascular endothelial cells. *Neurosci Lett* 371 6-11.

- Glenner GG & Wong CW 1984 Alzheimer's disease and Down's syndrome: sharing of a unique cerebrovascular amyloid fibril protein. *Biochem Biophys Res Commun* 122 1131-1135.
- Gloeckner SF, Meyne F, Wagner F, Heinemann U, Krasnianski A, Meissner B & Zerr I 2008 Quantitative analysis of transthyretin, tau and amyloid-beta in patients with dementia. *J Alzheimers Dis* 14 17-25.
- Goodman DS 1985 Retinoids and retinoid-binding proteins. *Harvey Lect* 81 111-132.
- Gottschall PE 1996 beta-Amyloid induction of gelatinase B secretion in cultured microglia: inhibition by dexamethasone and indomethacin. *Neuroreport* 7 3077-3080.
- Greenfield JP, Tsai J, Gouras GK, Hai B, Thinakaran G, Checler F, Sisodia SS, Greengard P & Xu H 1999 Endoplasmic reticulum and trans-Golgi network generate distinct populations of Alzheimer beta-amyloid peptides. *Proc Natl Acad Sci U S A* 96 742-747.
- Grundke-Iqbal I, Iqbal K, Tung YC, Quinlan M, Wisniewski HM & Binder LI 1986 Abnormal phosphorylation of the microtubule-associated protein tau (tau) in Alzheimer cytoskeletal pathology. *Proc Natl Acad Sci U S A* 83 4913-4917.
- Guan H, Liu Y, Daily A, Police S, Kim MH, Oddo S, LaFerla FM, Pauly JR, Murphy MP & Hersh LB 2009 Peripherally expressed neprilysin reduces brain amyloid burden: a novel approach for treating Alzheimer's disease. *J Neurosci Res* 87 1462-1473.
- Guo S, Wang S, Kim WJ, Lee SR, Frosch MP, Bacskai BJ, Greenberg SM & Lo EH 2006 Effects of apoE isoforms on beta-amyloid-induced matrix metalloproteinase-9 in rat astrocytes. *Brain Res* 1111 222-226.
- Haapasalo A & Kovacs DM 2011 The Many Substrates of Presenilin/gamma-Secretase. *J Alzheimers Dis*.
- Haass C 2004 Take five--BACE and the gamma-secretase quartet conduct Alzheimer's amyloid beta-peptide generation. *EMBO J* 23 483-488.
- Haass C & Selkoe DJ 1993 Cellular processing of beta-amyloid precursor protein and the genesis of amyloid beta-peptide. *Cell* 75 1039-1042.
- Hajjar IM, Keown M, Lewis P & Almor A 2008 Angiotensin converting enzyme inhibitors and cognitive and functional decline in patients with Alzheimer's disease: an observational study. *Am J Alzheimers Dis Other Demen* 23 77-83.
- Han SH, Jung ES, Sohn JH, Hong HJ, Hong HS, Kim JW, N DL, Kim M, Kim H, Ha HJ, Kim YH, Huh N, Jung MW, Mook-Jung I. 2011 Human Serum Transthyretin Levels Correlate Inversely with Alzheimer's Disease. *J Alzheimers Dis*.
- Hansson SF, Puchades M, Blennow K, Sjogren M & Davidsson P 2004 Validation of a prefractionation method followed by two-dimensional electrophoresis - Applied to cerebrospinal fluid proteins from frontotemporal dementia patients. *Proteome Sci* 2 7.
- Hardy J & Selkoe DJ 2002 The amyloid hypothesis of Alzheimer's disease: progress and problems on the road to therapeutics. *Science* 297 353-356.
- Hardy JA & Higgins GA 1992 Alzheimer's disease: the amyloid cascade hypothesis. *Science* 256 184-185.
- Heber S, Herms J, Gajic V, Hainfellner J, Aguzzi A, Rulicke T, von Kretschmar H, von Koch C, Sisodia S, Tremml P, et al. 2000 Mice with combined gene knock-outs reveal essential and partially redundant functions of amyloid precursor protein family members. *J Neurosci* 20 7951-7963.

- Hegde ML, Bharathi P, Suram A, Venugopal C, Jagannathan R, Poddar P, Srinivas P, Sambamurti K, Rao KJ, Scancar J, et al. 2009 Challenges associated with metal chelation therapy in Alzheimer's disease. *J Alzheimers Dis* 17 457-468.
- Helbecque N, Cotel D, Hermant X & Amouyel P 2007 Impact of the matrix metalloproteinase MMP-3 on dementia. *Neurobiol Aging* 28 1215-1220.
- Hemming ML & Selkoe DJ 2005 Amyloid beta-protein is degraded by cellular angiotensin-converting enzyme (ACE) and elevated by an ACE inhibitor. *J Biol Chem* 280 37644-37650.
- Hemming ML, Selkoe DJ & Farris W 2007 Effects of prolonged angiotensin-converting enzyme inhibitor treatment on amyloid beta-protein metabolism in mouse models of Alzheimer disease. *Neurobiol Dis* 26 273-281.
- Hickman SE, Allison EK & El Khoury J 2008 Microglial dysfunction and defective beta-amyloid clearance pathways in aging Alzheimer's disease mice. *J Neurosci* 28 8354-8360.
- Hidalgo J, Chung R, Penkowa M, Vařák M. 2009 Structure and function of vertebrate metallothioneins, in "Metallothioneins and Related Chelators", Vol. 5 of 'Metal Ions in Life Sciences'; Sigel, A., Sigel, H., Sigel, R.K.O., Eds. (2009) The Royal Society of Chemistry, Cambridge, UK, pp 279-317.
- Hirko AC, Meyer EM, King MA & Hughes JA 2007 Peripheral transgene expression of plasma gelsolin reduces amyloid in transgenic mouse models of Alzheimer's disease. *Mol Ther* 15 1623-1629.
- Holcomb L, Gordon MN, McGowan E, Yu X, Benkovic S, Jantzen P, Wright K, Saad I, Mueller R, Morgan D, et al. 1998 Accelerated Alzheimer-type phenotype in transgenic mice carrying both mutant amyloid precursor protein and presenilin 1 transgenes. *Nat Med* 4 97-100.
- Hook VY, Kindy M, Reinheckel T, Peters C & Hook G 2009 Genetic cathepsin B deficiency reduces beta-amyloid in transgenic mice expressing human wild-type amyloid precursor protein. *Biochem Biophys Res Commun* 386 284-288.
- Horstmann S, Budig L, Gardner H, Koziol J, Deuschle M, Schilling C & Wagner S 2010 Matrix metalloproteinases in peripheral blood and cerebrospinal fluid in patients with Alzheimer's disease. *Int Psychogeriatr* 22 966-972.
- Howells C, West AK & Chung RS 2010 Neuronal growth-inhibitory factor (metallothionein-3): evaluation of the biological function of growth-inhibitory factor in the injured and neurodegenerative brain. *Febs J* 277 2931-2939.
- Hsiao K, Chapman P, Nilsen S, Eckman C, Harigaya Y, Younkin S, Yang F & Cole G 1996 Correlative memory deficits, Abeta elevation, and amyloid plaques in transgenic mice. *Science* 274 99-102.
- Hu J, Igarashi A, Kamata M & Nakagawa H 2001 Angiotensin-converting enzyme degrades Alzheimer amyloid beta-peptide (A beta); retards A beta aggregation, deposition, fibril formation; and inhibits cytotoxicity. *J Biol Chem* 276 47863-47868.
- Hu J, Miyatake F, Aizu Y, Nakagawa H, Nakamura S, Tamaoka A, Takahash R, Urakami K & Shoji M 1999 Angiotensin-converting enzyme genotype is associated with Alzheimer disease in the Japanese population. *Neurosci Lett* 277 65-67.
- Huang Y 2010 Abeta-independent roles of apolipoprotein E4 in the pathogenesis of Alzheimer's disease. *Trends Mol Med* 16 287-294.

- Huang Y, Weisgraber KH, Mucke L & Mahley RW 2004 Apolipoprotein E: diversity of cellular origins, structural and biophysical properties, and effects in Alzheimer's disease. *J Mol Neurosci* 23 189-204.
- Hughes SR, Khorkova O, Goyal S, Knaeblein J, Heroux J, Riedel NG & Sahasrabudhe S 1998 Alpha2-macroglobulin associates with beta-amyloid peptide and prevents fibril formation. *Proc Natl Acad Sci U S A* 95 3275-3280.
- Huse JT, Liu K, Pijak DS, Carlin D, Lee VM & Doms RW 2002 Beta-secretase processing in the trans-Golgi network preferentially generates truncated amyloid species that accumulate in Alzheimer's disease brain. *J Biol Chem* 277 16278-16284.
- Ichai C, Chevallier N, Delaere P, Dournaud P, Epelbaum J, Hauw JJ, Vincent JP & Checler F 1994 Influence of region-specific alterations of neuropeptidase content on the catabolic fates of neuropeptides in Alzheimer's disease. *J Neurochem* 62 645-655.
- Irie, Y., and Keung, W. M. (2001) Metallothionein-III antagonizes the neurotoxic and neurotrophic effects of amyloid beta peptides. *Biochem Biophys Res Commun* 282, 416-420
- Irie, Y., and Keung, W. M. (2003) Anti-amyloid beta activity of metallothionein-III is different from its neuronal growth inhibitory activity: structure-activity studies. *Brain research* 960, 228-234
- Irizarry MC, McNamara M, Fedorchak K, Hsiao K & Hyman BT 1997a APPSw transgenic mice develop age-related A beta deposits and neuropil abnormalities, but no neuronal loss in CA1. *J Neuropathol Exp Neurol* 56 965-973.
- Irizarry MC, Soriano F, McNamara M, Page KJ, Schenk D, Games D & Hyman BT 1997b Abeta deposition is associated with neuropil changes, but not with overt neuronal loss in the human amyloid precursor protein V717F (PDAPP) transgenic mouse. *J Neurosci* 17 7053-7059.
- Isbir T, Agachan B, Yilmaz H, Aydin M, Kara I, Eker D & Eker E 2001 Interaction between apolipoprotein-E and angiotensin-converting enzyme genotype in Alzheimer's disease. *Am J Alzheimers Dis Other Demen* 16 205-210.
- Ito S, Ohtsuki S, Kamiie J, Nezu Y & Terasaki T 2007 Cerebral clearance of human amyloid-beta peptide (1-40) across the blood-brain barrier is reduced by self-aggregation and formation of low-density lipoprotein receptor-related protein-1 ligand complexes. *J Neurochem* 103 2482-2490.
- Jacobson DR, McFarlin DE, Kane I & Buxbaum JN 1992 Transthyretin Pro55, a variant associated with early-onset, aggressive, diffuse amyloidosis with cardiac and neurologic involvement. *Hum Genet* 89 353-356.
- Janmey PA, Chaponnier C, Lind SE, Zaner KS, Stossel TP & Yin HL 1985 Interactions of gelsolin and gelsolin-actin complexes with actin. Effects of calcium on actin nucleation, filament severing, and end blocking. *Biochemistry* 24 3714-3723.
- Ji L, Chauhan A & Chauhan V 2008 Cytoplasmic gelsolin in pheochromocytoma-12 cells forms a complex with amyloid beta-protein. *Neuroreport* 19 463-466.
- Jiang Q, Lee CY, Mandrekar S, Wilkinson B, Cramer P, Zelcer N, Mann K, Lamb B, Willson TM, Collins JL, et al. 2008 ApoE promotes the proteolytic degradation of Abeta. *Neuron* 58 681-693.
- Johnson KA, Bhushan S, Stahl A, Hallberg BM, Frohn A, Glaser E & Eneqvist T 2006 The closed structure of presequence protease PreP forms a unique 10,000 Angstroms³ chamber for proteolysis. *EMBO J* 25 1977-1986.

- Jorissen E, Prox J, Bernreuther C, Weber S, Schwanbeck R, Serneels L, Snellinx A, Craessaerts K, Thathiah A, Tesseur I, et al. 2010 The disintegrin/metalloproteinase ADAM10 is essential for the establishment of the brain cortex. *J Neurosci* 30 4833-4844.
- Jung SS, Zhang W & Van Nostrand WE 2003 Pathogenic A beta induces the expression and activation of matrix metalloproteinase-2 in human cerebrovascular smooth muscle cells. *J Neurochem* 85 1208-1215.
- Kang J & Muller-Hill B 1990 Differential splicing of Alzheimer's disease amyloid A4 precursor RNA in rat tissues: PreA4(695) mRNA is predominantly produced in rat and human brain. *Biochem Biophys Res Commun* 166 1192-1200.
- Kehoe PG & Wilcock GK 2007 Is inhibition of the renin-angiotensin system a new treatment option for Alzheimer's disease? *Lancet Neurol* 6 373-378.
- Kim J, Basak JM & Holtzman DM 2009 The role of apolipoprotein E in Alzheimer's disease. *Neuron* 63 287-303.
- Kimberly WT & Wolfe MS 2003 Identity and function of gamma-secretase. *J Neurosci Res* 74 353-360.
- Koistinaho M, Lin S, Wu X, Esterman M, Koger D, Hanson J, Higgs R, Liu F, Malkani S, Bales KR, et al. 2004 Apolipoprotein E promotes astrocyte colocalization and degradation of deposited amyloid-beta peptides. *Nat Med* 10 719-726.
- Konings CH, Kuiper MA, Scheltens P, Grijpma AM, van Pelt W & Wolters EC 1993 Re-evaluation of cerebrospinal fluid angiotensin-converting enzyme activity in patients with 'probable' Alzheimer's disease. *Eur J Clin Chem Clin Biochem* 31 495-497.
- Korenberg JR, Pulst SM, Neve RL & West R 1989 The Alzheimer amyloid precursor protein maps to human chromosome 21 bands q21.105-q21.05. *Genomics* 5 124-127.
- Kovacs DM 2000 alpha2-macroglobulin in late-onset Alzheimer's disease. *Exp Gerontol* 35 473-479.
- Kowalska A 2003 Amyloid precursor protein gene mutations responsible for early-onset autosomal dominant Alzheimer's disease. *Folia Neuropathol* 41 35-40.
- Kuhn PH, Wang H, Dislich B, Colombo A, Zeitschel U, Ellwart JW, Kremmer E, Rossner S & Lichtenthaler SF 2010 ADAM10 is the physiologically relevant, constitutive alpha-secretase of the amyloid precursor protein in primary neurons. *EMBO J* 29 3020-3032.
- Kurochkin IV & Goto S 1994 Alzheimer's beta-amyloid peptide specifically interacts with and is degraded by insulin degrading enzyme. *FEBS Lett* 345 33-37.
- Kwiatkowski DJ, Janmey PA, Mole JE & Yin HL 1985 Isolation and properties of two actin-binding domains in gelsolin. *J Biol Chem* 260 15232-15238.
- Lammich S, Kojro E, Postina R, Gilbert S, Pfeiffer R, Jasionowski M, Haass C & Fahrenholz F 1999 Constitutive and regulated alpha-secretase cleavage of Alzheimer's amyloid precursor protein by a disintegrin metalloprotease. *Proc Natl Acad Sci U S A* 96 3922-3927.
- Lanni C, Nardinocchi L, Puca R, Stanga S, Uberti D, Memo M, Govoni S, D'Orazi G & Racchi M 2010 Homeodomain interacting protein kinase 2: a target for Alzheimer's beta amyloid leading to misfolded p53 and inappropriate cell survival. *PLoS One* 5 e10171.

- Ledesma MD, Da Silva JS, Crassaerts K, Delacourte A, De Strooper B & Dotti CG 2000 Brain plasmin enhances APP alpha-cleavage and Abeta degradation and is reduced in Alzheimer's disease brains. *EMBO Rep* 1 530-535.
- Lee JY, Kweon HS, Cho E, Byun HR, Kim DH, Kim YH, Han PL & Koh JY 2007 Upregulation of tPA/plasminogen proteolytic system in the periphery of amyloid deposits in the Tg2576 mouse model of Alzheimer's disease. *Neurosci Lett* 423 82-87.
- Leissring MA, Farris W, Chang AY, Walsh DM, Wu X, Sun X, Frosch MP & Selkoe DJ 2003 Enhanced proteolysis of beta-amyloid in APP transgenic mice prevents plaque formation, secondary pathology, and premature death. *Neuron* 40 1087-1093.
- LePage RN, Fosang AJ, Fuller SJ, Murphy G, Evin G, Beyreuther K, Masters CL & Small DH 1995 Gelatinase A possesses a beta-secretase-like activity in cleaving the amyloid protein precursor of Alzheimer's disease. *FEBS Lett* 377 267-270.
- Li NC, Lee A, Whitmer RA, Kivipelto M, Lawler E, Kazis LE & Wolozin B 2010 Use of angiotensin receptor blockers and risk of dementia in a predominantly male population: prospective cohort analysis. *BMJ* 340 b5465.
- Li P, Kuo WL, Yousef M, Rosner MR & Tang WJ 2006 The C-terminal domain of human insulin degrading enzyme is required for dimerization and substrate recognition. *Biochem Biophys Res Commun* 343 1032-1037.
- Li R, Lindholm K, Yang LB, Yue X, Citron M, Yan R, Beach T, Sue L, Sabbagh M, Cai H, et al. 2004 Amyloid beta peptide load is correlated with increased beta-secretase activity in sporadic Alzheimer's disease patients. *Proc Natl Acad Sci U S A* 101 3632-3637.
- Li W, Poteet E, Xie L, Liu R, Wen Y & Yang SH 2011 Regulation of matrix metalloproteinase 2 by oligomeric amyloid beta protein. *Brain Res*.
- Liao A, Nitsch RM, Greenberg SM, Finckh U, Blacker D, Albert M, Rebeck GW, Gomez-Isla T, Clatworthy A, Binetti G, et al. 1998 Genetic association of an alpha2-macroglobulin (Val1000Ile) polymorphism and Alzheimer's disease. *Hum Mol Genet* 7 1953-1956.
- Link CD 1995 Expression of human beta-amyloid peptide in transgenic *Caenorhabditis elegans*. *Proc Natl Acad Sci U S A* 92 9368-9372.
- Liu L & Murphy RM 2006 Kinetics of Inhibition of beta-Amyloid Aggregation by Transthyretin. *Biochemistry* 45 15702-15709.
- Mahley RW & Huang Y 2006 Apolipoprotein (apo) E4 and Alzheimer's disease: unique conformational and biophysical properties of apoE4 can modulate neuropathology. *Acta Neurol Scand Suppl* 185 8-14.
- Mahley RW, Weisgraber KH & Huang Y 2006 Apolipoprotein E4: a causative factor and therapeutic target in neuropathology, including Alzheimer's disease. *Proc Natl Acad Sci U S A* 103 5644-5651.
- Malito E, Hulse RE & Tang WJ 2008 Amyloid beta-degrading cryptidases: insulin degrading enzyme, presequence peptidase, and neprilysin. *Cell Mol Life Sci* 65 2574-2585.
- Manelli AM, Bulfinch LC, Sullivan PM & LaDu MJ 2007 Abeta42 neurotoxicity in primary co-cultures: effect of apoE isoform and Abeta conformation. *Neurobiol Aging* 28 1139-1147.
- Martinho A, Goncalves I, Cardoso I, Almeida MR, Quintela T, Saraiva MJ & Santos CR 2010 Human metallothioneins 2 and 3 differentially affect amyloid-beta binding by transthyretin. *Febs J* 277 3427-3436.


- Matsumoto N, Kitayama H, Kitada M, Kimura K, Noda M & Ide C 2003 Isolation of a set of genes expressed in the choroid plexus of the mouse using suppression subtractive hybridization. *Neuroscience* 117 405-415.
- Matsuoka Y, Saito M, LaFrancois J, Gaynor K, Olm V, Wang L, Casey E, Lu Y, Shiratori C, Lemere C, et al. 2003 Novel therapeutic approach for the treatment of Alzheimer's disease by peripheral administration of agents with an affinity to beta-amyloid. *J Neurosci* 23 29-33.
- McDermott JR & Gibson AM 1997 Degradation of Alzheimer's beta-amyloid protein by human and rat brain peptidases: involvement of insulin-degrading enzyme. *Neurochem Res* 22 49-56.
- Meilandt WJ, Cisse M, Ho K, Wu T, Esposito LA, Scarce-Levie K, Cheng IH, Yu GQ & Mucke L 2009 Nephrilysin overexpression inhibits plaque formation but fails to reduce pathogenic A β oligomers and associated cognitive deficits in human amyloid precursor protein transgenic mice. *J Neurosci* 29 1977-1986.
- Melchor JP, Pawlak R & Strickland S 2003 The tissue plasminogen activator-plasminogen proteolytic cascade accelerates amyloid-beta (A β) degradation and inhibits A β -induced neurodegeneration. *J Neurosci* 23 8867-8871.
- Meloni G, Sonois V, Delaine T, Guilloureau L, Gillet A, Teissie J, Faller P & Vasak M 2008 Metal swap between Zn²⁺-metallothionein-3 and amyloid-beta-Cu protects against amyloid-beta toxicity. *Nat Chem Biol* 4 366-372.
- Merlos-Suarez A, Ruiz-Paz S, Baselga J & Arribas J 2001 Metalloprotease-dependent protransforming growth factor- α ectodomain shedding in the absence of tumor necrosis factor- α -converting enzyme. *J Biol Chem* 276 48510-48517.
- Miners JS, Ashby E, Van Helmond Z, Chalmers KA, Palmer LE, Love S & Kehoe PG 2008a Angiotensin-converting enzyme (ACE) levels and activity in Alzheimer's disease, and relationship of perivascular ACE-1 to cerebral amyloid angiopathy. *Neuropathol Appl Neurobiol* 34 181-193.
- Miners JS, Baig S, Palmer J, Palmer LE, Kehoe PG & Love S 2008b A β -degrading enzymes in Alzheimer's disease. *Brain Pathol* 18 240-252.
- Monji A, Yoshida I, Tashiro K, Hayashi Y, Matsuda K & Tashiro N 2000 Inhibition of A β fibril formation and A β -induced cytotoxicity by senile plaque-associated proteins. *Neurosci Lett* 278 81-84.
- Narita M, Holtzman DM, Schwartz AL & Bu G 1997 Alpha₂-macroglobulin complexes with and mediates the endocytosis of beta-amyloid peptide via cell surface low-density lipoprotein receptor-related protein. *J Neurochem* 69 1904-1911.
- Palha JA, Moreira P, Wisniewski T, Frangione B & Saraiva MJ 1996 Transthyretin gene in Alzheimer's disease patients. *Neurosci Lett* 204 212-214.
- Palmer JC, Kehoe PG & Love S 2010 Endothelin-converting enzyme-1 in Alzheimer's disease and vascular dementia. *Neuropathol Appl Neurobiol* 36 487-497.
- Penkowa M, Tio L, Giralt M, Quintana A, Molinero A, Atrian S, Vasak M & Hidalgo J 2006 Specificity and divergence in the neurobiologic effects of different metallothioneins after brain injury. *J Neurosci Res* 83 974-984.
- Perez A, Morelli L, Cresto JC & Castano EM 2000 Degradation of soluble amyloid beta-peptides 1-40, 1-42, and the Dutch variant 1-40Q by insulin degrading enzyme from Alzheimer disease and control brains. *Neurochem Res* 25 247-255.

- Pitas RE, Boyles JK, Lee SH, Foss D & Mahley RW 1987 Astrocytes synthesize apolipoprotein E and metabolize apolipoprotein E-containing lipoproteins. *Biochim Biophys Acta* 917 148-161.
- Qiu WQ, Borth W, Ye Z, Haass C, Teplow DB & Selkoe DJ 1996 Degradation of amyloid beta-protein by a serine protease-alpha2-macroglobulin complex. *J Biol Chem* 271 8443-8451.
- Qiu WQ, Walsh DM, Ye Z, Vekrellis K, Zhang J, Podlisny MB, Rosner MR, Safavi A, Hersh LB & Selkoe DJ 1998 Insulin-degrading enzyme regulates extracellular levels of amyloid beta-protein by degradation. *J Biol Chem* 273 32730-32738.
- Ray I, Chauhan A, Wegiel J & Chauhan VP 2000 Gelsolin inhibits the fibrillization of amyloid beta-protein, and also defibrillizes its preformed fibrils. *Brain Res* 853 344-351.
- Reitz C, van Rooij FJ, de Maat MP, den Heijer T, Hofman A, Witteman JC & Breteler MM 2008 Matrix metalloproteinase 3 haplotypes and dementia and Alzheimer's disease. The Rotterdam Study. *Neurobiol Aging* 29 874-881.
- Reitz C, van Rooij FJ, Soares HD, de Maat MP, Hofman A, Witteman JC & Breteler MM 2010 Matrix metalloproteinase 3 haplotypes and plasma amyloid beta levels: the Rotterdam Study. *Neurobiol Aging* 31 715-718.
- Roher AE, Kasunic TC, Woods AS, Cotter RJ, Ball MJ & Fridman R 1994 Proteolysis of A beta peptide from Alzheimer disease brain by gelatinase A. *Biochem Biophys Res Commun* 205 1755-1761.
- Saraiva MJ, Birken S, Costa PP & Goodman DS 1984 Family studies of the genetic abnormality in transthyretin (prealbumin) in Portuguese patients with familial amyloidotic polyneuropathy. *Ann N Y Acad Sci* 435 86-100.
- Schochet SS, Jr., Lampert PW & McCormick WF 1973 Neurofibrillary tangles in patients with Down's syndrome: a light and electron microscopic study. *Acta Neuropathol* 23 342-346.
- Schwarzman AL & Goldgaber D 1996 Interaction of transthyretin with amyloid beta-protein: binding and inhibition of amyloid formation. *Ciba Found Symp* 199 146-160; discussion 160-144.
- Schwarzman AL, Gregori L, Vitek MP, Lyubski S, Strittmatter WJ, Enghilde JJ, Bhasin R, Silverman J, Weisgraber KH, Coyle PK, et al. 1994 Transthyretin sequesters amyloid beta protein and prevents amyloid formation. *Proc Natl Acad Sci U S A* 91 8368-8372.
- Schwarzman AL, Tsiper M, Gregori L, Goldgaber D, Frakowiak J, Mazur-Kolecka B, Taraskina A, Pchelina S & Vitek MP 2005 Selection of peptides binding to the amyloid b-protein reveals potential inhibitors of amyloid formation. *Amyloid* 12 199-209.
- Schwarzman AL, Tsiper M, Wenthe H, Wang A, Vitek MP, Vasiliev V & Goldgaber D 2004 Amyloidogenic and anti-amyloidogenic properties of recombinant transthyretin variants. *Amyloid* 11 1-9.
- Serot JM, Christmann D, Dubost T & Couturier M 1997 Cerebrospinal fluid transthyretin: aging and late onset Alzheimer's disease. *J Neurol Neurosurg Psychiatry* 63 506-508.
- Seubert P, Vigo-Pelfrey C, Esch F, Lee M, Dovey H, Davis D, Sinha S, Schlossmacher M, Whaley J, Swindlehurst C, et al. 1992 Isolation and quantification of soluble Alzheimer's beta-peptide from biological fluids. *Nature* 359 325-327.

- Shah K, Qureshi SU, Johnson M, Parikh N, Schulz PE & Kunik ME 2009 Does use of antihypertensive drugs affect the incidence or progression of dementia? A systematic review. *Am J Geriatr Pharmacother* 7 250-261.
- Shirahama T, Skinner M, Westermark P, Rubinow A, Cohen AS, Brun A & Kemper TL 1982 Senile cerebral amyloid. Prealbumin as a common constituent in the neuritic plaque, in the neurofibrillary tangle, and in the microangiopathic lesion. *Am J Pathol* 107 41-50.
- Sluck JM, Lin RC, Katolik LI, Jeng AY & Lehmann JC 1999 Endothelin converting enzyme-1-, endothelin-1-, and endothelin-3-like immunoreactivity in the rat brain. *Neuroscience* 91 1483-1497.
- Solans A, Estivill X & de La Luna S 2000 A new aspartyl protease on 21q22.3, BACE2, is highly similar to Alzheimer's amyloid precursor protein beta-secretase. *Cytogenet Cell Genet* 89 177-184.
- Soprano DR, Herbert J, Soprano KJ, Schon EA & Goodman DS 1985 Demonstration of transthyretin mRNA in the brain and other extrahepatic tissues in the rat. *J Biol Chem* 260 11793-11798.
- Stabilini R, Vergani C, Agostoni A & Agostoni RP 1968 Influence of age and sex on prealbumin levels. *Clin Chim Acta* 20 358-359.
- Stahl A, Moberg P, Ytterberg J, Panfilov O, Brockenhuus Von Lowenhielm H, Nilsson F & Glaser E 2002 Isolation and identification of a novel mitochondrial metalloprotease (PreP) that degrades targeting presequences in plants. *J Biol Chem* 277 41931-41939.
- Stein TD, Anders NJ, DeCarli C, Chan SL, Mattson MP & Johnson JA 2004 Neutralization of transthyretin reverses the neuroprotective effects of secreted amyloid precursor protein (APP) in APPSW mice resulting in tau phosphorylation and loss of hippocampal neurons: support for the amyloid hypothesis. *J Neurosci* 24 7707-7717.
- Stein TD & Johnson JA 2002 Lack of neurodegeneration in transgenic mice overexpressing mutant amyloid precursor protein is associated with increased levels of transthyretin and the activation of cell survival pathways. *J Neurosci* 22 7380-7388.
- Takasugi N, Tomita T, Hayashi I, Tsuruoka M, Niimura M, Takahashi Y, Thinakaran G & Iwatsubo T 2003 The role of presenilin cofactors in the gamma-secretase complex. *Nature* 422 438-441.
- Talamagas AA, Efthimiopoulos S, Tsilibary EC, Figueiredo-Pereira ME & Tzinia AK 2007 Abeta(1-40)-induced secretion of matrix metalloproteinase-9 results in sAPPalpha release by association with cell surface APP. *Neurobiol Dis* 28 304-315.
- Tang BL 2009 Neuronal protein trafficking associated with Alzheimer disease: from APP and BACE1 to glutamate receptors. *Cell Adh Migr* 3 118-128.
- Tillement L, Lecanu L & Papadopoulos V 2011 Alzheimer's disease: effects of beta-amyloid on mitochondria. *Mitochondrion* 11 13-21.
- Tucker HM, Kihiko-Ehmann M & Estus S 2002 Urokinase-type plasminogen activator inhibits amyloid-beta neurotoxicity and fibrillogenesis via plasminogen. *J Neurosci Res* 70 249-255.
- Tucker HM, Kihiko-Ehmann M, Wright S, Rydel RE & Estus S 2000a Tissue plasminogen activator requires plasminogen to modulate amyloid-beta neurotoxicity and deposition. *J Neurochem* 75 2172-2177.
- Tucker HM, Kihiko M, Caldwell JN, Wright S, Kawarabayashi T, Price D, Walker D, Scheff S, McGillis JP, Rydel RE, et al. 2000b The plasmin system is induced by and degrades amyloid-beta aggregates. *J Neurosci* 20 3937-3946.

- Vahlquist A, Rask L, Peterson PA & Berg T 1975 The concentrations of retinol-binding protein, prealbumin, and transferrin in the sera of newly delivered mothers and children of various ages. *Scand J Clin Lab Invest* 35 569-575.
- Vargas T, Antequera D, Ugalde C, Spuch C & Carro E 2010 Gelsolin restores A beta-induced alterations in choroid plexus epithelium. *J Biomed Biotechnol* 2010 805405.
- Vassar R, Bennett BD, Babu-Khan S, Kahn S, Mendiaz EA, Denis P, Teplow DB, Ross S, Amarante P, Loeloff R, et al. 1999 Beta-secretase cleavage of Alzheimer's amyloid precursor protein by the transmembrane aspartic protease BACE. *Science* 286 735-741.
- Vassar R & Citron M 2000 Abeta-generating enzymes: recent advances in beta- and gamma-secretase research. *Neuron* 27 419-422.
- Vepsalainen S, Hiltunen M, Helisalmi S, Wang J, van Groen T, Tanila H & Soininen H 2008 Increased expression of Abeta degrading enzyme IDE in the cortex of transgenic mice with Alzheimer's disease-like neuropathology. *Neurosci Lett* 438 216-220.
- Vigo-Pelfrey C, Lee D, Keim P, Lieberburg I & Schenk DB 1993 Characterization of beta-amyloid peptide from human cerebrospinal fluid. *J Neurochem* 61 1965-1968.
- von Koch CS, Zheng H, Chen H, Trumbauer M, Thinakaran G, van der Ploeg LH, Price DL & Sisodia SS 1997 Generation of APLP2 KO mice and early postnatal lethality in APLP2/APP double KO mice. *Neurobiol Aging* 18 661-669.
- Wang DS, Dickson DW & Malter JS 2006 beta-Amyloid degradation and Alzheimer's disease. *J Biomed Biotechnol* 2006 58406.
- Wang S, Wang R, Chen L, Bennett DA, Dickson DW & Wang DS 2010 Expression and functional profiling of neprilysin, insulin-degrading enzyme, and endothelin-converting enzyme in prospectively studied elderly and Alzheimer's brain. *J Neurochem* 115 47-57.
- Wati H, Kawarabayashi T, Matsubara E, Kasai A, Hirasawa T, Kubota T, Harigaya Y, Shoji M & Maeda S 2009 Transthyretin accelerates vascular Abeta deposition in a mouse model of Alzheimer's disease. *Brain Pathol* 19 48-57.
- Weeraratna AT, Kalehua A, Deleon I, Bertak D, Maher G, Wade MS, Lustig A, Becker KG, Wood W, 3rd, Walker DG, et al. 2007 Alterations in immunological and neurological gene expression patterns in Alzheimer's disease tissues. *Exp Cell Res* 313 450-461.
- Weisgraber KH & Mahley RW 1996 Human apolipoprotein E: the Alzheimer's disease connection. *FASEB J* 10 1485-1494.
- Weisner B & Kauerz U 1983 The influence of the choroid plexus on the concentration of prealbumin in CSF. *J Neurol Sci* 61 27-35.
- Wen D, Corina K, Chow EP, Miller S, Janmey PA & Pepinsky RB 1996 The plasma and cytoplasmic forms of human gelsolin differ in disulfide structure. *Biochemistry* 35 9700-9709.
- Wisniewski T & Frangione B 1996 Molecular biology of brain aging and neurodegenerative disorders. *Acta Neurobiol Exp (Wars)* 56 267-279.
- Woeber KA & Ingbar SH 1968 The contribution of thyroxine-binding prealbumin to the binding of thyroxine in human serum, as assessed by immunoadsorption. *J Clin Invest* 47 1710-1721.
- Xie L, Helmerhorst E, Taddei K, Plewright B, Van Bronswijk W & Martins R 2002 Alzheimer's beta-amyloid peptides compete for insulin binding to the insulin receptor. *J Neurosci* 22 RC221.

- Yamada T, Yoshiyama Y, Sato H, Seiki M, Shinagawa A & Takahashi M 1995 White matter microglia produce membrane-type matrix metalloprotease, an activator of gelatinase A, in human brain tissues. *Acta Neuropathol* 90 421-424.
- Yan P, Hu X, Song H, Yin K, Bateman RJ, Cirrito JR, Xiao Q, Hsu FF, Turk JW, Xu J, et al. 2006 Matrix metalloproteinase-9 degrades amyloid-beta fibrils in vitro and compact plaques in situ. *J Biol Chem* 281 24566-24574.
- Yang JD, Feng G, Zhang J, Lin ZX, Shen T, Breen G, St Clair D & He L 2000 Association between angiotensin-converting enzyme gene and late onset Alzheimer's disease in Han chinese. *Neurosci Lett* 295 41-44.
- Yerbury JJ, Kumita JR, Meehan S, Dobson CM & Wilson MR 2009 alpha2-Macroglobulin and haptoglobin suppress amyloid formation by interacting with prefibrillar protein species. *J Biol Chem* 284 4246-4254.
- Yin KJ, Cirrito JR, Yan P, Hu X, Xiao Q, Pan X, Bateman R, Song H, Hsu FF, Turk J, et al. 2006 Matrix metalloproteinases expressed by astrocytes mediate extracellular amyloid-beta peptide catabolism. *J Neurosci* 26 10939-10948.
- Zhang YW, Thompson R, Zhang H & Xu H 2011 APP processing in Alzheimer's disease. *Mol Brain* 4 3.
- Zhao J & Pei G 2008 Evoking plasmin for beta-amyloid clearance. *Cell Res* 18 803-804.
- Zhao Z, Xiang Z, Haroutunian V, Buxbaum JD, Stetka B & Pasinetti GM 2007 Insulin degrading enzyme activity selectively decreases in the hippocampal formation of cases at high risk to develop Alzheimer's disease. *Neurobiol Aging* 28 824-830.
- Zheng H & Koo EH 2006 The amyloid precursor protein: beyond amyloid. *Mol Neurodegener* 1 5.
- Zhong N & Weisgraber KH 2009a Understanding the association of apolipoprotein E4 with Alzheimer disease: clues from its structure. *J Biol Chem* 284 6027-6031.
- Zhong N & Weisgraber KH 2009b Understanding the basis for the association of apoE4 with Alzheimer's disease: opening the door for therapeutic approaches. *Curr Alzheimer Res* 6 415-418.
- Zlokovic BV 2004 Clearing amyloid through the blood-brain barrier. *J Neurochem* 89 807-811.
- Zou K, Maeda T, Watanabe A, Liu J, Liu S, Oba R, Satoh Y, Komano H & Michikawa M 2009 Abeta42-to-Abeta40- and angiotensin-converting activities in different domains of angiotensin-converting enzyme. *J Biol Chem* 284 31914-31920.
- Zou K, Yamaguchi H, Akatsu H, Sakamoto T, Ko M, Mizoguchi K, Gong JS, Yu W, Yamamoto T, Kosaka K, et al. 2007 Angiotensin-converting enzyme converts amyloid beta-protein 1-42 (Abeta(1-42)) to Abeta(1-40), and its inhibition enhances brain Abeta deposition. *J Neurosci* 27 8628-8635.
- Zubenko GS, Volicer L, Drenfeld LK, Freeman M, Langlais PJ & Nixon RA 1985 Cerebrospinal fluid levels of angiotensin-converting enzyme in Alzheimer's disease, Parkinson's disease and progressive supranuclear palsy. *Brain Res* 328 215-221.



Alzheimer's Disease Pathogenesis-Core Concepts, Shifting Paradigms and Therapeutic Targets

Edited by Dr. Suzanne De La Monte

ISBN 978-953-307-690-4

Hard cover, 686 pages

Publisher InTech

Published online 12, September, 2011

Published in print edition September, 2011

Alzheimer's Disease Pathogenesis: Core Concepts, Shifting Paradigms, and Therapeutic Targets, delivers the concepts embodied within its title. This exciting book presents the full array of theories about the causes of Alzheimer's, including fresh concepts that have gained ground among both professionals and the lay public. Acknowledged experts provide highly informative yet critical reviews of the factors that most likely contribute to Alzheimer's, including genetics, metabolic deficiencies, oxidative stress, and possibly environmental exposures. Evidence that Alzheimer's resembles a brain form of diabetes is discussed from different perspectives, ranging from disease mechanisms to therapeutics. This book is further energized by discussions of how neurotransmitter deficits, neuro-inflammation, and oxidative stress impair neuronal plasticity and contribute to Alzheimer's neurodegeneration. The diversity of topics presented in just the right depth will interest clinicians and researchers alike. This book inspires confidence that effective treatments could be developed based upon the expanding list of potential therapeutic targets.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Cecília R A Santos, Isabel Cardoso and Isabel Gonçalves (2011). Key Enzymes and Proteins in Amyloid-Beta Production and Clearance, Alzheimer's Disease Pathogenesis-Core Concepts, Shifting Paradigms and Therapeutic Targets, Dr. Suzanne De La Monte (Ed.), ISBN: 978-953-307-690-4, InTech, Available from: <http://www.intechopen.com/books/alzheimer-s-disease-pathogenesis-core-concepts-shifting-paradigms-and-therapeutic-targets/key-enzymes-and-proteins-in-amyloid-beta-production-and-clearance>

INTECH
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2011 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the [Creative Commons Attribution-NonCommercial-ShareAlike-3.0 License](#), which permits use, distribution and reproduction for non-commercial purposes, provided the original is properly cited and derivative works building on this content are distributed under the same license.