

Psychopathology and Quality of Life in Children of Mentally Ill Parents

Silke Wiegand-Grefe, Susanne Halverscheid,
Franz Petermann and Angela Plass
*University Medical Center Hamburg-Eppendorf
Center for Clinical Psychology and Rehabilitation at Bremen University
Germany*

1. Introduction

Children of mentally ill parents are considered a high-risk population for the development of psychological disorders although the reported rates of affected children vary between studies. Whether or not a child will develop a mental illness himself depends on risk factors as well as protective factors. This chapter presents findings concerning the mental health status and the quality of life in children of mentally ill parents (N = 86). The results reveal an increased relative risk to develop psychopathological symptoms being 3 to 7 times higher than in children of the general population. At the same time, the quality of life in children with a mentally ill parent is substantially decreased. The psychological disturbances of the children are associated with the parents' perceived impairments caused by their mental illness. The results support the need for prevention programs for this special group of children.

2. Psychopathology in children of mentally ill parents

Children of mentally ill parents represent a high-risk population for the development of psychiatric disorders (Ramchandani & Psychogiou, 2009). The majority of studies examined the mental health status of children with a parent suffering from depression. The results reveal an increased symptomatology being 3 to 4 times higher compared to control samples (Weissman, Fendrich, Warner & Wickramaratne, 1992, 2005; Fergusson, Lynskey & Horwood, 1993; Gelfand & Teti, 1990). A study with children of parents suffering from diverse psychiatric disorders indicates a three- to sevenfold rate of psychopathology compared to the general population (Wiegand-Grefe, Geers, Plass, Petermann & Riedesser, 2009). Due to the psychiatric illness, psycho-social risk factors accumulate and determine each other (Mattejat, Lenz & Wiegand-Grefe 2011). Financial problems, resulting from an illness-related job loss, for example, constitute an additional burden on the family. In addition, the parent-child-interaction is often impaired, being highly associated with mental health problems in the children and a decreased quality of life. These risk configurations do not directly lead to developmental problems and psychiatric disturbances in the children, but rather interact with the children's vulnerability and resilience (Noeker & Petermann, 2008). The term resilience characterizes a hardiness against extreme stresses and strains that

may emerge within the children's environments. Resilient children overcome the sometimes potential traumatic conditions in a compensatory manner that allows them to surmount the adversities, displaying a healthy psychological functioning in the long term. Resilient children exhibit a balanced temper, high self-esteem, active coping mechanisms, good problem solving skills and an effective and competent social behavior. They have a secure attachment to at least one attachment figure and exhibit a clear value orientation (Noeker & Petermann, 2008). Resilient children are also very capable of showing empathy and expressing their own emotional states (Lenz & Kuhn, 2011). Which moderating factors can foster the children's mental health despite adverse living conditions is a central question for the development of effective prevention programs. In families with a mentally ill parent, the parental style of coping with the illness as well as reliable family relations are considered to be crucial protective factors for the children's development (Wiegand-Grefe et al., 2010a; Wiegand-Grefe, Halverscheid & Plass, 2011).

3. Quality of life in children of mentally ill parents

To date, little research has been conducted to assess the quality of life in children of mentally ill parents (Wiegand-Grefe et al. 2010b; Bullinger, 2011). Research in this area focuses rather on children's and adolescents' own somatic or psychiatric illness (Ravens-Sieberer & Bullinger, 1998; Ravens-Sieberer et al. 2006, 2008; Bullinger et al., 2008; Mattejat et al. 2005). Studies have shown that mentally ill children have a lower health-related quality of life (HRQL) than healthy or somatically ill children (Mattejat & Remschmidt, 2003; Ravens-Sieberer et al., 2008). In several Australian cities, N = 3597 children at the age of 6-17 years were assessed in their quality of life (Sawyer et al., 2002). The sample consisted of three psychiatric subsamples (ADHD, major depression and conduct disorder), two groups of children with physical illnesses, and a healthy control group of children. All children were assessed through parental evaluations. Children with mental health problems had decreased levels of quality of life in comparison to healthy children. In four out of five scales, they also displayed poorer rates in quality of life compared to the children with physical illnesses.

In a study with depressive and healthy children, aged 7 to 15 years, HRQL proved to be higher in healthy children (N = 1695) than in children suffering from depression (N = 248) (Kiss et al., 2009). In relation to the children's self-evaluations, maternal ratings of HRQL in depressive children was lower than in the children's self-assessments, while mothers of healthy children came up with a higher estimation of HRQL than the children themselves. The children's age and gender had no influence on the agreement between self-assessments and maternal ratings, neither was the mother's level of education influential.

In the representative German BELLA study with N = 1843 children and adolescents, adolescents with eating disorders showed lower HRQL scores than adolescents with normal eating habits (Herpertz-Dahlmann et al., 2008).

In a Dutch study, therapists rated the quality of life of 310 children and adolescents, aged 6 to 18 years, who were either healthy or suffered from one of five different psychiatric disorders (ADHD, anxiety disorders, pervasive developmental disorders, affective disorders or other diagnoses). Overall, children with pervasive developmental disorders were assessed to have the lowest HRQL (Bastiaansen et al., 2004).

Another study conducted by the Dutch research group assessed how quality of life changes relative to fluctuations in psychopathology. It was hypothesized that HRQL can increase

even when the severity of psychopathological symptoms does not change (Bastiaansen et al., 2005a). Thirty-three percent of the 126 children and adolescents aged 7 to 19 years showed improved rates in psychopathology as well as in their quality of life, 28 % improved in either one of the two areas, while 38 % of the children revealed no changes in both areas. However, 11 % of the children and adolescents showed increased rates in their HRQL within the given time frame, while the severity of their psychopathology did not change. Thus, psychotherapy should aim at improving patients' quality of life even though symptom reduction might not be possible.

In an outpatient-study, various factors besides psychopathology were assessed in their impact on the quality of life in 252 children and adolescents with mental health problems (Bastiaansen et al., 2005b). Within the examined child-related factors, decreased quality of life was associated with low self-esteem, poor social skills and comorbid somatic illnesses. Regarding the parent-related factors, maternal psychopathology and stress were linked with low HRQL in the children. In respect to environmental factors, a slight association between decreased quality of life was found in combination with a low socioeconomic status, little social support, poor family functioning, and stressful life events. The authors conclude that the treatment of psychological disturbances should be complemented by enhancing social competence, self-esteem, family functioning and social support.

Children who experience mental illnesses in their family are exposed to various stressors that are likely to influence their quality of life (Wiegand-Grefe et al., 2010b). Several studies of our work group document reduced levels of HRQL in affected children in contrast to comparative samples (Pollak et al., 2008; Jeske et al., 2009). Studies showed also relationships between children's HRQL and parental coping mechanisms (Jeske et al., 2009), family functioning (Pollak et al., 2008; Jeske et al., 2010) and parental attachment styles (Jeske et al., 2011).

This chapter presents the results of several studies examining psychopathology and quality of life in children of mentally ill parents. It is expected that the children are assessed by their parents to deal with more emotional and behavioral problems, the higher the parental subjective impairment is. It is also presumed that the quality of life in children of mentally ill parents is significantly lower than in children of the general population. Furthermore, it is expected that the HRQL is decreased with growing severity of parental symptomatology and that children's quality of life is reduced when the children suffer from emotional and behavioral problems themselves.

4. Study design

In a 9-month pilot study conducted within the project "CHIMPs - Children of mentally ill parents" at the University Medical Center Hamburg-Eppendorf, Germany, all patients referred to the Clinic of Psychiatry and Psychotherapy were registered from August 2005 to May 2006. For all patients, age, sex and diagnoses were recorded. Patients with minor children were examined further when they fulfilled the following inclusion criteria: aged 18-60 years, having at least one minor child between 0-18 years, receiving stationary treatment for at least five days, being sufficiently fluent in German language and giving informed consent for participating in the study. Exclusion criteria were: previous participation in the study in case of repeated hospitalization, a short duration of stationary treatment less than five days, and severe psychiatric or cognitive impairments. The patients were asked to answer questionnaires about their own mental health status and their children's situation.

When having several children, parents were asked to give only information on one randomly selected child in order to ensure the independence of observations.

5. Sample

Overall, 964 patients were registered. Among this sample, 167 (17%) patients had at least one minor child. Further 558 patients had no children (58 %), 104 patients had grown-up children above 18 years (11 %) and 135 patients were older than 60 years (14 %). Out of these 167 parents, 42 (25%) were not willing to participate in the study and further 39 parents (23 %) did not fulfill the inclusion criteria or had to be excluded from data analysis due to a large number of missing data. Overall, $N = 86$ parents answered the standardized questionnaires about the children's health-related quality of life, while a reduced number of $N = 67$ parents assessed the mental health status of their children.

The sample consists of 43 fathers and 43 mothers, aged between 23 and 59 years ($M = 41$ years, $SD = 7.9$). Forty-four patients are married (51 %), 19 parents are single or divorced (22 %), two parents are separated (2 %) and one patient is widowed (1 %). The most frequent school leaving certificate is a graduation from senior high-school (41 %), followed by graduations from intermediate secondary school (31%) and secondary general school (20%). Two parents are without school degree, 2 parents have another school leaving certificate. Regarding the qualifications, vocational trainings (39%) and University degrees (24%) are the most prevalent qualifications, 15% of the parents have no further qualification, 7 % hold another qualification. Thirty-one percent are white-collar employees, 20 % blue-collar workers, 16 % homemakers, 12 % self-employed, 3 % retired, and 2 % each are students or trainees.

Fifty-five parents (63 %) live together with their child, the remaining 31 patients (36 %) do not live with their children but have frequent contact to them (at least every two weeks). Here, a striking gender-specific effect can be observed: while 88% of the psychiatrically ill mothers live together with their children, only 53% of the fathers do so ($\chi^2=9.46$, $p=.002$). The children who are not living with their mentally ill parent live either with the other parent ($N = 21$, 68 %), with relatives ($N = 2$, 6 %), in foster- or adoption families ($N = 2$, 6 %), in institutional care ($N = 3$, 10 %) or in their own apartment. The examined children are aged between 4 to 18 years ($M = 11$ years, $SD = 4.49$), the distribution of boys and girls is equally balanced (50 % each).

As primary diagnosis according ICD-10 (WHO, 2000), 27 patients (32 %) are diagnosed with an affective disorder, 23 parents (26 %) suffer from either anxiety disorders, an obsessive-compulsive disorder or PTSD, further 20 parents (23 %) engage in substance abuse (mostly alcohol) and 14 patients (16 %) are affected by schizophrenia. Two parents have a personality disorder (PD), namely a Borderline PD and an emotional-unstable PD. Overall, 61% of the here examined parents have a comorbid psychiatric diagnosis.

6. Measurements

The mentally ill parents were asked to which extent they feel impaired by their illness through the German version of the Symptom-Checklist-14-R (SCL-14-R, Prinz et al., 2008). This self-report constitutes a short version of the SCL-90-R (Franke, 1995) and is used as a measure of general psychiatric symptomatology. It comprises 14 items on a 5-point Likert Scale, indicating the degree of symptomatology in the last seven days (not at all (0), a little

bit (1), moderately (2), quite a bit (3), extremely (4)). The SCL-14-R includes three subscales: Depressiveness, Somatoform Complaints and Phobic Anxiety. The total score is measured with the Global Severity Index (GSI). The GSI has proven to be a good indicator for the actual symptom severity.

The psychiatric diagnoses of the parents were given by the attending psychiatrists in accordance with the diagnostic criteria of the ICD-10 (WHO, 2005).

For the assessment of emotional and behavioral problems in the children and adolescents, parents were administered the Child Behavior Checklist/4-18 (Working Group German Child Behavior Checklist, 1994). On the CBCL, parents rate their children on 113 specific problem items using a 3-point scale (not true (0), somewhat/sometimes true (1), very/often true (2)). The resulting scales comprise a total problem score, two broadband scores (internalizing and externalizing problems) and eight syndrome scales (social withdrawal, somatic complaints, anxious/depressed, social problems, thought problems, attention problems, delinquent behavior, aggressive behavior).

The health-related quality of life of the children and adolescents was assessed by their parents with the KINDL^R questionnaire (Ravens-Sieberer & Bullinger, 2000). The KINDL^R comprehends 24 items on a 5-point Likert scale. The resulting subscales are: physical well-being, emotional well-being, self-esteem, family, friends, and everyday functioning (school or nursery/kindergarten). Psychometric results revealed a high degree of reliability (Cronbach's $\alpha \geq 0.70$ for most of the sub-scales and samples) and a satisfactory convergent validity.

7. Results

7.1 Parents' subjective impairment by the mental illness

All parents indicate to feel at least "a little bit" impaired by their illness. The Global Severity Index (GSI) lies around $M = 1.7$ ($SD = 0.79$). While "depressiveness" represents the greatest impact for the patients, phobic anxiety influences the here examined parents to a lesser degree. In comparison to the clinical norm sample (Prinz et al., 2008), the mentally ill parents display higher mean values on all scales, reaching significant differences in t-tests on the GSI, "depressiveness" and "phobic anxiety". On the scale "somatization", the parents differ only in comparison to the general population (Geers, 2006).

SCL-14-R Scales	Study Sample		Comparative clinical sample		General population	
	M	SD	M	SD	M	SD
Global Symptom Severity Index (GSI)	1.66	0.79	1.15	0.77	#	#
Depressiveness	2.26	1.05	1.49	1.05	0.45	0.58
Somatoform Complaints	1.38	1.07	1.20	1.01	0.46	0.63
Phobic Anxiety	1.05	1.13	0.59	0.88	0.17	0.39

Table 1. SCL-14-R values of the study sample ($N=62$), a comparative clinical sample and a representative sample of the general population (from Wiegand-Grefe, Geers, Plass, Petermann & Riedesser, 2009).

Annotation: M = mean value, SD = standard deviation, # = not specified.

7.2 Psychopathology in children of mentally ill parents

The children's emotional and behavioral problems, assessed through the CBCL scales, are listed in table 2 as T-values, taking age- and gender-specific aspects of emotional and behavioral development into account.

CBCL Scales	M	SD	Minimum	Maximum	N
Social Withdrawal	58.11	9.08	50	88	61
Somatic Complaints	57.62	9.32	50	81	60
Anxious/Depressed	58.90	9.17	50	83	60
Social Problems	58.00	8.29	50	84	58
Thought Problems	56.37	9.36	50	80	60
Attention Problems	58.90	8.98	50	83	60
Delinquent Behavior	57.33	6.95	50	75	61
Aggressive Behavior	59.31	9.68	50	95	62
Internalizing Problems	57.33	11.83	38	82	61
Externalizing Problems	57.58	10.42	35	84	62
Total	58.92	11.47	38	86	61

Table 2. Distribution of T-Values of the CBCL subscales and main scales (N=62) (from Wiegand-Grefe et al., 2009). Annotation: M = mean value, SD = standard deviation, N = sample size.

In order to assess how many children display psychological problems in the clinical ($T \geq 64$), subclinical (T between 60-63) or normal range ($T \leq 59$), the T-scores have been categorized accordingly. On the syndrome scales, 14 % of the children and adolescents are seen to have somatic complaints, 13 % suffer from clinical or subclinical levels of anxiety and depression. On the main CBCL scales, 31 % of the parents report internalizing problems and further 9 % see subclinical internalizing problems. Regarding externalizing problems, 29 % of the parents report behavioral problems in the clinical range, another 16 % see externalizing symptomatology in the subclinical range. Overall, 40 % of the parents report at least subclinical problems in the internalizing domain, and 45 % of the parents see externalizing behavior to an at least subclinical degree. Considering the total problem score, 52 % of the children are rated to lie within the normal range of emotional and behavioral difficulties, while 32 % are seen to display subclinical symptoms and 14 % to exhibit clinically relevant symptomatology. Sixty-six percent of the examined children are reported not to display any clinically relevant symptoms on the syndrome scales. The remaining 32 % of children and adolescents suffer from clinically relevant symptoms on at least one dimension measured by the CBCL.

When comparing the relative frequencies of children of mentally ill parents, lying in the subclinical or clinical range on the CBCL syndrome scales, with those of the general child population (with the top 5 % constituting the subclinical range, and the top 2 % defining the clinical range), it can be stated that on all syndrome scales, children of mentally ill parents display 1.5- to 7-times heightened rates of clinically relevant symptomatology (table 3). It is striking that the children are especially prone to suffer from somatic symptoms (relative risk, $RR = 7.26$). Besides, they have a 6-fold risk to display thought and attention problems ($RR = 6.45$) and are five times more likely to engage in social withdrawal in comparison to the general child population ($RR = 5.65$).

CBCL Subscales	Subclinical and clinical range		Clinical range	
	Reference value in the general population: 5 %		Reference value in the general population: 2 %	
	%	relative risk	%	relative risk
Social Withdrawal	16.13	3.23	11.29	5.65
Somatic Complaints	19.35	3.87	14.52	7.26
Anxious/Depressed	24.19	4.84	12.90	6.45
Social Problems	17.74	3.55	6.45	3.23
Thought Problems	20.97	4.19	12.90	6.45
Attention Problems	17.74	3.55	12.90	6.45
Delinquent Behavior	14.52	2.90	3.23	1.61
Aggressive Behavior	24.19	4.84	9.68	4.84

Table 3. Relative frequencies and relative risk in children of mentally ill parents (N = 62) displaying CBCL scores in the subclinical or clinical range in comparison with children of the general population lying in the clinical range (top 2 %) and subclinical/clinical range (top 5 %) (from Wiegand-Grefe et al., 2009).

7.3 Parents' subjective impairments and the psychological difficulties of their children

Since a positive linear association is expected between the parents' subjective impairments and the children's psychopathology, SCL-14-R and CBCL scale values are correlated (table 4). Significant positive correlations exist between the parent's global severity index (GSI), the SCL-14-R dimensions somatization and phobic anxiety, and all main scales of the CBCL. The correlative association between parental "somatization" and the children's "internalizing problems" represents the highest association with $r = .302$ indicating a small to medium sized linkage between the variables. No association has been found between parental "depressiveness" and psychopathology in children.

Mean Values of SCL-14-R-Scales		Internalizing CBCL-Scale (T-values)		Externalizing CBCL-Scale (T-values)		CBCL Total Score (T-values)	
		r	*	r	*	r	*
Global Symptom Severity Index (GSI)	r	.297	*	.275	*	.287	*
	p	.010		.015		.012	
	N	61		62		61	
Depressiveness	r	.204		.176		.182	
	p	.06		.09		.08	
	N	61		62		61	
Somatoform Complaints	r	.302	**	.233	*	.264	*
	p	.009		.03		.02	
	N	61		62		61	
Phobic Anxiety	ρ	.221	*	.279	*	.299	**
	p	.043		.014		.010	
	N	61		62		61	

Table 4. Correlative association between the SCL-14-R-Scales and the main CBCL-Scales (N=62) (from Wiegand-Grefe et al., 2009). Annotation: r = Pearson product-moment correlation coefficient; ρ = Spearman rank correlation coefficient; p= level of significance; * p <.05; ** p <.01; N = sample size.

7.4 Health-related quality of life in the children

The total score of the KINDL^R was transformed with “0” indicating the poorest and “100” indicating the maximum value of health-related quality of life (HRQL). In the examined population of children of mentally ill parents, the total score of HRQL lies by 70.91 (SD = 12.88). The children’s overall score on the dimension physical well-being lies by 71.01 (SD = 18.64), the “emotional well-being” lies around 71.28 (SD = 18.55). The KINDL^R subscales range between 67.50 on the lowest end (self-esteem, SD = 17.22) and 74.26 constituting the maximum value (friends, SD = 13.77). The scales family (M = 68.51, SD = 19.33) and school (M = 70.00, SD = 18.24) rank in between. In comparison to the reference sample of the general population, the here examined children of mentally ill parents display low values in the total HRQL-score that can be explained and specified by poor results on the scales emotional well-being, family and school (Jeske et al., 2010, Wiegand-Grefe et al., 2010). When comparing the ratings of mentally ill mothers and fathers, it can be stated that no differences were found in their assessment of the HRQL in their children.

7.5 Subjective parental impairment and quality of life in the children

Slight negative associations have been found between the subjective parental impairment, as measured by the SCL-14-R, and the children’s HRQL dimensions emotional well-being, self-esteem, family, school and the overall total score (table 5).

Mean Values of SCL-14-R-Scales		KINDL ^R Total Score	Physical well-being	Emotional well-being	Self-esteem	Family	Friends	School
Global Symptom Severity Index (GSI)	r	-.274		-.189		-.267		-.253
	ρ		-.149		-.259		-.038	
	p	.008**	.094	.048*	.012*	.009**	.373	.017*
	N	77	80	79	75	78	76	70
Depressive-ness	r	-.288		-.216		-.289		-.305
	ρ		-.179		-.160		-.027	
	p	.006**	.059	.028*	.086	.005**	.408	.005**
	N	77	80	79	75	75	76	70
Somatoform Complaints	r	-.261		-.157		-.248		-.207
	ρ		-.169		-.325		.049	
	p	.011*	.068	.084	.002**	.014*	.337	.043*
	N	77	80	79	75	75	76	70
Phobic Anxiety	r							
	ρ	-.081	.010	-.055	-.164	-.047	-.040	-.133
	p	.243	.465	.315	.080	.340	.365	.137
	N	77	80	79	75	78	76	70

Table 5. Correlative association between SCL-14-R and KINDL^R (N=62) (from Wiegand-Grefe et al. 2010). Annotation: r = Pearson product-moment correlation coefficient, ρ = Spearman rank correlation coefficient; p= level of significance; * p <.05; ** p <.01 (1-sided); N = sample size.

Further significant correlations exist between parental „depressiveness“ and the children’s HRQL in the total score and the areas of school functioning, family, and emotional well-

being. Somatoform complaints of the parents are associated with poor “self-esteem” and decreased values on the dimensions school, family and the total score. No linear association was found between the HRQL in children and parental “phobic anxiety”.

7.6 Psychopathology and health-related quality of life in the children

When regarding the association between the children’s psychopathology and their health-related quality of life, considerable negative associations can be found between the main CBCL scales and all KINDL^R dimensions. According to the results, parents see substantial losses in their children’s HRQL with increasing emotional and behavioral problems (table 6).

CBCL Scales (T-values)		KINDL ^R Total Score	Physical Well-being	Emotional Well-being	Self-esteem	Family	Friends	School
Total Problem Score	r	-.611		-.522		-.347		-.442
	ρ		-.405		-.597		-.291	
	p	.000***	.000***	.000***	.000***	.002**	.009**	.000***
	N	67	67	66	64	66	66	65
Externalizing Problems	r	-.492		-.445		-.396		-.296
	ρ		-.320		-.453		-.216	
	p	.000***	.004**	.000***	.000***	.000***	.040*	.008**
	N	66	67	66	64	66	66	65
Internalizing Problems	r	-.694		-.667		-.272		-.561
	ρ		-.561		-.744		-.278	
	p	.000***	.000***	.000***	.000***	.014*	.012*	.000***
	N	66	66	66	64	65	65	66

Table 6. Correlative association between CBCL and KINDL^R (N=67) (from Wiegand-Grefe et al., 2010) Annotation: r = Pearson product-moment correlation coefficient, ρ = Spearman rank correlation coefficient; p = level of significance; * p <.05; ** p <.01; N = sample size

8. Discussion

In comparison to the general population, children of mentally ill parents have a substantial risk to develop psychopathological symptomatology. On the main scales and the syndrome scales of the CBCL, the children’s relative risk to develop clinically relevant symptomatology is three- to sevenfold. In the existing literature, children of mentally ill parents are reported to be assessed up to five times more likely to exhibit psychological difficulties (Kölch, Schielke, Becker, Fegert & Schmid, 2008). In the study at hand, 32 % of the parents see clinically relevant symptoms in their children. When including the cases of subclinical symptomatology, even 47 % of the examined children are regarded to be somehow affected by emotional and behavioral problems. This rate goes along with reported frequencies in comparable studies (Beidel & Turner, 1997; Hill, Locke, Lowers & Connolly, 1999; Kelley & Fals-Stewart, 2004; Lapalme, Hodgins & La Roche, 1997; Merikangas, Dierker & Szatmari, 1998). It may be argued that the percentage of affected children might even be higher considering the fact that many parents undergo feelings of guilt and shame and therefore tend to minimize difficulties in clinical evaluations. It is

striking, for example, that adolescents often report more difficulties in self-reports than the parents indicate in the according ratings (Najman et al., 2000). Thus, it might be expected that adolescents with a mentally ill parent might even exhibit higher rates of symptomatology than in the examined parental assessments.

In respect to the characteristics of the parental illness, positive associations were found between the severity of the illness, parental anxiety and somatic complaints and children's psychopathology. This result is analogous to other studies showing that the severity of parental symptomatology goes along with increased psychopathology in the children (Brennan et al., 2000; Hammen, Burge, Burney & Adrian, 1990; Keller et al., 1986; Weissman et al., 2005). It is rather astonishing that, contrary to other studies (Mattejat et al., 2000), no association was found between depressive symptoms in parents and emotional and behavioral difficulties in the here examined population of children. It can be assumed that parents with phobic anxieties and somatic complaints are especially sensitive for their children's difficulties. It is interesting that parental somatization is related to psychological problems in children on all CBCL main scales. This finding underlines that somatic complaints and illnesses in parents stand in a close relationship with children's mental health (Romer & Hagen), although somatizations have to be distinguished from severe somatic illnesses. Since patients with somatizations and anxieties tend to be over-sensitive towards potential difficulties, the results, indicating increased symptomatology in the children, should be cautiously interpreted.

In the present study, children with a mentally ill parent display a decreased quality of life in comparison to the general population. The overall quality of life in children is especially associated with the global severity of parental symptomatology, as well as parental somatization and depressiveness in specific. With increasing degree of perceived impairment, parents rated their children's HRQL to be poorer. In a previous study, it was found that children, who were exposed to a parental mental illness for a longer time frame, exhibited higher quality of life scores on the KINDL^R scale "friends". Presumably, these children might have mastered their situation by compensating their familial difficulties with peer group activities.

The psychopathology of the examined children is substantially linked to the quality of life, as indicated by high correlations between the KINDL^R and the CBCL scales.

In respect to methodological constrictions, it has to be underlined that the assessment of the children's psychopathology and quality of life is solely based on the parents' view. The informative validity of assessments by mentally ill parents has not yet been explored sufficiently. Hitherto, studies have examined response patterns in parents with depression and anxiety disorders. Overall, the findings do not allow drawing specific conclusions in respect to potential response biases. In general, the most credible source for internalizing disorders are self-assessments of children and adolescents, while externalizing problems are best assessed by parents (Fombonne, 2002). In the consequent CHIMPs project, families will be examined as a whole, which will allow contrasting the evaluations of mentally ill parents and partners with the children's self-reports. The suggestion to include teacher reports (Thiels & Schmitz, 2008) would certainly be promising and beneficial, but forms an obstacle in the shame-related area of parental mental illness.

9. Implications for clinical practice

The present chapter deals with the mental health status and the quality of life of children with a mentally ill parent. The impairments resulting from the mental illness are associated

with the children's mental health and their quality of life. The heightened rates of psychopathology in the children underline the need to implement appropriate prevention programs. In the development of such preventive measures, the improvement of the children's quality of life should constitute a central component next to the reduction of psychopathological symptoms.


10. References

- Achenbach, T. M. (1991). Manual for the Child Behavior Checklist / 4-18 and 1991 Profile. Burlington: University of Vermont, Department of Psychiatry.
- Arbeitsgruppe Deutsche Child Behavior Checklist (1994). CBCL/4-18. Elternfragebogen über das Verhalten von Kindern und Jugendlichen. Deutsche Bearbeitung der Child Behavior Checklist. Köln: Arbeitsgruppe Kinder-, Jugend- und Familiendiagnostik (KJFD).
- Bastiaansen, D., Koot, H.M., Ferdinand, R.F. & Verhulst, F.C. (2004). Quality of life in children with psychiatric disorders: Self-, parent and clinician report. *American Academy of Child and Adolescent Psychiatry*, 43, 221-230.
- Bastiaansen, D., Koot, H.M. & Ferdinand, R.F. (2005a). Psychopathology in children: Improvement of quality of life without psychiatric symptom reduction? *European Child and Adolescent Psychiatry*, 14, 364-370.
- Bastiaansen, D., Koot, H.M. & Ferdinand, R.F. (2005b). Determinants of quality of life in children with psychiatric disorders. *Quality of Life Research*, 14, 1599-1612.
- Beidel, D.C. & Turner, S.M. (1997). At risk for anxiety. I. Psychopathology in the offspring of anxious parents. *Journal of the American Academy of Child and Adolescent Psychiatry*, 36, 918-924.
- Brennan, P.A., Hammen, C., Andersen, M.J., Bor, W., Najman, J.M. & Williams, G.M. (2000). Chronicity, severity and timing of maternal depressive symptoms: Relationship with child outcomes at age 5. *Developmental Psychology*, 36, 759-766.
- Bullinger, M. (2011) Lebensqualität von Kindern und Jugendlichen im Kontext der Gesundheit ihrer Eltern. In: S. Wiegand-Grefe, F. Matthejat & A. Lenz (Hrsg.). *Kinder mit psychisch kranken Eltern. Klinik und Forschung*. (S. 401-415). Göttingen: Vandenhoeck & Ruprecht.
- Fergusson, M., Lynskey, M.T. & Horwood, L.J. (1993). The effect of maternal depression on maternal ratings of child behavior. *Journal of Abnormal Child Psychology*, 21, 245-269.
- Fombonne, E. (2002). Case identification in an epidemiological context. In M. Rutter & E. Taylor (Eds.), *Child and adolescent psychiatry* (4th ed., pp. 52-86). Oxford: Blackwell.
- Franke, G. H. (1995). SCL-90-R. Die Symptomcheckliste von Derogatis - Deutsche Version - Manual. Göttingen: Hogrefe
- Geers, P. (2006). Psychische Gesundheit der Kinder von Eltern mit psychischen Störungen. Bremen: Unveröffentlichte Diplomarbeit, Studiengang Psychologie, Universität Bremen.
- Gelfand, D.M. & Teti, D.M. (1990). The effects of maternal depression on children. *Clinical Psychology Review*, 10, 329-353.

- Goodman, S.H. & Gotlib, I.H. (1999). Risk for psychopathology in the children of depressed mothers. A developmental model for understanding mechanisms of transmission. *Psychological Review*, 106, 458-490.
- Hammen, C., Burge, D., Burney, E. & Adrian, C. (1990). Longitudinal study of diagnoses in children of woman with unipolar and bipolar affective disorder. *Archives of General Psychiatry*, 47, 1112-1117.
- Herpertz-Dahlmann, B., Wille, N., Hölling, H., Vloet, T.D., Ravens-Sieberer, U. & BELLA study group (2008). Disordered eating behavior and attitudes, associated psychopathology and health-related quality of life: results of the BELLA study. *European Child and Adolescent Psychiatry*, 17, 82-91.
- Hill, S.Y., Locke, J., Lowers, L. & Connolly, J. (1999). Psychopathology and Achievement in children at high risk for developing alcoholism. *Journal of the American Academy of Child and Adolescent Psychiatry*, 38, 883-891.
- ICD-10-GM Version (2005) Internationale statistische Klassifikation der Krankheiten und verwandter Gesundheitsprobleme (ICD)- German Modification. Band 1: Systematisches Verzeichnis. Kooperationsausgabe. Stuttgart: Kohlhammer. Deutsche Krankenhaus Verlagsgesellschaft mbH
- Jeske, J., Bullinger, M., Plass, A., Petermann, F. & Wiegand-Grefe, S. (2009). Risikofaktor Krankheitsverarbeitung. Zusammenhänge zwischen der Krankheitsverarbeitung einer elterlichen psychischen Erkrankung und der gesundheitsbezogenen Lebensqualität der Kinder. *Zeitschrift für Psychiatrie, Psychologie und Psychotherapie*, 57, 207-213
- Jeske, J., Bullinger, M. & Wiegand-Grefe, S. (2010). Familien mit psychisch kranken Eltern. Zusammenhang von Familienfunktionalität und gesundheitsbezogener Lebensqualität der Kinder. *Familiendynamik* 35: 338-347.
- Jeske, J., Bullinger, M. & Wiegand-Grefe, S. (2011). Do attachment patterns of parents with a mental illness have an impact upon how they view the quality of life of their children? *Vulnerable Children and Youth Studies* 6, 39-50.
- Keller, M.B., Beardslee, W.R., Dorer, D.J., Lavori, P.W., Samuelson, H. & Klerman, G.R. (1986). Impact of severity and chronicity of parental affective illness on adaptive functioning and psychopathology in children. *Archives of General Psychiatry*, 43, 930-937.
- Kelley, M.L. & Fals-Stewart, W. (2004). Psychiatric disorders of children living with drug-abusing, alcohol-abusing, and non-substance-abusing fathers. *Journal of the American Academy of Child and Adolescent Psychiatry*, 43, 621-628.
- Kiss, E., Kapornai, K., Baji, I., Mayer, L. & Vetró, Á. (2009). Assessing quality of life: mother-child agreement in depressed Hungarian. *European Child and Adolescent Psychiatry*, 18, 265-273.
- Kölch, M., Schielke, A., Becker, T., Fegert, J.M. & Schmid, M. (2008). Belastung Minderjähriger aus Sicht der psychisch kranken Eltern. Ergebnisse einer Befragung stationär behandelter Patienten mit dem SDQ. *Nervenheilkunde*, 27, 527-532.
- Lapalme, M., Hodgins, S. & La Roche, C. (1997). Children of parents with bipolar disorder : A metaanalysis of risk for mental disorders. *Canadian Journal of Psychiatry*, 42, 623-631.
- Lee, C. & Gotlib, I.H. (1991). Adjustment of Children of depressed mothers: A 10-month follow-up. *Journal of Abnormal Psychology*, 100, 473-477.

- Lenz, A., Kuhn, J. (2011). Was stärkt Kinder psychisch kranker Eltern und fördert ihre Entwicklung? Überblick über Ergebnisse der Resilienz- und Copingforschung. In: S. Wiegand-Grefe, F. Mattejat & A. Lenz (Hrsg.). *Kinder mit psychisch kranken Eltern. Klinik und Forschung*. (S. 269-298). Göttingen: Vandenhoeck & Ruprecht.
- Mattejat, F., Remschmidt, H. (2003) Zur Erfassung der Lebensqualität bei psychisch gestörten Kindern und Jugendlichen- Eine Übersicht. (Online), 30/07/2011, Available from <http://www.kjp.uni-marburg.de/lq/index.php?include=publikat>
- Mattejat, F., Wüthrich, C. & Remschmidt, H. (2000). Kinder psychisch kranker Eltern. Forschungsperspektiven am Beispiel von Kindern depressiver Eltern. *Nervenarzt*, 71, 164-172.
- Mattejat, F., Lenz, A. & Wiegand-Grefe, S. (2011). Kinder psychisch kranker Eltern - Eine Einführung in die Thematik. In: S. Wiegand-Grefe, A- Lenz & F. Mattejat (Hrsg.) *Kinder mit psychisch kranken Eltern. Klinik und Forschung*. (S. 13-24). Göttingen: Vandenhoeck und Ruprecht.
- Mattejat, F., Koenig, U., Barchewitz, C., Felbel, D., Herpertz-Dahlmann, B., Hoehne, D., Janthur, B., Jungmann, J., Katzenski, B., Kirchner, J., Naumann, A., Nölkel, P., Schaff, C., Schulz, E., Warnke, A., Wienand, F. & Remschmidt, H. (2005). Zur Lebensqualität von psychisch kranken Kindern und ihren Eltern: Ergebnisse der ersten multizentrischen Studie mit der Elternversion des Inventars zur Erfassung der Lebensqualität bei Kindern und Jugendlichen (ILK). *Kindheit und Entwicklung*, 14, 39-47
- Merikangas, K.R., Dierker, L.C. & Szatmari, P. (1998). Psychopathology among offspring of parents with substance abuse and/or anxiety disorders: A high-risk study. *The Journal of Child Psychology and Psychiatry*, 39, 711-720.
- Najman, J.M., Williams, G.M., Nikles, J., Spence, S., Bor, W., O'Callaghan, M., LeBroque, R. & Andersen, M.J. (2000). Mothers' mental illness and child behavior problems: cause-effect association or observation bias? *Journal of the American Academy of Child and Adolescent Psychiatry*, 39, 592-602.
- Noeker, M. & Petermann, F. (2008). Resilienz: Funktionale Adaptation an widrige Umgebungsbedingungen. *Zeitschrift für Psychiatrie, Psychologie und Psychotherapie*, 56, 255-263.
- Pollak, E., Bullinger, M. Jeske, J. Wiegand-Grefe, S (2008): How do mentally ill parents evaluate their childrens quality of life? Associations with the parent's illness and family functioning. *Praxis der Kinderpsychologie und Kinderpsychiatrie*, 57, 301-314.
- Prinz, U., Nutzinger, D.O., Schulz, H., Petermann, F. Braukhaus, C. & Andreas, S. (2008). Die Symptom-Checkliste-90-R und ihre Kurzversion: Psychometrische Analysen bei Patienten mit psychischen Krankheiten. *Physikalische Medizin, Rehabilitationsmedizin, Kurortmedizin*, 18, 337-343.
- Ramchandani, P. & Psychogiou, L. (2009). Paternal psychiatric disorders and children's psychosocial development. *Lancet Journal*, 374, 646-653
- Ravens-Sieberer, U. & Bullinger, M. (1998). Assessing health-related quality of life in chronically ill children with the German KINDL: first psychometric and content analytical results. *Quality of Life Research*, 7, 399-407
- Ravens-Sieberer, U. & Bullinger, M. (2000). KINDL-R. Fragebogen zur Erfassung der gesundheitsbezogenen Lebensqualität bei Kindern und Jugendlichen. Revidierte Form. Manual. (Online), 30/07/2011, Available from <http://www.kindl.org/daten/pdf/ManGerman.pdf>

- Ravens-Sieberer, U., Erhart, M., Wille, N., Wetzel, R., Nickel, J. & Bullinger, M. (2006). Generic Health-Related Quality-of-Life Assessment in Children and Adolescents. *Pharmacoeconomics*, 24, 1199-1220
- Ravens-Sieberer, U., Erhart, M., Wille, N., Bullinger, M. & BELLA study group (2008). Health-related quality of life in children and adolescents in Germany: results of the BELLA study. *European Child and Adolescent Psychiatry*, 17, 148-156.
- Romer, G. & Haagen, M. (2007). *Kinder körperlich kranker Eltern*. Göttingen: Hogrefe.
- Sawyer, M.G., Whaites, L., Rey, J.M., Hazell, P.L., Graetz, B.W. & Baghurst, P. (2002). Health-related quality of life of children and adolescents with mental disorders. *Journal of the American Academy of Child and Adolescent Psychiatry*, 41, 530-537.
- Thiels, C. & Schmitz, G.S. (2008). Selbst- und Fremdbeurteilung von Verhaltensauffälligkeiten bei Kindern und Jugendlichen. Zur Validität von Eltern- und Lehrerurteilen. *Kindheit und Entwicklung*, 17, 118-125.
- Weissman M.M., Leckman, J.F., Merikangas, K.R., Gammon, G.D. & Prusoff, B.A. (1984). Depression and Anxiety Disorders in Parents and Children. Results From the Yale Family Study. *Archives of General Psychiatry*, 41, 845-852.
- Weissman, M.M., Wickramaratne, P., Nomura, Y., Warner, V., Verdeli, H., Pilowsky, D.J., Grillon, C. & Bruder, G. (2005). Families at high and low risk for depression. A 3-Generation Study. *Archives of General Psychiatry*, 62, 29-36.
- Wiegand-Grefe, S., Geers, P., Halverscheid, S., Petermann, F. & Plass, A. (2010a). Kinder psychisch kranker Eltern. Zusammenhänge zwischen der Krankheitsbewältigung einer elterlichen psychischen Erkrankung und der Gesundheit der Kinder. *Zeitschrift für Klinische Psychologie und Psychotherapie*, 39, 13-23.
- Wiegand-Grefe, S., Geers, P., Petermann, F. & Plass, A. (2011). Kinder psychisch kranker Eltern: Merkmale elterlicher psychiatrischer Erkrankung und Gesundheit der Kinder aus Elternsicht. *Fortschritte der Neurologie und Psychiatrie*, 79, 32-40.
- Wiegand-Grefe, S., Geers, P., Plass, A., Petermann, F. & Riedesser, P. (2009). Kinder psychisch kranker Eltern. Zusammenhänge zwischen subjektiver elterlicher Beeinträchtigung und psychischer Auffälligkeit der Kinder aus Elternsicht. *Kindheit und Entwicklung*, 18, 111-121.
- Wiegand-Grefe, S., Halverscheid, S. & Plass, A. (2011). Kinder und ihre psychisch kranken Eltern. Familienorientierte Prävention – Der CHIMPs-Beratungsansatz. Göttingen: Hogrefe.
- Wiegand-Grefe, S., Jeske, J., Bullinger, M., Plass, A. & Petermann, F. (2010b). Lebensqualität von Kindern psychisch kranker Eltern. Zusammenhänge zwischen Merkmalen elterlicher Erkrankung und gesundheitsbezogener Lebensqualität der Kinder aus Elternsicht. *Zeitschrift für Psychiatrie, Psychologie, und Psychotherapie*, 58, 315-322.
- Wolke, D. (2008). Von Null bis Drei: Entwicklungsrisiken und Entwicklungsabweichungen. In F. Petermann (Hrsg.), *Lehrbuch der Klinischen Kinderpsychologie* (6., völlig veränd. Aufl., S.65-80. Göttingen: Hogrefe.
- Ziegenhain, U., Derksen, B. & Dreisörner, R. (2004). Frühe Förderung von Resilienz bei jungen Müttern und ihren Säuglingen. *Kindheit und Entwicklung*, 13, 226-234.



Mental Illnesses - Evaluation, Treatments and Implications

Edited by Prof. Luciano LAbate

ISBN 978-953-307-645-4

Hard cover, 476 pages

Publisher InTech

Published online 13, January, 2012

Published in print edition January, 2012

In the book "Mental Illnesses - Evaluation, Treatments and Implications" attention is focused on background factors underlying mental illness. It is crucial that mental illness be evaluated thoroughly if we want to understand its nature, predict its long-term outcome, and treat it with specific rather than generic treatment, such as pharmacotherapy for instance. Additionally, community-wide and cognitive-behavioral approaches need to be combined to decrease the severity of symptoms of mental illness. Unfortunately, those who should profit the most by combination of treatments, often times refuse treatment or show poor adherence to treatment maintenance. Most importantly, what are the implications of the above for the mental health community? Mental illness cannot be treated with one single form of treatment. Combined individual, community, and socially-oriented treatments, including recent distance-writing technologies will hopefully allow a more integrated approach to decrease mental illness world-wide.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Silke Wiegand-Grefe, Susanne Halverscheid, Franz Petermann and Angela Plass (2012). Psychopathology and Quality of Life in Children of Mentally Ill Parents, *Mental Illnesses - Evaluation, Treatments and Implications*, Prof. Luciano LAbate (Ed.), ISBN: 978-953-307-645-4, InTech, Available from: <http://www.intechopen.com/books/mental-illnesses-evaluation-treatments-and-implications/psychopathology-and-quality-of-life-in-children-of-mentally-ill-parents>

INTECH

open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2012 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.