

Coumarin-Derived Fluorescent Chemosensors

Hongqi Li, Li Cai and Zhen Chen
College of Chemistry,
Chemical Engineering and Biotechnology,
Donghua University, Shanghai,
China

1. Introduction

Fluorescent chemosensors are highly valuable in a variety of fields including environmental chemistry, analytical chemistry, and bio-medicinal science. They have provided accurate, on-line, and low-cost detection of toxic heavy metal ions, anions, and enzymes with high selectivity and sensitivity. Coumarins, with the structure of benzopyrone, have many advantages including high fluorescence quantum yield, large Stokes shift, excellent light stability, and less toxicity. Therefore coumarins have been widely used in the fields of biology, medicine, perfumes, cosmetics, and fluorescent dyes. By far coumarin derivatives have been used as fluorescent probes of pH, for detection of nitric oxide, nitroxide, and hydrogen peroxide. Moreover, coumarin derivatives have served as good chemosensors of anions including cyanide, fluoride, pyrophosphate, acetate, benzoate, and dihydrogenphosphate as well as various metal ions comprised of Hg(II), Cu(II), Zn(II), Ni(II), Ca(II), Pb(II), Mg(II), Fe(III), Al(III), Cr(III), and Ag(I). Several systems containing coumarin exhibited simultaneous sensitivity toward two or more different metal ions, e.g. Ca(II) and Mg(II), Ni(II) and Co(II), Cu(II) and Hg(II), Na(I) and K(I), Cu(II) and Ni(II), Hg(II) and Ag(I), Cu(II)/Ni(II)/Cd(II), Zn(II)/Cd(II)/Pb(II), or Ni(II)/Pd(II)/Ag(I). Herein a brief review of fluorescent chemical sensors derived from coumarins is presented.

2. Structural characteristics of coumarin

The fusion of a pyrone ring with a benzene ring gives rise to a class of heterocyclic compounds known as benzopyrones, of which two distinct types are recognized, namely benzo- α -pyrones, commonly called coumarins, and benzo- γ -pyrones, called chromones, the latter differing from the former only in the position of the carbonyl group in the heterocyclic ring as shown in Fig. 1 (Sethna & Shah, 1945). It is well known that stilbene with a *trans* conformation is highly fluorescent. From the viewpoint of molecular structure, coumarins bear a carbon-carbon double bond which is fixed as *trans* conformation as in *trans*-stilbene through a lactone structure. This can help to avoid the *trans-cis* transformation of the double bond under ultraviolet (UV) irradiation as observed in stilbene compounds and results in strong fluorescence and high fluorescence quantum yield and photostability in most of coumarin derivatives.




Fig. 1. Structures and numbering scheme of coumarin and related compounds

It was shown in the late 1950s that substitutions on the coumarin structure shifted the fluorescence band. For instance, adding a methyl group to the 4-position of 7-hydroxy- or 7-methoxycoumarin red shifts the fluorescence spectra. Addition of electron-repelling groups in the 4-, 6-, or 7-position or electron-attracting groups in the 3-position all shifts the fluorescence band to longer wavelengths. When the carbonyl is substituted with a thione, the absorbance was red shifted and the fluorescence was quenched (Trenor et al., 2004). The reduction of the acceptor strength of the substituent at the 3-position, as in a 7-diethylaminocoumarin dye **1** (the structure of which is shown in Fig. 1) does not always result in sustained fluorescence in polar, aprotic solvents. Thus 7-diethylamino-3-styrylcoumarin dyes are not technically important as laser dyes; however, the extreme sensitivity of coumarin **1** to the medium polarity could provide an opportunity to probe the microenvironment experienced by the molecule (Bangar Raju & Varadarajan, 1995).

Changing the solvent or the solution pH also affected the fluorescence spectra. Study on the effect of solution pH on 7-hydroxy-4-methylcoumarin showed that increasing the solution pH raised the fluorescence intensity. Studies on the effect of changing the solvent polarity on 13 coumarin derivatives revealed that increasing solvent polarity red shifted the absorbance as well as red shifted and broadened the emission of the coumarins due to increased hydrogen bonding. Studies on the excited-state properties of 4- and 7-substituted coumarin derivatives revealed that solvent polarity shifted both the emission and absorption peaks, with a greater shift observed in the emission spectra, indicating that the excited-state dipole moment of the solute molecule was greater than the ground-state dipole moment (Trenor et al., 2004).

3. Coumarin-derived fluorescent chemosensors

3.1 Coumarin-derived fluorescent chemosensors for metal ions

Considering the threat of mercury to the environment and human health, a great effort has been devoted to the utilization of fluorescent methods for detection of Hg^{2+} ions. More than ten coumarin-derived fluorescent chemosensors for Hg^{2+} ions have been reported. The structures and references of these chemical sensors are listed in Table 1.

The recognition mechanisms of these chemosensors mainly involve photoinduced electron transfer (PET), intramolecular charge transfer (ICT), fluorescence resonance energy transfer (FRET), coordination, and desulfurization. For instance, the fluorescence detection of **2** upon Hg^{2+} addition is promoted by a Hg^{2+} -induced desulfurization of the thiourea moiety, leading to a decrease in an ICT character of the excited-state coumarin moiety (Shiraishi et al., 2010). Coumarin-thiazolobenzo-crown ether based chemosensor **3** has been developed for $\text{Hg}(\text{II})$ ions that utilizes the strong coordination of $\text{Hg}(\text{II})$ ions on the crown oxygen and thiazole nitrogen. The complexation of $\text{Hg}(\text{II})$ disrupts the ICT from the oxygen donor to the

















Structure of sensor	Reference	Structure of sensor	Reference
	Shiraishi et al., 2010		W. Ma et al., 2010
	S. H. Lee et al., 2010	 10: R = acetylene  11: R = 1,3-dithiane-2-yl 	D.-N. Lee et al., 2009 J. H. Kim et al., 2009
	Wang et al., 2006		M. G. Choi et al., 2009
	Ryu et al., 2010		H. J. Kim et al., 2010
	H. J. Kim et al., 2009a		Voutsadakis et al., 2010
	Q.-j. Ma et al., 2010		Jiang & Wang, 2009
	R. Sheng et al., 2008		Cho & Ahn, 2010

Table 1. Structures and references of coumarin-derived chemosensors for Hg(II) ions

coumarin fluorophore resulting in blue-shift in absorption and quenching of the fluorescence (S. H. Lee et al., 2010). The fluorescent ratiometric Hg²⁺ ion sensor **4**, based on a coumarin platform coupled with a tetraamide receptor, can specifically detect Hg²⁺ ions through the ICT mechanism (Wang et al., 2006). Fluorescein-coumarin chemodosimeter **5** for

signaling Hg^{2+} ions is designed based on FRET arising from the interaction between a pair of fluorophores (Ryu et al., 2010). Rhodamine-coumarin conjugate **7** was developed as a probe for $\text{Hg}(\text{II})$ ions. The fluorescence response to $\text{Hg}(\text{II})$ ions is attributed to the 1:1 complex formation between probe **7** and Hg^{2+} (Q.-J. Ma et al., 2010). Chemosensor **9** based on the coumarin thiosemicarbazide displays a selective fluorescence enhancement for Hg^{2+} , which is attributed to the transformation of thiosemicarbazide unit to 1,3,4-oxadiazoles via Hg^{2+} -induced desulfurization reaction in aqueous media (W. Ma et al., 2010). Coumarinyldithiane **11** and thiocoumarin **12** selectively sense Hg^{2+} also due to the Hg^{2+} -induced desulfurization reaction. Probe **14** belongs to the turn-on class of sensors, functioning via a PET process (Voutsadaki et al., 2010).

Chemosensors **15** and **16** are not coumarin derivatives but in the presence of $\text{Hg}(\text{II})$ ions the weakly fluorescent precursor **15** can be transformed to strongly fluorescent coumarin **17** via a desulfurization-lactonization cascade reaction as shown in Fig. 2 (Jiang & Wang, 2009). Similarly, **16** selectively senses inorganic mercury in the turn-on mode through a $\text{Hg}(\text{II})$ ion-promoted hydrolysis-cyclization reaction that leads to coumarin **17** as shown in Fig. 3 (Cho & Ahn, 2010).




Fig. 2. Transformation of sensor **14** to fluorescent coumarin **16** via $\text{Hg}(\text{II})$ -induced desulfurization-lactonization




Fig. 3. The reaction of probe **16** with HgCl_2

Detection of trace amount of Cu^{2+} is important not only for environmental applications, but also for toxicity determination in living organs. Following the report that a new cavitaand bearing four coumarin groups acts as fluorescent chemosensor for Cu^{2+} (Jang et al., 2006), over ten more coumarin-derived fluorescent chemosensors for $\text{Cu}(\text{II})$ ions have been envisaged. The structures and references of these chemical sensors are listed in Table 2.













Structure of sensor	Reference	Structure of sensor	Reference
 <p>18</p>	G. He et al., 2010a	 <p>24</p>	X. Chen et al., 2011
 <p>19</p>	M. H. Kim et al., 2009	 <p>25</p>	Ko et al., 2011
 <p>20</p>	Chandrasekhar et al., 2009	 <p>26</p>	N. Li et al., 2010
 <p>21</p>	Chandrasekhar et al., 2009	 <p>27</p>	Ciesiński et al., 2010
 <p>22</p>	J. R. Sheng et al., 2008	 <p>28</p>	Helal et al., 2011
 <p>23</p>	Zhao et al., 2010	 <p>29</p>	Jung et al., 2009

Table 2. Structures and references of coumarin-derived chemosensors for $\text{Cu}(\text{II})$ ions

A turn-on fluorescent probe **18** for Cu^{2+} is presented by incorporating coumarin fluorophores within the benzyl dihydrazone moiety. It is among the brightest Cu^{2+} binding sensors in aqueous media reported to date (G. He et al., 2010a). Coumarin **19** is a highly effective turn-on fluorescent sensor that is catalytically hydrolyzed by Cu^{2+} leading to a

large increase in the fluorescence intensity (M. H. Kim et al., 2009). Studies on interaction of phosphorus-supported multidentate coumarin-containing fluorescent sensors **20** and **21** with various transition metal ions reveal substantial fluorescence enhancement upon interaction with Cu^{2+} enabling a selective detection mechanism for Cu^{2+} (Chandrasekhar et al., 2009). Biscoumarin **22** linked by a C=N double bond is highly sensitive and selective to Cu^{2+} and the fluorescent sensing mechanism is based on C=N isomerization (J. R. Sheng et al., 2008). Coumarin probe **23** is highly selective for Cu^{2+} over biologically relevant alkali metals, alkaline earth metals and the first row transition metals due to the formation of a 1:2 complex between Cu^{2+} and **23** (Zhao et al., 2010). Fluorescent biscoumarin **24** linked by a piperazine unit shows high selectivity towards Cu^{2+} (Chen et al., 2011). Rationally designed iminocoumarin fluorescent sensor **25** displays high selectivity for Cu^{2+} over a variety of competing metal ions in aqueous solution with a significant fluorescence increase (Ko et al., 2011). Nonfluorescent coumarin derivative **26** is synthesized as an efficient turn-on fluorescent chemodosimeter for Cu^{2+} in water. Mechanism studies suggest that **26** forms a complex with Cu^{2+} in a 1:2 metal-to-ligand ratio, and a 50-fold fluorescence enhancement is observed when the complex simultaneously undergoes Cu^{2+} -promoted hydrolysis (N. Li et al., 2010). Sensor **27** relies on a coumarin-tagged ligand that selectively binds Cu^{2+} over other biometals to induce fluorescence quenching, which is subsequently relieved upon UV irradiation to provide the turn-on response (Ciesieski et al., 2010). For chemosensor **28** the mechanism of fluorescence is based on ICT, which is modified by the introduction of an electron-donating diethylamino group making it chromogenic and increasing the binding affinity (Helal et al., 2011). Coumarin **29** appending 2-picolyamide enables efficient tridentate complexation for Cu(II) in preference to a variety of other common heavy and toxic metal ions (Jung et al., 2009).

Owing to the important role of zinc, the second most abundant transition metal in the human body, more and more attention has been paid to development of Zn^{2+} -specific chemosensors including coumarin-derived fluorescent chemosensors for Zn(II) ions, the structures and references of which are listed in Table 3.

Study on the sensory capabilities of two novel di(2-picoly)amine (DPA)-substituted coumarins **30** and **31** shows that the variation of the point of attachment of the DPA group to the coumarin framework controls their sensing behavior: the 4-substituted system **30** is a chelation-enhanced fluorescence (CHEF)-type sensor which shows a significant increase in fluorescence intensity upon Zn^{2+} binding, whereas the 3-substituted coumarin **31** is a ratiometric sensor (N. C. Lim & Brückner, 2004). Coumarin Schiff-base **32** acts as a turn-on fluorescent chemosensor for Zn(II) ions (H. Li et al., 2009) and **33** does not show any two-photon activity in the wavelength range 760-860 nm but in the presence of Zn(II) **33** exhibits large two-photon absorption as well as emission in the same wavelength range (Ray et al., 2010). Another coumarin Schiff-base **43** is a highly sensitive and selective fluorescent probe for Zn^{2+} in tetrahydrofuran (THF) (Yan et al., 2011). In coumarin Schiff-base **34** the fluorescence quenching is dominant because of the nitrogen lone pair orbital contribution to the excitation. Upon Zn^{2+} coordination **34** shows a significant fluorescence enhancement due to the blocking of the nitrogen lone pair orbital by metal coordination (Jung et al., 2010). A series of coumarin-based fluorescent probes **35-38** for detecting Zn^{2+} with high affinities show the ratiometric fluorescent properties (Mizukami et al., 2009). Another series of coumarin-derived chemosensors **39-42** belong to the CHEF-type and have been showed to













Structure of sensor	Reference	Structure of sensor	Reference
 30	N. C. Lim & Brückner, 2004	 38	Mizukami et al., 2009
 31	N. C. Lim & Brückner, 2004	 39	N. C. Lim et al., 2005
 32: R = OH 33: R = NEt₂	H. Li et al., 2009 Ray et al., 2010	 40 • 4 HCl 41: R = H 42: R = CH₂CO₂Et	N. C. Lim et al., 2005
 34	Jung et al., 2010	 41: R = H 42: R = CH₂CO₂Et	N. C. Lim et al., 2005
 35	Mizukami et al., 2009	 43	Yan et al., 2011
 36: R = H 37: R = Cl	Mizukami et al., 2009	 44	Su et al., 2010

Table 3. Structures and references of coumarin-derived chemosensors for Zn(II) ions

be competent for detecting zinc pools in cultured rat pituitary (GH3) and hepatoma (H4IIE) cell lines (N. C. Lim et al., 2005). Coumarin **44** is a fluorescent sensor for Zn²⁺ and exhibits lower background fluorescence due to intramolecular PET but upon mixing with Zn²⁺ in aqueous ethanol, a turn-on fluorescence emission is observed (Su et al., 2010). Recently it was reported that a biscoumarin linked by bi-thiazole acted as a colorimetric receptor selectively for Zn²⁺ (Upadhyay & Mishra, 2010).

Structures of several coumarin-derived fluorescent chemosensors for iron(III) are shown in Fig. 4. Squarate hydroxamate-coumarin conjugate **45** is designed as a CHEF-type sensor for Fe(III). Due to a PET process, **45** possesses a low fluorescence yield but upon exposure of **45** to Fe(III), an irreversible 9-fold fluorescence intensity increase is observed as the result of an oxidation/hydrolysis reaction (N. C. Lim et al., 2009). Coumarin derivative **46** exhibits high selectivity for Fe³⁺ and the selectivity is not affected by the presence of representative alkali metals, alkali earth metals and other transition metal salts (Yao et al., 2009). Mugineic acid-coumarin derivative **47** synthesized by click chemistry acts as a fluorescent probe for Fe³⁺ (Namba et al., 2010). Coumarin-based hexadentate fluorescent probes for selective quantification of iron(III) have also been designed and synthesized (Y. M. Ma & Hider, 2009).




Fig. 4. Structures of coumarin-derived fluorescent chemosensors for iron(III) ion

Structures of coumarin-derived fluorescent chemosensors for Mg(II) are shown in Fig. 5. Coumarin-based two-photon probe **48** is developed for the detection of free Mg²⁺ ions in living cells and living tissues. The probe can be excited by 880 nm laser photons, emits strong two-photon excited fluorescence in response to Mg²⁺ ions (H. M. Kim et al., 2007). Coumarin Schiff-base **49**, without two-photon activity in the wavelength range 760-860 nm, exhibits large two-photon absorption as well as emission in the presence of Mg²⁺ (Ray et al., 2010). Coumarin-based chromoionophore **50** implemented in a transparent membrane can be used as an optical one-shot sensor for Mg²⁺ (Capitán-Vallvey, 2006). Two coumarin salen-based sensors **51** and **52** exhibit a pronounced fluorescence enhancement response toward Mg²⁺ in the presence of Na⁺ as a synergic trigger (Dong et al., 2011). Coumarin-derived fluorescent molecular probes **53** and **54** can be used for highly selective detection of Mg²⁺ versus Ca²⁺ by means of monitoring the absorption and fluorescence spectral change (Suzuki et al., 2002).




Fig. 5. Structures of coumarin-derived fluorescent chemosensors for Mg(II) ion

Fig. 6 shows the structures of coumarin-derived fluorescent chemosensors for Ag(I). A highly sensitive and selective fluorescent chemosensor **55** for Ag⁺ based on a coumarin-Se₂N chelating conjugate is developed. Due to inhibiting a PET quenching pathway, a fluorescent enhancement of 4-fold is observed under the binding of the Ag⁺ cation to **55** with a detection limit down to the 10⁻⁸ M range (S. Huang et al., 2011). Coumarin **56** and **57** are highly silver ion selective fluorescence ionophores (Sakamoto et al., 2010). By a microwave-assisted dual click reaction, fluorogenic 3-azidocoumarin can be rapidly introduced onto a 3,4-dipropargylglucosyl or galactosyl scaffold with restored fluorescence. Subsequent desilylation leads to water soluble sugar-bis-triazolocoumarin conjugates which are applicable toward selective Ag⁺ detection in aqueous media via fluorescence spectroscopy (X.-P. He et al., 2011).




Fig. 6. Structures of coumarin-derived fluorescent chemosensors for silver(I) ion

Structures of coumarin-derived fluorescent chemosensors for lead(II) are shown in Fig. 7. Fluorescent chemosensor **58** based on a coumarin-crown ether conjugate exhibits a high affinity and selectivity for Pb²⁺ (C.-T. Chen & W.-P. Huang, 2002). Coumarin dyes **59-61** seem to fulfill most of the criteria required for intracellular lead indicators, as they exhibit high selectivity for Pb²⁺ (Roussakis et al., 2008). Ion competition studies and fluorescence experiments show that a fullerene-coumarin dyad is selective for Pb²⁺ complexation (Pagona et al., 2010).




Fig. 7. Structures of coumarin-derived fluorescent chemosensors for lead(II) ion

Apart from the above-mentioned metal ions, coumarin-derived fluorescent chemosensors have been used for detection of other metal ions including Cd²⁺ (Taki et al., 2008), Al³⁺ (Maity & Govindaraju, 2010), and Cr³⁺ (Hu et al., 2011). Some coumarin-derived fluorescent chemosensors exhibit simultaneous sensitivity toward two or more different metal ions, e.g. Cu(II) and Hg(II) based on FRET mechanism (G. He et al., 2010b) or via selective anion-induced demetallation (Lau et al., 2011), Cu²⁺ and Ni²⁺ (H. Li et al., 2011), Ni²⁺ and Zn²⁺ (Chattopadhyay et al., 2006), Ni²⁺ and Co²⁺ (Lin et al., 2009), Ca²⁺ and Mg²⁺ (Suresh & Das, 2009), Na⁺ and K⁺ (Ast et al., 2011), Hg²⁺ and Ag⁺ (Tsukamoto et al., 2011), Cu²⁺/Ni²⁺/Cd²⁺ (Lin et al., 2008), Zn²⁺/Cd²⁺/Pb²⁺ (Kulatileke et al., 2006), or Ni²⁺/Pd²⁺/Ag⁺ (Santos et al., 2009).

3.2 Coumarin-derived fluorescent chemosensors for anions

Development of highly efficient chemosensors for cyanide is of extreme significance due to the detrimental aspect of cyanide. Much attention has been paid to the utilization of fluorescent methods for the detection of cyanide. Several coumarin-derived fluorescent chemosensors for CN⁻ have been reported, the structures of which are shown in Fig. 8.




Fig. 8. Structures of coumarin-derived fluorescent chemosensors for cyanide

Cobalt(II)-coumarinylsalen complex **62** exhibits selective and tight binding to a cyanide anion and displays a significant fluorescence enhancement upon the addition of cyanide owing to the interruption of PET from the coumarin fluorophore to the cobalt(II)-salen moiety (J. H. Lee et al., 2010). Coumarin-spiropyran conjugate **63** is a highly sensitive chemosensor for CN⁻ and shows a CN⁻-selective fluorescence enhancement under UV irradiation (Shiraishi et al., 2011). An indole conjugated coumarin **64** for KCN chemodosimeter displays considerable dual changes in both absorption (blue-shift) and emission (turn-on) bands exclusively for KCN. The fluorescence enhancement of the **64**-KCN is mainly due to blocking of the ICT process (H. J. Kim et al., 2011). Doubly activated coumarin **65** acts as a colorimetric and fluorescent chemodosimeter for cyanide (G.-J. Kim & H.-J. Kim, 2010a). A simple aldehyde-functionalized coumarin **66** has been utilized as a doubly activated Michael acceptor for cyanide (G.-J. Kim & H.-J. Kim, 2010b). Coumarin-based fluorescent chemodosimeter **67** with a salicylaldehyde functionality as a binding site has shown selectivity for cyanide anions over other anions in water at biological pH (K.-S. Lee et al., 2008a).

Recognition and detection of fluoride, the smallest anion with unique chemical properties is of growing interest. Several coumarin-derived fluorescent chemosensors for F⁻ have been developed, the structures of which are shown in Fig. 9.

Coumarin derivative **68** has been developed as a fluorescent probe for detection of F⁻ ion in water and bioimaging in A549 human lung carcinoma cells (S. Y. Kim et al., 2009). Coumarin **69** is a simple, highly selective, and sensitive chemosensor for fluoride anions in organic and aqueous media based on the specific affinity of fluoride anion to silicon (Sokkalingam & Lee, 2011). Coumarin-derived chemosensor **70** shows an obvious color change from yellow to blue upon addition of F⁻ ion with a large red shift of 145 nm in acetonitrile (Zhuang et al., 2011). Coumarin-based hydrazone **71** is an ICT probe for fluoride in aqueous medium (Upadhyay et al., 2010a). Coumarin-based system **72** has been developed as a novel turn-on fluorescent and colorimetric sensor for fluoride anions (J. Li et al., 2009).




Fig. 9. Structures of coumarin-derived fluorescent chemosensors for F^- ion




Fig. 10. Structures of coumarin-derived fluorescent chemosensors for anions other than CN^- and F^-

Structures of coumarin-derived fluorescent chemosensors for detection of anions other than CN^- and F^- have been shown in Fig. 10. Coumarin-based fluorescent probes **73-76** with salicylaldehyde functionality as recognition unit have been developed for selective detection of bisulfite anions in water (K. Chen et al., 2010). Coumarin Schiff-base **77** is a highly selective and sensitive turn-on fluorogenic probe for detection of HSO_4^- ions in aqueous solution (H. J. Kim et al., 2009b). Coumarin-based hydrazone **78** acts as an ICT probe for detection of acetate, benzoate, and dihydrogenphosphate (Upadhyay et al., 2010b). Another coumarin-based hydrazone **79** has been utilized as both a colorimetric and ratiometric chemosensor for acetate and a selective fluorescence turn-on probe for iodide (Mahapatra et al., 2011). Compound **80** is a colorimetric and fluorescence anion sensor with the urea group as binding site and the coumarin moiety as signal unit. The sensor displays significant fluorescence enhancement response to anions such as acetate, because of complex formation (Shao, 2010). Coumarin-derived fluorescent chemosensors have also been used for detection of pyrophosphate (S. K. Kim et al., 2009), $H_2PO_4^-$ and $PhPO_3H^-$ (K. Choi & Hamilton, 2001), and multiple anions including pyrophosphate, citrate, ATP and ADP (Mizukami et al., 2002). A new Zn^{II} -2,2':6',2''-terpyridine complex, derivatized with a coumarin moiety, acts as a fluorescent chemosensor for different biologically important phosphates like PPI, AMP and ADP in mixed aqueous media (Das et al., 2011). A strapped calix[4]pyrrole-coumarin conjugate has been developed as a fluorogenic anions (Cl^- , Br^- and AcO^-) receptor modulated by cation and anion binding (Miyaji et al., 2005).

3.3 Coumarin derivatives as fluorescent probes of pH

Structures of coumarin-derived fluorescent chemosensors for sensing pH are shown in Fig. 11. Several iminocoumarin (**81**) derivatives **82-84** have been synthesized as a new type of fluorescent pH indicator. The indicators possess moderate to high brightness, excellent photostability and compatibility with light-emitting diodes. The indicators can be covalently immobilized on the surface of amino-modified polymer microbeads which in turn are incorporated into a hydrogel matrix to afford novel pH-sensitive materials. When a mixture of two different microbeads is used, the membranes are capable of optical pH sensing over a very wide range comparable to the dynamic range of the glass electrode (pH 1-11) (Vasylevska et al., 2007). Coumarin derivative **85** containing piperazine and imidazole moieties has been developed as a fluorophore for hydrogen ions sensing. The fluorescence enhancement of the sensor with an increase in hydrogen ions concentration is based on the hindering of PET from the piperazinyl amine and the imidazolyl amine to the coumarin fluorophore by protonation. The sensor **85** has a novel molecular structure design of fluorophore-spacer-receptor(1)-receptor(2) format and therefore is proposed to sense two range of pH from 2.5 to 5.5 and from 10 to 12 instead of sensing one pH range (Saleh et al., 2008). By using rational molecular design, two molecular functions, the transport by vesicular monoamine transporter (VMAT) and ratiometric optical pH sensing, have been integrated to develop ratiometric pH-responsive fluorescent false neurotransmitter (FFN) probes (M. Lee et al., 2010). A FRET sensor with a donor and an acceptor attached to each end of pH-sensitive polysulfoamides exhibits an instantaneous conformation change from coil to globule at a specific pH, which results in the drastic on-and-off FRET efficiency. To detect a specific pH region, sulfadimethoxine and sulfamethizole are selected among various sulfonamides since their pK_a values are in the physiological pH. For tuning the emission color arising from FRET, 7-hydroxy-4-bromomethylcoumarin and coumarin 343 are used as a FRET donor and an acceptor, respectively, for a blue-to-green FRET sensor (Hong & Jo, 2008).




Fig. 11. Structures of coumarin-derived fluorescent chemosensors for sensing pH

3.4 Coumarin-derived fluorescent chemosensors for thiols

Biothiols such as cysteine (Cys), homocysteine (Hcy), and glutathione (GSH) are of great significance because they are involved in myriad vital cellular processes including redox homeostasis and cellular growth. Alteration of the cellular biothiols is also implicated in

cancer and AIDS. Study on fluorescent and colorimetric probes for detection of thiols has received much attention and many coumarin-derived fluorescent chemosensors for detection of thiols have been reported. The structures of these chemical sensors are shown in Fig. 12.




Fig. 12. Structures of coumarin-derived chemosensors for detection of thiols

Generally detection of thiols by optical probes is based on two features of thiols, their strong nucleophilicity and high binding affinity toward metal ions. Accordingly, most of the fluorescent probes for thiols are in fact chemodosimeters, which involve specific reactions between probes and thiols, such as Michael addition, cyclization with aldehyde (or ketone), cleavage of disulfide by thiols, metal complexes-displace coordination, demetalization from Cu-complex, thiolysis of dinitrophenyl ether, and Staudinger ligation. For instance, detection of thiols by chemosensors **87-91** and **93** involves Michael addition between the probes and thiols. For sensors **67** (Fig. 8), **86** and **98**, cyclization reactions occur between the sensors and thiols in the detection process. Coumarin **86** is a ratiometric fluorescent probe for specific detection of Cys over Hcy and GSH based on the drastic distinction in the kinetic profiles (Yuan et al., 2011). Nonfluorescent coumarin-malonitrile conjugate **87** can be transformed into a strongly fluorescent molecule through the Michael addition and thus exhibits a highly selective fluorescence response toward biothiols including Cys, Hcy and GSH with micromolar sensitivity (Kwon et al., 2011). Similarly, nonfluorescent **88** displays a highly selective fluorescence enhancement with thiols and has been successfully applied to thiols determination in intracellular, in human urine and blood samples (Zuo et al., 2010). Coumarin **89** has been developed as a water-soluble, fast-response, highly sensitive and selective fluorescence thiol quantification probe (Yi et al., 2009). Compound **90** (G.-J. Kim et al., 2011) and **91** (S. Y. Lim et al., 2011) with a hydrogen bond act as highly selective ratiometric fluorescence turn-on probes for GSH. Structure **92** has been judiciously designed and synthesized as a new type of selective benzenethiol fluorescent probe based on the thiolysis of dinitrophenyl ether (Lin et al., 2010a). Coumarin-based chemodosimeter **93** effectively and selectively recognizes thiols based on a Michael type reaction, showing a preference for Cys over other biological materials including Hcy and GSH (Jung et al.,

2011a). Iminocoumarin-Cu(II) ensemble-based chemodosimeter **94** sensitively senses thiols followed by hydrolysis to give a marked fluorescence enhancement over other amino acids based on demetalization from Cu-complex (Jung et al., 2011b). Nonfluorescent coumarin-phosphine dye **95** reacts with S-nitrosothiols (RSNOs) to form a fluorescent coumarin derivative and thus may be used as a tool in the detection of RSNOs. The reaction mechanism is similar to the well-known Staudinger ligation (Pan et al., 2009). 7-Mercapto-4-methylcoumarin **96** is a reporter of thiol binding to the CdSe quantum dot surface (González-Béjar et al., 2009). Coumarin-derived complex **97** has been developed as a reversible fluorescent probe for highly selective and sensitive detection of mercapto biomolecules such as Cys, Hcy and GSH (J. Wu et al., 2011). A simple coumarin derivative **98** is the first fluorescence turn-on probe for thioureas by the double functional group transformation strategy. The probe exhibits high sensitivity and selectivity for thioureas over other structurally and chemically related species including urea and thiophenol (Lin et al., 2010b). The simple coumarin sensor **67** (Fig. 8) has shown fluorescence selectivity for not only cyanide anions but also Hcy and Cys in water (K.-S. Lee et al., 2008b). A new coumarin-containing zinc complex has been developed as a colorimetric turn-on and fluorescence turn-off sensor which shows high selectivity for hydrogen sulfide in the presence of additional thiols like Cys or GSH (Galardon et al., 2009).

3.5 Coumarin-derived fluorescent chemosensors for H₂O₂, O₂, hydroxyl radicals or chemical warfare agents

Structures of coumarin-derived fluorescent chemosensors for hydrogen peroxide, oxygen, hydroxyl radicals or chemical warfare agents are shown in Fig. 13. Water-soluble umbelliferone-based fluorescent probe **99** shows very large increase (up to 100-fold) in fluorescent intensity upon reaction with hydrogen peroxide, and good selectivity over other reactive oxygen species (Du et al., 2008). Another water-soluble fluorescent hydrogen peroxide probe **100** based on a 'click' modified coumarin fluorophore shows significant intensity increases (up to fivefold) in near-green fluorescence upon reaction with H₂O₂, and good selectivity over other reactive oxygen species (Du Y et al., 2010). More recently a simple and highly sensitive fluorometric method was proposed for the determination of H₂O₂ in milk samples. In this method, nonfluorescent coumarin was oxidized to highly fluorescent 7-hydroxycoumarin by hydroxyl radicals generated in a Fenton reaction, and the oxidation product had strong fluorescence with a maximum intensity at 456 nm and could be used as a fluorescent probe for H₂O₂ (Abbas et al., 2010). Thiazo-coumarin ligand directly cyclometallated Pt(II) complex **101** has been used for luminescent O₂ sensing (W. Wu et al., 2011). A hybrid coumarin-cyanine platform **102** has been developed as the first ratiometric fluorescent probe for detection of intracellular hydroxyl radicals (Yuan et al., 2010). More recently a coumarin-neutral red (CONER) nanoprobe was developed for detection of hydroxyl radical based on the ratiometric fluorescence signal between 7-hydroxy coumarin 3-carboxylic acid and neutral red dyes. Biocompatible poly lactide-co-glycolide nanoparticles containing encapsulated neutral red were produced using a coumarin 3-carboxylic acid conjugated poly(sodium *N*-undecylenyl-*Nε*-lysinate) as moiety reactive to hydroxyl radicals. The response of the CONER nanoprobe was dependent on various parameters such as reaction time and nanoparticle concentration. The probe was selective for hydroxyl radicals as compared with other reactive oxygen species including O₂^{•-}, H₂O₂, ¹O₂ and OCl⁻ (Ganea et al., 2011).

The current rise in international concern over criminal terrorist attacks using chemical warfare agents has brought about the need for reliable and affordable detection methods of toxic gases. One of the applicable technologies is the design of fluorogenic chemosensors for the specific detection of nerve agents (Royo et al., 2007). A coumarin oximate **103** has been developed for detection of chemical warfare simulants based on the PET mechanism that gives an "off-on" fluorescent response with a half-time of approximately 50 ms upon phosphorylation of a reactive oximate functionality (Wallace et al., 2006). Coumarin-derived hydroxy oxime **104** serves as a nerve agent sensor based on the reaction of β -hydroxy oxime with organophosphorus agent mimics (Dale & Rebek, 2009). A FRET approach towards potential detection of phosgene has been developed as shown in Fig. 14. When both coumarins **105** and **106** are mixed together with triphosgene in the presence of Et_3N in CHCl_3 , hybrid urea **107** forms in a statistical yield. Significant fluorescence enhancement is detected which is particularly important since the acceptor unit alone does not emit under the same condition. The fluorescence increase is obviously due to the formation of urea **107**. Simultaneously, the fluorescence from the donor unit decreases due to the quenching, indicating that efficient energy transfer takes place from the donor to the acceptor. This system is selective, since other gases/agents rarely can serve for cross-linking (Zhang & Rudkevich, 2007).




Fig. 13. Structures of coumarin-derived fluorescent sensors for H_2O_2 , O_2 , hydroxyl radicals or chemical warfare agents




Fig. 14. Coumarins **105** and **106** react with phosgene to form urea **107**

3.6 Coumarin-derived fluorescent chemosensors for amines, amino acids or other organic compounds

Structures of coumarin-derived fluorescent chemosensors for amines, amino acids or other organic compounds are shown in Fig. 15. Butyl-substituted coumarin aldehyde **108** is an excellent chemosensor for detection of amines and unprotected amino acids in aqueous conditions by formation of highly fluorescent iminium ions (Feuster & Glass, 2003). Boronic acid-containing coumarin aldehyde **109** binds to primary catecholamines with good affinity and acts as an effective colorimetric sensor for dopamine and norepinephrine with excellent selectivity over epinephrine, amino acids, and glucose. In the fluorescence manifold, sensor **109** responds differentially to catechol amines over simple amines, giving a fluorescence decrease in response to catechol-containing compounds and a fluorescence increase with other amines (Secor & Glass, 2004). Coumarin-based fluorescent functional monomers containing a carboxylic acid functionality, **110** and **111** have been synthesized, which allow for the preparation of fluorescent imprinted polymer sensors for chiral amines (Nguyen & Ansell, 2009). Coumarin aldehyde **66** (Fig. 8) can be utilized as not only a doubly activated Michael acceptor for cyanide but also a highly selective and sensitive fluorescence turn-on probe for proline (G.-J. Kim & H.-J. Kim, 2010c). Coumarin-azacrown ether conjugate **112** has been developed as a fluorescent probe for identifying melamine (Xiong et al., 2010). Thiocoumarin **113** can be efficiently desulfurized to its corresponding coumarin by the reaction with mCPBA, and results in a pronounced fluorescence turn-on type signaling. The conversion also provides a significant change in absorption behavior which allows a ratiometric analysis, providing a convenient detection method for mCPBA in aqueous environment (Cha et al., 2010). Polymers containing 4,8-dimethylcoumarin have been developed for detection of 2,4-dinitrotoluene (DNT) and 2,4,6-trinitrotoluene (TNT).




Fig. 15. Structures of coumarin-derived fluorescent chemosensors for amines, amino acids or other organic compounds

The fluorescence quenching of these copolymers in solution can be attributed to the collisional quenching. The response of these polymeric sensors is promising and can easily detect DNT and TNT at few parts per billion levels (Kumar et al., 2010). A novel kind of luminescent vesicular chemosensors for the recognition of biologically important ions and molecules such as imidazoles has been developed by the self-assembly of lipids, amphiphilic binding sites, and fluorescent coumarin reporter dyes that are sensitive to their environment (Gruber et al., 2010). Two hybrid compounds **114** and **115**, linked via an ester-bond between the 7-hydroxyl residue of an umbelliferone and a carboxylic acid residue of two nitroxide radicals, act as fluorescence and spin-label probes. The ESR intensities of **114** and **115** are proportionally reduced after the addition of ascorbic acid sodium salt, and their fluorescence intensities are increased maximally by eight- and nine-fold, respectively (Sato et al., 2008).

3.7 Coumarin-derived fluorescent chemosensors for TiO₂, monolayer, polymerization or polymeric micelles

Structures of coumarin-derived fluorescent chemosensors for detection of TiO₂, monolayer, or photopolymerization are shown in Fig. 16. A novel acac-coumarin chromophore linker **116** for robust sensitization of TiO₂ has been developed to find molecular chromophores with suitable properties for solar energy conversion. The synthesis and spectroscopic characterization confirms that **116** yields improved sensitization to solar light and provides robust attachment to TiO₂ even in aqueous conditions (Xiao et al., 2011). A new amphiphilic coumarin dye, 7-aminocoumarin-4-acetic acid octadecylamide (**117**) forms a stable monolayer at the air-water interface and may be utilized as an efficient fluorescent probe for monolayer studies (Kele et al., 2001). Performance of amidocoumarins **118-120** as probes for monitoring of cationic photopolymerization of monomers by fluorescence probe technology has been investigated. 7-Diethylamino-4-methylcoumarin **118** can be used for monitoring cationic photopolymerization of monomers using the fluorescence intensity ratio as an indicator of the polymerization process. The replacement of diethylamino group in **118** with benzamido or acetamido groups eliminates the effect of the probe protonation on kinetics of cationic photopolymerization. 7-Benzamido-4-methylcoumarin **119** and 7-acetamido-4-methylcoumarin **120** can be used as fluorescent probes for monitoring progress of cationic polymerization of vinyl ethers under stationary measurement conditions, using normalized fluorescence intensity as an indicator of the polymerization progress (Ortyl et al., 2010). Coumarin 153 has been used as a fluorescent probe molecule to monitor the possible micellization of several amphiphilic block copolymers (Basu et al., 2009).




Fig. 16. Structures of coumarin-derived fluorescent sensors for detection of TiO₂, monolayer, or photopolymerization

3.8 Coumarin-derived fluorescent chemosensors for enzymes

Structures of coumarin-derived fluorescent chemosensors for enzymes are shown in Fig. 17. Hemicyanine-coumarin hybrid **121** represents a new class of far-red emitting fluorogenic dyes whose fluorescence is unveiled through an enzyme-initiated domino reaction and thus acts as a fluorogenic probe for penicillin G acylase (Richard et al., 2008). Similarly novel self-immolative spacer systems **122** and **123** have been developed and are utilized as fluorogenic probes for sensing penicillin amidase (Meyer et al., 2008). A library of 6-arylcoumarins has been developed as candidate fluorescent sensors of which **124** has the strongest fluorescence intensity, whose quantum yield is similar to that of ethyl 7-diethylaminocoumarin-3-carboxylate, a well-known fluorophore as labeling or sensing biomolecule. The transformation of the methoxy group (**125**) to a hydroxyl group (**126**) induces a change of fluorescence intensity, which suggests that **125** may be useful as a fluorescent sensor for dealkylating enzymes such as glycosidase. Coumarin **127** shows 50% decrease of the fluorescence intensity at pH 8.0 compared with that at pH 6.0 and this decrease may be derived from the deprotonation of the triazole ring. Thus **127** may be used as a fluorescent sensor for nitric oxide (Hirano et al., 2007). Histone deacetylases are intimately involved in epigenetic regulation and, thus, are one of the key therapeutic targets for cancer. Coumarin-suberoylanilide hydroxamic acid **128** is a fluorescent probe for determining binding affinities and off-rates of histone deacetylase inhibitors (Singh et al., 2011). A quinone-methide-rearrangement reaction as the off-on optical switch has been successfully implemented into the design of the first long-wavelength latent fluorogenic substrate **129** which is a sensitive fluorimetric indicator for analyte determination in salicylate hydroxylase-coupled dehydrogenase assay (S.-T. Huang et al., 2010). Another switch-on long-wavelength latent fluorogenic substrate **130** is a fluorescent probe for nitroreductase (H.-C. Huang et al., 2011).




Fig. 17. Structures of coumarin-derived fluorescent chemosensors for enzymes

Apart from the above-mentioned coumarin-derived fluorescent chemosensors for enzymes, clickable biocompatible nanoparticles have been prepared in a one-pot process by microemulsion polymerization, which are then readily modified by the Huisgen Cu(I)-catalyzed azide-alkyne cycloaddition reaction to afford a coumarin-containing subtilisin responsive nanosensor (Welser et al., 2009). A coumarin-containing time-resolved fluorescence probe for dipeptidyl peptidase 4 has also been reported (Kawaguchi et al., 2010). A coumarin-derived triple-signaling fluorescent probe has been successfully applied for intracellular measurement of different enzyme activity (Y. Li et al., 2011). As shown in Fig. 18, a sensitive, selective, and fluorogenic probe **131** has been developed for monoamine oxidases (MAO A and B). Nonfluorescent aminocoumarin **131** can be converted to fluorescent pyrrolocoumarin **132** in the presence of MAO A and B (G. Chen et al., 2005). A new fluorogenic transformation based on a quinone reduction/lactonization sequence as shown in Fig. 19 has been developed and evaluated as a tool for probing redox phenomena in a biochemical context.



Fig. 18. Systematic mapping of aminocoumarin **131** and the corresponding pyrrolocoumarin **132**

3.9 Coumarin-derived fluorescent chemosensors for proteins, DNA, RNA and other uses

Structures of coumarin-derived fluorescent chemosensors for DNA, RNA, nitroxyl and proteins are shown in Fig. 20. A novel coumarin C-riboside **133** is designed and synthesized based on the well-known photoprobe Coumarin 102. The coumarin C-glycoside **133** has been incorporated synthetically into DNA oligomers, and has been used to probe ultrafast dynamics of duplex DNA using time-resolved Stokes shift methods (Coleman et al., 2007). Coumarin-triazole **134** reacts with CuCl_2 to form a chelated Cu(II)-134 complex which shows highly selective turn-on type fluorogenic behavior upon addition of Angeli's salt ($\text{Na}_2\text{N}_2\text{O}_3$) and can be used for detection of nitroxyl in living cells (Zhou et al., 2011). A simple coumarin derivative 7-diethylaminocoumarin-3-carboxylic acid **135** has been used as an acceptor to construct a useful and effective FRET system for detection of RNA-small molecule binding (Xie et al., 2009). Coumarin dye bearing an indolenine substituent **136** displays high emission and bright fluorescence and offers promise as a fluorescent chemosensor for protein detection (Kovalska et al., 2010). Coumarin-containing trifunctional probe **137**, assembled using a cleavable linker, is useful for efficient enrichment and detection of glycoproteins (Tsai et al., 2010). Coumarin 6 has been used as a fluorescent probe to monitor protein aggregation and can distinguish between both amorphous and fibrillar aggregates (Makwana et al., 2011). An $\text{S}_{\text{N}}\text{Ar}$ reaction-triggered fluorescence probe is developed using a new fluorogenic compound derivatized from 7-aminocoumarin for oligonucleotides detection (Shibata et al., 2009). Coumarin C343 has been conjugated to silica nanoparticles and entrapped in a sol-gel matrix to produce a nanosensor capable of monitoring lipid peroxidation (Baker et al., 2007). Coumarin-containing dual-emission chemosensors for nucleoside polyphosphates have been developed based on a new

mechanism involving binding-induced recovery of FRET. These sensors demonstrate that binding-induced modulation of spectral overlap is a powerful strategy for the rational design of FRET-based chemosensors (Kurishita et al., 2010).




Fig. 19. Structures of coumarin-derived fluorescent chemosensors for DNA, RNA, nitroxyl and proteins

4. Conclusions

Coumarin-derived fluorescent chemosensors have been extensively applied in a variety of fields. Though these sensors are effective for detection of many species, their performance toward different species might decrease in the following order: metal ions, anions, biothiols, enzymes, pH, amines and amino acids, chemical warfare agents, proteins, hydrogen peroxide, hydroxyl radicals, polymerization and polymeric micelles, DNA and RNA, oxygen, titania. Continuous efforts will be devoted to development of fluorescent chemosensors with higher selectivity and sensitivity for more single target or simultaneously for multiple targets, thus providing practical fluorescent chemosensors for application in environmental chemistry, analytical chemistry, and bio-medical science.

5. References

- Abbas, M. E.; Luo, W.; Zhu, L.; Zou, J. & Tang, H. (2010). Fluorometric determination of hydrogen peroxide in milk by using a Fenton reaction system. *Food Chem.*, vol.120, No.1, (May 2010), pp. 327-331, ISSN 0308-8146
- Ast, S.; Müller, H.; Flehr, R.; Klamroth, T.; Walz, B. & Holdt, H.-J. (2011). High Na⁺ and K⁺-induced fluorescence enhancement of a π -conjugated phenylaza-18-crown-6-triazol-substituted coumarin fluoroionophore. *Chem. Commun.*, vol.47, No.16, (April 2011), pp. 4685-4687, ISSN 1359-7345
- Baker, N.; Greenway, G. M.; Wheatley, R. A. & Wiles, C. (2007). A chemiluminescence nanosensor to monitor lipid peroxidation. *Analyst*, vol.132, No.2, (January 2007), pp. 104-106, ISSN 0003-2654
- Bangar Raju, B. & Varadarajan, T. S. (1995). Spectroscopic studies of 7-diethylamino-3-styrylcoumarins. *J. Photochem. Photobio. A*, vol.85, No.3, (January 1995), pp. 263-267, ISSN 1010-6030

- Basu, S.; Chatterjee, D. P.; Chatterjee, U.; Mondal, S. & Mandal, D. (2009). Fluorescence probing of block copolymeric micelles using Coumarin 153. *Colloids Surf. A*, vol.341, No.1-3, (June 2009), pp. 13-20, ISSN 0927-7757
- Capitán-Vallvey, L. F.; Fernández-Ramos, M. D.; Lapresta-Fernández, A.; Brunet, E.; Rodríguez-Ubis, J. C. & Juanes, O. (2006). Magnesium optical one-shot sensor based on a coumarin chromoionophore. *Talanta*, vol.68, No.5, (February 2006), pp. 1663-1670, ISSN 0039-9140
- Cha, S.; Hwang, J.; Choi, M. G. & Chang, S.-K. (2010). Dual signaling of *m*-chloroperbenzoic acid by desulfurization of thiocoumarin. *Tetrahedron Lett.*, vol.51, No.50, (December 2010), pp. 6663-6665, ISSN 0040-4039
- Chandrasekhar, V.; Bag, P. & Pandey, M. D. (2009). Phosphorus-supported multidentate coumarin-containing fluorescence sensors for Cu²⁺. *Tetrahedron*, vol.65, No.47, (November 2009), pp. 9876-9883, ISSN 0040-4020
- Chattopadhyay, N.; Mallick, A. & Sengupta, S. (2006). Photophysical studies of 7-hydroxy-4-methyl-8-(4'-methylpiperazin-1'-yl) methylcoumarin: a new fluorescent chemosensor for zinc and nickel ions in water. *J. Photochem. Photobio. A*, vol.177, No.1, (January 2006), pp. 55-60, ISSN 1010-6030
- Chen, C.-T. & Huang, W.-P. (2002). A highly selective fluorescent chemosensor for lead ions. *J. Am. Chem. Soc.*, vol.124, No.22, (June 2002), pp. 6246-6247, ISSN 0002-7863
- Chen, G.; Yee, D. J.; Gubernator, N. G. & Sames, D. (2005). Design of optical switches as metabolic indicators: new fluorogenic probes for monoamine oxidases (MAO A and B). *J. Am. Chem. Soc.*, vol.127, No.13, (April 2005), pp. 4544-4545, ISSN 0002-7863
- Chen, K.; Guo, Y.; Lu, Z.; Yang, B. & Shi, Z. (2010). Novel coumarin-based fluorescent probe for selective detection of bisulfite anion in water. *Chin. J. Chem.*, vol.28, No.1, (January 2010), pp. 55-60, ISSN 1001-604X
- Chen, X.; Xi, H.; Sun, X.; Zhao, T.; Meng, Q. & Jiang, Y. (2011). Synthesis and fluorescent probes properties of a coumarin-based piperazine containing fluorine. *Chin. J. Org. Chem.*, vol.31, No.4, (April 2011), pp. 544-547, ISSN 0253-2786
- Cho, Y.-S. & Ahn, K. H. (2010). A 'turn-on' fluorescent probe that selectively responds to inorganic mercury species. *Tetrahedron Lett.*, vol.51, No.29, (July 2010), pp. 3852-3854, ISSN 0040-4039
- Choi, K. & Hamilton, A. D. (2001). A dual channel fluorescence chemosensor for anions involving intermolecular excited state proton transfer. *Angew. Chem. Int. Ed.*, vol.40, No.20, (October 2001), pp. 3912-3915, ISSN 1433-7851
- Choi, M. G.; Kim, Y. H.; Namgoong, J. E. & Chang, S.-K. (2009). Hg²⁺-selective chromogenic and fluorogenic chemodosimeter based on thiocoumarins. *Chem. Commun.*, No.24, (June 2009), pp. 3560-3562, ISSN 1359-7345
- Ciesinski, K. L.; Hyman, L. M.; Derisavifard, S. & Franz, K. J. (2010). Toward the detection of cellular copper(II) by a light-activated fluorescence increase. *Inorg. Chem.*, vol.49, No.15, (August 2010), pp. 6808-6810, ISSN 0020-1669
- Coleman, R. S.; Berg, M. A. & Murphy, C. J. (2007). Coumarin base-pair replacement as a fluorescent probe of ultrafast DNA dynamics. *Tetrahedron*, vol.63, No.17, (April 2007), pp. 3450-3456, ISSN 0040-4020
- Dale, T. J. & Rebek, J., Jr. (2009). Hydroxy oximes as organophosphorus nerve agent sensors. *Angew. Chem. Int. Ed.*, vol.48, No.42, (October 2009), pp. 7850-7852, ISSN 1433-7851
- Das, P.; Ghosh, A.; Kesharwani, M. K.; Ramu, V.; Ganguly, B. & Das, A. (2011). Zn^{II}-2,2':6',2''-Terpyridine-based complex as fluorescent chemosensor for PPI, AMP and ADP. *Eur. J. Inorg. Chem.*, No.20, (July 2011), pp. 3050-3058, ISSN 1434-1948

- Dong, Y.; Li, J.; Jiang, X.; Song, F.; Cheng, Y. & Zhu, C. (2011). Na⁺ triggered fluorescence sensors for Mg²⁺ detection based on a coumarin salen moiety. *Org. Lett.*, vol.13, No.9, (May 2011), pp. 2252-2255, ISSN 1523-7060
- Du, L.; Li, M.; Zheng, S. & Wang, B. (2008). Rational design of a fluorescent hydrogen peroxide probe based on the umbelliferone fluorophore. *Tetrahedron Lett.*, vol.49, No.19, (May 2008), pp. 3045-3048, ISSN 0040-4039
- Du Y, L.; Ni Y, N.; Li, M. & Wang, B. (2010). A fluorescent hydrogen peroxide probe based on a 'click' modified coumarin fluorophore. *Tetrahedron Lett.*, vol.51, No.8, (February 2010), pp. 1152-1154, ISSN 0040-4039
- Feuster, E. K. & Glass, T. E. (2003). Detection of amines and unprotected amino acids in aqueous conditions by formation of highly fluorescent iminium ions. *J. Am. Chem. Soc.*, vol.125, No.52, (December 2003), pp. 16174-16175, ISSN 0002-7863
- Galardon, E.; Tomas, A.; Roussel, P. & Artaud, I. (2009). New fluorescent zinc complexes: towards specific sensors for hydrogen sulfide in solution. *Dalton Trans.*, No.42, (November 2009), pp. 9126-9130, ISSN 1477-9226
- Ganea, G. M.; Kolic, P. E.; El-Zahab, B. & Warner, I. M. (2011). Ratiometric coumarin-neutral red (CONER) nanoprobe for detection of hydroxyl radicals. *Anal. Chem.*, vol.83, No.7, (April 2011), pp. 2576-2581, ISSN 0003-2700
- González-Béjar, M.; Frenette, M.; Jorge, L. & Scaiano, J. C. (2009). 7-Mercapto-4-methylcoumarin as a reporter of thiol binding to the CdSe quantum dot surface. *Chem. Commun.*, No.22, (June 2009), pp. 3202-3204, ISSN 1359-7345
- Gruber, B.; Stadlbauer, S.; Späth, A.; Weiss, S.; Kalinina, M. & König, B. (2010). Modular chemosensors from self-assembled vesicle membranes with amphiphilic binding sites and reporter dyes. *Angew. Chem. Int. Ed.*, vol.49, No.39, (September 2010), pp. 7125-7128, ISSN 1433-7851
- He, G.; Zhang, X.; He, C.; Zhao, X. & Duan, C. (2010b). Ratiometric fluorescence chemosensors for copper(II) and mercury(II) based on FRET systems. *Tetrahedron*, vol.66, No.51, (December 2010), pp. 9762-9768, ISSN 0040-4020
- He, G.; Zhao, X.; Zhang, X.; Fan, H.; Wu, S.; Li, H.; He, C. & Duan, C. (2010a). A turn-on PET fluorescence sensor for imaging Cu²⁺ in living cells. *New J. Chem.*, vol.34, No.6, (June 2010), pp. 1055-1058, ISSN 1144-0546
- He, X.-P.; Song, Z.; Wang, Z.-Z.; Shi, X.-X. & Chen, K. (2011). Creation of 3,4-bis-triazolocoumarin-sugar conjugates via fluorogenic dual click chemistry and their quenching specificity with silver(I) in aqueous media. *Tetrahedron*, vol.67, No.19, (May 2011), pp. 3343-3347, ISSN 0040-4020
- Helal, A.; Rashid, M. H. O.; Choi, C.-H. & Kim, H.-S. (2011). Chromogenic and fluorogenic sensing of Cu²⁺ based on coumarin. *Tetrahedron*, vol.67, No.15, (April 2011), pp. 2794-2802, ISSN 0040-4020
- Hirano, T.; Hiromoto, K. & Kagechika, H. (2007). Development of a library of 6-aryl coumarins as candidate fluorescent sensors. *Org. Lett.*, vol.9, No.7, (March 2007), pp. 1315-1318, ISSN 1523-7060
- Hong, S. W. & Jo, W. H. (2008). A fluorescence resonance energy transfer probe for sensing pH in aqueous solution. *Polymer*, vol.49, No. 19, (September 2008), pp. 4180-4187, ISSN 0032-3861
- Hu, X.; Zhang, X.; He, G.; He, C. & Duan, C. (2011). A FRET approach for luminescence sensing Cr³⁺ in aqueous solution and living cells through functionalizing glutathione and glucose moieties. *Tetrahedron*, vol.67, No.6, (February 2011), pp. 1091-1095, ISSN 0040-4020

- Huang, H.-C.; Wang, K.-L.; Huang, S.-T.; Lin, H.-Y. & Lin, C.-M. (2011). Development of a sensitive long-wavelength fluorogenic probe for nitroreductase: a new fluorimetric indicator for analyte determination by dehydrogenase-coupled biosensors. *Biosensors Bioelectronics*, vol.26, No.8, (April 2011), pp. 3511-3516, ISSN 0956-5663
- Huang, S.; He, S.; Lu, Y.; Wei, F.; Zeng, X. & Zhao, L. (2011). Highly sensitive and selective fluorescent chemosensor for Ag⁺ based on a coumarin-Se₂N chelating conjugate. *Chem. Commun.*, vol.47, No.8, (February 2011), pp. 2408-2410, ISSN 1359-7345
- Huang, S.-T.; Teng, C.-J.; Lee, Y.-H.; Wu, J.-Y.; Wang, K.-L. & Lin, C.-M. (2010). Design and synthesis of a long-wavelength latent fluorogenic substrate for salicylate hydroxylase: a useful fluorimetric indicator for analyte determination by dehydrogenase-coupled biosensors. *Anal. Chem.*, vol.82, No.17, (September 2010), pp. 7329-7334, ISSN 0003-2700
- Jang, Y. J.; Moon, B.-S.; Park, M. S.; Kang, B.-G.; Kwon, J. Y.; Hong, J. S. J.; Yoon, Y. J.; Lee, K. D. & Yoon, J. (2006). New cavitand derivatives bearing four coumarin groups as fluorescent chemosensors for Cu²⁺ and recognition of dicarboxylates utilizing Cu²⁺ complex. *Tetrahedron Lett.*, vol.47, No.16, (April 2006), pp. 2707-2710, ISSN 0040-4039
- Jiang, W. & Wang, W. (2009). A selective and sensitive "turn-on" fluorescent chemodosimeter for Hg²⁺ in aqueous media via Hg²⁺ promoted facile desulfurization-lactonization reaction. *Chem. Commun.*, No.26, (July 2009), pp. 3913-3915, ISSN 1359-7345
- Jung, H. S.; Han, J. H.; Habata, Y.; Kang, C. & Kim, J. S. (2011b). An iminocoumarin-Cu(II) ensemble-based chemodosimeter toward thiols. *Chem. Commun.*, vol.47, No.18, (May 2011), pp. 5142-5144, ISSN 1359-7345
- Jung, H. S.; Ko, K. C.; Kim, G.-H.; Lee, A.-R.; Na, Y.-C.; Kang, C.; Lee, J. Y. & Kim, J. S. (2011a). Coumarin-based thiol chemosensor: synthesis, turn-on mechanism, and its biological application. *Org. Lett.*, vol.13, No.6, (March 2011), pp. 1498-1501, ISSN 1523-7060
- Jung, H. S.; Ko, K. C.; Lee, J. H.; Kim, S. H.; Bhuniya, S.; Lee, J. Y.; Kim, Y.; Kim, S. J. & Kim, J. S. (2010). Rationally designed fluorescence turn-on sensors: a new design strategy based on orbital control. *Inorg. Chem.*, vol.49, No.18, (September 2010), pp. 8552-8557, ISSN 0020-1669
- Jung, H. S.; Kwon, P. S.; Lee, J. W.; Kim, J. I.; Hong, C. S.; Kim, J. W.; Yan, S.; Lee, J. Y.; Lee, J. H.; Joo, T. & Kim, J. S. (2009). Coumarin-derived Cu²⁺-selective fluorescence sensor: synthesis, mechanisms, and applications in living cells. *J. Am. Chem. Soc.*, vol.131, No.5, (February 2009), pp. 2008-2012, ISSN 0002-7863
- Kawaguchi, M.; Okabe, T.; Terai, T.; Hanaoka, K.; Kojima, H.; Minegishi, I. & Nagano, T. (2010). A time-resolved fluorescence probe for dipeptidyl peptidase 4 and its application in inhibitor screening. *Chem. Eur. J.*, vol.16, No.45, (December 2010), pp. 13479-13486, ISSN 0947-6539
- Kele, P.; Orbulescu, J.; Mello, S. V.; Mabrouki, M. & Leblanc, R. M. (2001). Langmuir-Blodgett film characterization of a new amphiphilic coumarin derivative. *Langmuir*, vol.17, No.23, (November 2001), pp. 7286-7290, ISSN 0743-7463.
- Kim, G.-J. & Kim, H.-J. (2010a). Doubly activated coumarin as a colorimetric and fluorescent chemodosimeter for cyanide. *Tetrahedron Lett.*, vol.51, No.1, (January 2010), pp. 185-187, ISSN 0040-4039
- Kim, G.-J. & Kim, H.-J. (2010b). Coumarinyl aldehyde as a Michael acceptor type of colorimetric and fluorescent probe for cyanide in water. *Tetrahedron Lett.*, vol.51, No.21, (May 2010), pp. 2914-2916, ISSN 0040-4039

- Kim, G.-J. & Kim, H.-J. (2010c). Highly selective and sensitive fluorescence turn-on probe for proline. *Tetrahedron Lett.*, vol.51, No.35, (September 2010), pp. 4670-4672, ISSN 0040-4039
- Kim, G.-J.; Lee, K.; Kwon, H. & Kim, H.-J. (2011). Ratiometric fluorescence imaging of cellular glutathione. *Org. Lett.*, vol.13, No.11, (June 2011), pp. 2799-2801, ISSN 1523-7060
- Kim, H. J.; Bhuniya, S.; Mahajan, R. K.; Puri, R.; Liu, H.; Ko, K. C.; Lee, J. Y. & Kim, J. S. (2009b). Fluorescence turn-on sensors for HSO_4^- . *Chem. Commun.*, No.46, (December 2009), pp. 7128-7130, ISSN 1359-7345
- Kim, H. J.; Kim, Y.; Kim, S. J.; Park, S. Y.; Lee, S. Y.; Kim, J. H.; No, K. & Kim, J. S. (2010). Iminocoumarin-based Hg(II) ion probe. *Bull. Korean Chem. Soc.*, vol.31, No.1, (January 2010), pp. 230-233, ISSN 0253-2964
- Kim, H. J.; Ko, K. C.; Lee, J. H.; Lee, J. Y. & Kim, J. S. (2011). KCN sensor: unique chromogenic and 'turn-on' fluorescent chemodosimeter: rapid response and high selectivity. *Chem. Commun.*, vol.47, No.10, (March 2011), pp. 2886-2888, ISSN 1359-7345
- Kim, H. J.; Park, J. E.; Choi, M. G.; Ahn, S. & Chang, S.-K. (2009a). Selective chromogenic and fluorogenic signalling of Hg^{2+} ions using a fluorescein-coumarin conjugate. *Dyes Pigments*, vol.84, No.1, (January 2010), pp. 54-58, ISSN 0143-7208
- Kim, H. M.; Yang, P. R.; Seo, M. S.; Yi, J.-S.; Hong, J. H.; Jeon, S.-J.; Ko, Y.-G.; Lee, K. J. & Cho, B. R. (2007). Magnesium ion selective two-photon fluorescent probe based on a benzo[h]chromene derivative for in vivo imaging. *J. Org. Chem.*, vol.72, No.6, (March 2007), pp. 2088-2096, ISSN 0022-3263
- Kim, J. H.; Kim, H. J.; Kim, S. H.; Lee, J. H.; Do, J. H.; Kim, H.-J.; Lee, J. H. & Kim, J. S. (2009). Fluorescent coumarinyldithiane as a selective chemodosimeter for mercury(II) ion in aqueous solution. *Tetrahedron Lett.*, vol.50, No.43, (October 2009), pp. 5958-5961, ISSN 0040-4039
- Kim, M. H.; Jang, H. H.; Yi, S.; Chang, S.-K. & Han, M. S. (2009). Coumarin-derivative-based off-on catalytic chemodosimeter for Cu^{2+} ions. *Chem. Commun.*, No.32, (August 2009), pp. 4838-4840, ISSN 1359-7345
- Kim, S. K.; Lee, D. H.; Hong, J.-I. & Yoon, J. (2009). Chemosensors for pyrophosphate. *Acc. Chem. Res.*, vol.42, No.1, (January 2009), pp. 23-31, ISSN 0001-4842
- Kim, S. Y.; Park, J.; Koh, M.; Park, S. B. & Hong, J.-I. (2009). Fluorescent probe for detection of fluoride in water and bioimaging in A549 human lung carcinoma cells. *Chem. Commun.*, No.31, (August 2009), pp. 4735-4737, ISSN 1359-7345
- Ko, K. C.; Wu, J.-S.; Kim, H. J.; Kwon, P. S.; Kim, J. W.; Bartsch, R. A.; Lee, J. Y. & Kim, J. S. (2011). Rationally designed fluorescence 'turn-on' sensor for Cu^{2+} . *Chem. Commun.*, vol.47, No.11, (March 2011), pp. 3165-3167, ISSN 1359-7345
- Kovalska, V. B.; Volkova, K. D.; Manaev, A. V.; Losytskyy, M. Y.; Okhrimenko, I. N.; Traven, V. F. & Yarmoluk, S. M. (2010). 2-Quinolone and coumarin polymethines for the detection of proteins using fluorescence. *Dyes Pigments*, vol.84, No.2, (February 2010), pp. 159-164, ISSN 0143-7208
- Kulatilleke, C. P.; de Silva, S. A. & Eliav, Y. (2006). A coumarin based fluorescent photoinduced electron transfer cation sensor. *Polyhedron*, vol.25, No.13, (September 2006), pp. 2593-2596, ISSN 0277-5387
- Kumar, A.; Pandey, M. K.; Anandakathir, R.; Mosurkal, R.; Parmar, V. S.; Watterson, A. C. & Kumar, J. (2010). Sensory response of pegylated and siloxanated 4,8-dimethylcoumarins: a fluorescence quenching study by nitro aromatics. *Sensors Actuators B*, vol.147, No.1, (May 2010), pp. 105-110, ISSN 0925-4005

- Kurishita, Y.; Kohira, T.; Ojida, A. & Hamachi, I. (2010). Rational design of FRET-based ratiometric chemosensors for in vitro and in cell fluorescence analyses of nucleoside polyphosphates. *J. Am. Chem. Soc.*, vol.132, No.38, (September 2010), pp. 13290-13299, ISSN 0002-7863
- Kwon, H.; Lee, K. & Kim, H.-J. (2011). Coumarin-malonitrile conjugate as a fluorescent turn-on probe for biothiols and its cellular expression. *Chem. Commun.*, vol.47, No.6, (February 2011), pp. 1773-1775, ISSN 1359-7345
- Lau, Y. H.; Price, J. R.; Todd, M. H. & Rutledge, P. J. (2011). A click fluorophore sensor that can distinguish Cu^{II} and Hg^{II} via selective anion-induced demetallation. *Chem. Eur. J.*, vol.17, No.10, (March 2011), pp. 2850-2858, ISSN 0947-6539
- Lee, D.-N.; Kim, G.-J. & Kim, H.-J. (2009). A fluorescent coumarinylalkyne probe for the selective detection of mercury(II) ion in water. *Tetrahedron Lett.*, vol.50, No.33, (August 2009), pp. 4766-4768, ISSN 0040-4039
- Lee, J. H.; Jeong, A. R.; Shin, I.-S.; Kim, H.-J. & Hong, J.-I. (2010). Fluorescence turn-on sensor for cyanide based on a cobalt(II)-coumarinylsalen complex. *Org. Lett.*, vol.12, No.4, (February 2010), pp. 764-767, ISSN 1523-7060
- Lee, K.-S.; Kim, H.-J.; Kim, G.-H.; Shin, I. & Hong, J.-I. (2008a). Fluorescent chemodosimeter for selective detection of cyanide in water. *Org. Lett.*, vol.10, No.1, (January 2008), pp. 49-51, ISSN 1523-7060
- Lee, K.-S.; Kim, T.-K.; Lee, J. H.; Kim, H.-J. & Hong, J.-I. (2008b). Fluorescence turn-on probe for homocysteine and cysteine in water. *Chem. Commun.*, No.46, (December 2008), pp. 6173-6175, ISSN 1359-7345
- Lee, M.; Gubernator, N. G.; Sulzer, D. & Sames, D. (2010). Development of pH-responsive fluorescent false neurotransmitters. *J. Am. Chem. Soc.*, vol.132, No.26, (July 2010), pp. 8828-8830, ISSN 0002-7863
- Lee, S. H.; Helal, A. & Kim, H.-S. (2010). Fluorescence sensing properties of thiazolobenzo-crown ether incorporating coumarin. *Bull. Korean Chem. Soc.*, vol.31, No.3, (March 2010), pp. 615-619, ISSN 0253-2964
- Li, H.; Cai, L.; Li, J.; Hu, Y.; Zhou, P. & Zhang, J. (2011). Novel coumarin fluorescent dyes: synthesis, structural characterization and recognition behavior towards Cu(II) and Ni(II). *Dyes Pigments*, vol.91, No.3, (December 2011), pp. 309-316, ISSN 0143-7208
- Li, H.; Gao, S. & Xi, Z. (2009). A colorimetric and "turn-on" fluorescent chemosensor for Zn(II) based on coumarin Schiff-base derivative. *Inorg. Chem. Commun.*, vol.12, No.4, (April 2009), pp. 300-303, ISSN 1387-7003
- Li, J.; Lin, H.; Cai, Z. & Lin, H. (2009). A novel coumarin-based switching-on fluorescent and colorimetric sensor for F⁻. *J. Lumin.*, vol.129, No.5, (May 2009), pp. 501-505, ISSN 0022-2313
- Li, N.; Xiang, Y. & Tong, A. (2010). Highly sensitive and selective "turn-on" fluorescent chemodosimeter for Cu²⁺ in water via Cu²⁺-promoted hydrolysis of lactone moiety in coumarin. *Chem. Commun.*, vol.46, No.19, (May 2010), pp. 3363-3365, ISSN 1359-7345
- Li, Y.; Wang, H.; Li, J.; Zheng, J.; Xu, X. & Yang, R. (2011). Simultaneous intracellular β -D-glucosidase and phosphodiesterase I activities measurements based on a triple-signaling fluorescent probe. *Anal. Chem.*, vol.83, No.4, (February 2011), pp. 1268-1274, ISSN 0003-2700
- Lim, N. C. & Brückner, C. (2004). DPA-substituted coumarins as chemosensors for zinc(II): modulation of the chemosensory characteristics by variation of the position of the chelate on the coumarin. *Chem. Commun.*, No.9, (May 2004), pp. 1094-1095, ISSN 1359-7345

- Lim, N. C.; Pavlova, S. V. & Brückner, C. (2009). Squaramide hydroxamate-based chemodosimeter responding to iron(III) with a fluorescence intensity increase. *Inorg. Chem.*, vol.48, No.3, (February 2009), pp. 1173-1182, ISSN 0020-1669
- Lim, N. C.; Schuster, J. V.; Porto, M. C.; Tanudra, M. A.; Yao, L.; Freake, H. C. & Brückner, C. (2005). Coumarin-based chemosensors for zinc(II): toward the determination of the design algorithm for CHEF-type and ratiometric probes. *Inorg. Chem.*, vol.44, No.6, (March 2005), pp. 2018-2030, ISSN 0020-1669
- Lim, S.-Y.; Lee, S.; Park, S. B. & Kim, H.-J. (2011). Highly selective fluorescence turn-on probe for glutathione. *Tetrahedron Lett.*, vol.52, No.30, (July 2011), pp. 3902-3904, ISSN 0040-4039
- Lin, W.; Cao, X.; Yuan, L. & Ding, Y. (2010b). Double functional group transformations for fluorescent probe construction: a fluorescence turn-on probe for thioureas. *Chem. Eur. J.*, vol.16, No.22, (June 2010), pp. 6454-6457, ISSN 0947-6539
- Lin, W.; Long, L. Tan, W. (2010a). A highly sensitive fluorescent probe for detection of benzenethiols in environmental samples and living cells. *Chem. Commun.*, vol.46, No.9, (March 2010), pp. 1503-1505, ISSN 1359-7345
- Lin, W.; Yuan, L.; Cao, X.; Tan, W. & Feng, Y. (2008). A coumarin-based chromogenic sensor for transition-metal ions showing ion-dependent bathochromic shift. *Eur. J. Org. Chem.*, No.29, (October 2008), pp. 4981-4987, ISSN 1434-193X
- Lin, W.; Yuan, L.; Cao, Z.; Feng, J. & Feng, Y. (2009). Fluorescence enhancement of coumarin-quinoline by transition metal ions: detection of paramagnetic Ni²⁺ and Co²⁺. *Dyes Pigments*, vol.83, No.1, (October 2009), pp. 14-20, ISSN 0143-7208
- Ma, Q.-J.; Zhang, X.-B.; Zhao, X.-H.; Jin, Z.; Mao, G.-J.; Shen, G.-L. & Yu, R.-Q. (2010). A highly selective fluorescent probe for Hg²⁺ based on a rhodamine-coumarin conjugate. *Anal. Chim. Acta*, vol.663, No.1, (March 2010), pp. 85-90, ISSN 0003-2670
- Ma, W.; Xu, Q.; Du, J.; Song, B.; Peng, X.; Wang, Z.; Li, G. & Wang, X. (2010). A Hg²⁺-selective chemodosimeter based on desulfurization of coumarin thiosemicarbazide in aqueous media. *Spectrochim. Acta A*, vol.76, No.2, (July 2010), pp. 248-252, ISSN 1386-1425
- Ma, Y. M. & Hider, R. C. (2009). The selective quantification of iron by hexadentate fluorescent probes. *Bioorg. Med. Chem.*, vol.17, No.23, (December 2009), pp. 8093-8101, ISSN 0960-894X
- Mahapatra, A. K.; Hazra, G.; Roy, J. & Sahoo, P. (2011). A simple coumarin-based colorimetric and ratiometric chemosensor for acetate and a selective fluorescence turn-on probe for iodide. *J. Lumin.*, vol.131, No.7, (July 2011), pp. 1255-1259, ISSN 0022-2313
- Maity, D. & Govindaraju, T. (2010). Conformationally constrained (coumarin-triazolyl-bipyridyl) click fluoroionophore as a selective Al³⁺ sensor. *Inorg. Chem.*, vol.49, No.16, (August 2010), pp. 7229-7231, ISSN 0020-1669
- Makwana, P. K.; Jethva, P. N. & Roy, I. (2011). Coumarin 6 and 1,6-diphenyl-1,3,5-hexatriene (DPH) as fluorescent probes to monitor protein aggregation. *Analyst*, vol.136, No.10, (May 2011), pp. 2161-2167, ISSN 0003-2654
- Meyer, Y.; Richard, J.-A.; Massonneau, M.; Renard, P.-Y. & Romieu, A. (2008). Development of a new nonpeptidic self-immolative spacer. Application to the design of protease sensing fluorogenic probes. *Org. Lett.*, vol.10, No.8, (April 2008), pp. 1517-1520, ISSN 1523-7060
- Miyaji, H.; Kim, H.-K.; Sim, E.-K.; Lee, C.-K.; Cho, W.-S.; Sessler, J. L. & Lee, C.-H. (2005). Coumarin-strapped calyx[4]pyrrole: a fluorogenic anion receptor modulated by

- cation and anion binding. *J. Am. Chem. Soc.*, vol.127, No.36, (September 2005), pp. 12510-12512, ISSN 0002-7863
- Mizukami, S.; Nagano, T.; Urano, Y.; Odani, A. & Kikuchi, K. (2002). A fluorescent anion sensor that works in neutral aqueous solution for bioanalytical application. *J. Am. Chem. Soc.*, vol.124, No.15, (April 2002), pp. 3920-3925, ISSN 0002-7863
- Mizukami, S.; Okada, S.; Kimura, S. & Kikuchi, K. (2009). Design and synthesis of coumarin-based Zn²⁺ probes for ratiometric fluorescence imaging. *Inorg. Chem.*, vol.48, No.16, (August 2009), pp. 7630-7638, ISSN 0020-1669
- Namba, K.; Kobayashi, K.; Murata, Y.; Hirakawa, H.; Yamagaki, T.; Iwashita, T.; Nishizawa, M.; Kusumoto, S. & Tanino, K. (2010). Mugineic acid derivatives as molecular probes for the mechanistic elucidation of iron acquisition in barley. *Angew. Chem. Int. Ed.*, vol.49, No.51, (December 2010), pp. 9956-9959, ISSN 1433-7851
- Nguyen, T. H. & Ansell, R. J. (2009). Fluorescent imprinted polymer sensors for chiral amines. *Org. Biomol. Chem.*, vol.7, No.6, (March 2009), pp. 1211-1220, ISSN 1477-0520
- Ortyl, J.; Sawicz, K. & Popielarz, R. (2010). Performance of amidocoumarins as probes for monitoring of cationic photopolymerization of monomers by fluorescence probe technology. *J. Polym. Sci. A*, vol.48, No.20, (October 2010), pp. 4522-4528, ISSN 1099-0518
- Pagona, G.; Economopoulos, S. P.; Tsikalas, G. K.; Katerionopoulos, H. E. & Tagmatarchis, N. (2010). Fullerene-coumarin dyad as a selective metal receptor: synthesis, photophysical properties, electrochemistry and ion binding studies. *Chem. Eur. J.*, vol.16, No.39, (October 2010), pp. 11969-11976, ISSN 0947-6539
- Pan, J.; Downing, J. A.; McHale, J. L. & Xian, M. (2009). A fluorogenic dye activated by S-nitrosothiols. *Mol. Biosyst.*, vol.5, No.9, (September 2009), pp. 918-920, ISSN 1742-206X
- Ray, D.; Nag, A.; Jana, A.; Goswami, D. & Bharadwaj, P. K. (2010). Coumarin derived chromophores in the donor-acceptor-donor format that give fluorescence enhancement and large two-photon activity in presence of specific metal ions. *Inorg. Chim. Acta*, vol.363, No.12, (October 2010), pp. 2824-2832, ISSN 0020-1693
- Richard, J.-A.; Massonneau, M.; Renard, P.-Y. & Romieu, A. (2008). 7-Hydroxycoumarin-hemicyanine hybrids: a new class of far-red emitting fluorogenic dyes. *Org. Lett.*, vol.10, No.19, (October 2008), pp. 4175-4178, ISSN 1523-7060
- Roussakis, E.; Pergantis, S. A. & Katerinopoulos, H. E. (2008). Coumarin-based ratiometric fluorescent indicators with high specificity for lead ions. *Chem. Commun.*, No.46, (December 2008), pp. 6221-6223, ISSN 1359-7345
- Royo, S.; Martínez-Mañez, R.; Sancenón, F.; Costero, A. M.; Parra, M. & Gil, S. (2007). Chromogenic and fluorogenic reagents for chemical warfare nerve agents' detection. *Chem. Commun.*, No.46, (December 2007), pp. 4839-4847, ISSN 1359-7345
- Ryu, D. H.; Noh, J. H. & Chang, S.-K. (2010). Selective ratiometric signaling of Hg²⁺ ions by a fluorescein-coumarin chemodosimeter. *Bull. Korean Chem. Soc.*, vol.31, No.1, (January 2010), pp. 246-249, ISSN 0253-2964
- Sakamoto, H.; Ishikawa, J.; Osuga, H.; Doi, K. & Wada, H. (2010). Highly silver ion selective fluorescent ionophore: fluorescent properties of polythiazaalkane derivatives bearing 8-(7-hydroxy-4-methyl)coumarinyl moiety in aqueous solution and in liquid-liquid extraction systems. *Analyst*, vol.135, No.3, (February 2010), pp. 550-558, ISSN 0003-2654

- Saleh, N.; Al-Soud, Y. A. & Nau, W. M. (2008). Novel fluorescent pH sensor based on coumarin with piperazine and imidazole substituents. *Spectrochim. Acta A*, vol.71, No.3, (December 2008), pp. 818-822, ISSN 1386-1425
- Santos, H. M.; Pedras, B.; Tamayo, A.; Casabó, J.; Escriche, L.; Covelo, B.; Capelo, J. L. & Lodeiro, C. (2009). New chemosensors based on thiomacrocyclic-containing coumarin-343 fluoroionophor: X-ray structures and previous results on the effect of cation binding on the photophysical properties. *Inorg. Chem. Commun.*, vol.12, No.11, (November 2009), pp. 1128-1134, ISSN 1387-7003
- Sato, S.; Suzuki, M.; Soma, T. & Tsunoda, M. (2008). Synthesis and properties of umbelliferone-nitroxide radical hybrid compounds as fluorescence and spin-label probes. *Spectrochim. Acta A*, vol.70, No.4, (September 2008), pp. 799-804, ISSN 1386-1425
- Secor, K. E. & Glass, T. E. (2004). Selective amine recognition: development of a chemosensor for dopamine and norepinephrine. *Org. Lett.*, vol.6, No.21, (October 2004), pp. 3727-3730, ISSN 1523-7060
- Sethna, S. M. & Shah, N. M. (1945). The chemistry of coumarins. *Chem. Rev.*, vol.36, No.1, (February 1945), pp. 1-62, ISSN 0009-2665
- Shao, J. (2010). A novel colorimetric and fluorescence anion sensor with a urea group as binding site and a coumarin group as signal unit. *Dyes Pigments*, vol.87, No.3, (November 2010), pp. 272-276, ISSN 0143-7208
- Sheng, J. R.; Feng, F.; Qiang, Y.; Liang, F. G.; Sen, L. & Wei, F.-H. (2008). A coumarin-derived fluorescence chemosensors selective for copper(II). *Anal. Lett.*, vol.41, No.12, (August 2008), pp. 2203-2213, ISSN 0003-2719
- Sheng, R.; Wang, P.; Liu, W.; Wu, X. & Wu, S. (2008). A new colorimetric chemosensor for Hg²⁺ based on coumarin azine derivatives. *Sensors Actuators B*, vol.128, No.2, (January 2008), pp. 507-511, ISSN 0925-4005
- Shibata, A.; Abe, H.; Ito, M.; Kondo, Y.; Shimizu, S.; Aikawa, K. & Ito, Y. (2009). DNA templated nucleophilic aromatic substitution reactions for fluorogenic sensing of oligonucleotides. *Chem. Commun.*, No.43, (November 2009), pp. 6586-6588, ISSN 1359-7345
- Shiraishi, Y.; Sumiya, S. & Hirai, T. (2010). A coumarin-thiourea conjugate as a fluorescent probe for Hg(II) in aqueous media with a broad pH range 2-12. *Org. Biomol. Chem.*, vol.8, No.6, (March 2010), pp. 1310-1314, ISSN 1477-0520
- Shiraishi, Y.; Sumiya, S. & Hirai, T. (2011). Highly sensitive cyanide anion detection with a coumarin-spiropyran conjugate as a fluorescent receptor. *Chem. Commun.*, vol.47, No.17, (May 2011), pp. 4953-4955, ISSN 1359-7345
- Singh, R. K.; Mandal, T.; Balasubramanian, N.; Cook, G. & Srivastava, D. K. (2011). Coumarin-suberoylanilide hydroxamic acid as a fluorescent probe for determining binding affinities and off-rates of histone deacetylase inhibitors. *Anal. Biochem.*, vol.408, No.2, (January 2011), pp. 309-315, ISSN 0003-2697
- Sokkalingam, P. & Lee, C.-H. (2011). Highly sensitive fluorescence "turn-on" indicator for fluoride anion with remarkable selectivity in organic and aqueous media. *J. Org. Chem.*, vol.76, No.10, (May 2011), pp. 3820-3828, ISSN 0022-3263
- Su, Z.; Chen, K.; Guo, Y.; Qi, H.; Yang, X.-F. & Zhao, M. (2010). A coumarin-based fluorescent chemosensor for Zn²⁺ in aqueous ethanol media. *J. Fluoresc.*, vol.20, No.4, (July 2010), pp. 851-856, ISSN 1053-0509
- Suresh, M. & Das, A. (2009). New coumarin-based sensor molecule for magnesium and calcium ions. *Tetrahedron Lett.*, vol.50, No.42, (October 2009), pp. 5808-5812, ISSN 0040-4039

- Suzuki, Y.; Komatsu, H.; Ikeda, T.; Saito, N.; Araki, S.; Citterio, D.; Hisamoto, H.; Kitamura, Y.; Kubota, T.; Nakagawa, J.; Oka, K. & Suzuki, K. (2002). Design and synthesis of Mg^{2+} -selective fluoroionophores based on a coumarin derivative and application for Mg^{2+} measurement in a living cell. *Anal. Chem.*, vol.74, No.6, (March 2002), pp. 1423-1428, ISSN 0003-2700
- Taki, M.; Desaki, M.; Ojida, A.; Iyoshi, S.; Hirayama, T.; Hamachi, I. & Yamamoto, Y. (2008). Fluorescence imaging of intracellular cadmium using a dual-excitation ratiometric chemosensor. *J. Am. Chem. Soc.*, vol.130, No.38, (September 2008), pp. 12564-12565, ISSN 0002-7863
- Trenor, S. R.; Shultz, A. R.; Love, B. J. & Long, T. E. (2004). Coumarins in polymers: from light harvesting to photo-cross-linkable tissue scaffolds. *Chem. Rev.*, vol.104, No.6, (June 2004), pp. 3059-3077, ISSN 0009-2665
- Tsai, C.-S.; Liu, P.-Y.; Yen, H.-Y.; Hsu, T.-L. & Wong, C.-H. (2010). Development of trifunctional probes for glycoproteomic analysis. *Chem. Commun.*, vol.46, No.30, (August 2010), pp. 5575-5577, ISSN 1359-7345
- Tsakamoto, K.; Shinohara, Y.; Iwasaki, S. & Maeda, H. (2011). A coumarin-based fluorescent probe for Hg^{2+} and Ag^{+} with an *N*-acetylthioureido group as a fluorescence switch. *Chem. Commun.*, vol.47, No.17, (May 2011), pp. 5073-5075, ISSN 1359-7345
- Upadhyay, K. K. & Mishra, R. K. (2010). Zn^{2+} specific colorimetric receptor based on coumarin. *Bull. Chem. Soc. Jpn.*, vol.83, No.10, (October 2010), pp. 1211-1215, ISSN 0009-2673
- Upadhyay, K. K.; Mishra, R. K.; Kumar, V. & Chowdhury, P. K. R. (2010a). A coumarin based ICT probe for fluoride in aqueous medium with its real application. *Talanta*, vol.82, No.1, (June 2010), pp. 312-318, ISSN 0039-9140
- Upadhyay, K. K.; Mishra, R. K.; Kumar, A.; Zhao, J. & Prasad, R. (2010b). Self assembled pseudo double helix architecture and anion sensing behavior of a coumarin based ICT probe. *J. Mol. Struct.*, vol.963, No.2-3, (January 2010), pp. 228-233, ISSN 0022-2860
- Vasylevska, A. S.; Karasyov, A. A.; Borisov, S. M. & Krause, C. (2007). Novel coumarin-based fluorescent pH indicators, probes and membranes covering a broad pH range. *Anal. Bioanal. Chem.*, vol.387, No.6, (March 2007), pp. 2131-2141, ISSN 1618-2642
- Voutsadaki, S.; Tsikalas, G. K.; Klontzas, E.; Froudakis, G. E. & Katerinopoulos, H. E. (2010). A "turn-on" coumarin-based fluorescent sensor with high selectivity for mercury ions in aqueous media. *Chem. Commun.*, vol.46, No.19, (May 2010), pp. 3292-3294, ISSN 1359-7345
- Wallace, K. J.; Fagbemi, R. I.; Folmer-Andersen, F. J.; Morey, J.; Lynth, V. M. & Anslyn, E. V. (2006). Detection of chemical warfare simulants by phosphorylation of a coumarin oximate. *Chem. Commun.*, No.37, (October 2006), pp. 3886-3888, ISSN 1359-7345
- Wang, J.; Qian, X. & Cui, J. (2006). Detecting Hg^{2+} ions with an ICT fluorescent sensor molecule: remarkable emission spectra shift and unique selectivity. *J. Org. Chem.*, vol.71, No.11, (May 2006), pp. 4308-4311, ISSN 0022-3263
- Welser, K.; Perera, M. D. A.; Aylott, J. W. & Chan, W. C. (2009). A facile method to clickable sensing polymeric nanoparticles. *Chem. Commun.*, No.43, (November 2009), pp. 6601-6603, ISSN 1359-7345
- Wu, J.; Sheng, R.; Liu, W.; Wang, P.; Ma, J.; Zhang, H. & Zhuang, X. (2011). Reversible fluorescent probe for highly selective detection of mercapto biomolecules. *Inorg. Chem.*, vol.50, No.14, (July 2011), pp. 6543-6551, ISSN 0020-1669

- Wu, W.; Wu, W.; Ji, S.; Guo, H. & Zhao, J. (2011). Accessing the long-lived emissive ^3IL triplet excited states of coumarin fluorophores by direct cyclometallation and its application for oxygen sensing and upconversion. *Dalton Trans.*, vol.40, No.22, (June 2011), pp. 5953-5963, ISSN 1477-9226
- Xiao, D.; Martini, L. A.; Snoeberger III, R. C.; Crabtree, R. H. & Batista, V. S. (2011). Inverse design and synthesis of acac-coumarin anchors for robust TiO_2 sensitization. *J. Am. Chem. Soc.*, vol.133, No.23, (June 2011), pp. 9014-9022, ISSN 0002-7863
- Xie, Y.; Dix, A. V. & Tor, Y. (2009). FRET enabled real time detection of RNA-small molecule binding. *J. Am. Chem. Soc.*, vol.131, No.48, (December 2009), pp. 17605-17614, ISSN 0002-7863
- Xiong, M.; Xi, H.; Fu, Y. & Sun, X. (2010). Synthesis and properties of novel coumarin-based fluorescent probes for identifying melamine. *Chin. J. Org. Chem.*, vol.30, No.6, (June 2010), pp. 908-911, ISSN 0253-2786
- Yan, M.; Li, T. & Yang, Z. (2011). A novel coumarin Schiff-base as a Zn(II) ion fluorescent sensor. *Inorg. Chem. Commun.*, vol.14, No.3, (March 2011), pp. 463-465, ISSN 1387-7003
- Yao, J.; Dou, W.; Qin, W. & Liu, W. (2009). A new coumarin-based chemosensor for Fe^{3+} in water. *Inorg. Chem. Commun.*, vol.12, No.2, (February 2009), pp. 116-118, ISSN 1387-7003
- Yi, L.; Li, H.; Sun, L.; Liu, L.; Zhang, C. & Xi, Z. (2009). A highly sensitive fluorescence probe for fast thiol-quantification assay of glutathione reductase. *Angew. Chem. Int. Ed.*, vol.48, No.22, (May 2009), pp. 4034-4037, ISSN 1433-7851
- Yuan, L.; Lin, W. & Song, J. (2010). Ratiometric fluorescent detection of intracellular hydroxyl radicals based on a hybrid coumarin-cyanine platform. *Chem. Commun.*, vol.46, No.42, (November 2010), pp. 7930-7932, ISSN 1359-7345
- Yuan, L.; Lin, W. & Yang, Y. (2011). A ratiometric fluorescent probe for specific detection of cysteine over homocysteine and glutathione based on the drastic distinction in the kinetic profiles. *Chem. Commun.*, vol.47, No.22, (June 2011), pp. 6275-6277, ISSN 1359-7345
- Zhang, H. & Rudkevich, D. M. (2007). A FRET approach to phosgene detection. *Chem. Commun.*, No.12, (March 2007), pp. 1238-1239, ISSN 1359-7345
- Zhao, X.; Zhang, Y.; He, G. & Zhou, P. (2010). Highly sensitive fluorescent coumarin-based probes for selective detection of copper ion. *Chin. J. Lumin.*, vol.31, No.3, (June 2010), pp. 433-438, ISSN 1000-7032
- Zhou, Y.; Liu, K.; Li, J.-Y.; Fang, Y.; Zhao, T.-C. & Yao, C. (2011). Visualization of nitroxyl in living cells by a chelated copper(II) coumarin complex. *Org. Lett.*, vol.13, No.6, (March 2011), pp. 1290-1293, ISSN 1523-7060
- Zhuang, X.; Liu, W.; Wu, J.; Zhang, H. & Wang, P. (2011). A novel fluoride ion colorimetric chemosensor based on coumarin. *Spectrochim. Acta A*, vol.79, No.5, (September 2011), pp. 1352-1355, ISSN 1386-1425
- Zuo, Q.-P.; Li, B.; Pei, Q.; Li, Z. & Liu, S.-K. (2010). A highly selective fluorescent probe for detection of biological samples thiol and its application in living cells. *J. Fluoresc.*, vol.20, No.6, (November 2010), pp. 1307-1313, ISSN 1053-0509



Advances in Chemical Sensors

Edited by Prof. Wen Wang

ISBN 978-953-307-792-5

Hard cover, 358 pages

Publisher InTech

Published online 20, January, 2012

Published in print edition January, 2012

The chemical sensor plays an essential role in the fields of environmental conservation and monitoring, disaster and disease prevention, and industrial analysis. A typical chemical sensor is a device that transforms chemical information in a selective and reversible way, ranging from the concentration of a specific sample component to total composition analysis, into an analytically useful signal. Much research work has been performed to achieve a chemical sensor with such excellent qualities as quick response, low cost, small size, superior sensitivity, good reversibility and selectivity, and excellent detection limit. This book introduces the latest advances on chemical sensors. It consists of 15 chapters composed by the researchers active in the field of chemical sensors, and is divided into 5 sections according to the classification following the principles of signal transducer. This collection of up-to-date information and the latest research progress on chemical sensor will provide valuable references and learning materials for all those working in the field of chemical sensors.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Hongqi Li, Li Cai and Zhen Chen (2012). Coumarin-Derived Fluorescent Chemosensors, *Advances in Chemical Sensors*, Prof. Wen Wang (Ed.), ISBN: 978-953-307-792-5, InTech, Available from: <http://www.intechopen.com/books/advances-in-chemical-sensors/coumarin-derived-fluorescent-chemosensors>

INTECH

open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2012 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.