

Therapeutic Targets in Colorectal Cancer

Rajunor Ettarh¹, Alvis Calamai² and Anthony Cullen²

¹*Department of Structural and Cellular Biology,*

Tulane University School of Medicine, New Orleans,

²*School of Medicine and Medical Sciences, University College, Dublin,*

¹USA

²Ireland

1. Introduction

Colon cancer is common worldwide: nearly a million people develop the disease every year and in the United States, colorectal cancer ranks third for frequency of occurrence and mortality in both men and women, with projected estimates for 2011 for occurrence and mortality put respectively at approximately 140,000 and 49,000 (American Cancer Society, 2011; Jemal et al, 2005). The projection for total deaths from all cancers in 2010 was 569,490 (Aliperti et al, 2011).

Significant progress in understanding colon cancer has produced a wealth of information that has aided improvements in aspects of diagnosis and disease management, contributing in the process to reduced mortality rates. The mechanisms that facilitate colorectal carcinogenesis and sustain progression and metastatic spread have been extensively investigated. The cause of colorectal cancer is multi-factorial. Notwithstanding the various contributing elements to the disease, the primary manifestation of colorectal carcinoma is the relentless and uncontrolled proliferation of cells and tissues in the intestinal mucosal epithelium. This pattern of abnormal proliferation is a disruption of the normal balance between new cell production by the epithelial cells in the mucosal crypts, and the release and loss of epithelial cells into the intestinal lumen i.e. cell-producing proliferation is normally finely and properly counter-balanced by regulated apoptotic and physical cell loss (Raz, 2002).

Given the multistep, multifactor origins of colorectal cancer, the rationale for targeted therapies and the identification of therapeutic targets is that the disease can be (a) prevented prior to initiation (b) obstructed in its progression by blocking or inhibiting mechanisms that sustain progression and facilitate metastasis (c) reversed. The list of potential targets include microbes and bacteria that facilitate tumor initiation, molecular targets such as adenomatous polyposis coli (APC), and cancer stem cells (CSCs) where targeted destruction is thought to be central to preventing metastatic tumor spread.

As with all cancers, finding and delivering therapeutic targets in colorectal cancer is based on the premise that there is one originating cell type (van der Flier & Clevers, 2009). If this population of mutant originating cells is eliminated, the ability for new initiation, progression and distant seeding of tumor cells should be impaired and eventually abolished. Several therapeutic approaches have shown promising results in experimental

studies. However, this chapter will focus largely on molecular targets in Wnt signaling, the nuclear receptor peroxisome proliferator-activated receptor (PPAR), and cancer stem cells (also known as cancer initiating cells).

2. Colonic epithelial cell renewal

The colon is the distal part of the intestinal tract and is lined internally by a simple layer of columnar epithelial cells (colonocytes) that send tube-like extensions called crypts into the mucosal layer of the intestinal wall. The crypts provide a conducive environment for the regulation and renewal of the epithelial covering of the colonic mucosa.

The epithelial cells in the crypt divide continuously and rapidly, achieving a turnover rate of epithelial renewal of between 5-6 days in mammals, with much shorter cell kinetic data reported for rodents (Di Garbo et al, 2010; Hall et al, 1994; Heath, 1996; Giles et al, 2003; Li et al, 1994; Loeffler et al, 1986; Potten & Loeffler, 1990; Okamoto & Watanabe, 2004; Wright & Alison, 1984). In the small intestine, between 8-9 cells are produced by each crypt epithelium every hour in mice; 2-3 dividing cells per crypt support cell production in the proximal intestine while up to 5 dividing cells are required to maintain cell production in the distal intestine (McGarvey et al, 2007a, 2007b). The renewal mechanism is sustained by a hierarchical arrangement of epithelial cells within the crypts, exemplified by the model described by Tomlinson and Bodmer (1995), with stem cells thought to reside in the lower part of the crypts, while differentiated cells populate the upper part of the crypt. By dividing and supplying transit (semi-differentiated) cells that migrate up the crypts, the stem cells are capable of and responsible for producing the various cell types that are found in the colonic epithelium. Differentiated cells at the top of the crypt and colonic mucosal surface eventually undergo spontaneous apoptosis and are released into the intestinal lumen (Fig 1).




Fig. 1. Schematic diagram of colonic crypt, illustrating the three zones and cell categories that constitute the kinetic framework for cell production and regeneration.

The maintenance of functional and structural integrity and viability of the enteric mucosal epithelium depends on the preservation of the crypt cell renewing and emigration mechanisms for repopulating the continuously shedding epithelial cell cover (McGarvey et al, 2007a, 2007b). Several models for investigating the dynamics of colon cell regulation have been described (Boman et al, 2001; Hardy & Stark, 2002; Lander, 2009; Michor et al, 2004; Paulus et al, 1992; van Leeuwen et al, 2006; Wodarz, 2007). Many of these models have been employed in studies of the mechanisms that underlie normal colonic epithelial cell regulation and regeneration, as well as the dysregulated proliferation in colorectal cancer.

3. Apoptosis

All of the new cells that are produced by proliferation of the cells in the stem cell compartment of the crypt, and numerically amplified in the semi-differentiated compartment, are distributed to the colonic mucosal epithelium to provide functionally important roles in absorption and secretion as well as providing a selectively permeable surface cover (Hall et al, 1994). The supply of new cells towards the upper crypt and surface epithelium is designed to satisfy the losses caused by cell injury, loss and programmed death (apoptosis). Surface cover cells are therefore removed or shed by processes that are as controlled and as balanced as the crypt-mediated cell renewal mechanism, and involves a cessation of proliferative processes in conjunction with the initiation of disposal and cell loss pathways (Leblond, 1964; Wright & Alison, 1984; Hall et al, 1994). Because the enteric epithelium is associated with underlying connective tissue fibroblasts, the accompanying fluxes in these cells are also correspondingly regulated in a controlled manner for proliferation and for cell loss (Marsh & Trier, 1974a, 1974b; Parker et al, 1974; Pascal et al, 1968a, 1968b). Together, the careful balance of cell production and cell loss maintains homeostasis in the colonic epithelium. Apoptosis does not occur randomly, rather it is seen towards the distal end of the cell migration route up the crypt (Hall et al, 1994). In colon cancer, proliferation is elevated and apoptosis is dysregulated, making the restoration of apoptosis an attractive proposition for therapeutic control of colon cancer growth (Evan & Vousden, 2001, Johnstone et al, 2002).

A number of cyclooxygenase (COX) inhibitors induce apoptosis by activating mechanisms that are either upstream (via the lipid metabolite 13-S-hydroxyoctadecadienoic acid) or downstream (via 14-3-3 ϵ proteins) of the nuclear hormone receptor PPAR δ (Liou et al, 2007; Shureiqi et al, 2003), indicating that the pro-apoptotic effect of COX inhibitors on cancer cells is dependent on down-regulation of PPAR δ . In APC min mice, short-term treatment with nitric-oxide-donating aspirin (NO-ASA) induces apoptosis in differentiated intestinal epithelial cells while prolonged treatment with sulindac reverses the anti-apoptotic effect of APC (Mahmoud et al, 1998; Ouyang et al, 2006). In contrast, celecoxib administration produces no effect on apoptosis (Williams et al, 2000).

Other agents that have been shown to reduce colorectal cancer growth *in vitro* include CDDO-Me, an oleanane synthetic triterpenoid that achieves its apoptotic effect partly through the generation of reactive oxygen (ROS) and the activation of procaspases (Gao et al, 2011), green tea polyphenols that achieve their apoptotic effect through the induction of caspases (Oz & Ebersole, 2010), and tocotrienol, a member of the vitamin E family of compounds that induces morphological changes similar to apoptosis (paraptosis) and an accompanying reduction in Wnt signaling and its down-stream genes (Zhang et al, 2011).

4. Wnt and colorectal cancer

One of the primary regulators of epithelial cell proliferation is Wnt signaling (Di Garbo et al, 2010). This signaling pathway involves the intermediate elements beta catenin, glycogen synthase kinase 3 beta (GSK3 β), casein kinase I (CKI), axin, adenomatous polyposis coli (APC) and T-cell factor/lymphoid enhancer factor (TCF/LEF). Inappropriate activation or disruption of Wnt signaling upsets the careful regulatory balance in epithelial kinetics, leads to disorderly proliferation, and is an important contributor to the process of colorectal carcinogenesis. Wnt signaling helps to control the levels of cytoplasmic beta catenin, between pools bound to APC and to the cell adhesion molecule E-cadherin. The APC-bound pool of beta catenin is held in a stable complex of axin, GSK3 β , CKI and APC that serves to regulate its cytoplasmic levels via targeted ubiquitin-mediated proteasomal degradation (Kikuchi et al, 2003; Pinto & Clevers, 2005). Wnt ligand signaling via membrane receptor proteins triggers a cascade that alters the relationship between the scaffold protein axin and GSK3 β , interrupts regulated destruction of beta catenin, and leads to accumulation of non-phosphorylated beta catenin in the cytoplasm that then reaches the nucleus. Translocation of beta catenin into the nucleus after binding with TCF/LEF leads to the activation of target genes that regulate proliferation, differentiation and apoptosis (Araki et al, 2003; Coghlan et al, 2000; DiGarbo et al, 2010; Fagotto et al, 1998; He et al, 1998; Kishida et al, 1999; Shtutman et al, 1999; Tetsu & McCormick, 1999; van der Flier & Clevers, 2009; Yamamoto et al, 1999; Yanagawa et al, 1995; Yost et al, 1996). Direct binding of TCF to regulatory elements in downstream genes have aided identification of target genes and suggest that Wnt-activated gene expression shows a gradient-wise concentration of activity in intestinal crypts with the highest expression in the bottom of the crypt (Gregorieff et al, 2005). Most of these target genes are expressed in normal crypts and in adenomas (van der Flier et al, 2007; van der Wetering et al, 2002).

5. Wnt and COX inhibition

Colon cancer is associated with dysregulation and overexpression of COX, a key enzyme in the biosynthetic conversion of arachidonic acid to eicosanoids (Botting, 2006). Increased levels of expression of COX-2 are seen in up to 85% of colorectal adenomas and carcinomas (Eberhart et al, 1994; Fujita et al, 1998; Rigas et al, 1993; Sheng et al, 1997).

COX inhibitors demonstrate an ability to disrupt proliferation in several CRC cell lines. In HT29 colorectal adenoma cell lines, suppression of proliferation is evident as early as 48 hours after treatment with naproxen and piroxicam and at later timepoints with aspirin, indomethacin, aspirin and NS398 (Shiff et al, 1996; Shureiqi et al, 2000). But in some studies, naproxen and salicylic acid showed no effect on proliferation in the same cell lines pointing to differing potencies for inhibition of COX as well as effects on growth and apoptosis (Piazza et al, 1997). Although anti-proliferative effects have been reported in studies using HCA7, HT115 and SW620 cell lines which all express COX, the non-COX expressing cell line HT116 also shows reduced growth when treated with celecoxib for 72 hours (Shureiqi et al, 2003). Most of the evidence allows the conclusion that the anti-proliferative effects of COX inhibitors on colon cancer cell lines are not related to COX expression or activity.

When COX inhibitors are administered to APC min mice, initiation and progression of intestinal and colonic polyps is inhibited and polyp load is reduced (Jacoby et al, 1996, 2000; Kohno et al, 2005; Mahmoud et al, 1998, Moorghen et al, 1988, 1998; Narisawa et al, 1983;

Rao et al, 1995, 2009; Reddy et al, 1993). Prevention of tumorigenesis or tumor load reduction reflects either decreased cell proliferation or increased cell death but findings from animal studies are inconsistent (Table 1). For example, celecoxib treatment reduces tumor numbers and inhibits cell proliferation but data from studies using various sulindac preparations point to a variability that may be rodent species dependent (Jacoby et al, 2000; Mahmoud et al, 1998; Moorghen et al, 1988, 1998; Rao et al 1995, 2009).

| Model | Inhibitor | Dose & duration (wks) | | Inhibition effect | Reference |
|-------------|-------------------------|-----------------------|------|-------------------|----------------------------|
| APC | | | | | |
| mouse | sulindac | 160ppm | 10 | none | <i>Shiff et al 1996</i> |
| mouse | sulindac S ₂ | 20mg/kg | 11 | none | <i>Swamy et al 2006</i> |
| mouse | celecoxib | 1500ppm | 6 | tumor number | <i>Han et al 2008</i> |
| DMH | | | | | |
| mouse | sulindac | 5mg/kg | 24 | tumorigenesis | <i>Shureiqi et al 2000</i> |
| mouse | sulindac | 5mg/kg | 18 | n/a | <i>Kim et al 2009</i> |
| AOM | | | | | |
| mouse | nimesulide | 0.04%w/w | 14 | n/a | <i>Shureiqi et al 2003</i> |
| rat | celecoxib | 300ppm | 46 | n/a | <i>Guo et al 2009</i> |
| rat | aspirin | 200-400ppm | 52 | tumorigenesis | <i>Piazza et al 1997</i> |
| NMNU | | | | | |
| rat | indomethacin | 10ppm | 1-30 | tumorigenesis | <i>Hanif et al 1996</i> |

Table 1. Effect of COX inhibitors on initiation and progression of experimental colon cancer in vivo. S₂ = sulfide, NMNU = n-methyl-N-nitrosourea, AOM = azoxymethane, DMH = 1,2-dimethylhydrazine, APC = adenomatous polyposis coli, n/a = not measured

Some of the inconsistency in findings from animal studies is reflected in the results from clinical investigations in patients. Treatment with aspirin and celecoxib shows beneficial prevention of colorectal cancer in patients, and treatment with 150mg sulindac twice daily for nine months reduces number and size of colorectal adenomas. However, treatment with standard sulindac doses (25-150 mg twice daily) for 48 months did not prevent adenomas in patients (Giardiello et al, 1993, 2002; Giovannucci et al, 1994; Lanas & Fernandez, 2009; Thun et al, 1991).

6. PPAR and COX inhibition

Peroxisome proliferator-activated receptors (PPAR) are part of the nuclear hormone receptor superfamily. While PPAR α and PPAR γ have been shown to be involved in various aspects of dietary lipid and glucose metabolism, PPAR δ is implicated in the control of cell proliferation, differentiation and colorectal carcinogenesis (Desvergne & Wahil, 1999; Michalik et al, 2003; Wang & Dubois, 2010). Ligand activation of PPAR δ is associated with suppressed induction of colon cancer (genetic and chemical treatment models) in mice via mechanisms that are linked to colonocyte differentiation and apoptosis (Harman et al, 2004;

Marin et al, 2006). Conversely, inactivation of PPAR δ in APC-min mice enhances predisposition to multiple intestinal and colorectal polyps (Harman et al, 2004; Reed et al, 2004). Such evidence suggests that PPAR δ attenuates colon cancer. However, Park and colleagues found a reduction in the ability of PPAR δ -/(null) cells to form tumors in nude mice and they concluded that PPAR δ might function to assist the tumor-suppressing function of adenomatous polyposis coli (APC) protein (Park et al, 2001).

Despite significant insights into the role of PPAR δ in colorectal cancer, the physiological role of PPAR δ in epithelia is still not completely understood. The unresolved nature of the available data has not prevented studies that have explored the possibility of targeting PPAR δ therapeutically in colorectal cancer. Prostacyclin I₂ can act as a natural ligand for PPAR δ (Gupta et al, 2000), and because COX-2 inhibitors can suppress carcinogenesis and reduce intestinal polyposis (Hollingshead et al, 2008; Jacoby et al, 1996; Mahmoud et al, 1998), a number of studies examined the use of COX inhibition to influence PPAR δ activity. Sulindac and indomethacin inhibit colorectal carcinogenesis in vitro by rapidly downregulating transcriptional activity of PPAR δ via disruption of DNA binding to PPAR δ -response elements (He et al, 1999). A similar effect on PPAR δ is also observed following administration of sulindac and celecoxib but this is preceded by induction of the enzyme 15-lipoxygenase-1 (Shureiqi et al, 2003). Administration of nitric-oxide-donating aspirin reduces PPAR δ expression and intestinal polyp numbers in mice but neither nimesulide nor GW0742 (a PPAR δ ligand) has an effect on PPAR δ mRNA levels, despite the fact that both agents reduce intestinal polyp numbers (Gupta et al, 2004; Hollingshead et al, 2008; Kohno et al, 2005).

COX-2 inhibitors and PPAR δ ligands can separately attenuate cancer growth, however combinatorial protocols have so far failed to produce potentiated inhibition of colon cancer indicating that COX-inhibitory and PPAR δ pathways are mechanistically separate (Hollingshead et al, 2008). In addition, concurrent expression of PPAR δ and COX-2 in colorectal tumors has poor prognostic implications for patients (Yoshinaga et al, 2011).

Ligand activation of PPAR γ is also anti-neoplastic in several tissues, but the data regarding its role in colorectal cancer is just as conflicting as the data for PPAR δ . PPAR γ activation inhibits colon cancer cell growth in vitro whereas a mutation-dependent pro-tumorigenic effect has been reported in vivo (Girnun et al, 2002; Yoshizumi et al, 2004). The mechanistically interrelated and inter-dependent nature of colorectal cancer is illustrated by the finding that PPAR γ agonists induce apoptosis by suppressing activation of NF κ B and GSK3 β (Ban et al, 2010). Other investigators have shown that PPAR γ induces apoptosis via inactivation of survivin and activation of caspase-3 in colorectal cancer cell lines and were able to inhibit PPAR γ -ligand induced apoptosis by activating PPAR δ (Wang et al, 2011).

7. Clones and stem cells

The crypt structure of the colonic epithelium is maintained by the putative presence of pluripotent intestinal crypt stem cells (Schmidt et al, 1988). Initially crypts are polyclonal and subsequently become monoclonal. Two kinetic models of the stem-cell-sustained intestinal crypt have been described. In the classic model, intestinal stem cells are thought to reside in the 4th cell position from the bottom of the crypt (the +4 cell). These stem cells supply daughter cells to the proliferative, transit-amplifying zone of the crypt; stem cells can be replaced by these daughter cells if necessary (Marshman et al, 2002; Potten, 1977; Potten et al, 1974, 2002). The zone model localizes stem cells to the bottom of the crypt; these cells are proposed to be the undifferentiated crypt base columnar (CBC) cells (Bjerknes & Cheng

1981a, 1981b, 1999, 2006). On the basis of modelling studies, it is proposed that stem cells and crypts can suffer losses and be replaced (Cairnie & Millen, 1975; Nicolas et al, 2007; Yatabe et al, 2001).

Unequivocal stem cell identification has long remained elusive but, using genetic lineage tracing experiments, Barker et al (2007) showed that Lgr5, a G-protein-coupled receptor, is expressed in CBC cells. The study followed Lgr5-positive daughter cells up intestinal crypts and on to the intestinal villous epithelium, where all differentiated epithelial cell types could be demonstrated. The ability of Lgr5-positive stem cells in the crypt to give rise to crypt-villus units appear to be dependent on proximity to CD24+ cells at the bottom of the crypt (Sato et al, 2011). Stem cells have also been identified in mammalian epidermal hair follicles where they express Lgr6 (Snippert et al, 2010). Deletion of the APC gene in crypt stem cells in Lgr5 knock-in mice facilitates intestinal microadenoma growth; deletion of APC in transit-amplifying, semi-differentiated crypt cells in Lgr5 knock-in mice significantly reduces the growth of intestinal adenomas. Together this suggests that APC loss needs to be stem cell specific to propagate unrestrained tumor growth (Barker et al, 2009). The finding that single isolated Lgr5-positive stem cells can give rise to self-organizing crypt-villus units (Sato et al, 2009) raises the possibility that these cells may be useful in treatment strategies that aim to repopulate enteric epithelia.

There is experimental evidence for several proposed colon cancer stem cell markers including CD133, CD44, CD166, the extracellular matrix protein olfactomedin-4 (OLFM4), aldehyde dehydrogenase (ALDH1A1), Lgr5, and pleckstrin homology-like domain family A member 1 (PHLDA1). Some of these markers are associated with IL6-STAT3-JAK2 signaling (Becker et al, 2008; Dalerba et al, 2007; O'Brien et al, 2007; Ricci-Vitani et al, 2007; Sakthianandeswaren et al, 2011; Sanders & Majumdar, 2011; Shmelkov et al, 2008; Tsai et al, 2011; Uchida et al, 2010; van der Flier et al, 2009).

In contrast to the idea that carcinogenic mutations can occur in any cell, the cancer stem cell model (first described in 1997 for hematologic malignancies) proposes that tumor transformation, progression and metastatic initiation is driven by the acquisition of oncogenic self-renewal properties by tissue stem cells, contributing to differentiation and the cellular heterogeneity of tumors (Chen et al, 2011; Sanders & Majumdar, 2011). This has led to the idea that conventional cancer therapies that target only proliferating cells in tumors may not necessarily be effective against cancer stem cells that mediate metastasis (Abdul Khalek et al, 2010, Sanders & Majumdar, 2011; Soltanian & Matin, 2011), and that these therapies may therefore be ineffective in producing long-term remissions. CSCs have greater DNA repair capacity and expression of ABC transporter genes, both of which contribute to relatively higher resistance to chemotherapy and radiation (Bao et al, 2006; Cho & Clarke, 2008; Hirschmann-Jax et al, 2004; Zhou et al, 2009). GO-Y030, a curcumin analogue has been shown to inhibit STAT3 phosphorylation signaling in colon cancer stem cells, offering the possibility of targeting STAT3 signaling in colon CSCs (Lin et al, 2011). The clonogenic and proliferative properties of CSCs are significantly interrupted by histone deacetylase (HDAC) inhibitors and this effect is associated with apoptotic cell death and modified Wnt signalling (Sikandar et al 2010).

8. Conclusion

1. When applied to colorectal cancer, the concept of hierarchical compartmentalization (as described in crypt kinetic models) offers target environments for stemness, proliferation

- and differentiation. Potential targets in each compartment include dividing cells, apoptotic mechanisms and cancer stem cells.
2. Wnt signalling has been targeted for inhibition because of its relationship with proliferation. Activity in this pathway is highest in the stem zone which provides the source of new cells.
 3. COX inhibitors have variable effects on proliferation that may be related to differing potencies, and the evidence suggests that these effects may not be due to any inhibitory action by the compounds on COX. Inconsistencies remain in trying to reproduce in patients the experimental outcomes on tumor loads seen following treatment with COX inhibitors.
 4. A range of compounds, including nutritional and synthetic substances, induce apoptosis in colorectal cancer cell lines. Not all COX inhibitors induce apoptosis.
 5. Some COX inhibitors down-regulate PPAR δ , other inhibitors do not. However, combination treatments do not produce the expected potentiation effect. The conflicting evidence of the roles of PPAR δ and PPAR γ in colorectal cancer remains unresolved.
 6. Stem cells markers are increasingly being identified and involvement in signalling pathways such as IL6-STAT3 point to new targets that may be modulated using therapeutic agents or genetic manipulations.

9. References

- Abdul Khalek, F. J.; Gallicano, G. I. & Mishra, L. (2010). Colon cancer stem cells. *Gastrointest Cancer Res* (Suppl 1), pp. S16-23
- Aliperti, L. A.; Predina J. D.; Vachani, A. & Singhal, S. (2011). Local and systemic recurrence is the Achilles heel of cancer surgery. *Ann Surg Oncol*, Vol.18, No.3 pp. 603-7.
- American Cancer Society. (2011). *Colorectal Cancer Facts & Figures 2011-2013*, American Cancer Society, Atlanta, United States
- Araki, Y.; Okamura, S.; Hussain, S. P.; Nagashima, M.; He, P.; Shiseki, M.; Miura, K. & Harris, C. C. (2003). Regulation of cyclooxygenase-2 expression by the Wnt and ras pathways. *Cancer Res*, Vol.63, pp. 728-34.
- Ban, J. O.; Kwak, D. H.; Oh, J. H.; Park, E. J.; Cho, M. C.; Song, H. S.; Song, M. J.; Han, S. B.; Moon, D. C.; Kang, K. W. & Hong, J. T. (2010). Suppression of NF-kappaB and GSK-3beta is involved in colon cancer cell growth inhibition by the PPAR agonist troglitazone. *Chem Biol Interact*, Vol.188, No.1, pp. 75-85
- Bao, S.; Wu, Q.; McLendon, R. E.; Hao, Y.; Shi, Q.; Hjelmeland, A. B.; Dewhirst, M. W.; Bigner, D. D. & Rich, J. N. (2006). Glioma stem cells promote radioresistance by preferential activation of the DNA damage response. *Nature*, Vol.444, No.7120, pp. 756-60.
- Barker, N.; Ridgway, R. A.; van Es, J. H.; van de Wetering, M.; Begthel, H.; van den Born, M.; Danenberg, E.; Clarke, A. R.; Sansom, O. J. & Clevers, H. (2009). Crypt stem cells as the cells-of-origin of intestinal cancer. *Nature*, Vol.457(7229), pp. 608-11
- Barker, N.; van Es, J. H.; Kuipers, J.; Kujala, P.; van den Born, M.; Cozijnsen, M.; Haegebarth, A.; Korving, J.; Begthel, H.; Peters, P. J. & Clevers, H. (2007). Identification of stem cells in small intestine and colon by marker gene *Lgr5*. *Nature*, Vol.449, pp. 1003-7
- Becker, L.; Huang, Q. & Mashimo, H. (2008). Immunostaining of *lgr5*, an intestinal stem cell marker, in normal and premalignant human gastrointestinal tissue. *Sci. World J*, Vol.8, pp. 1168-76.

- Bjerknes, M. & Cheng, H. (1981a). The stem-cell zone of the small intestinal epithelium. I. Evidence from Paneth cells in the adult mouse. *Am. J. Anat*, Vol.160, pp. 51-63
- Bjerknes, M. & Cheng, H. (1981b). The stem-cell zone of the small intestinal epithelium. III. Evidence from columnar, enteroendocrine, and mucous cells in the adult mouse. *Am. J. Anat*, Vol.160, pp. 77-91
- Bjerknes, M. & Cheng, H. (1999). Clonal analysis of mouse intestinal epithelial progenitors. *Gastroenterology*, Vol.116, pp. 7-14
- Bjerknes, M. & Cheng, H. (2006). Intestinal epithelial stem cells and progenitors. *Methods Enzymol*, Vol.419, pp. 337- 83
- Boman, B. M.; Fields, J. Z.; Bonham-Carter, O. & Runquist, O. A. (2001). Computer modeling implicates stem cell overproduction in colon cancer initiation. *Cancer Res*, Vol.61, pp. 8408-11
- Botting, R. M. (2006). Inhibitors of cyclooxygenases: mechanisms, selectivity and uses. *J Physiol Pharmacol*, Vol.57, Suppl 5, pp. 113-24.
- Cairnie, A. B. & Millen, B. H. (1975). Fission of crypts in the small intestine of the irradiated mouse. *Cell Tissue Kinet*, Vol.8, pp. 189-96
- Chen, S. Y.; Huang, Y. C.; Liu, S. P.; Tsai, F. J.; Shyu, W. C. & Lin, S. Z. (2011). An overview of concepts for cancer stem cells. *Cell Transplant*, Vol.20, No.1, pp. 113-20
- Cho, R. W. & Clarke, M. F. (2008). Recent advances in cancer stem cells. *Curr Opin Genet Dev*, Vol.18, No.1, pp. 48-53.
- Coghlan, M. P.; Culbert, A. A.; Cross, D. A.; Corcoran, S. L.; Yates, J. W.; Pearce, N. J.; Rausch, O. L; Murphy, G. J.; Carter, P. S.; Roxbee Cox, L.; Mills, D.; Brown, M. J.; Haigh, D.; Ward, R. W.; Smith, DG.; Murray, K. J.; Reith, A. D. & Holder, J. C. (2000). Selective small molecule inhibitors of glycogen synthase kinase-3 modulate glycogen metabolism and gene transcription. *Chem Biol*, Vol.7, pp. 793-803.
- Dalerba, P.; Dylla, S. J.; Park, I. K.; Liu, R.; Wang, X.; Cho, R. W.; Hoey, T.; Gurney, A.; Huang, E. H.; Simeone, D. M.; Shelton, A. A.; Parmiani, G.; Castelli, C. & Clarke, M. F. (2007). Phenotypic characterization of human colorectal cancer stem cells. *Proc. Natl. Acad. Sci. USA*, Vol.104, pp. 10158-63.
- Desvergne, B. & Wahli, W. (1999). Peroxisome proliferator-activated receptors: nuclear control of metabolism. *Endocr Rev*, Vol.20, pp. 649-88.
- Di Garbo, A.; Johnston, M. D.; Chapman, S. J. & Maini, P. K. (2010). Variable renewal rate and growth properties of cell populations in colon crypts. *Physical Review*, Vol.81, pp. 061909.1-12
- Eberhart, C. E.; Coffey, R. J.; Radhika, A.; Giardiello, F. M.; Ferrenbach, S. & DuBois, R. N. (1994). Up-regulation of cyclooxygenase-2 gene expression in human colorectal adenomas and adenocarcinomas. *Gastroenterology*, Vol.107, pp. 1183-88.
- Evan, G. I. & Vousden, K. H. (2001). Proliferation, cell cycle and apoptosis in cancer. *Nature*, Vol.411, pp. 342-48.
- Fagotto, F.; Gluck, U. & Gumbiner, B. M. (1998). Nuclear localization signal-independent and importin/karyopherin-independent nuclear import of β -catenin. *Curr Biol*, Vol.8, pp. 181-90.
- Fujita, T.; Matsui, M.; Takaku, K.; Uetake, H.; Ichikawa, W.; Taketo, M. M. & Sugihara, K. (1998). Size- and invasion-dependent increase in cyclooxygenase-2 levels in human colorectal carcinomas. *Cancer Res*, Vol.58, pp. 4823-26.

- Gao, X.; Deeb, D.; Liu, P.; Liu, Y.; Arbab-Ali, S.; Dulchavsky, S. A. & Gautam, S. C. (2011). Role of reactive oxygen species (ROS) in CDDO-Me-mediated growth inhibition and apoptosis in colorectal cancer cells. *J Exp Ther Oncol*, Vol.9, No.2, pp. 119-27.
- Giardiello, F. M.; Hamilton, S. R.; Krush, A. J.; Piantadosi, S.; Hyland, L. M.; Celano, P.; Booker, S. V.; Robinson, C. R. & Offerhaus, G. J. (1993). Treatment of colonic and rectal adenomas with sulindac in familial adenomatous polyposis. *N Engl J Med*, Vol.328, pp. 1313-16
- Giardiello, F. M.; Yang, V. W.; Hyland, L. M.; Krush, A. J.; Petersen, G. M.; Trimbath, J. D.; Piantadosi, S.; Garrett, E.; Geiman, D. E.; Hubbard, W.; Offerhaus, G. J. & Hamilton, S. R. (2002). Primary chemoprevention of familial adenomatous polyposis with sulindac. *N Engl J Med*, Vol.346, pp. 1054-59
- Giles, R. H.; van Es, J. H. & Clevers, H. (2003). Caught up in a Wnt storm: Wnt signaling in cancer. *Biochim Biophys Acta*, Vol.1653, pp. 1-24.
- Giovannucci, E.; Rimm, E. B.; Stampfer, M. J.; Colditz, G. A.; Ascherio, A. & Willett, W. C. (1994). Aspirin use and the risk for colorectal cancer and adenoma in male health professionals. *Ann Intern Med*, Vol.121, pp. 241-6
- Girnun, G. D.; Smith, W. M.; Drori, S.; Sarraf, P.; Mueller, E.; Eng, C.; Nambiar, P.; Rosenberg, D. W.; Bronson, R. T.; Edelmann, W.; Kucherlapati, R.; Gonzalez, F. J. & Spiegelman, B. M. (2002). APC-dependent suppression of colon carcinogenesis by PPARgamma. *Proc Natl Acad Sci USA*, Vol.99, pp. 13771-76.
- Gregorieff, A.; Pinto, D.; Begthel, H.; Destree, O.; Kileman, M. & Clevers, H. (2005). Expression patterns of Wnt signaling components in the adult intestine. *Gastroenterology*, Vol.129, pp. 626-38
- Guo, Q.; Wu, M.; Lian, P.; Liao, M.; Xiao, Z.; Wang, X. & Shen, S. (2009). Synergistic effect of indomethacin and NGX6 on proliferation and invasion by human colorectal cancer cells through modulation of the Wnt/beta-catenin signaling pathway. *Mol Cell Biochem*, Vol.330, pp. 71-81.
- Gupta, R. A.; Tan, J.; Krause, W. F.; Geraci, M. W.; Willson, T. M.; Dey, S. K. & DuBois, R. N. (2000). Prostacyclin-mediated activation of peroxisome proliferator-activated receptor delta in colorectal cancer. *Proc. Natl. Acad. Sci. USA*, Vol.97, pp. 13275-80.
- Gupta, R. A.; Wang, D.; Katkuri, S.; Wang, H.; Dey, S.K. & DuBois, R. N. (2004). Activation of nuclear hormone receptor peroxisome proliferator-activated receptor-delta accelerates intestinal adenoma growth. *Nat Med*, Vol.10, pp. 245-7.
- Hall, P. A.; Coates, P. J.; Ansari, B. & Hopwood, D. (1994). Regulation of cell number in the mammalian gastrointestinal tract: the importance of apoptosis. *J. Cell Sci*, Vol.107(Pt. 12), pp. 3569-77
- Han, A.; Song, Z.; Tong, C.; Hu, D.; Bi, X.; Augenlicht, L. H. & Yang, W. (2008). Sulindac suppresses beta-catenin expression in human cancer cells. *Eur J Pharmacol*, Vol.583, pp. 26-31.
- Hanif, R.; Pittas, A.; Feng, Y.; Koutsos, M. I.; Qiao, L.; Staiano-Coico, L.; Shiff, S. I. & Rigas, B. (1996). Effects of nonsteroidal anti-inflammatory drugs on proliferation and on induction of apoptosis in colon cancer cells by a prostaglandin-independent pathway. *Biochemical pharmacology*, Vol.52, pp. 237-45.
- Hardy, K. & Stark, J. (2002). Mathematical models of the balance between apoptosis and proliferation. *Apoptosis*, Vol.7, pp. 373-81.

- Harman FS, Nicol CJ, Marin HE, Ward JM, Gonzalez FJ, Peters JM. (2004). Peroxisome proliferator-activated receptor-delta attenuates colon carcinogenesis. *Nat. Med.*, Vol.10, pp. 481-83.
- He, T. C.; Chan, T. A.; Vogelstein, B. & Kinzler, K. W. (1999). PPARdelta is an APC-regulated target of nonsteroidal anti-inflammatory drugs. *Cell*, Vol.99, pp. 335-45.
- He, T. C.; Sparks, A. B.; Rago, C.; Hermeking, H.; Zawel, L.; da Costa, L. T.; Morin, P. J.; Vogelstein, B. & Kinzler, K. W. (1998). Identification of c-MYC as a target of the APC pathway. *Science*, Vol.281, pp. 1509-12.
- Heath, J. P. (1996). Epithelial cell migration in the intestine. *Cell Biol Int*, Vol.20, pp. 139-46.
- Hirschmann-Jax, C.; Foster, A. E.; Wulf, G. G.; Nuchtern, J. G.; Jax, T. W.; Gobel, U.; Goodell, M. A. & Brenner, M. K. (2004). A distinct "side population" of cells with high drug efflux capacity in human tumor cells. *Proc Natl Acad Sci USA*, Vol.101, No.39, pp. 14228-33
- Hollingshead, H. E.; Borland, M. G.; Billin, A. N.; Willson, T. M.; Gonzalez, F. J. & Peters, J. M. (2008). Ligand activation of peroxisome proliferator-activated receptor-beta/delta (PPARbeta/delta) and inhibition of cyclooxygenase 2 (COX2) attenuate colon carcinogenesis through independent signaling mechanisms. *Carcinogenesis*, Vol.29, pp. 169-76.
- Jacoby, R. F.; Marshall, D. J.; Newton, M. A.; Novakovic, K.; Tutsch, K.; Cole, C. E.; Lubet, R. A.; Kelloff, G. J.; Verma, A.; Moser, A. R. & Dove, W. F. (1996). Chemoprevention of spontaneous intestinal adenomas in the Apc Min mouse model by the nonsteroidal anti-inflammatory drug piroxicam *Cancer Res*, Vol.56, pp. 710-14
- Jacoby, R. F.; Seibert, K.; Cole, C. E.; Kelloff, G. & Lubet, R. A. (2000). The cyclooxygenase-2 inhibitor celecoxib is a potent preventive and therapeutic agent in the min mouse model of adenomatous polyposis *Cancer Res*, Vol.60, pp. 5040-44
- Jemal, A.; Murray, T.; Ward, E.; Samuels, A.; Tiwari, R. C.; Ghafoor, A.; Feuer, E. J. & Thun, M. J. (2005). Cancer statistics, 2005. *CA Cancer J Clin*, Vol.55, No.1, pp. 10-30.
- Johnstone, R. W.; Ruefli, A. A. & Lowe, S. W. (2002). Apoptosis: a link between cancer genetics and chemotherapy. *Cell*, Vol.108, pp. 153-64.
- Kikuchi, A. (2003). Tumor formation by genetic mutations in the components of the Wnt signaling pathway. *Cancer Sci*, Vol.94, pp. 225-29.
- Kim, Y. H.; Kim, M. H.; Kim, B. J.; Kim, J. J.; Chang, D. K.; Son, H. J.; Rhee, P. L. & Rhee, J. C. (2009). Inhibition of cell proliferation and invasion in a human colon cancer cell line by 5-aminosalicylic acid. *Dig Liver Dis*, Vol.41, pp. 328-37.
- Kishida, S.; Yamamoto, H.; Hino, S.; Ikeda, S.; Kishida, M. & Kikuchi, A. (1999). DIX domains of Dvl and axin are necessary for protein interactions and their ability to regulate beta-catenin stability. *Mol. Cell. Biol*, Vol.19, pp. 4414-22.
- Kohno, H.; Suzuki, R.; Sugie, S. & Tanaka, T. (2005). Suppression of colitis-related mouse colon carcinogenesis by a COX-2 inhibitor and PPAR ligands. *BMC Cancer*, Vol.5, pp. 46
- Lanas, A. & Ferrandez, A. (2009). NSAIDs and the colon. *Curr Opin Gastroenterol*, Vol.25, pp. 44-49
- Lander, A. D. (2009). The 'stem cell' concept: is it holding us back? *J Biol*, Vol.8, No.8, pp. 70.
- Leblond, C. P. (1964). Classification of cell populations on the basis of their proliferative behaviour. *J. Nat. Cancer Inst. Monograph*, Vol.14, pp. 119-48.
- Li, Y. Q.; Roberts, S. A.; Paulus, U.; Loeffler, M. & Potten, C. S. (1994). The crypt cycle in mouse small intestinal epithelium. *J Cell Sci*, Vol.107, pp. 3271-79.


- Lin, L.; Liu, Y.; Li, H.; Li, P. K.; Fuchs, J.; Shibata, H.; Iwabuchi, Y. & Lin, J. (2011). Targeting colon cancer stem cells using a new curcumin analogue, GO-Y030. *Br J Cancer*, Vol.105, No.2, pp. 212-20
- Liou, J. Y.; Ghelani, D.; Yeh, S. & Wu, K. K. (2007). Nonsteroidal anti-inflammatory drugs induce colorectal cancer cell apoptosis by suppressing 14-3-3epsilon. *Cancer Res*, Vol.67, pp. 3185-91.
- Loeffler, M.; Stein, R.; Wichmann, H-E.; Potten, C. S.; Kaur, P. & Chwalinski, S. (1986). *Cell Tissue Kinet*, Vol.19, pp. 627- 45.
- Mahmoud, N. N.; Boolbol, S. K.; Dannenberg, A. J.; Mestre, J. R.; Bilinski, R. T.; Martucci, C.; Newmark, H. L.; Chadburn, A. & Bertagnolli, M. M. (1998). The sulfide metabolite of sulindac prevents tumors and restores enterocyte apoptosis in a murine model of familial adenomatous polyposis. *Carcinogenesis*, Vol.19, pp. 87-91
- Marin, H. E.; Peraza, M. A.; Billin, A. N.; Willson, T. M.; Ward, J. M.; Kennett, M. J.; Gonzalez, F. J. & Peters, J. M. (2006). Ligand activation of peroxisome proliferator-activated receptor beta inhibits colon carcinogenesis. *Cancer Res*, Vol.66, pp. 4394-401.
- Marsh, M. N. & Trier, J. S. (1974a). Morphology and cell proliferation of subepithelial fibroblasts in adult mouse jejunum. I. Structural features. *Gastroenterol*, Vol.67, pp. 622-35.
- Marsh, M. N. & Trier, J. S. (1974b). Morphology and cell proliferation of subepithelial fibroblasts in adult mouse jejunum. II. Radioautographic studies. *Gastroenterology*, Vol.67, pp. 636-45.
- Marshman, E.; Booth, C. & Potten C. S. (2002). The intestinal epithelial stem cell. *Bioessays* Vol.24, pp. 91-8
- McGarvey, M. A.; Bass, G. & Ettarh, R. R. (2007a). Nimesulide alters cell recruitment into mitosis in murine intestinal crypts without influencing the cell production rate. *Dig. Dis. Sci*, Vol.52, pp. 1471-78.
- McGarvey, M. A.; O'Kelly, F. & Ettarh R. R. (2007b). Nimesulide inhibits crypt epithelial cell proliferation at 6 hours in the small intestine in CD-1 mice. *Dig. Dis. Sci*, Vol.52, pp. 2087-94.
- Michalik, L.; Desvergne, B. & Wahli, W. (2003). Peroxisome proliferator-activated receptors beta/delta: emerging roles for a previously neglected third family member. *Curr Opin Lipidol*, Vol.14, pp. 129-35.
- Michor, F.; Iwasa, Y.; Rajagopalan, H.; Lengauer, C. & Nowak, M. (2004). Linear model of colon cancer initiation. *Cell Cycle*, Vol.3, pp. 358-62.
- Moorghen, M.; Ince, P.; Finney, K. J.; Sunter, J. P.; Appleton, D. R. & Watson, A. J. (1988). A protective effect of sulindac against chemically-induced primary colonic tumours in mice. *J Pathol*, Vol.156, pp. 341-47
- Moorghen, M.; Orde, M.; Finney, K. J.; Appleton, D. R. & Watson, A. J. (1998). Sulindac enhances cell proliferation in DMH-treated mouse colonic mucosa. *Cell Prolif*, Vol.31, pp. 59-70
- Narisawa, T.; Satoh, M.; Sano, M. & Takahashi, T. (1983). Inhibition of initiation and promotion by N-methylnitrosourea-induced colon carcinogenesis in rats by non-steroid anti-inflammatory agent indomethacin. *Carcinogenesis*, Vol.4, pp. 1225-27
- Nicolas, P.; Kim, K. M.; Shibata, D. & Tavare, S. (2007). The stem cell population of the human colon crypt: analysis via methylation patterns. *PLoS Comput. Biol*, Vol.3, pp. e28

- O'Brien, C. A.; Pollett, A.; Gallinger, S. & Dick, J. E. (2007). A human colon cancer cell capable of initiating tumour growth in immunodeficient mice. *Nature*, Vol.445, pp. 106-10.
- Okamoto, R. & Watanabe, M. (2004). Molecular and clinical basis for the regeneration of human gastrointestinal epithelia. *J Gastroenterol*, Vol.39, pp. 1-6.
- Ouyang, N.; Williams, J. L. & Rigas, B. (2006). NO-donating aspirin isomers downregulate peroxisome proliferator-activated receptor (PPAR)delta expression in APC(min/+) mice proportionally to their tumor inhibitory effect: Implications for the role of PPARdelta in carcinogenesis. *Carcinogenesis*, Vol.27, pp. 232-39.
- Oz, H. S. & Ebersole, J. L. (2010). Green tea polyphenols mediated apoptosis in intestinal epithelial cells by a FADD-dependent pathway. *J Cancer Ther*, Vol.1, No.3, pp. 105-13.
- Park, B. H.; Vogelstein, B. & Kinzler, K. W. (2001). Genetic disruption of PPARdelta decreases the tumorigenicity of human colon cancer cells. *Proc. Natl. Acad. Sci. USA*, Vol.98, pp. 2598-603.
- Parker, F. G.; Barnes, E. N. & Kaye, G. I. (1974). The pericryptal fibroblast sheath. IV. Replication, migration and differentiation of the subepithelial fibroblasts of the crypt and villus of the rabbit jejunum. *Gastroenterology*, Vol.67, pp. 607-21.
- Pascal, R. R.; Kaye, G. I. & Lane, N. (1968a). Colonic pericryptal fibroblast sheath: replication, migration and cytodifferentiation of a mesenchymal cell system in adult tissue. I. Autoradiographic studies of normal rabbit colon. *Gastroenterology*, Vol.54, pp. 835-51.
- Pascal, R. R.; Kaye, G. I. & Lane, N. (1968b). Colonic pericryptal fibroblast sheath: replication, migration and cytodifferentiation of a mesenchymal cell system in adult tissue. II. Fine structural aspects of normal rabbit and human colon. *Gastroenterology*, Vol.54, pp. 852-65.
- Paulus, U.; Potten, C. S. & Loeffler, M. (1992). A model of the control of cellular regeneration in the intestinal crypt after perturbation based solely on local stem cell regulation. *Cell Prolif*, Vol.25, pp. 559-78
- Piazza, G. A.; Rahm, A. K.; Finn, T. S.; Fryer, B. H.; Li, H.; Stoumen, A. L.; Pamukcu, R. & Ahnen, D. J. (1997). Apoptosis primarily accounts for the growth-inhibitory properties of sulindac metabolites and involves a mechanism that is independent of cyclooxygenase inhibition, cell cycle arrest, and p53 induction. *Cancer Res*, Vol.57, pp. 2452-59.
- Pinto, D. & Clevers, H. (2005). Wnt control of stem cells and differentiation in the intestinal epithelium. *Exp. Cell Res*, Vol.306, pp. 357-63.
- Potten, C. S. (1977). Extreme sensitivity of some intestinal crypt cells to X and gamma irradiation. *Nature*, Vol.269, pp. 518-21
- Potten, C. S. & Loeffler, M. (1990). *Development*, Vol.110, pp. 1001-20.
- Potten, C. S.; Kovacs, L. & Hamilton, E. (1974). Continuous labelling studies on mouse skin and intestine. *Cell Tissue Kinet*, Vol.7, pp. 271-83
- Potten, C. S.; Owen, G. & Booth, D. (2002). Intestinal stem cells protect their genome by selective segregation of template DNA strands. *J. Cell Sci*, Vol.115, pp. 2381-88
- Rao, C. V.; Rivenson, A.; Simi, B.; Zang, E.; Kelloff, G.; Steele, V. & Reddy, B. S. (1995). Chemoprevention of colon carcinogenesis by sulindac a nonsteroidal anti-inflammatory agent. *Cancer Res*, Vol.55, pp. 1464-72

- Rao, C. V.; Steele, V. E.; Swamy, M. V.; Patlolla, J. M.; Guruswamy, S. & Kopelovich, L. (2009). Inhibition of azoxymethane-induced colorectal cancer by CP-31398 a TP53 modulator alone or in combination with low doses of celecoxib in male F344 rats. *Cancer Res*, Vol.69, pp. 8175–82
- Raz, A. (2002). Is inhibition of cyclooxygenase required for the anti-tumorigenic effects of nonsteroidal, anti-inflammatory drugs (NSAIDs)? In vitro versus in vivo results and the relevance for the prevention and treatment of cancer. *Biochem. Pharmacol*, Vol.63, pp. 343–47.
- Reddy, B. S.; Rao, C. V.; Rivenson, A. & Kelloff, G. (1993). Inhibitory effect of aspirin on azoxymethane- induced colon carcinogenesis in F344 rats. *Carcinogenesis*, Vol.14, pp. 1493–97
- Reed, K. R.; Sansom, O. J.; Hayes, A. J.; Gescher, A. J.; Winton, D. J.; Peters, J. M. & Clarke, A. R. (2004). PPARdelta status and Apc-mediated tumourigenesis in the mouse intestine. *Oncogene*, Vol.23, pp. 8992-96.
- Ricci-Vitiani, L.; Lombardi, D. G.; Pilozzi, E.; Biffoni, M.; Todaro, M.; Peschle, C. & De Maria, R. (2007). Identification and expansion of human colon-cancer-initiating cells. *Nature*, Vol.445, pp. 111–15.
- Rigas, B.; Goldman, I. S. & Levine, L. (1993). Altered eicosanoid levels in human colon cancer. *J. Lab. Clin. Med*, Vol.122, pp. 518–23
- Sakthianandeswaren, A.; Christie, M.; D'Andreti, C.; Tsui, C.; Jorissen, R. N.; Li, S.; Fleming, N. I.; Gibbs, P.; Lipton, L.; Malatterre, J.; Ramsay, R. G.; Pheese, T. J.; Ernst, M.; Jeffery, R. E.; Poulson, R.; Leedham, S. J.; Segditsas, S.; Tomlinson, I. P.; Bernhard, O. K.; Simpson, R. J.; Walker, F.; Faux, M. C.; Church, N.; Catimel, B.; Flanagan, D. J.; Vincan, E. & Sieber, O. M. (2011). PHLDA1 expression marks the putative epithelial stem cells and contributes to intestinal tumorigenesis. *Cancer Res*, Vol.71, No.10, pp. 3709-19
- Sanders, M. A. & Majumdar, A. P. (2011). Colon cancer stem cells: implications in carcinogenesis. *Front Biosci*, Vol.16, pp. 1651-62
- Sato, T.; van Es, J. H.; Snippert, H. J.; Stange, D. E.; Vries, R. G.; van den Born, M.; Barker, N.; Shroyer, N. F.; van de Wetering, M. & Clevers, H. (2011). Paneth cells constitute the niche for Lgr5 stem cells in intestinal crypts. *Nature*, Vol.469(7330), pp. 415-8.
- Sato, T.; Vries, R. G.; Snippert, H. J.; van de Wetering, M.; Barker, N.; Stange, D. E.; van Es, J. H.; Abo, A.; Kujala, P.; Peters, P. J. & Clevers, H. (2009). Single Lgr5 stem cells build crypt-villus structures in vitro without a mesenchymal niche. *Nature*, Vol.459(7244), pp. 262-65.
- Schmidt, G. H.; Winton, D. J. & Ponder, B. A. (1988). Development of the pattern of cell renewal in the crypt-villus unit of chimaeric mouse small intestine. *Development*, Vol.103, No.4, pp. 785-90.
- Sheng, H.; Shao, J.; Kirkland, S. C.; Isakson, P.; Coffey, R. J.; Morrow, J.; Beauchamp, R. D. & DuBois, R. N. (1997). Inhibition of human colon cancer cell growth by selective inhibition of cyclooxygenase-2. *J. Clin. Invest*, Vol.99, pp. 2254–59.
- Shiff, S. J.; Koutsos, M. I.; Qiao, L. & Rigas, B. (1996). Nonsteroidal antiinflammatory drugs inhibit the proliferation of colon adenocarcinoma cells: effects on cell cycle and apoptosis. *Exp Cell Res*, Vol.222, pp. 179-88.
- Shmelkov, S. V.; Butler, J. M.; Hooper, A. T.; Hormigo, A.; Kushner, J.; Milde, T.; St Clair, R.; Baljevic, M.; White, I.; Jin, D. K.; Chadburn, A.; Murphy, A. J.; Valenzuela, D. M.; Gale, N. W.; Thurston, G.; Yancopoulos, G. D.; D'Angelica, M.; Kemeny, N.; Lyden,

- D. & Rafii, S. (2008). Cd133 expression is not restricted to stem cells, and both cd133+ and cd133- metastatic colon cancer cells initiate tumors. *J. Clin. Invest.*, Vol.118, pp. 2111-20.
- Shtutman, M.; Zhurinsky, J.; Simcha, I.; Albanese, C.; D'Amico, M.; Pestell, R. & Ben-Ze'ev, A. (1999). The cyclin D1 gene is a target of the β -catenin/LEF-1 pathway. *Proc. Nat. Acad. Sci. USA*, Vol.96, pp. 5522-27.
- Shureiqi, I.; Chen, D.; Lotan, R.; Yang, P.; Newman, R. A.; Fischer, S. M. & Lippman, S. M. (2000). 15-Lipoxygenase-1 mediates nonsteroidal anti-inflammatory drug-induced apoptosis independently of cyclooxygenase-2 in colon cancer cells. *Cancer Res*, Vol.60, pp. 6846-50
- Shureiqi, I.; Jiang, W.; Zuo, X.; Wu, Y.; Stimmel, J. B.; Leesnitzer, L. M.; Morris, J. S.; Fan, H. Z.; Fischer, S. M. & Lippman, S. M. (2003). The 15-lipoxygenase-1 product 13-S-hydroxyoctadecadienoic acid down-regulates PPAR- δ to induce apoptosis in colorectal cancer cells. *Proc. Nat. Acad. Sci. USA*, Vol.100, pp. 9968-73.
- Sikandar, S.; Dizon, D.; Shen, X.; Li, Z.; Besterman, J. & Lipkin, S. M. (2010). The class I HDAC inhibitor MGCD0103 induces cell cycle arrest and apoptosis in colon cancer initiating cells by upregulating Dickkopf-1 and non-canonical Wnt signaling. *Oncotarget*, Vol.1, No.7, pp. 596-605.
- Snippert, H. J.; Haegbarth, A.; Kasper, M.; Jaks, V.; van Es, J. H.; Barker, N.; van de Wetering, M.; van den Born, M.; Begthel, H.; Vries, R. G.; Stange, D. E.; Toftgård, R. & Clevers, H. (2010). Lgr6 marks stem cells in the hair follicle that generate all cell lineages of the skin. *Science*, Vol.327(5971), pp. 1385-89.
- Soltanian, S. & Matin, M. M. (2011). Cancer stem cells and cancer therapy. *Tumour Biol*, Vol.32, No.3, pp. 425-40
- Swamy, M. V.; Patlolla, J. M.; Steele, V. E.; Kopelovich, L.; Reddy, B. S. & Rao, C. V. (2006). Chemoprevention of familial adenomatous polyposis by low doses of atorvastatin and celecoxib given individually and in combination to APCmin mice. *Cancer research*, Vol.66, pp. 7370-77.
- Tetsu, O. & McCormick, F. (1999). β -catenin regulates expression of cyclin D1 in colon carcinoma cells. *Nature*, Vol.398, pp. 422-26.
- Thun, M. J.; Namboodiri, M. M.; Heath, C.W. Jr. (1991). Aspirin use and reduced risk of fatal colon cancer. *N Engl J Med*, Vol.325, pp. 1593-96
- Tomlinson, I. P. & Bodmer, W. F. (1995). Failure of programmed cell death and differentiation as causes of tumors: some simple mathematical models. *Proc Natl Acad Sci USA*, Vol.92, No.24, pp. 11130-34.
- Tsai, K. S.; Yang, S. H.; Lei, Y. P.; Tsai, C. C.; Chen, H. W.; Hsu, C. Y.; Chen, L. L.; Wang, H. W.; Miller, S. A.; Chiou, S. H.; Hung, M. C. & Hung, S. C. (2011). Mesenchymal Stem Cells Promote Formation of Colorectal Tumors in Mice. *Gastroenterology*, doi:10.1053/j.gastro.2011.05.045
- Uchida, H.; Yamazaki, K.; Fukuma, M.; Yamada, T.; Hayashida, T.; Hasegawa, H.; Kitajima, M.; Kitagawa, Y. & Sakamoto, M. (2010). Overexpression of leucine-rich repeat-containing g protein-coupled receptor 5 in colorectal cancer. *Cancer Sci*, Vol.101, pp. 1731-37
- van der Flier, L. G. & Clevers, H. (2009). Stem Cells, Self-Renewal, and Differentiation in the Intestinal Epithelium. *Annu Rev Physiol*, 71:241-60

- van der Flier, L. G.; Haegbarth, A.; Stange, D. E.; van de Wetering, M. & Clevers, H. (2009). Olfm4 is a robust marker for stem cells in human intestine and marks a subset of colorectal cancer cells. *Gastroenterology*, 137:15-17
- van Leeuwen, I.; Byrne, H.; Jensen, O. & King, J. (2006). Crypt dynamics and colorectal cancer: advances in mathematical modelling. *Cell Prolif*, 39, 157-81.
- Wang, D. & DuBois, R. N. (2010). Therapeutic potential of peroxisome proliferator-activated receptors in chronic inflammation and colorectal cancer. *Gastroenterol Clin North Am*, Vol.39, No.3, pp. 697-707
- Wang, D.; Ning, W.; Xie, D.; Guo, L. & Dubois, R. N. (2011). Peroxisome proliferator-activated receptor δ confers resistance to peroxisome proliferator-activated receptor γ -induced apoptosis in colorectal cancer cells. *Oncogene*, doi: 10.1038/onc.2011.299
- Williams, C. S.; Watson, A. J.; Sheng, H.; Helou, R.; Shao, J. & DuBois, R. N. (2000). Celecoxib prevents tumor growth in vivo without toxicity to normal gut: lack of correlation between in vitro and in vivo models. *Cancer Res*, Vol.60, pp. 6045-51.
- Wodarz, D. (2007). Effect of stem cell turnover rates on protection against cancer and aging. *J. Theor. Biol*, Vol.245, pp. 449-58.
- Wright, N. A. & Alison, M. (1985). *The Biology of Epithelial Cell Populations*, Vol.2, ISBN 978-0-19-857615-0, Oxford University Press, United States
- Yamamoto, H.; Kishida, S.; Kishida, M.; Ikeda, S.; Takada, S. & Kikuchi, A. (1999). Phosphorylation of axin, a Wnt signal negative regulator, by glycogen synthase kinase-3 β regulates its stability. *J. Biol. Chem*, Vol.274, pp. 10681-84.
- Yanagawa, S.; van Leeuwen, F.; Wodarz, A.; Klingensmith, J. & Nusse, R. (1995). The dishevelled protein is modified by wingless signaling in Drosophila. *Gene Develop*, Vol.9, pp. 1087-97.
- Yatabe, Y.; Tavaré, S. & Shibata, D. (2001). Investigating stem cells in human colon by using methylation patterns. *Proc. Natl. Acad. Sci. USA*, Vol.98, pp. 10839-44
- Yoshinaga, M.; Taki, K.; Somada, S.; Sakiyama, Y.; Kubo, N.; Kaku, T.; Tsuruta, S.; Kusumoto, T.; Sakai, H.; Nakamura, K.; Takayanagi, R. & Muto, Y. (2011). The expression of both peroxisome proliferator-activated receptor delta and cyclooxygenase-2 in tissues is associated with poor prognosis in colorectal cancer patients. *Dig Dis Sci*, Vol.56, No.4, pp. 1194-200.
- Yoshizumi, T.; Ohta, T.; Ninomiya, I.; Terada, I.; Fushida, S.; Fujimura, T.; Nishimura, G.; Shimizu, K.; Yi, S. & Miwa, K. (2004). Thiazolidinedione, a peroxisome proliferator-activated receptor-gamma ligand, inhibits growth and metastasis of HT-29 human colon cancer cells through differentiation-promoting effects. *Int J Oncol*, Vol.25, pp. 631-39
- Yost, C.; Torres, M.; Miller, J. R.; Huang, E.; Kimelman, D. & Moon, R. T (1996). The axis-inducing activity, stability, and subcellular distribution of beta-catenin is regulated in Xenopus embryos by glycogen synthase kinase 3. *Gene. Develop*. Vol.10, pp. 1443-54.
- Zhang, J. S.; Li, D. M.; He, N.; Liu, Y. H.; Wang, C. H.; Jiang, S. Q.; Chen, B. Q. & Liu, J. R. (2011). A paraptosis-like cell death induced by δ -tocotrienol in human colon carcinoma SW620 cells is associated with the suppression of the Wnt signaling pathway. *Toxicology*, Vol.285, pp. 8-17.
- Zhou, B. B.; Zhang, H.; Damelin, M.; Geles, K. G.; Grindley, J. C. & Dirks, P. B. (2009). Tumour-initiating cells: challenges and opportunities for anticancer drug discovery. *Nat Rev Drug Discov*, Vol.8, No.10, pp. 806-23



Colorectal Cancer - From Prevention to Patient Care

Edited by Dr. Rajunor Ettarh

ISBN 978-953-51-0028-7

Hard cover, 538 pages

Publisher InTech

Published online 17, February, 2012

Published in print edition February, 2012

The projections for future growth in the number of new patients with colorectal cancer in most parts of the world remain unfavorable. When we consider the substantial morbidity and mortality that accompanies the disease, the acute need for improvements and better solutions in patient care becomes evident. This volume, organized in five sections, represents a synopsis of the significant efforts from scientists, clinicians and investigators towards finding improvements in different patient care aspects including nutrition, diagnostic approaches, treatment strategies with the addition of some novel therapeutic approaches, and prevention. For scientists involved in investigations that explore fundamental cellular events in colorectal cancer, this volume provides a framework for translational integration of cell biological and clinical information. Clinicians as well as other healthcare professionals involved in patient management for colorectal cancer will find this volume useful.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Rajunor Ettarh, Alvis Calamai and Anthony Cullen (2012). Therapeutic Targets in Colorectal Cancer, Colorectal Cancer - From Prevention to Patient Care, Dr. Rajunor Ettarh (Ed.), ISBN: 978-953-51-0028-7, InTech, Available from: <http://www.intechopen.com/books/colorectal-cancer-from-prevention-to-patient-care/therapeutic-targets-in-colorectal-cancer>

INTECH
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2012 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.