

The ATF/CREB Family of Transcription Factors in Breast Cancer

Jeremy K. Haakenson, Mark Kester and David X. Liu
*Pennsylvania State University College of Medicine,
USA*

1. Introduction

Transcription factors are proteins that bind DNA and either promote or block gene transcription. The activating transcription factor/cyclic AMP response element binding (ATF/CREB) family of transcription factors are involved in various cellular processes, including cell stress responses, cell survival, and cell growth. In this chapter, we will first give an overview of the transcriptional regulation of oncogenesis, followed by a brief summary of the roles of ATF/CREB family members in breast cancer. After this, we will describe the structure of ATF/CREB family members and then go into detail concerning each ATF/CREB family member that has a function relevant to breast cancer. Finally, we will end the chapter with some forward-looking remarks on ATF/CREB-based breast cancer therapeutics.

2. Transcriptional regulation in breast cancer

Transcription factors play an important role in breast cancer tumorigenesis and progression. For example, ETS1, ELF3, PDEF, PEA3, HIF-1, and MYC are all transcription factors that are overexpressed in breast cancer (Kurpios and others 2003). Overexpression of MYC in MCF7 human non-metastatic breast cancer cells causes those cells to display a metastatic-like phenotype (de Launoit and others 2000). The transcriptional targets of PEA3 include MMP1, MMP3, MMP9, vimentin, and ICAM-1, genes that are involved in breast cancer cell invasion and migration (de Launoit and others 2000). Other transcription factors involved in breast cancer include snail, which blocks E-cadherin transcription (Cano and others 2000); CBP/p300, a transcriptional co-activator involved in HER2 expression (Wang and others 2001); STAT6, which blocks the immune response to breast cancer (Sinha and others 2005); and Fra-1, whose overexpression leads to increased invasion in breast cancer (Mizutani and others 2004). Like PEA3, Runx2 is also involved in breast cancer metastasis, specifically, metastasis to bone (Shore 2005). Its targets include collagenase-3 and BSP, which are upregulated in metastatic breast cancer cells (Shore 2005). Since it is well known that tumors can become hypoxic, it is not surprising that HIF-1 is overexpressed in primary breast tumors (Kimbrow and Simons 2006). Its best known target is VEGF, an inducer of angiogenesis (Kimbrow and Simons 2006). MYC is amplified and overexpressed in breast cancer (Chen and Olopade 2008). Its amplification is correlated with tumor progression and poor prognosis (Chen and Olopade 2008). MYC binds MAX to form a heterodimer that

induces the transcription of genes such as MTA1, which is involved in transformation; PEG10, which plays a role in proliferation; hTERT; and VEGF (Chen and Olopade 2008). In a transgenic mouse model, overexpression of MYC in the mammary epithelium caused breast cancer to develop after eight months (Rose-Hellekant and Sandgren 2000). Similar to MYC, MYB is amplified in hereditary breast cancer (Kauraniemi and others 2000). MYB is involved in ER-dependent breast cancer cell proliferation (Ramsay and Gonda 2008). Its targets are anti-apoptotic genes, such as MYC, cyclin A1, and BCL2 (Ramsay and Gonda 2008).

Several transcription factors are constitutively active in breast cancer, including the aryl hydrocarbon receptor, STAT3, and NF- κ B (Pensa, Watson, Poli 2009; Schlezinger and others 2006; Vogel and Matsumura 2009). The aryl hydrocarbon receptor targets CYP1B1, a P450 enzyme (Schlezinger and others 2006). STAT3 is correlated with increased breast cancer cell survival, cytoskeletal reorganization, and migration (Pensa, Watson, Poli 2009). It activates HGF (Elliott and others 2002), cyclin D1, MYC, and bcl-XL (Silva and Shupnik 2007). In addition, it blocks the transcription of p21, which is an inhibitor of cell cycle progression (Silva and Shupnik 2007). Constitutively active NF- κ B, which is found in more than 90% of breast cancer (Vogel and Matsumura 2009), leads to breast cancer tumors that are hormone-independent (Wysocki and Wierusz-Wysocka 2010) and drug resistant (Garg and others 2003).

Many transcription factors act as tumor suppressors, including FoxP3, KLF10, p53, and GATA3. FoxP3, an X-linked breast cancer tumor suppressor, blocks HER2 expression (Medema and Burgering 2007). KLF10 is a transcription factor whose expression is inversely correlated with breast cancer stage (Subramaniam and others 2010). Its targets are BARD1 and Smad2, which are tumor suppressors (Subramaniam and others 2010). The tumor suppressor, p53, is mutated in about 26% of all breast cancers (Patocs and others 2007). It transactivates p21 and can also transcriptionally activate Bax1, leading to apoptosis (Ingvarsson 1999). GATA3, the most highly expressed transcription factor in the mammary epithelium (Kouros-Mehr and others 2008), is normally involved in luminal epithelial cell differentiation (Chou, Provot, Werb 2010). However, it is lost during cancer progression (Chou, Provot, Werb 2010). Indeed, decreased GATA3 expression is correlated with a worse prognosis, characterized by less differentiated, ER- cancer (Chou, Provot, Werb 2010).

The steroid hormone receptors, which are ligand-activated nuclear receptors that act as transcription factors, deserve special mention when it comes to breast cancer. The estrogen receptor (ER) and the progesterone receptor (PR) are overexpressed in many breast cancers. This has led to the development of anti-estrogens and aromatase inhibitors for breast cancer treatment. Besides leading to breast cancer tumorigenesis, ER also induces breast cancer progression (Carroll and Brown 2006), and it is not only overexpression, but mutations and alternative splicing that can cause breast cancer (Toran-Allerand 2004). ER targets include PR, c-jun (Fang, Chen, Weigel 2009), p52/TFF1, MYC, cyclin D1, and BCL2 (Welboren and others 2009). Like ER, PR also contributes to breast cancer. The progesterone receptor causes cells to enter S phase and mediates anchorage-independent growth in response to progesterin (Daniel, Knutson, Lange 2009). PR targets include MYC (Lange 2008), p21, EGFR, SGK, Tissue Factor, Muc-1, HB-EGF, and IRS-1 (Daniel, Knutson, Lange 2009). In addition to ER and PR in breast cancer, it is interesting to note that the androgen receptor (AR) protects against breast cancer. AR expression is correlated with a better prognosis (Yeap, Wilce, Leedman 2004), and mutations in the AR DNA-binding domain have been found in male breast cancer (MacLean, Warne, Zajac 1995).

3. ATF/CREB transcription factors in breast cancer

Like the transcription factors described in the previous section, the ATF/CREB family of transcription factors plays a role in breast cancer and may prove to be an effective therapeutic target. Some family members, such as ATF1, are protective against breast cancer, while the roles of others, such as ATF2 and ATF3, remain controversial. Still other family members, such as ATF4, ATF5, and CREB, promote breast cancer pathology (Table 1). These latter may be the most promising targets for intervention, since inhibiting their production or activation could block tumor growth and metastasis. The following sections will focus on the structure and function of individual ATF/CREB transcription factors.

4. Structure of ATF/CREB family members

The ATF/CREB family consists of a group of transcription factors that all contain an N-terminal DNA-binding domain and a C-terminal basic leucine zipper (B-ZIP) domain that binds other B-ZIP transcription factors to form homo- and heterodimers (Vinson and others 2002). ATF and CREB were both originally named in 1987 and were later found to bind to the same consensus sequence (Hai and Hartman 2001). The DNA-binding consensus sequence for this family is GTGACGT A/C A/G, which is found in the promoter region of target genes (Hai and Hartman 2001). Specificity is achieved by homo- and heterodimerization and epigenetic mechanisms, such as DNA methylation (Hai and Hartman 2001). For example, ATF1 forms heterodimers with CREM Ia and CREM Iib (Newman and Keating 2003). Likewise, ATF2, ATF4, and ATF5 all form heterodimers with C/EBP γ (Newman and Keating 2003). Based on the dimers that they form, ATF/CREB transcription factors are able to regulate the transcription of many target genes involved in breast cancer pathology and suppression.

5. ATF1

Several groups have identified ATF1 targets in both mice and humans. For example, Torti's group at Wake Forest found that ATF1 aids in the transcription of H ferritin in mouse fibroblasts (Tsuji, Torti, Torti 1998). Perhaps more relevant to breast cancer, ATF1 increases the transcription of H-2D^d, a major histocompatibility complex (MHC) class I gene (Ishiguro, Brown, Meruelo 1997). The transcription of such a gene could allow the immune system to tag breast cancer cells for destruction. Besides its effects on the immune system, ATF1 may also play a role in breast cancer via its regulation of steroidal hormone synthesis. Clem et al. (Clem, Hudson, Clark 2005) showed that ATF1 binds the steroidal acute regulatory protein (StAR) promoter, although they failed to determine whether this association activates or blocks transcription. Regardless of ATF1's effect, regulation of StAR, an enzyme necessary for estradiol synthesis, may play an important role in preventing breast cancer because a longer time of exposure to estradiol (menarche to menopause) is correlated with increased breast cancer risk. In addition to its actions on H-2D^d and StAR in mice, ATF1 blocks thrombospondin1 transcription in human thyroid cancer cells (Ghoneim and others 2007). Whether or not this effect translates to breast cancer cells has yet to be confirmed. Most relevant to breast cancer is that BRCA1 activates ATF1 (Houvras and others 2000). BRCA1 and BRCA2 act as tumor suppressor genes, and they often contain mutations in breast cancer. Houvras et al. (Houvras and others 2000) found that wild type (wt) BRCA1 bound ATF1, causing transcription of a luciferase reporter gene. Such a finding strongly suggests

ATF/CREB Family Member	Role in Breast Cancer	Targets
ATF1	Most likely suppresses tumor formation	H ferritin (Tsuiji, 1998) H-2D ^d (Ishiguro, 1997) StAR (Clem, 2005) Thrombospondin 1 (Ghoneim, 2007)
ATF2	Unclear	Collagen, type X, $\alpha 1$ (Tsuchimochi, 2010) MMP13 (Tsuchimochi, 2010) Cyclin A (van Dam, 2001) Aromatase (Deb, 2006) MMP2 (Song, 2006) COX2 (Subbaramaiah, 2002) Cyclin D1 (Lee, 1999) Maspin (Maekawa, 2008) IFN γ (Xue, 2005) FoxP3 (Liu, 2009)
ATF3	Unclear	Fibronectin (Yin, 2010) Snail (Yin, 2010) Twist (Yin, 2010) Slug (Yin, 2010) αv integrin (Yin, 2010) $\beta 6$ integrin (Yin, 2010) E-cadherin (Yin, 2010)
ATF4	Increased malignancy Angiogenesis	Osteocalcin (St-Arnaud, 2007) Type I collagen (Ameri, 2008) TRB3 (Ameri, 2008) E-selectin (Ameri, 2008) Asparagine synthetase (Ameri, 2008) RANKL (Ameri, 2008) VEGF (Ameri, 2008) GADD34 (Fels, 2006) CHOP (Fels, 2006) Gamma-synuclein (Hua, 2009) LAMP3 (Mujcic, 2009)
ATF5	Cell Survival	Phosphoenolpyruvate carboxykinase 2 (Pascual, 2008) Aldolase B (Pascual, 2008) ID1 (Gho, 2008) Egr-1 (Li, 2009; Liu, 2011) Mcl-1 (Sheng, 2010) Bcl-2 (Dluzen, 2011) Hsp27 (Wang, 2008) CYP2B6 (Pascual, 2008)
CREB	Malignancy	Aromatase (Brown, 2010) Bcl-2 (Dong, 1999)

Table 1. The roles of ATF/CREB family members in breast cancer.

that ATF1 protects against breast cancer. The fact that ATF1 associates with BRCA1, activates H-2D^d, and may block StAR synthesis indicates that it acts as a tumor suppressor in breast cancer. On the other hand, its blockage of thrombospondin would suggest that it plays a pro-tumorigenic role. However, the work with thrombospondin has not been verified *in vivo*, and such an effect may only occur in thyroid cancer cells. Unfortunately,

very few studies have been performed on the role of ATF1 in breast cancer. Important future studies will elucidate the effects of ATF1 on immunity and hormone synthesis in both normal and neoplastic breast cells, as well as its role in breast cancer *in vivo*.

6. ATF2

The exact role of ATF2 in breast cancer is unclear. Targets for ATF2 have been found in both chickens and mice. One such target, Col10a1 (collagen, type X, $\alpha 1$), mediates chondrocyte differentiation (Tsuchimochi and others 2010). ATF2 also increases the transcription of matrix metalloproteinase 13 (MMP13), which may help facilitate breast cancer metastasis (Tsuchimochi and others 2010). Matrix metalloproteinases degrade the basement membrane during the metastatic cascade. So the fact that ATF2 enhances the transcription of MMP13 indicates that it may increase the likelihood of tumor metastasis. In addition, Jun-ATF2 dimers have been shown to lead to the transcription of cyclin A, which increases cell proliferation (van Dam and Castellazzi 2001), providing further evidence for a possible oncogenic role for ATF2.

In primary human adipose fibroblasts obtained from breast cancer patients, co-culture with malignant epithelial cells increased the levels of phosphorylated ATF2 (pATF2) found at the I.3/II promoter of aromatase, the enzyme responsible for estrogen synthesis (Deb and others 2006). Co-culture with the malignant cells also increased binding of pATF2 to itself, C/EBP β , and CBP (Deb and others 2006). The fact that ATF2 aids in the transcription of a gene that increases estrogen levels makes it seem pro-tumorigenic. Furthermore, pATF2 has been shown to aid in the transcription of matrix metalloproteinase 2 (MMP2), which increases migration in H-Ras-transformed MCF10A human breast epithelial cells (Song and others 2006). Such activity indicates that ATF2 may play a role in breast cancer metastasis, if not oncogenesis. ATF2 also forms a complex with c-Jun and c-Fos that mediates HER2's induction of cyclooxygenase-2 (COX2), which itself may be carcinogenic (Subbaramaiah and others 2002). The fact that ATF2 is downstream of HER2 lends strong support to the notion that it is involved in some breast cancers. In addition, Lee et al. (Lee and others 1999) found that v-src causes ATF2 and CREB to bind the CRE/ATF site of the cyclin D1 gene, leading to transcription of cyclin D1 in MCF7 human breast cancer cells. Cyclin D1 inactivates the retinoblastoma tumor suppressor (RB), predisposing cells to malignancy (Hunter and Pines 1994; Sherr 1994). So ATF2's role here would again indicate that it acts as an oncogene.

In contrast to the above-mentioned studies, which describe ATF2 as contributing to a malignant phenotype, several groups have characterized it as protective against breast cancer. For instance, Maekawa et al. found that knockout of ATF2 increased cell number, decreased apoptosis, and increased the number of v-K-ras-induced colonies in mouse embryonic fibroblasts (MEF's) (Maekawa and others 2007). Plus, knockdown of ATF2 decreased levels of one of its targets, the breast cancer suppressor, maspin, in the mammary tumors of ATF2^{+/-} mice (Maekawa and others 2007; Maekawa and others 2008). Such findings would support the argument that ATF2 inhibits breast cancer formation. Along those lines, ATF2 has been shown to bind to the proximal element of interferon gamma (IFN γ) in MCF7 cells (Xue, Firestone, Bjeldanes 2005). Increased production and secretion of IFN γ by breast cancer cells could help the immune system destroy malignant cells before they spread, thereby inhibiting tumor growth and preventing metastasis. In a similar vein, decreased ATF2 levels have been shown to diminish the level of FoxP3, a breast cancer tumor suppressor (Liu and others 2009), lending further support to the idea that ATF2

inhibits breast cancer. However, studies in a wide range of human breast cancer cell lines are needed in order to definitively determine ATF2's role in breast cancer.

7. ATF3

Like ATF2, ATF3's role in breast cancer remains poorly delineated, although recent evidence characterizes it as an oncogene that may be involved in metastasis. Yin et al. (Yin and others 2010) found that ATF3 mediates the TGF β -induced increase in the expression of fibronectin, twist, snail, and slug in MCF10C1a1 human breast cancer cells. In addition, ATF3 overexpression led to an increase in α v and β 6 integrins, as well as increased migration in MCF10C1a1 cells (Yin and others 2010). Furthermore, ATF3 caused a decrease in E-cadherin expression, as well as other genetic and morphological changes characteristic of the epithelial to mesenchymal transition (EMT) (Yin and others 2010). All of these findings support the idea that ATF3 mediates EMT and metastasis in breast cancer. In addition, ATF3 also supports oncogenesis, as overexpression of ATF3 in MCF10C1a1 cells that were injected into nude mice increased the incidence of tumor formation (Yin and others 2010). Similarly, overexpression of ATF3 in the mammary gland myoepithelial cells of transgenic mice caused squamous metaplasia in nulliparous females and led to the formation of mammary tumors in animals that had given birth twice (Wang and others 2008). This is perhaps the most convincing evidence that ATF3 acts as an oncogene in breast cancer.

All of the above findings on ATF3 seem to establish it as an oncogene and metastasis promoter, yet some doubt remains. Einbond et al. (Einbond and others 2007) found that actin decreased cell proliferation in mouse embryonic fibroblasts (MEF's) in an ATF3-dependent manner. However, this finding may pertain only to MEF's and not to breast cancer cells. Other groups have found that the anti-neoplastic agents, doxorubicin and gemcitabine, increase ATF3 levels in human breast cancer cell lines (Hernandez-Vargas and others 2007; Mallory and others 2005). However, it remains unclear whether these drugs are acting through ATF3 or ATF3 is upregulated as part of an intracellular compensatory response. The best evidence that we have today suggests that ATF3 acts to promote tumorigenesis and metastasis. However, more studies must be conducted in order to confirm these results.

8. ATF4

Unlike ATF2 and ATF3, ATF4 is clearly pro-tumorigenic in breast cancer. Perhaps most telling, increased ATF4 levels have been found in human tumors compared to normal tissue (Ameri and Harris 2008), and several ATF4 targets indicate an oncogenic phenotype. Targets of ATF4 include osteocalcin (St-Arnaud and Elchaarani 2007), type I collagen, TRB3, E-selectin, and asparagine synthetase (Ameri and Harris 2008). More relevant to cancer, ATF4 acts as an activating transcription factor for RANKL, which activates Akt/PKB, an anti-apoptotic enzyme (Ameri and Harris 2008). Such activity in the breast would favor the formation of cancer. Besides RANKL expression, ATF4 also induces VEGF transcription (Ameri and Harris 2008), which can lead to angiogenesis at the sites of primary and secondary tumors, allowing hypoxic tumors to reach normoxia. In addition, ATF4 likely protects tumor cells from hypoxia by increasing the transcription of GADD34 and CHOP, which protect cells from hypoxic damage (Fels and Koumenis 2006). The center of a tumor often becomes hypoxic, and ATF4 may play a role in preventing hypoxia-induced cancer

cell death. Although ATF4 benefits the organism during development, bone formation, and stress, it can also act in a sinister fashion by aiding in tumor formation.

Indeed, several studies have shown a link between ATF4 levels and a more malignant genotype in human breast cancer cells. For instance, decreased ATF4 levels have been shown to cause a decrease in gamma-synuclein levels in T47D breast cancer cells (Hua and others 2009). Such a correlation with gamma-synuclein, which is up-regulated in advanced breast cancer (Bruening and others 2000), may indicate that ATF4 mediates breast tumor progression. Furthermore, hypoxia has been shown to increase LAMP3 levels in MCF7 cells via the action of ATF4 (Mujcic and others 2009). This further implicates ATF4 in tumor progression and metastasis, as LAMP3 has been shown to play a role in metastasis (Mujcic and others 2009). Interestingly, this finding suggests that hypoxic tumors may be more likely to metastasize than those with an adequate blood supply. Finally, ATF4 has been shown to mediate the osteopontin-induced increase in VEGF expression in MDA-MB-231 cells (Chakraborty, Jain, Kundu 2008). Since VEGF secretion leads to the recruitment of endothelial cells as part of the process of angiogenesis, this would suggest that ATF4 is able to help primary and secondary breast tumors survive, at least once osteopontin has activated it. Clearly, overexpression or constitutive activation of ATF4 causes breast cancer cells to become more malignant.

9. ATF5

Like ATF4, ATF5 contributes to a malignant phenotype. Targets of ATF5 include aldolase B (Pascual and others 2008), ID1 (Gho and others 2008), Egr-1 (Li and others 2009; Liu and others 2011), Mcl-1 (Sheng and others 2010), Bcl-2 (Dluzen and others 2011), and phosphoenolpyruvate carboxykinase 2 (PEPCK) (Pascual and others 2008). Up-regulation of PEPCK, a glycolytic enzyme, could indicate that ATF5 plays a role in the Warburg effect, in which cancer cells use aerobic glycolysis to generate ATP rather than oxidative phosphorylation. It is thought that this use of a less efficient means of producing ATP allows proliferating cells to more efficiently produce other needed metabolites, such as acetyl-CoA and NADPH (Vander Heiden, Cantley, Thompson 2009). In addition to increasing PEPCK production, ATF5 activates the transcription of heat shock protein 27 (Hsp27), which blocks apoptosis (Wang, Lin, Zhang 2007). Such an anti-apoptotic mechanism may lead to breast cancer oncogenesis. Furthermore, ATF5 has been shown to increase the level of CYP2B6 in human hepatoma cells (Pascual and others 2008). This P450 enzyme metabolizes cyclophosphamide, which is used to treat breast cancer. Thus, by inducing an enzyme that degrades cyclophosphamide, ATF5 may contribute not only to oncogenesis, but also to drug resistance. Based on the target genes that it up-regulates, such as Hsp27 and CYP2B6, ATF5 should be pro-tumorigenic in breast cancer.

Indeed, our recent studies indicate that ATF5 induces transcription of the pro-mitogenic early growth response factor (Egr-1) gene in MCF7 human breast cancer cells (Li and others 2009; Liu and others 2011). In addition, ATF5 also binds the Bcl-2 P2 promoter and transactivates Bcl-2, an anti-apoptotic gene, leading to breast cancer cell survival (Dluzen and others 2011). In the future, we may see ATF5 inhibitors used to treat breast cancer, as inhibition of ATF5 leads to cell death in breast cancer cells but not human breast epithelial cells (HBEC's) (Dluzen and others 2011). To this end, we have recently found that Hsp70 interacts with the N-terminal of ATF5 and protects ATF5 from both caspase- and

proteasome-dependent protein degradation (Li and others, 2011). The next step will involve using *in vivo* studies to determine the efficacy and toxicity of ATF5 inhibitors as treatments for breast cancer.

10. CREB

Similar to ATF5, CREB can contribute to malignancy of the breast. Importantly, CREB induces the transcription of aromatase in breast adipose mesenchymal cells (Brown and Simpson 2010). Increased levels of aromatase will lead to increased estrogen levels, which have been implicated in breast cancer. In fact, aromatase inhibitors, such as exemestane and anastrozole, are currently used to treat breast cancer (Goodman and others 2006). As an example of positive feedback regulation, estrogen causes CREB to bind and activate the cyclin D1 promoter (Castro-Rivera, Samudio, Safe 2001). By activating cyclin D1, which causes cells to progress through the cell cycle, activation of CREB may further contribute to carcinogenesis. In addition, dominant negative CREB has been shown to block the transcription of bcl-2 in MCF7 cells (Dong and others 1999). Since bcl-2 blocks apoptosis, this implicates CREB as being proto-oncogenic. Although good *in vitro* studies have been done on the role of CREB in breast cancer, *in vivo* studies are needed to confirm its status as a transcription factor that supports tumor growth.

11. Conclusions

A host of transcription factors regulate breast cancer, acting as both oncogenes and tumor suppressors. The ATF/CREB family is an example of a group of transcription factors involved in breast cancer development, with some family members leading to breast cancer pathogenesis and some blocking it. Therapeutic approaches that target members that are oncogenic may lead to the next generation of breast cancer therapies. A drug that inhibits those three genes without affecting other members of the ATF/CREB family could be especially powerful because it could block many downstream targets at once. An appropriate drug discovery screen would use luciferase assays of ATF 1-5 and CREB. Compounds that inhibit ATF4, ATF5, and CREB while not affecting ATF1, ATF2, and ATF3 would be considered hits. The screen could initially compare a highly metastatic breast cancer cell line, such as 410.4 cells to a normal mammary epithelial cell line, such as HC11 cells. The screen would then be confirmed in both ER⁺ (MCF7) and ER⁻ (MDA-MB-231) cells. Such a screen could lead to efficacious new breast cancer therapeutics with relatively little toxicity.

12. Acknowledgements

This work was supported in part by NIH grant R01HL076789 and Tobacco Settlement Fund, State of Pennsylvania to M.K. and American Cancer Society Research Scholar Award RSG-08-288-01-GMC and Department of Defense grant BC085617 to D.X.L.

13. Abbreviations

Acetyl-CoA, acetyl coenzyme A; ATF/CREB, activating transcription factor/cyclic AMP response element binding; BARD1, BRCA1 associated RING domain 1; Bcl-2, B-cell

CLL/lymphoma 2; BRCA1, breast cancer 1, early onset; BRCA2, breast cancer 2, early onset; BSP, integrin-binding sialoprotein; C/EBP β , CCAAT/enhancer binding protein, beta; C/EBP γ , CCAAT/enhancer binding protein, gamma; CHOP, DNA-damage-inducible transcript 3; CREM, cyclic AMP responsive element modulator; CYP1B1, cytochrome P450, family 1, subfamily B, polypeptide 1; CYP2B6, cytochrome P450, family 2, subfamily B, polypeptide 6; EGFR, epidermal growth factor receptor; Egr-1, early growth response 1; ELF3, E74-like factor 3 (ets domain transcription factor, epithelial specific); ER, estrogen receptor; ETS1, v-ets erythroblastosis virus E26 oncogene homolog 1 (avian); GADD34, protein phosphatase 1, regulatory (inhibitor) subunit 15A; GATA3, GATA binding protein 3; HB-EGF, heparin-binding EGF-like growth factor; HER2, v-erb-b2 erythroblastic leukemia viral oncogene homolog 2, neuro/glioblastoma derived oncogene homolog (avian); HIF-1, hypoxia inducible factor 1; Hsp27, heat shock protein 27; Hsp70, heat shock protein 70; hTERT, human telomerase reverse transcriptase; ICAM-1, intercellular adhesion molecule 1; ID1, inhibitor of DNA binding 1, dominant negative helix-loop-helix protein; IRS-1, insulin receptor substrate 1; KLF10, Kruppel-like factor 10; LAMP3, lysosomal-associated membrane protein 3; MAX, MYC associated factor X; Mcl-1, myeloid cell leukemia sequence 1 (BCL2-related); MMP1, matrix metalloproteinase 1 (interstitial collagenase); MMP2, matrix metalloproteinase 2 (gelatinase A, 72 kDa gelatinase, 72 kDa type IV collagenase); MMP3, matrix metalloproteinase 3 (stromelysin 1, progelatinase); MMP9, matrix metalloproteinase 9 (gelatinase B, 92kDa gelatinase, 92kDa type IV collagenase); MTA1, metastasis associated 1; MYB, v-myb myeloblastosis viral oncogene homolog (avian); MYC, v-myc myelocytomatosis viral oncogene homolog (avian); NADPH, nicotinamide adenine dinucleotide phosphate; NF- κ B, nuclear factor of kappa light polypeptide gene enhancer in B cells; PDEF, SAM pointed domain containing ets transcription factor; PEA3, ets variant 4; PEG10, paternally expressed 10; RANKL, tumor necrosis factor (ligand) superfamily, member 11; SGK, serum/glucocorticoid regulated kinase; STAT3, signal transducer and activator of transcription 3 (acute-phase response factor); STAT6, signal transducer and activator of transcription 6, interleukin 4-induced; TGF β , transforming growth factor, beta; TRB3, tribbles homolog 3 (*Drosophila*); VEGF, vascular endothelial growth factor.

14. References

- Ameri K and Harris AL. 2008. Activating transcription factor 4. *Int J Biochem Cell Biol* 40(1):14-21.
- Brown KA and Simpson ER. 2010. Obesity and breast cancer: Progress to understanding the relationship. *Cancer Res* 70(1):4-7.
- Bruening W, Giasson BI, Klein-Szanto AJ, Lee VM, Trojanowski JQ, Godwin AK. 2000. Synucleins are expressed in the majority of breast and ovarian carcinomas and in preneoplastic lesions of the ovary. *Cancer* 88(9):2154-63.
- Cano A, Perez-Moreno MA, Rodrigo I, Locascio A, Blanco MJ, del Barrio MG, Portillo F, Nieto MA. 2000. The transcription factor snail controls epithelial-mesenchymal transitions by repressing E-cadherin expression. *Nat Cell Biol* 2(2):76-83.
- Carroll JS and Brown M. 2006. Estrogen receptor target gene: An evolving concept. *Mol Endocrinol* 20(8):1707-14.
- Castro-Rivera E, Samudio I, Safe S. 2001. Estrogen regulation of cyclin D1 gene expression in ZR-75 breast cancer cells involves multiple enhancer elements. *J Biol Chem* 276(33):30853-61.


- Chakraborty G, Jain S, Kundu GC. 2008. Osteopontin promotes vascular endothelial growth factor-dependent breast tumor growth and angiogenesis via autocrine and paracrine mechanisms. *Cancer Res* 68(1):152-61.
- Chen Y and Olopade OI. 2008. MYC in breast tumor progression. *Expert Rev Anticancer Ther* 8(10):1689-98.
- Chou J, Provot S, Werb Z. 2010. GATA3 in development and cancer differentiation: Cells GATA have it! *J Cell Physiol* 222(1):42-9.
- Clem BF, Hudson EA, Clark BJ. 2005. Cyclic adenosine 3',5'-monophosphate (cAMP) enhances cAMP-responsive element binding (CREB) protein phosphorylation and phospho-CREB interaction with the mouse steroidogenic acute regulatory protein gene promoter. *Endocrinology* 146(3):1348-56.
- Daniel AR, Knutson TP, Lange CA. 2009. Signaling inputs to progesterone receptor gene regulation and promoter selectivity. *Mol Cell Endocrinol* 308(1-2):47-52.
- de Launoit Y, Chotteau-Lelievre A, Beaudoin C, Coutte L, Netzer S, Brenner C, Huvent I, Baert JL. 2000. The PEA3 group of ETS-related transcription factors. role in breast cancer metastasis. *Adv Exp Med Biol* 480:107-16.
- Deb S, Zhou J, Amin SA, Imir AG, Yilmaz MB, Lin Z, Bulun SE. 2006. A novel role of sodium butyrate in the regulation of cancer-associated aromatase promoters I.3 and II by disrupting a transcriptional complex in breast adipose fibroblasts. *J Biol Chem* 281(5):2585-97.
- Dluzen D, Li G, Tancelosky D, Moreau M, Liu DX. 2011. BCL-2 is a downstream target of ATF5 that mediates the prosurvival function of ATF5 in a cell type-dependent manner. *J Biol Chem* 286(9):7705-13.
- Dong L, Wang W, Wang F, Stoner M, Reed JC, Harigai M, Samudio I, Kladden MP, Vyhlidal C, Safe S. 1999. Mechanisms of transcriptional activation of bcl-2 gene expression by 17beta-estradiol in breast cancer cells. *J Biol Chem* 274(45):32099-107.
- Einbond LS, Su T, Wu HA, Friedman R, Wang X, Ramirez A, Kronenberg F, Weinstein IB. 2007. The growth inhibitory effect of actein on human breast cancer cells is associated with activation of stress response pathways. *Int J Cancer* 121(9):2073-83.
- Elliott BE, Hung WL, Boag AH, Tuck AB. 2002. The role of hepatocyte growth factor (scatter factor) in epithelial-mesenchymal transition and breast cancer. *Can J Physiol Pharmacol* 80(2):91-102.
- Fang SH, Chen Y, Weigel RJ. 2009. GATA-3 as a marker of hormone response in breast cancer. *J Surg Res* 157(2):290-5.
- Fels DR and Koumenis C. 2006. The PERK/eIF2alpha/ATF4 module of the UPR in hypoxia resistance and tumor growth. *Cancer Biol Ther* 5(7):723-8.
- Garg AK, Hortobagyi GN, Aggarwal BB, Sahin AA, Buchholz TA. 2003. Nuclear factor-kappa B as a predictor of treatment response in breast cancer. *Curr Opin Oncol* 15(6):405-11.
- Gho JW, Ip WK, Chan KY, Law PT, Lai PB, Wong N. 2008. Re-expression of transcription factor ATF5 in hepatocellular carcinoma induces G2-M arrest. *Cancer Res* 68(16):6743-51.
- Ghoneim C, Soula-Rothhut M, Blanchevoys C, Martiny L, Antonicelli F, Rothhut B. 2007. Activating transcription factor-1-mediated hepatocyte growth factor-induced down-regulation of thrombospondin-1 expression leads to thyroid cancer cell invasion. *J Biol Chem* 282(21):15490-7.

- Goodman LS, Gilman A, Brunton LL, Lazo JS, Parker KL. 2006. Goodman & Gilman's the pharmacological basis of therapeutics. 11th ed. New York: McGraw-Hill.
- Hai T and Hartman MG. 2001. The molecular biology and nomenclature of the activating transcription factor/cAMP responsive element binding family of transcription factors: Activating transcription factor proteins and homeostasis. *Gene* 273(1):1-11.
- Hernandez-Vargas H, Rodriguez-Pinilla SM, Julian-Tendero M, Sanchez-Rovira P, Cuevas C, Anton A, Rios MJ, Palacios J, Moreno-Bueno G. 2007. Gene expression profiling of breast cancer cells in response to gemcitabine: NF-kappaB pathway activation as a potential mechanism of resistance. *Breast Cancer Res Treat* 102(2):157-72.
- Houvras Y, Benezra M, Zhang H, Manfredi JJ, Weber BL, Licht JD. 2000. BRCA1 physically and functionally interacts with ATF1. *J Biol Chem* 275(46):36230-7.
- Hua H, Xu L, Wang J, Jing J, Luo T, Jiang Y. 2009. Up-regulation of gamma-synuclein contributes to cancer cell survival under endoplasmic reticulum stress. *J Pathol* 217(4):507-15.
- Hunter T and Pines J. 1994. Cyclins and cancer. II: Cyclin D and CDK inhibitors come of age. *Cell* 79(4):573-82.
- Ingvarsson S. 1999. Molecular genetics of breast cancer progression. *Semin Cancer Biol* 9(4):277-88.
- Ishiguro N, Brown GD, Meruelo D. 1997. Activation transcription factor 1 involvement in the regulation of murine H-2Dd expression. *J Biol Chem* 272(25):15993-6001.
- Kauraniemi P, Hedenfalk I, Persson K, Duggan DJ, Tanner M, Johannsson O, Olsson H, Trent JM, Isola J, Borg A. 2000. MYB oncogene amplification in hereditary BRCA1 breast cancer. *Cancer Res* 60(19):5323-8.
- Kimbro KS and Simons JW. 2006. Hypoxia-inducible factor-1 in human breast and prostate cancer. *Endocr Relat Cancer* 13(3):739-49.
- Kouros-Mehr H, Kim JW, Bechis SK, Werb Z. 2008. GATA-3 and the regulation of the mammary luminal cell fate. *Curr Opin Cell Biol* 20(2):164-70.
- Kurpios NA, Sabolic NA, Shepherd TG, Fidalgo GM, Hassell JA. 2003. Function of PEA3 ets transcription factors in mammary gland development and oncogenesis. *J Mammary Gland Biol Neoplasia* 8(2):177-90.
- Lange CA. 2008. Integration of progesterone receptor action with rapid signaling events in breast cancer models. *J Steroid Biochem Mol Biol* 108(3-5):203-12.
- Lee RJ, Albanese C, Stenger RJ, Watanabe G, Inghirami G, Haines GK, 3rd, Webster M, Muller WJ, Brugge JS, Davis RJ, et al. 1999. pp60(v-src) induction of cyclin D1 requires collaborative interactions between the extracellular signal-regulated kinase, p38, and jun kinase pathways. A role for cAMP response element-binding protein and activating transcription factor-2 in pp60(v-src) signaling in breast cancer cells. *J Biol Chem* 274(11):7341-50.
- Li G, Li W, Angelastro JM, Greene LA, Liu DX. 2009. Identification of a novel DNA binding site and a transcriptional target for activating transcription factor 5 in c6 glioma and mcf-7 breast cancer cells. *Mol Cancer Res* 7(6):933-43.
- Li G, Xu Y, Guan D, Liu Z, Liu DX. 2011. HSP70 protein promotes survival of C6 and U87 glioma cells by inhibition of ATF5 degradation. *J Biol Chem* 286(23):20251-9.
- Liu DX, Qian D, Wang B, Yang JM, Lu Z. 2011. p300-dependent ATF5 acetylation is essential for Egr-1 gene activation and cell proliferation and survival. *Mol Cell Biol* 31(18):3906-16.

- Liu Y, Wang Y, Li W, Zheng P, Liu Y. 2009. Activating transcription factor 2 and c-jun-mediated induction of FoxP3 for experimental therapy of mammary tumor in the mouse. *Cancer Res* 69(14):5954-60.
- MacLean HE, Warne GL, Zajac JD. 1995. Defects of androgen receptor function: From sex reversal to motor neurone disease. *Mol Cell Endocrinol* 112(2):133-41.
- Maekawa T, Sano Y, Shinagawa T, Rahman Z, Sakuma T, Nomura S, Licht JD, Ishii S. 2008. ATF-2 controls transcription of maspin and GADD45 alpha genes independently from p53 to suppress mammary tumors. *Oncogene* 27(8):1045-54.
- Maekawa T, Shinagawa T, Sano Y, Sakuma T, Nomura S, Nagasaki K, Miki Y, Saito-Ohara F, Inazawa J, Kohno T, et al. 2007. Reduced levels of ATF-2 predispose mice to mammary tumors. *Mol Cell Biol* 27(5):1730-44.
- Mallory JC, Crudden G, Oliva A, Saunders C, Stromberg A, Craven RJ. 2005. A novel group of genes regulates susceptibility to antineoplastic drugs in highly tumorigenic breast cancer cells. *Mol Pharmacol* 68(6):1747-56.
- Medema RH and Burgering BM. 2007. The X factor: Skewing X inactivation towards cancer. *Cell* 129(7):1253-4.
- Mizutani N, Luo Y, Mizutani M, Reisfeld RA, Xiang R. 2004. DNA vaccines suppress angiogenesis and protect against growth of breast cancer metastases. *Breast Dis* 20:81-91.
- Mujcic H, Rzymiski T, Rouschop KM, Koritzinsky M, Milani M, Harris AL, Wouters BG. 2009. Hypoxic activation of the unfolded protein response (UPR) induces expression of the metastasis-associated gene LAMP3. *Radiother Oncol* 92(3):450-9.
- Newman JR and Keating AE. 2003. Comprehensive identification of human bZIP interactions with coiled-coil arrays. *Science* 300(5628):2097-101.
- Pascual M, Gomez-Lechon MJ, Castell JV, Jover R. 2008. ATF5 is a highly abundant liver-enriched transcription factor that cooperates with constitutive androstane receptor in the transactivation of CYP2B6: Implications in hepatic stress responses. *Drug Metab Dispos* 36(6):1063-72.
- Patocs A, Zhang L, Xu Y, Weber F, Caldes T, Mutter GL, Platzer P, Eng C. 2007. Breast-cancer stromal cells with TP53 mutations and nodal metastases. *N Engl J Med* 357(25):2543-51.
- Pensa S, Watson CJ, Poli V. 2009. Stat3 and the inflammation/acute phase response in involution and breast cancer. *J Mammary Gland Biol Neoplasia* 14(2):121-9.
- Ramsay RG and Gonda TJ. 2008. MYB function in normal and cancer cells. *Nat Rev Cancer* 8(7):523-34.
- Rose-Hellekant TA and Sandgren EP. 2000. Transforming growth factor alpha- and c-myc-induced mammary carcinogenesis in transgenic mice. *Oncogene* 19(8):1092-6.
- Schlezinger JJ, Liu D, Farago M, Seldin DC, Belguise K, Sonenshein GE, Sherr DH. 2006. A role for the aryl hydrocarbon receptor in mammary gland tumorigenesis. *Biol Chem* 387(9):1175-87.
- Sheng Z, Li L, Zhu LJ, Smith TW, Demers A, Ross AH, Moser RP, Green MR. 2010. A genome-wide RNA interference screen reveals an essential CREB3L2-ATF5-MCL1 survival pathway in malignant glioma with therapeutic implications. *Nat Med* 16(6):671-7.
- Sherr CJ. 1994. The ins and outs of RB: Coupling gene expression to the cell cycle clock. *Trends Cell Biol* 4(1):15-8.

- Shore P. 2005. A role for Runx2 in normal mammary gland and breast cancer bone metastasis. *J Cell Biochem* 96(3):484-9.
- Silva CM and Shupnik MA. 2007. Integration of steroid and growth factor pathways in breast cancer: Focus on signal transducers and activators of transcription and their potential role in resistance. *Mol Endocrinol* 21(7):1499-512.
- Sinha P, Clements VK, Miller S, Ostrand-Rosenberg S. 2005. Tumor immunity: A balancing act between T cell activation, macrophage activation and tumor-induced immune suppression. *Cancer Immunol Immunother* 54(11):1137-42.
- Song H, Ki SH, Kim SG, Moon A. 2006. Activating transcription factor 2 mediates matrix metalloproteinase-2 transcriptional activation induced by p38 in breast epithelial cells. *Cancer Res* 66(21):10487-96.
- St-Arnaud R and Elchaarani B. 2007. Identification of additional dimerization partners of FIAT, the factor inhibiting ATF4-mediated transcription. *Ann N Y Acad Sci* 1116:208-15.
- Subbaramaiah K, Norton L, Gerald W, Dannenberg AJ. 2002. Cyclooxygenase-2 is overexpressed in HER-2/neu-positive breast cancer: Evidence for involvement of AP-1 and PEA3. *J Biol Chem* 277(21):18649-57.
- Subramaniam M, Hawse JR, Rajamannan NM, Ingle JN, Spelsberg TC. 2010. Functional role of KLF10 in multiple disease processes. *Biofactors* 36(1):8-18.
- Toran-Allerand CD. 2004. Minireview: A plethora of estrogen receptors in the brain: Where will it end? *Endocrinology* 145(3):1069-74.
- Tsuchimochi K, Otero M, Dragomir CL, Plumb DA, Zerbini LF, Libermann TA, Marcu KB, Komiya S, Ijiri K, Goldring MB. 2010. GADD45beta enhances Col10a1 transcription via the MTK1/MKK3/6/p38 axis and activation of C/EBPbeta-TAD4 in terminally differentiating chondrocytes. *J Biol Chem* 285(11):8395-407.
- Tsuji Y, Torti SV, Torti FM. 1998. Activation of the ferritin H enhancer, FER-1, by the cooperative action of members of the AP1 and Sp1 transcription factor families. *J Biol Chem* 273(5):2984-92.
- van Dam H and Castellazzi M. 2001. Distinct roles of jun : Fos and jun : ATF dimers in oncogenesis. *Oncogene* 20(19):2453-64.
- Vander Heiden MG, Cantley LC, Thompson CB. 2009. Understanding the warburg effect: The metabolic requirements of cell proliferation. *Science* 324(5930):1029-33.
- Vinson C, Myakishev M, Acharya A, Mir AA, Moll JR, Bonovich M. 2002. Classification of human B-ZIP proteins based on dimerization properties. *Mol Cell Biol* 22(18):6321-35.
- Vogel CF and Matsumura F. 2009. A new cross-talk between the aryl hydrocarbon receptor and RelB, a member of the NF-kappaB family. *Biochem Pharmacol* 77(4):734-45.
- Wang A, Arantes S, Yan L, Kiguchi K, McArthur MJ, Sahin A, Thames HD, Aldaz CM, Macleod MC. 2008. The transcription factor ATF3 acts as an oncogene in mouse mammary tumorigenesis. *BMC Cancer* 8:268.
- Wang H, Lin G, Zhang Z. 2007. ATF5 promotes cell survival through transcriptional activation of Hsp27 in H9c2 cells. *Cell Biol Int* 31(11):1309-15.
- Wang SC, Zhang L, Hortobagyi GN, Hung MC. 2001. Targeting HER2: Recent developments and future directions for breast cancer patients. *Semin Oncol* 28(6 Suppl 18):21-9.

- Welboren WJ, Sweep FC, Span PN, Stunnenberg HG. 2009. Genomic actions of estrogen receptor alpha: What are the targets and how are they regulated? *Endocr Relat Cancer* 16(4):1073-89.
- Wysocki PJ and Wierusz-Wysocka B. 2010. Obesity, hyperinsulinemia and breast cancer: Novel targets and a novel role for metformin. *Expert Rev Mol Diagn* 10(4):509-19.
- Xue L, Firestone GL, Bjeldanes LF. 2005. DIM stimulates IFN γ gene expression in human breast cancer cells via the specific activation of JNK and p38 pathways. *Oncogene* 24(14):2343-53.
- Yeap BB, Wilce JA, Leedman PJ. 2004. The androgen receptor mRNA. *Bioessays* 26(6):672-82.
- Yin X, Wolford CC, Chang YS, McConoughey SJ, Ramsey SA, Aderem A, Hai T. 2010. ATF3, an adaptive-response gene, enhances TGF β signaling and cancer-initiating cell features in breast cancer cells. *J Cell Sci* 123(Pt 20):3558-65.



Targeting New Pathways and Cell Death in Breast Cancer

Edited by Dr. Rebecca Aft

ISBN 978-953-51-0145-1

Hard cover, 190 pages

Publisher InTech

Published online 29, February, 2012

Published in print edition February, 2012

This book presents novel and interesting findings by multiple accomplished investigators in breast cancer. These chapters elucidate new mechanisms of breast cancer cell death as well as discuss new pathways for therapeutic targeting.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Jeremy K. Haakenson, Mark Kester and David X. Liu (2012). The ATF/CREB Family of Transcription Factors in Breast Cancer, Targeting New Pathways and Cell Death in Breast Cancer, Dr. Rebecca Aft (Ed.), ISBN: 978-953-51-0145-1, InTech, Available from: <http://www.intechopen.com/books/targeting-new-pathways-and-cell-death-in-breast-cancer/the-atf-creb-family-of-transcription-factors-in-breast-cancer>

INTECH

open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2012 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.