

Inequality of Oral Health in a Life-Course Perspective

Dorthe Holst¹ and Annemarie A. Schuller²

¹*Section of Community Dentistry, Dental Faculty, University of Oslo,*

²*TNO Prevention and Health*

¹*Norway*

²*The Netherlands*

1. Introduction

Viewing health and disease in a life-course perspective has gained scientific interest recently (Kuh & Ben-Shlomo, 1997). Panel studies are scarce but other designs come close to being able to follow health and disease through life (Kuh & Ben-Shlomo, 1997). There are three major perspectives in life-course research: One line emphasizes the importance of life-style and deprivation in childhood for adult chronic disease. This research investigates environmental conditions and experiences through prenatal life, infancy, childhood and adolescence that may make individuals more susceptible to developing adult chronic disease. Pearce et al., 2004 studied the effect of birth-weight, early diet, use of comforter and social status on oral health of young adults, but found only effect of social status in the expected direction. Nicolau and co-workers found a relationship between several biological factors and caries among adolescents (Nicolau et al., 2003, 2007). A second line of research assumes biological programming during critical periods of development either during pregnancy or in early life (Barker, 1994). A third line of research suggests an accumulation of risk through the life course. Accumulation of risk is different from programming in that it does not require the notion of a critical period. This approach explicitly places more emphasis on a greater range of biological and social experiences in childhood, adolescence and in early adulthood than either the life style or programming models. There are reasons to believe that adult oral health is affected through a range of life-course mechanisms. The present work leans on the third perspective arguing that oral health is continuously exposed to environmental and behavioral risks that lead to accumulated plaque in the mouth and diseases in the dental tissues (Fejerskov & Kidd, 2008).

On a population basis the vast majority of children are born with a good oral health. Exposure to different life-styles and nutritional and hygienic conditions appears as a threat to oral health through life (Holst & al., 2007). The global picture of oral health is patterned by variation in living conditions and variation in life-course patterns of oral health (Petersen et al., 2006).

A number of studies have described oral health of populations in repeated cross-sectional studies (Schuller & Holst, 1998, Kelly et al., 2000; Skudutyte-Rysstad & Eriksen 2007, Krustrup et al., 2008). These studies provide valuable information about background related changes in oral health conditions at certain points of time. It has for example been shown

that edentulousness has a much lower incidence now than 30 years ago (Petersen et al., 2004; Holst, 2008). The main explanations for this are improved standard of living, availability of fluoride toothpaste and more accessible dental services. Despite the improvement, social status still affects oral health even though recent research indicates that this relationship has become weaker in some countries (Holst, 2008). There is reason to believe that avoiding edentulousness and maintaining oral health requires a life-long attention to healthy diet, oral health promotion, oral hygiene and preventive dental services.

In this chapter the influence of social status on clinical aspects of oral health is assessed from childhood through adolescence to adulthood in the same birth-cohorts in Norway. This unique possibility was made possible through a careful design of a series of cross-sectional studies in the counties of Trøndelag in Norway (Schuller & Holst, 1998). On a population basis the vast majority of children are born with a good oral health. Exposure to different life-styles and nutritional and hygienic conditions forms a threat to oral health through life (Holst & al., 2007). The global picture of oral health is patterned by variation in living conditions and variation in life-course patterns of oral health (Petersen et al., 2006).

A number of studies have described oral health of populations in repeated cross-sectional studies (Schuller & Holst, 1998, Kelly et al., 2000; Skudutyte-Rysstad & Eriksen 2007, Krustrup et al., 2008). These studies provide valuable information about background related changes in oral health conditions at certain points of time. It has for example been shown that edentulousness has a much lower incidence now than 30 years ago (Petersen et al., 2004; Holst, 2008). The main explanations for this are improved standard of living, availability of fluoride toothpaste and more accessible dental services. Despite the improvement, social status still affects oral health even though recent research indicates that this relationship has become weaker in some countries (Holst, 2008). There is reason to believe that avoiding edentulousness and maintaining oral health requires a life-long attention to healthy diet, oral health promotion, oral hygiene and preventive dental services.

2. Material and methods

The material comprised data from independent random samples of three birth-cohorts living in the counties of Sør- and Nord-Trøndelag in 1983. The birth-cohorts were 1929-1938, 1939-1948 and 1959-1960, and they were 45-54-, 34-44 and 23-24-years old in 1983 (Table 1, sample a). In 2006 two samples were drawn from the 1929-1938 and 1959-1960 birth-cohorts in Nord-Trøndelag only, who were then 67-78-year-old and 46-47-year old (Table 1). The age specific sample size for each of the participating counties were 500 in 1983 and was reduced to 250 in 2006. The sample in the two-year age-group 46-47 was 100 persons.

The methods of data collection comprised standardized clinical measurements and self-administered questionnaires (Bærum et al., 1985; Schuller and Holst, 1998). In 1983 and in 2006 ten and two calibrated dental teams, respectively, collected the data. Two senior researchers (DH and AAS) followed and guided the procedures in order to secure standardized conditions and comparability among the surveys. The first Trøndelag study in 1973 started as part of the first WHO International Collaborative Study survey (Arnljot et al.,

1985). The study was also repeated in 1994 but not reported here. Calibration exercises were conducted each study year. Calibration was performed for paired examiners and intra-examiner variability was high for the DMF index ($r > 0.92$). Inter-examiner agreement was exercised until $r > 0.85$ between all pairs, and the results otherwise found satisfactory (Bærum et al., 1985; Holst et al., 2007). The examinations took place at the public dental clinics of the South- and Nord-Trøndelag counties. Permission was granted by public authorities and by the participants' informed consent. All necessary permissions were given throughout the study period and by the participants' informed consent. In 2006 the study was approved by the Regional ethical committee Middle of Norway and approved by the Norwegian Council of Research.

| Birth-cohort | Age | 1983 a | | 1983 b | | 2006 | |
|--------------|-------|--------|----|--------|----|------|----|
| | | n | % | n | % | n | % |
| 1959-1960 | 23-24 | 1000 | 84 | 500 | 81 | | |
| | 46-47 | | | | | 100 | 90 |
| 1929-1938 | 35-44 | 1000 | 82 | 500 | 80 | | |
| | 45-54 | 1000 | 74 | 500 | 72 | 350 | 71 |
| | 68-77 | | | | | 250 | 61 |

1983 a : The sample includes Nord- and Sør-Trøndelag

1983 b: The sample includes Nord-Trøndelag

Table 1. Trøndelagsstudies. Samples in 1983 and 2006 according to birth-cohort and age. Participation in percent

In the present study the outcome variables were number of sound teeth and toothsurfaces (ST, SS) and sound + filled teeth and toothsurfaces (SFT, SFS) and DMFT and DMFS index. DMFT and DMFS are the sums of DT/S, MT/S and FT/S, where DT/S is defined as the number of teeth/toothsurfaces with primary and secondary caries, including root and coronal caries. Only caries with a distinguishable break in the surface was recorded. Missing surfaces is the number of missing tooth surfaces irrespective of cause. FS is the number of surfaces filled, both root and coronal restorations, including all types of filling materials and crowns. The clinical examination comprised recording of the condition of the visible part of the tooth. The analyses were based on 28 teeth excluding third molars.

As part of the study in 1983 twelve questions were asked about social and dental conditions when the sampled persons were ten years old. Social status was measured in two ways: By father's and mother's number of natural teeth in three categories: Many own teeth (2), some own teeth (1) and no own teeth (0). The variables were summed and dummy variables constructed (Table 2, column 4). Eleven other questions were asked about oral health environment at age 10 (Table 2). The questions comprised whether the families had rules for eating sweets, tooth brushing habits, advice about oral health from teachers, school nurse/medical doctor and school dentist, fathers and mothers dental status, visits to a dentist during preschool and school age, parents control of tooth brushing, use of toothpicks

and dental floss. Advice from teachers, school nurse/medical doctor and school dentist were collapsed into an index called advice about oral health (Table 2, column 3). Visits to a dentist during pre-school and school age were summarized to yearly and not yearly. The questions comprising whether the families had rules for eating sweets, tooth brushing habits were combined into parents attention and dummy variables constructed. Sex was included in the meaning of a social construct assuming females to be more engaged in health and oral health behaviors.

Social status in early adulthood was measured as number of years with formal education. Length of education was divided into four quartiles each comprising 25 % of the samples. The first quartile comprised the 25 per cent of the sample with the shortest education, the second quartile the 25 per cent of the sample with the second shortest education, the third quartile the 25 per cent with the second longest education and the fourth quartile comprised the 25 per cent of the sample with the longest education. The quartiles thus represent the distribution of length of education in equally sized groups. Using quartiles eliminates the problem often faced with measuring length of education that the length of education the population changes over time. Length of education was transformed into dummy variables (Table 2, column 4). Four question of oral hygiene practices (1983) were added into oral health behavior index and dummy variables constructed.

| Variables | Categories and coding | Additive indices | Analytical categories/dummy variables |
|--|--|--|--|
| Rules for sweet consumption | Yes (1) no (0) | | |
| Toothbrushing habits | Twice a day (1) Once or less (0) | | |
| Advice from teacher | Yes (1) no (0) | Advice about oral health Advice (1-3) No advice (0+0+0) | Advice (1) No advice (0) |
| Advice from doctor/nurse | Yes (1) no (0) | | |
| Advice from school / district dentist | Yes (1) no (0) | | |
| Father's dental status | Many teeth (2), few (1) none (0) | Parents' dental status Many (4) Few (1-3) None (0) | Dummy parents' dental status many Dummy parents'dental status few Reference category |
| Mother's dental status | Many teeth (2), few (1) none (0) | | |
| | | | |
| Visited a dentist pre school | Yearly (2) a few times (1) never (0) | Dental care at age 10 Yearly (4) A few times (1-3) Never (0) | Yearly (1), Not yearly (0) |
| Visited a dentist during school | Yearly (2) a few times (1) never (0) | | |
| Parents controlled toothbrushing | Often/daily (2) a few times (1) never (0) | Parents' attention Daily (5-6) A few times (1-3) Never (0) | Dummy parents attention daily Dummy parents attention a few times Reference category |
| Used toothpicks | Often/daily (2) a few times (1) never (0) | | |
| Used dental floss | Often daily/ (2) a few times (1) never (0) | | |
| Gender | Female (1) Male (0) | | |
| Length of education | Highest quartile (3) | | Dummy highest quartile |
| | Second highest quartile (2) | | Dummy second highest quartile |
| | Second lowest quartile (1) | | Dummy second lowest quartile |
| | Lowest quartile (0) | | Reference category |
| Regular dental visits last three years | Regular each year (1)irregular (0) | | |
| Brushed yesterday | Yes (1) No (0) | Oral health behaviour Good (4) Middle (2-3) Bad (0-1) | Dummy oral behaviour good Dummy oral behaviour middle Reference category |
| Used dental floss yesterday | Yes (1) No (0) | | |
| Used toothpicks yesterday | Yes (1) No (0) | | |
| Had sweets yesterday | Yes (1) No (0) | | |

Table 2. Variables, categories, indices and analytical categories

2.1 Analysis

In the first part of the analysis social status together with the early oral health environment and sex were related to oral health variables in 1983 by multiple regressions. Since all the dependent variables were measured on the same scale (tooth surfaces, range 1-128), the regression coefficients can be interpreted directly as effects of the independent variables in number of surfaces. For the second part of the analysis the data files from 1983 and 2006 were combined to one data file in order to study whether the impact of social status changed during this period. Multiple regression analysis was used, and the level of significance was $p = 0.05$. Associations nearly reaching significance ($0.07 > p > 0.05$) are shown.

3. Results

The distribution of the independent variables according to age-groups is shown in Table 3. The table provides a picture of how the oral health environment at age ten years varied between the birth-cohorts. The youngest birth-cohort had the best level of oral health environment.

| | Age in 1983 | | |
|--|-------------|------------|------------|
| | 23-24 year | 35-44 year | 45-54 year |
| Oral health environment at age 10 | | | |
| Had rules for sweet consumption | 29.4 | 20.4 | 14.4 |
| Toothbrushing twice per day | 73.0 | 64.0 | 59.4 |
| Got advice about oral health | 98.2 | 70.1 | 53.8 |
| Parents' many teeth | 72.4 | 47.0 | 47.4 |
| Dental care yearly | 19.3 | 6.1 | 3.7 |
| Parents' attention high | 34.2 | 12.4 | 6.8 |
| Behaviour in 1983 | | | |
| Regular dental visits | 65.8 | 69.4 | 63.7 |
| Oral health behaviour good | 23.8 | 15.2 | 11.0 |

Table 3. Descriptive statistics for independent variables in 1983. Sample a. Percentage

Table 4 shows the means and standard deviations of the clinical variables. The table provides a clear picture of the variation between the age-groups in the condition of the tooth surfaces. The table also shows how different clinical indicators show very different results. The number of sound tooth-surfaces is absolutely highest in the youngest age group and the number of DMFS and MS highest in the oldest age-group.

Tables 5-7 show how the early oral health environment, the social variables and the oral health behaviours each and combined (R^2) affected oral health in the age-groups in 1983.

| Age-group | n | SS | | DS | | FS | | MS | | DMFS | | |
|-----------|-------|-------|-------|-------|------|-------|-------|-------|-------|-------|-------|-------|
| | | Mean | sd | Mean | sd | Mean | sd | Mean | sd | Mean | sd | |
| 23-24 | 773 | 84.10 | 17.41 | 1.09 | 2.52 | 37.89 | 16.05 | 4.89 | 7.34 | 43.89 | 17.36 | |
| 1983* | 35-44 | 773 | 48.54 | 21.51 | 2.25 | 6.23 | 48.64 | 22.86 | 28.55 | 30.77 | 79.45 | 21.51 |
| | 45-54 | 675 | 35.11 | 24.24 | 2.19 | 5.38 | 37.02 | 27.3 | 53.66 | 42.51 | 92.88 | 24.24 |
| 2006 | 46-47 | 96 | 79.08 | 21.02 | 0.80 | 1.9 | 38.59 | 16.73 | 5.38 | 7.2 | 44.77 | 18.83 |
| | 68-77 | 150 | 33.02 | 22.02 | 1.61 | 3.31 | 39.53 | 25.71 | 44.16 | 36.53 | 85.29 | 20.17 |

* Sample a

Table 4. Dental variables in 1883 and 2006. Mean and standard deviation (sd) (basis 124 tooth surfaces, 28 teeth)

3.1 Birth-cohort 1959-1960, 23-24 years in 1983

Table 5 shows the impact of the independent variables on the outcome variables. Having positive oral health behaviors at age ten increased the likelihood of more sound surfaces (SS), more filled surfaces (FS) and more surfaces with caries experience (DMFS) at age 23-24. If parents had many own teeth the 23-24-years-olds had 5.0 more sound surfaces (SS) and 5.0 less surfaces with caries experience (DMFS). Length of education was statistically significantly related to the D-M-F-S variables in the expected direction. Regular dental care was related to mean number of surfaces with untreated decay (DS). Good oral health behavior was statistically significant related to more sound surfaces (SS) and less filled surfaces (FS). The variables in the model explained from 4 -13 % of the variation in the dependent variables.

3.2 Birth-cohort 1939-1948, 35-44 years in 1983

Table 6 shows that parents' dental status and yearly dental visits at age ten had a statistically significant impact on several of the oral health variables. Having parents with many of their own teeth at age ten the 35-44 year olds had more functional surfaces (SFS), less missing surfaces (MS) and surfaces with caries experience (DMFS). Women had less sound surfaces (SS) and more filled surfaces (FS) and DMFS than men when they were 35-44-years-old in 1983. The longer the education, the better the values of the oral health indicators were; the differences between the quartiles of education were big. Dental care last year had a statistical significant influence on DS, SFS, MS and FS. The explained variation varied from 11 - 27 %.

3.3 Birth-cohort 1929-1938, 45-54 years in 1983

Table 7 shows that parents' dental status at age ten years had a statistically significant effect on the oral health variables. In addition tooth brushing and dental care at age 10 had a significant effect on untreated caries (DS). Women had more sound surfaces (SS), less untreated decay (DS), and more filled (FS) and DMFS than men had. Length of education had a significant effect on all oral health variables except untreated caries (DS). Oral health behavior had a similar effect, while regular dental care also affected untreated caries (DS).

Table 5. Relationship between DMFS and life-course variables and indices. Multiple regression. Birth-cohort 1959-60 in 1983

| Variables | Oral health | | | | Untreated disease | | FS | |
|------------------------------------|-------------------|------|--------------------|------|-------------------|------|-------------|------|
| | SS Coefficient | se | SFS Coefficient | se | DS Coefficient | se | Coefficient | se |
| Intercept | 76.13 | 4.62 | 118.92 | 2.11 | 4.07 | 0.66 | 42.79 | 4.33 |
| At age 10 | | | | | | | | |
| Rules for sweet consumption | 2.42 | 1.53 | 0.60 | 0.7 | -0.19 | 0.22 | -1.82 | 1.43 |
| Toothbrushing habits | 4.39 * | 1.78 | -0.71 | 0.81 | 0.21 | 0.25 | 3.68 * | 1.67 |
| Advice about oral health | -0.33 | 3.69 | -1.57 | 1.68 | -0.43 | 0.52 | -1.24 | 3.46 |
| Parents dental status few teeth | -0.56 | 2.77 | 1.21 | 1.27 | 0.03 | 0.39 | 1.77 | 2.60 |
| Parents dental status many teeth | 5.04 * | 2.51 | 2.14 ^ | 1.15 | -0.39 | 0.36 | -2.91 | 2.35 |
| Dental care at age 10 | -2.09 | 1.59 | -0.16 | 0.72 | -1.41 | 0.22 | 1.93 | 1.48 |
| Parents' attention a few times | -1.29 | 1.62 | -0.16 | 0.74 | -0.34 | 0.23 | 1.13 | 1.52 |
| Parents' attention daily | -0.37 | 1.96 | -0.22 | 0.9 | -0.31 | 0.28 | 0.15 | 1.85 |
| Early adult age | | | | | | | | |
| Gender | 0.06 | 1.37 | -0.56 | 0.62 | -0.57 * | 0.19 | -6.26 | 1.28 |
| Education-second lowest quartile | 3.65 * | 1.71 | 2.33 * | 0.78 | -0.92 * | 0.24 | -1.32 | 1.60 |
| Education-second highest quartile | 3.96 * | 1.82 | 1.91 * | 0.83 | -0.98 * | 0.26 | -2.05 | 1.70 |
| Education- second highest quartile | 8.84 * | 1.86 | 4.04 * | 0.85 | -1.28 * | 0.26 | -4.80 * | 1.74 |
| Last year | | | | | | | | |
| Regular dental care | -1.03 | 1.33 | 0.54 | 0.61 | -1.05 * | 0.19 | 1.58 | 1.24 |
| Oral health behaviour middle | 5.14 * | 2.13 | 1.08 | 0.97 | -0.58 * | 0.30 | -4.06 * | 2.00 |
| Oral health behaviour- good | 5.77 * | 2.44 | 1.36 | 0.12 | -0.45 | 0.35 | -4.39 * | 2.29 |
| R ² | 0.07 | | 0.05 | | 0.13 | | 0.04 | |

* = p < 0.05

^ = 0.05 < p < 0.07

Table 6. Relationship between DMFS and life-course variables and indices. Multiple regression. Birth-cohort 1939-1948 in 1983

| Variables | Oral health | | | | Untreated disease | | FS | | |
|------------------------------------|-------------|------|-------------|------|-------------------|------|-------------|----------|--|
| | SS | | SFS | | DS | | FS | | |
| | Coefficient | se | Coefficient | se | Coefficient | se | Coefficient | se | |
| Intercept | 37.86 | 2.77 | 58.28 | 3.77 | 7.52 | 0.81 | 24.14 | 2.61 | |
| At age 10 | | | | | | | | | |
| Rules for sweet consumption | 0.45 | 2.35 | -1.87 | 3.20 | -0.07 | 0.67 | -1.83 | 2.16 | |
| Toothbrushing habits | -0.38 | 2.08 | -3.57 | 2.84 | -0.61 | 0.60 | -2.94 | 1.93 | |
| Advice about oral health | 1.84 | 1.66 | -0.38 | 2.26 | -0.08 | 0.47 | -2.15 | 1.53 | |
| Parents dental status few teeth | -0.12 | 2.14 | 2.59 | 2.92 | 1.28 * | 0.61 | 1.84 | 1.98 | |
| Parents dental status many teeth | 3.94 * | 2.05 | 6.25 * | 2.80 | 0.63 | 0.59 | 1.65 | 1.89 | |
| Dental care at age 10 | -7.60 * | 3.32 | -1.06 | 4.53 | 0.87 | 0.93 | 5.58 ^ | 3.01 | |
| Parents' attention a few times | -0.59 | 1.89 | 2.11 | 2.58 | -0.03 | 0.54 | 2.93 | 1.74 | |
| Parents' attention daily | 0.21 | 3.23 | 1.26 | 4.40 | -0.06 | 0.92 | 0.77 | 2.97 | |
| Early adult age | | | | | | | | | |
| Gender | -7.23 * | 1.66 | -1.06 | 2.26 | -0.79 | 0.47 | 6.31 * | 1.52 | |
| Education-second lowest quartile | 5.49 * | 2.06 | 10.95 * | 2.82 | -1.06 | 0.59 | 4.78 * | 1.91 | |
| Education-second highest quartile | 6.78 * | 2.14 | 20.97 * | 2.92 | -1.65 * | 0.61 | 13.04 * | 1.97 | |
| Education- second highest quartile | 11.98 * | 2.26 | 27.80 * | 3.08 | -1.55 * | 0.64 | 14.60 * | 2.07 | |
| Last year | | | | | | | | | |
| Regular dental care | 2.58 | 1.74 | 17.38 * | 2.38 | -4.73 * | 0.50 | 12.97 * | 1.62 | |
| Oral health behaviour middle | 6.43 * | 2.28 | 14.05 * | 3.11 | -0.65 | 0.66 | 6.88 * | 2.13 | |
| Oral health behaviour good | 3.15 | 2.86 | 15.72 * | 3.90 | -1.11 | 0.83 | 12.56 * | 2.66 | |
| R ² | 0.11 | | 0.26 | | 0.17 | | 0.27 | | |
| | | | | | | | | ^p=0.064 | |

* = p < 0.05

^ = 0.05 < p < 0.07

Table 7. Relationship between DMFS-and life-course variables and indices. Multiple regression. Birth-cohort 1929-38 in 1983

| Variables | Oral health | | | | Untreated disease | | | |
|------------------------------------|-------------|------|-------------|------|-------------------|------|-------------|------|
| | SS | | SFS | | DS | | FS | |
| | Coefficient | se | Coefficient | se | Coefficient | se | Coefficient | se |
| Intercept | 18.46 | 2.70 | 17.95 | 3.99 | 8.19 | 0.82 | 6.61 | 2.82 |
| At age 10 | | | | | | | | |
| Rules for sweet consumption | -4.20 | 2.85 | -7.65 ^ | 4.20 | 1.35 | 0.79 | -1.68 | 2.72 |
| Toothbrushing habits | -0.01 | 2.26 | 2.41 | 3.34 | -1.51 * | 0.65 | 3.51 | 2.25 |
| Advice about oral health | 0.47 | 1.73 | 0.91 | 2.55 | -0.30 | 0.48 | 0.89 | 1.65 |
| Parents dental status few teeth | 3.45 | 2.29 | 5.15 | 3.38 | -0.78 | 0.65 | 0.69 | 2.25 |
| Parents dental status many teeth | 5.86 * | 2.26 | 8.53 * | 3.34 | -0.79 | 0.64 | 1.65 | 2.22 |
| Dental care at age 10 | -7.91 | 3.59 | 1.45 | 5.29 | 2.06 * | 0.94 | 1.30 | 3.24 |
| Parents' attention a few times | -0.02 | 2.06 | 1.94 | 3.04 | 0.69 | 0.56 | 1.68 | 1.95 |
| Parents' attention daily | 0.76 | 3.78 | 6.12 | 5.57 | -0.02 | 1.03 | 3.95 | 3.54 |
| Early adult age | | | | | | | | |
| Gender | -8.24 * | 1.86 | -3.47 | 2.74 | -1.27 * | 0.52 | 7.39 * | 1.80 |
| Education-second lowest quartile | 3.65 | 2.23 | 8.35 * | 3.29 | -0.76 | 0.64 | 5.11 * | 2.22 |
| Education-second highest quartile | 4.68 ^ | 2.58 | 19.27 * | 3.81 | -0.52 | 0.73 | 17.50 * | 2.51 |
| Education- second highest quartile | 13.36 * | 2.48 | 32.93 * | 3.66 | -0.93 | 0.67 | 18.53 * | 2.32 |
| Last year | | | | | | | | |
| Regular dental care | 10.17 * | 1.88 | 36.16 * | 2.78 | -3.23 * | 0.56 | 19.92 * | 1.92 |
| Oral health behaviour middle | 8.58 * | 2.32 | 15.52 * | 3.42 | -0.94 | 0.67 | 4.45 * | 2.32 |
| Oral health behaviour good | 9.10 * | 3.16 | 16.56 * | 4.65 | -0.46 | 0.90 | 4.70 | 3.09 |
| R ² | 0.191 | | 0.434 | | 0.135 | | 0.388 | |
| | ^p=0.070 | | ^p=0.069 | | | | | |

* = p< 0.05

^ =0.05< p< 0.07

3.4 The influence of social status from 1983 to 2006

For this part of the analysis data files from 1983 and 2006 were combined into one datafile. An interaction term between social status measured as length of education and study year was included in order to see if the effect of length of education was important both years or only one of the years.

3.4.1 Birth-cohort 1959-60, age 23-24 and 46-47 in the combined file

Table 8 shows that the interaction between length of education and study year was not significant with the exception of the effect on sound surfaces, where the persons in the second lowest education group had kept nearly 15 more sound surfaces than the lowest group. Gender and regular dental care had an independent effect on mean number of sound surfaces (DS). Oral health behavior had a significant effect on oral health variables.

3.4.2 Birth-cohort 1929-38, age 45-54 and 68-77 in the combined file

Table 9 shows that the effect of social status measured as length of education was not dependent upon which year it was measured except that the persons in the second highest education quartile on average had more surfaces with untreated decay compared with persons in the lowest quartile. In this birth-cohort gender and particularly regular dental care had an effect on several of the outcome variables. Functional surfaces increased by nearly 34 surfaces among persons with visits to the dentist compared to those without. Likewise the average number of FS and MS were much higher and lower, respectively, among those with regular dental care.

4. Discussion

This study has shown that social status and oral health environment in childhood was important for adults' oral health during the most of the 20. century. The attention from parents and the local environment lead to a better oral health outcome in adulthood. Social status measured by length of education was also a personal resource that guided choices leading to better oral health. The longer the education the better the oral health was. Regular dental visits were important especially for the eldest birth-cohort. Good oral health behaviors early and during adulthood were also important for oral health. Effects of more than 30 surfaces were found on indicators like missing and functional tooth surfaces. When the birth-cohorts were followed from 1983 to 2006 social status had an effect in both 1983 and 2006. Judged by the number of tooth surfaces the difference between social status groups had not increased by 2006. The latter observation deserves a critical comment. The cumulative DMFS measure is sensitive to increased levels of risk factors in the sense that more surfaces can be affected, until saturation is reached. When lower risk levels occur, the DMFS figures cannot decline within the same birth-cohort. The Missing, Filled and Sound indicators (the DMF index) cannot reverse. Only the number of decayed surfaces can reverse (Holst and Schuller, 2000). In the present study the average number of decayed tooth-surfaces was significantly reduced and indicated a lower level of recent risk (Holst et al., 2007). This is a serious limitation. With regard to estimating the influence of social status and other explanatory variables, a reduced effect can thus not be shown, and it can only be concluded that the effect of social status did not increase from 1983 to 2006. In a cohort analysis of the relationship between social status and average number of DMFT in 35-44-

Table 8. Relationship between DMFS variables and independent variables. Combined datafile 1983 and 2006. Birth-cohort 1959-1960

| Variables | Oral health | | | | Untreated disease | | | |
|-----------------------------------|-------------|------|-------------|------|-------------------|------|-------------|------|
| | SS | | SFS | | DS | | FS | |
| | Coefficient | se | Coefficient | se | Coefficient | se | Coefficient | se |
| Intercept | 73.24 | 2.97 | 117.84 | 1.42 | 4.47 | 0.42 | 44.37 | 2.68 |
| Gender | 0.36 | 1.81 | -1.23 | 0.89 | -0.68 * | 0.26 | -1.09 | 1.63 |
| Regular dental care | -0.61 | 2.05 | -2.24 | 0.96 | -0.74 * | 0.29 | 0.14 | 1.85 |
| Education-second lowest quartile | 5.09 | 2.79 | 3.16 * | 1.30 | -1.73 * | 0.40 | -1.97 | 2.52 |
| Education-second highest quartile | 1.06 | 3.00 | 2.65 ^ | 1.39 | -1.54 * | 0.43 | 1.49 | 2.71 |
| Education-highest quartile | 11.84 * | 2.95 | 6.40 * | 1.37 | -2.08 * | 0.42 | -5.64 * | 2.67 |
| Oral behaviour middle | 5.88 * | 2.45 | 0.91 | 1.23 | -1.12 * | 0.35 | -4.91 * | 2.22 |
| Oral behaviour good | 6.74 * | 2.81 | 1.22 | 1.42 | -1.30 * | 0.40 | -4.67 ^ | 2.54 |
| Year | 3.57 | 4.55 | -3.35 | 2.55 | -1.11 | 0.65 | -2.26 | 4.11 |
| Interact. Year/second lowest q. | -14.55 * | 5.84 | -2.55 | 3.30 | 1.08 | 0.83 | 7.43 | 5.28 |
| Interact. Year/second highest q. | -1.47 | 6.20 | -2.54 | 3.88 | 0.22 | 0.88 | -2.57 | 5.60 |
| Interact. Year/ highest q. | -1.10 | 6.68 | -3.71 | 3.78 | 0.66 | 0.95 | -3.40 | 6.03 |
| R ² | 0.11 | | 0.12 | | 0.16 | | 0.06 | |
| | | | ^p=0.058 | | | | ^p=0.067 | |

* = p < 0.05

^ = 0.05 < p < 0.07

Table 9. Relationship between DMFS- and independent variables. Combined datatiles 1983 and 2006. Birth-cohort 1929-1938

| Variables | Oral health | | | | Untreated disease | | | | |
|-----------------------------------|-------------|------|-------------|-------|-------------------|------|-------------|------|--|
| | SS | | SFS | | DS | | FS | | |
| | Coefficient | se | Coefficient | se | Coefficient | se | Coefficient | se | |
| Intercept | 20.45 | 2.96 | 23.64 | 4.42 | 5.20 | 0.71 | 0.02 | 2.77 | |
| Gender | -4.31 ^ | 2.20 | 1.57 | 3.30 | -1.23 * | 0.53 | 6.03 * | 2.06 | |
| Regular dental care | 12.02 * | 2.63 | 33.84 * | 4.05 | -1.29 * | 0.63 | 25.55 * | 2.46 | |
| Education-second lowest quartile | 1.38 | 3.30 | 6.57 | 4.80 | -1.06 | 0.79 | 4.79 | 3.08 | |
| Education-second highest quartile | 6.08 | 3.80 | 22.72 * | 5.54 | -1.10 | 0.91 | 15.64 * | 3.55 | |
| Education-highest quartile | 10.50 * | 4.08 | 33.98 * | 5.98 | -9.63 | 0.98 | 21.38 * | 3.82 | |
| Oral behaviour middle | 6.99 * | 2.63 | 14.31 * | 3.96 | -0.64 | 0.63 | 9.31 * | 2.46 | |
| Oral behaviour good | 0.87 | 3.28 | 6.98 | 4.98 | -0.63 | 0.79 | 8.52 * | 3.06 | |
| Year | -6.42 | 3.87 | 9.27 | 6.47 | -2.14 * | 0.93 | 2.65 | 3.62 | |
| Interact. Year/second lowest q. | 7.32 | 5.58 | 3.03 | 8.79 | 1.25 | 1.34 | 2.67 | 5.22 | |
| Interact. Year/second highest q. | 5.78 | 6.29 | -12.92 | 9.58 | 3.65 * | 1.51 | -6.91 | 5.88 | |
| Interact. Year/ highest q. | 9.56 | 6.79 | -12.23 | 10.63 | 2.02 | 1.63 | -10.29 | 6.35 | |
| R ² | 0.16 | | 0.36 | | 0.06 | | 0.43 | | |
| | ^p=0.050 | | | | | | | | |

* = p < 0.05

^ = 0.05 < p < 0.07

year-olds in 1983 and a new cohort of 35-44-year-olds in 2006 from the same material, the relationship between social status and number of present teeth had disappeared in 2006, and the relationship between social status and average number of DMFT and DMFS was significantly reduced (Holst et al., 2007). This shows that the DMF index can be used in cross-sectional research comparing birth-cohorts of the same age; the index has serious limitations in semi-longitudinal research.

It is important to draw attention to the different dimensions that the chosen oral health indicators reflect. The indicators SS and SFS reflect oral health and function and high and increasing values represent positive expressions of oral health. MF and FS are negative expressions of oral health and high and increasing values show reduced oral health. These treatment indicators have limitations since they do not include repeated treatment in the same teeth. DMFS (or DMFT) are a summarized expression of untreated and treated disease, and the values may be difficult to interpret because the indicators of the index move in contrasting directions over time. It is important that researchers in oral epidemiology engage in finding new ways measures of disease activity that are different from measures of treatment activity.

There are a number of threats to reliability and validity of the data when surveys are repeated and the same variables are used over time, and different birth-cohorts are exposed to the same procedures. Concepts of behavioral norms and interpretation of clinical symptoms change, and treatment criteria change (Gimmestad & Holst, 2003). Most of the questions in the present surveys, however, were about factual events and clear to the respondents; some memory bias among the respondents with regard to events at age 10 years should be expected. These are measurement errors that increase the variance of the variables and reduce the discriminative ability of the statistical tests. Even though one of the authors (DH) was present at all the surveys and has acted as the golden standard, it is difficult to avoid flow in the translation of the standardized criteria.

The results from the present study have a limited statistical inference with regard to the size of the population the results may be generalized to. On the other hand when it comes to modeling social processes generalization is based on how validly the model catches the specific underlying social processes. It was not the intention to explain all the variation in the dependent variables. It is interesting to notice that R^2 was high in the oldest cohort. It cannot be settled whether this is a cohort or an age effect. Probably it is both, assuming that age reflects the cumulative exposure to plaque during the life-course, and the later born birth-cohorts have experienced a different environment that will result in a better oral health. There are reasons to believe that our data and the model have caught some of those social processes that were important for oral health and its development over time. Other and nationally representative Norwegian data support the finding of a more equally distributed oral health (Holst et al., 2007; Skudutyte and Eriksen, 2007, Holst & Skau, 2010). The Trøndelag studies started at a time where data on oral health and its determinants were scarce. In hind-sight these studies have provided opportunities for valuable descriptions and explanations of the changes in oral health.

The Norway is considered to have had a homogeneous population compared to many other countries (Krokstad & Westin, 2004). Yet, the demography, the size of the country and the arctic location has resulted in cultural and distributional differences. Living conditions and social disparities have to a large extent affected oral health of the population previously in Norway (Arnljot et al., 1985). During the last decade larger income differences have been

observed that might have lead to increased social inequalities in both oral health and demand for dental services (Krokstad 2004). That seems not to have occurred. Cross-sectional data will typically focus on cross-sectional social differences and discuss these with limited insight over time. Often will the lead time between exposure and result be overlooked. Panel data and data with the present analytical potential can detect whether or not a social problem is increasing or decreasing. It cannot be ignored that the results of this study can be ascribed to welfare policies across a number of living conditions in Norway. The public dental service with a population responsibility and out-reach services in this country is an example of one such public policy that have contributed to increasing public awareness of oral health as a value. A high level of public awareness may be expected to influence both the promotion of oral health and accessible adequate dental care. A life-course perspective provides an opportunity understand oral health over time. The present work supports the assumption that oral health is continuously exposed to environmental and behavioral risks that affect life-time oral health.

5. Conclusion

The purpose of the work was to study the influence of 1) the oral health environment at age 10, 2) of adolescent and adulthood dental behaviors and of 3) social status on oral health in three birth-cohorts later in life (1983) and in 2006 in Norway.

The material comprised data from random samples of three birth-cohorts living in the counties of Sør- and Nord-Trøndelag in 1983. The birth-cohorts were 1929-1938, 1939-1948 and 1959-1960. In 2006 two samples were drawn from the 1929-1938 and 1959-1960 birth-cohort. The data collection comprised standardized clinical measurements and self-administered questionnaires. The early oral health environment and social status and sex were related to oral health in 1983 by multiple regressions. The impact of social status was studied in combined datafiles from 1983 and 2006.

The oral health environment in childhood was important for adults' oral health. The attention from parents and the local environment lead to a better oral health outcome in adulthood. Social status affected choices leading to better oral health. Regular dental visits were important especially for the eldest birth-cohort. Good oral health behaviors early and during adulthood were also important for oral health. Judged by number of tooth surfaces the difference between social status groups had not increased by 2006.

A life-course perspective provides an opportunity to understand oral health over time. The present study supports the assumption that oral health is continuously exposed to environmental and behavioral risks that lead to accumulated diseases in the dental tissues.

6. References

- Arnlot, H.A.; Barmes, D.E.; Cohen, L.K.; Hunter, P.B.V. & Ship, II.(1985). *Oral health care systems. An international collaborative study.* World Health Organization , Geneva.
- Barker, D.J.P. (1994). Fetal origin of adult disease. *The Fetal and Maternal Medicine Review*, Vol. 6, pp. 71-80.
- Bærum, P.; Holst, D. & Rise, J. (1985). *Dental health in Trøndelag 1983. Changes from 1973 - 1983.* Helsedirektoratet, Oslo.

- Fejerskov, O. & Kidd, E. (2008). *Dental caries. The disease and its clinical management*. 2nd edn. Blackwell Munksgaard, Oxford.
- Gimmestad, A.L. & Holst, D. (2003). Changes in restorative caries treatment in 15-year-olds in Oslo, Norway, 1979–1996. *Community Dentistry and Oral Epidemiology*, Vol. 31, pp. 246-251.
- Holst, D. (2008). Oral health equality during 30 years in Norway. *Community Dentistry and Oral Epidemiology*, Vol. 36, pp.326-334.
- Holst, D. & Schuller, A. A. (2000). Oral health changes in an adult Norwegian population: a cohort analytical approach. *Community Dentistry and Oral Epidemiology*, Vol. 28, pp. 102-111.
- Holst, D., Schuller, A.A. & Dahl, K.E. (2007). Bedre tannhelse for alle? Tannhelseutvikling i den voksne befolkning i Nord-Trøndelag fra 1973 til 2006. *Den norske tannlegeforenings Tidende*, Vol. 117, pp. 804-811.
- Holst D. & Skau, I. (2010). Tenner og tannstatus i den voksne befolkning i Norge *Den norske tannlegeforenings Tidende*, Vol. 129, pp. 164-169.
- Kelly, M., Steele, J., Nuttall, N., Bradnock, G., Morris, J., Nunn, J., Pine, C., Pitts, N., Treasure, E. & White, D. (2000). *Adult Dental Health Survey. Oral health in the United Kingdom 1998*. The Stationery Office, London.
- Krokstad, S. (2004). *Socioeconomic inequalities in health and disability. Social epidemiology in the Nord-Trøndelag Health Study (HUNT), Norway*. Norwegian University of Science and Technology, Verdal.
- Krokstad, S., Westin, S. (2004). Disability in society-medical and non-medical determinants for disability pension in a Norwegian total county population study. *Social Science & Medicine*, Vol. 58, pp. 1837-1848.
- Krustrup, U., Holm-Pedersen, P., Petersen, P.E., Lund, R. & Avlund, K. (2008). The overtime effect of social position on dental caries experience in a group of old-aged Danes born in 1914. *Journal of Public Health Dentistry*, Vol. 68, pp. 46-52.
- Kuh, D., & Ben-Shlomo, Y. (1997). *A life course approach to chronic disease epidemiology*. Oxford University Press, New York.
- Nicolau, B., Marcenes, W., Bartley, M. & Sheiham, A. (2003). A life course approach to assessing causes of dental caries experience: The relationship between biological, behavioural, socio-economic and psychological conditions and caries in adolescents. *Caries Research*, Vol. 37, pp. 319-326.
- Nicolau, B., Thomson, W.M., Steele, J.G. & Allison, P.J. (2007). Life-course epidemiology: concepts and theoretical models and its relevance to chronic oral conditions. *Community Dentistry and Oral Epidemiology*, Vol. 35, pp. 241-249.
- Pearce M.S., Steele J.G., Mason J., Walls A.W.G. & Parker L. (2004). Do circumstances in early life contribute to tooth retention in middle age? *Journal of Dental Research*, Vol. 83, pp. 562-566.
- Petersen, P.E., Kjølner, M. & Christensen, L.B. (2004). Changing dentate status of adults, use of dental health services, and achievement of national dental health goals in Denmark by the year 2000. *Journal of Public Health Dentistry*, Vol. 64, pp. 227-235.
- Petersen P.E, Burgeois D., Ogawa H., Estipunan-Day S. & Ndiaye C. (2005) The global burden of oral diseases and oral health. *Bulletin of the World Health Organization*, Vol. 83, pp. 661-669.

- Schuller, A.A. & Holst, D. (1998). Changes in the adult oral health of adults from Trøndelag, Norway 1973-1983-1994. *Community Dentistry and Oral Epidemiology*, Vol. 26, pp. 201-208.
- Skudutyte-Rysstad, R. & Eriksen, H. (2007). Changes in caries experience among 35-year-old Oslo citizens, 1973-2003. *Acta Odontologica Scandinavica*. Vol. 65, pp. 72-77.



Oral Health Care - Prosthodontics, Periodontology, Biology, Research and Systemic Conditions

Edited by Prof. Mandeep Virdi

ISBN 978-953-51-0040-9

Hard cover, 372 pages

Publisher InTech

Published online 29, February, 2012

Published in print edition February, 2012

Geriatric dentistry, or gerodontics, is the branch of dental care dealing with older adults involving the diagnosis, prevention, and treatment of problems associated with normal aging and age-related diseases as part of an interdisciplinary team with other healthcare professionals. Prosthodontics is the dental specialty pertaining to the diagnosis, treatment planning, rehabilitation, and maintenance of the oral function, comfort, appearance, and health of patients with clinical conditions associated with missing or deficient teeth and/or oral and maxillofacial tissues using biocompatible materials. Periodontology, or Periodontics, is the specialty of oral healthcare that concerns supporting structures of teeth, diseases, and conditions that affect them. The supporting tissues are known as the periodontium, which includes the gingiva (gums), alveolar bone, cementum, and the periodontal ligament. Oral biology deals with the microbiota and their interaction within the oral region. Research in oral health and systemic conditions concerns the effect of various systemic conditions on the oral cavity and conversely helps to diagnose various systemic conditions.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Dorthe Holst and Annemarie A. Schuller (2012). Inequality of Oral Health in a Life-Course Perspective, Oral Health Care - Prosthodontics, Periodontology, Biology, Research and Systemic Conditions, Prof. Mandeep Virdi (Ed.), ISBN: 978-953-51-0040-9, InTech, Available from: <http://www.intechopen.com/books/oral-health-care-prosthodontics-periodontology-biology-research-and-systemic-conditions/dynamic-changes-of-oral-health>

INTECH
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2012 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.