

Expression of the Histamine H₄ Receptor in Human Tissue

Katsunori Yamaura¹, Masahiko Suzuki², Takao Namiki³ and Koichi Ueno^{1,4}

¹*Department of Geriatric Pharmacology and Therapeutics, Graduate School of Pharmaceutical Sciences,*

²*Research Center for Frontier Medical Engineering,*

³*Department of Japanese-Oriental "KAMPO" Medicine, Graduate School of Medicine,*

⁴*Center for Preventive Medical Science, Chiba University, Japan*

1. Introduction

The histamine H₄ receptor (H₄R) is the most recently identified of the four histamine receptors (H₁R–H₄R), and belongs to the same G-protein-coupled receptor (GPCR) family. The amino acid sequence of H₄R shares approximately 26%, 27%, and 58% homology with H₁R, H₂R, and H₃R, respectively, in the transmembrane regions (Nguyen et al., 2001). Furthermore, H₄R couples to Gi/o proteins and shows 10³- to 10⁴-fold higher ligand affinity than H₁R and H₂R (Thurmond et al., 2008).

Several organs express H₄R, and immune tissues such as the spleen, thymus, bone marrow, and leukocytes have a wide range of expression levels (Oda et al., 2000). It has been reported that chemotaxis of mast cells and eosinophils is stimulated by histamine via H₄R; the receptor is therefore attractive as a new target of research into allergic diseases (de Esch et al., 2005).

2. Expression of H₄R in synovial tissue in rheumatoid arthritis

A role for histamine has been implicated in rheumatoid arthritis (RA). RA consists mainly of synovial tissue inflammation that may be dispersed throughout the body, but its molecular etiology remains unclear. Macrophage infiltration and excessive formation of fibroblasts cause a variety of cytokines to be secreted from synovial membranes in patients with RA, and this in turn stimulates osteolytic activity (Sweeney & Firestein, 2004). There is evidence of a significant increase in histamine concentration in synovial samples from patients with RA (Frewin et al., 1986). These observations suggest a potentially significant role of H₄R in the cause, progression, and treatment of RA.

The presence of H₁R and H₂R in human synovial cell culture (HSCC) has been clearly shown by ligand-binding experiments (Nagata, 1991). However, there has been no definitive evidence or conclusive reports of the similar presence of H₃R or H₄R. Therefore, utilizing our expertise in reverse transcription polymerase chain reaction (RT-PCR) techniques, we examined the H₄R-specific mRNA expression in HSCC obtained from 11 RA patients who underwent artificial knee-replacement surgery (Ikawa et al., 2005).

macrophage markers showed that human H₄R protein is expressed in both fibroblast-like and macrophage-like cells in RA synovial tissues (Fig. 3 and Fig. 4). Others have also reported identification of H₄R in synovial tissue of patients with RA (Grzybowska-Kowalczyk et al., 2007).




Fig. 2. Expression of H₄R-specific mRNA in HSCC from 11 RA patients. The gel was loaded with 5 μ L of amplified products. A 100-bp DNA ladder was used as a molecular weight marker. S: standard; C: control; RE: relative expression; sample from 11 different RA patients, RA1 to RA11. RE of mRNA was calculated by normalizing the separated sample intensity value, taking that of the corresponding internal control (GAPDH) as 100%. Intensity values were measured using an image analyzer (IX81, OLYMPUS). Figure reprinted with permission from the Pharmaceutical Society of Japan.

Next, we focused attention on the expression of H₄R mRNA in synovial tissues, and serum matrix metalloproteinase-3 (MMP-3) concentration in RA. We found a significant negative correlation between H₄R expression in synovial tissues and serum MMP-3 concentration, but no correlation between MMP-3 and H₁R or H₂R (Yamaura et al., 2011). These observations suggest that H₄R is a potential target of novel pharmacotherapeutic agents for RA, and H₄R functional analysis may be useful in developing such treatments. In addition, we detected H₄R expression in human cartilage chondrocytes and in a murine chondrocytic cell line, ATDC5 (Yamaura et al., in press). Further work is needed to determine the expression mechanism and function of H₄R on chondrocytes.

3. Expression of H₄R in human skin

Following detection of H₄R expression in synovial tissue, we also analyzed H₄R expression in human epidermal tissue (Yamaura et al., 2009) and fibroblast cell cultures (Ikawa et al., 2008). Our immunoassays revealed that H₄R is expressed in both human epidermal tissues and dermal fibroblasts.

Keratinocytes are the major constituent of the epidermis. We found that immunohistochemical staining showed strong H₄R expression in keratin (K)10-positive differentiated keratinocytes in the prickle cell and granular layers of the epidermis (Fig. 5a). By contrast, H₄R was weakly expressed in K14-positive proliferating keratinocytes in the basal layer (Fig. 5b).




Fig. 3. Co-expression of H₄R protein with fibroblast-specific marker proteins. (a) Mouse anti-PH followed by Cy2-conjugated anti-mouse (green); (b) rabbit anti-H₄R followed by Cy3-conjugated anti-rabbit (red); (c) superposition of a on b; (d) Nomarski phase contrast microscopy image (NPCMI); (e) rabbit anti-H₄R followed by Cy2-conjugated anti-rabbit (green); (f) mouse anti-CD55 followed by Cy3-conjugated anti-mouse (red); (g) superposition of e on f; (h) NPCMI. Scale bar: 50 μ m. Figure reproduced with permission from the Pharmaceutical Society of Japan.




Fig. 4. Co-expression of H₄R protein with macrophage-specific marker proteins. (a) Mouse anti-CD68 followed by Cy2-conjugated anti-mouse (green); (b) rabbit anti-H₄R followed by Cy3-conjugated anti-rabbit (red); (c) superposition of a on b; (d) NPCMI; (e) mouse anti-CD163 followed by Cy2-conjugated anti-mouse (green); (f) rabbit anti-H₄R followed by Cy3-conjugated anti-rabbit (red); (g) superposition of e on f; (h) NPCMI. Scale bar: 50 μ m. Figure reproduced with permission from the Pharmaceutical Society of Japan.




Fig. 5. H₄R expression in human epidermal tissues. Double immunofluorescence staining of human epidermal tissues with anti-human H₄R antibody followed by Cy2-conjugated anti-rabbit secondary antibody (red), and anti-K10 (a) or anti-K14 (b) antibody followed by Cy2-conjugated anti-mouse secondary antibody (green). (c) For the negative control, tissues were only exposed to the secondary antibody. Figure reprinted with permission from the Japanese Society of Toxicology.

Keratinization is the result of keratinocytes dividing in the basal lamina and moving to the upper layer as they mature. K10 is expressed in keratinocytes in the early stages following differentiation, whereas K14 is expressed in undifferentiated keratinocytes. Accordingly, our results suggest that keratinocytes increase expression of H₄R following differentiation; however, further work is necessary to determine the expression mechanism and the physiological role of the receptors.

Increased H₄R expression has been reported in CD4⁺ T cells of patients with atopic dermatitis (Gutzmer et al., 2009), and skin mast cells have been shown to express H₄R (Lippert et al., 2004). These findings suggest that dermal cells may play an important role, via H₄R, in skin disorders. Dermal fibroblasts are a major component of the dermis. When the skin is damaged, they perform important roles including production of extracellular matrix molecules such as collagens. We have demonstrated the expression of H₄R in human dermal fibroblast cells using immunohistochemical staining (Fig. 6). Furthermore, Western blot analysis showed enhancement of the expression level of H₄R in dermal fibroblasts by stimulation with dexamethasone (Fig. 7).




Fig. 6. Expression of H₄R on human dermal fibroblasts. Double immunofluorescence staining of dermal fibroblast cells treated with anti-human PH antibody followed by Cy2-conjugated anti-mouse secondary antibody (green), and anti-human H₄R antibody followed by Cy3-conjugated anti-rabbit secondary antibody (red). Magnification x 400. Figure reprinted with permission from the Japanese Society of Toxicology.

Ohnishi et al. (2008) reported that the levels of leukotriene B₄ receptor 1, which belongs to the GPCR family, were up-regulated by dexamethasone. This suggests that H₄R, which is part of the same family, may be up-regulated by dexamethasone too. This up-regulation of H₄R expression might be associated with itching that occurs as a rebound phenomenon after withdrawal of high-dose topical glucocorticoids. We confirmed that significant enhancement of pruritus occurred after chronic topical application of dexamethasone in

mice (Yamaura et al., 2011). However, further study is needed to investigate the relationship between the expression of H₄R in skin and pruritus, which are both enhanced by dexamethasone.




Fig. 7. Effects of dexamethasone on H₄R protein expression in human dermal fibroblast cell cultures. Expression of H₄R was analyzed by Western blot analysis. Protein levels of H₄R were normalized to the corresponding β-actin protein levels. The mean value of the non-treatment group was set to 1.0. Data are expressed as means (n=2-3). Figure reprinted with permission from the Japanese Society of Toxicology.

4. Effect of H₄R antagonists on pruritus model

Chronic pruritus, associated with chronic conditions such as skin, liver, and kidney diseases and metabolic disorders, is a major diagnostic and therapeutic problem and can have a profound impact on the quality of life of patients. Recent studies have raised the possibility that H₄R, in addition to the H₁R, may contribute to histamine-mediated pruritic responses in mice (Bell et al., 2004). Both specific H₄R agonists and histamine were shown to induce pruritic responses which could be blocked by pretreatment with H₄R antagonists; the response was also found to be markedly attenuated in H₄R-deficient mice. We thus examined the effectiveness of selective H₄R antagonists as antipruritic drugs by their effect on histamine H₁R antagonist-resistant acute pruritus induced by substance P in mice.

We investigated the effect of the H₁R antagonist fexofenadine and the H₄R antagonist JNJ7777120 on histamine-induced acute pruritus (Yamaura et al., 2009). Oral administration of fexofenadine caused a slight reduction in scratching, whereas JNJ7777120 showed a significant reduction (Fig. 8). We then examined the effect of these antagonists in substance P-mediated acute pruritus. Fexofenadine showed no reduction in substance P-induced scratching. By contrast, JNJ7777120 at 10 and 30 mg/kg doses reduced substance P-induced scratching in a dose-dependent manner (Fig. 9). Although JNJ7777120 crosses the blood-brain barrier, it does not cause sedation in rodents (Dunford et al., 2007); hence its antipruritic action is not a secondary effect of sedation. The results suggest that H₁R has only limited involvement in histamine-induced pruritus. By contrast, the significant effect of JNJ7777120 suggests that H₄R

has a much greater role. Substance P-induced pruritus is resistant to H₁R antagonists (Togashi et al., 2002); given its occurrence in mast cell-deficient mice (Andoh et al., 2001), histamine from mast cells is unlikely to be involved. The role of H₁R is also thought to be minor, with fexofenadine having no observable effect. However, the suppression of the pruritic response by JNJ7777120 suggests that histamine may act via H₄R rather than H₁R.




Fig. 8. Effect of H₄R on scratching behavior induced by histamine. Histamine (300 nmol) was injected intradermally into shaved skin on the back of each mouse. Immediately after the injection of pruritogen, scratching events were counted for 30 min using the MicroAct apparatus (Neuroscience Inc., Tokyo, Japan). Fexofenadine (a) or JNJ7777120 (b) was administered orally 20 min before the injection of pruritogen. Values represent the mean \pm SEM of four mice. * $p < 0.05$ vs. control (Dunnett's multiple comparisons). Figure reprinted with permission from the Japanese Society of Toxicology.




Fig. 9. Effect of H₄R on scratching behavior induced by substance P. Substance P (100 nmol) was injected intradermally into shaved skin on the back of each mouse. Immediately after the injection of pruritogen, scratching events were counted for 30 min using MicroAct. Fexofenadine (a) or JNJ7777120 (b) was administered orally 20 min before the injection of pruritogen. Values represent the mean ± SEM of four mice. * $p < 0.05$ vs. control (Dunnett's multiple comparisons). Figure reprinted with permission from the Japanese Society of Toxicology.

Further, we created a chronic itch model in which repeated application of 2,4,6-trinitrochlorobenzene to the back skin of HR-1 mice was seen to elicit frequent scratching behavior at 24 h after challenge. JNJ7777120 at 10 and 30 mg/kg doses reduced this scratching behavior, whereas fexofenadine had no such effect (Suwa et al., 2011). These results suggest that H₄R antagonists may be useful for treatment of H₁R antagonist-resistant chronic pruritus such as atopic dermatitis.

5. Conclusion

We have demonstrated the expression of H₄R in human synovial cells from patients with RA and found a significant negative correlation between H₄R expression in RA synovial tissues and serum MMP-3 concentration. Furthermore, we have shown expression of H₄R in human skin and demonstrated that an H₄R antagonist ameliorates both H₁R antagonist-resistant acute and chronic pruritus. Taken together, these results suggest that histamine H₄R could be a new drug target for therapeutic use in RA or pruritic skin disorders such as atopic dermatitis.

6. Acknowledgment


We thank Johnson & Johnson Pharmaceutical Research & Development, L.L.C., for generously providing JNJ7777120. This research was supported in part by Grants-in-Aid for Scientific Research from the Japan Society for the Promotion of Science.

7. References

- Andoh T., Katsube N., Maruyama M., & Kuraishi Y. (2001). Involvement of leukotriene B(4) in substance P-induced itch-associated response in mice. *J. Invest. Dermatol.* 117: 1621-1626, ISSN 0022-202X.
- Bell JK., McQueen DS., & Rees JL. (2004). Involvement of histamine H₄ and H₁ receptors in scratching induced by histamine receptor agonists in Balb C mice. *Br. J. Pharmacol.* 142: 374-380, ISSN 0007-1188.
- de Esch IJ., Thurmond RL., Jongejan A., & Leurs R. (2005). The histamine H₄ receptor as a new therapeutic target for inflammation. *Trends Pharmacol. Sci.* 26: 462-469, ISSN 0165-6147.
- Dunford PJ., Williams KN., Desai PJ., Karlsson L., McQueen D., & Thurmond RL. (2007). Histamine H₄ receptor antagonists are superior to traditional antihistamines in the attenuation of experimental pruritus. *J. Allergy Clin. Immunol.* 119: 176-183, ISSN 0091-6749.
- Fonseca JE., Cortez-Dias N., Francisco A. Sobral M., Canhão H., Resende C., Castelão W., Macieira C., Sequeira G., Saraiva F., da Silva JA., Carmo-Fonseca M., & Viana Queiroz M. (2005). Inflammatory cell infiltrate and RANKL/OPG expression in rheumatoid synovium: comparison with other inflammatory arthropathies and correlation with outcome. *Clin. Exp. Rheumatol.* 23: 185-192, ISSN 0392-856X.
- Frewin DB., Cleland LG., Jonsson JR., & Robertson PW. (1986). Histamine levels in human synovial fluid. *J. Rheumatol.* 13: 13-14, ISSN 0315-162X.
- Grzybowska-Kowalczyk A., Wojtecka-Lukasik E., Maslinska D., Gujski M., & Maslinski S. (2007). Distribution pattern of histamine H₄ receptor in human synovial tissue from patients with rheumatoid arthritis. *Inflamm. Res.* 56(Suppl 1): S59-S60, ISSN 1023-3830.
- Gutzmer R., Mommert S., Gschwandtner M., Zwingmann K., Stark H., & Werfel T. (2009). The histamine H₄ receptor is functionally expressed on T(H)₂ cells. *J. Allergy Clin. Immunol.* 123: 619-625, ISSN 0091-6749.

- Ikawa Y., Suzuki M., Shiono S., Ohki E., Moriya H., Negishi E., & Ueno K. (2005). Histamine H₄ receptor expression in human synovial cells obtained from patients suffering from rheumatoid arthritis. *Biol. Pharm. Bull.* 28: 2016-2018, ISSN 0918-6158.
- Ikawa Y., Shiba K., Ohki E., Mutoh N., Suzuki M., Sato H., & Ueno K. (2008). Comparative study of histamine H₄ receptor expression in human dermal fibroblasts. *J. Toxicol. Sci.* 33: 503-508, ISSN 0388-1350.
- Lippert U., Artuc M., Grützkau A., Babina M., Guhl S., Haase I., Blaschke V., Zachmann K., Knosalla M., Middel P., Krüger-Krasagakis S., & Henz BM. (2004). Human skin mast cells express H₂ and H₄, but not H₃ receptors. *J. Invest. Dermatol.* 123: 116-123, ISSN 0022-202X.
- Nagata Y. (1991). The role of histamine in human synovial fibroblasts. *Med. J. Kinki Univ.* 16: 117-126 (in Japanese), ISSN 0385-8367.
- Nguyen T., Shapiro DA., George SR., Setola V., Lee DK., Cheng R., Rauser L., Lee SP., Lynch KR., Roth BL., & O'Dowd BF. (2001). Discovery of a novel member of the histamine receptor family. *Mol. Pharmacol.* 59:427-433, ISSN 0026-895X.
- Oda T., Morikawa N., Saito Y., Masuho Y., & Matsumoto S. (2000). Molecular cloning and characterization of a novel type of histamine receptor preferentially expressed in leukocytes. *J. Biol. Chem.* 275: 36781-36786, ISSN 0021-9258.
- Ohnishi H., Miyahara N., Dakhama A., Takeda K., Mathis S., Haribabu B., & Gelfand EW. (2008). Corticosteroids enhance CD8+ T cell-mediated airway hyperresponsiveness and allergic inflammation by upregulating leukotriene B₄ receptor 1. *J. Allergy Clin. Immunol.* 121: 864-871, ISSN 0091-6749.
- Ohki E., Suzuki M., Aoe T., Ikawa Y., Negishi E., & Ueno K. (2007). Expression of histamine H₄ receptor in synovial cells from rheumatoid arthritic patients. *Biol. Pharm. Bull.* 30: 2217-2220, ISSN 0918-6158.
- Suwa E., Yamaura K., Oda M., Namiki T., & Ueno K. (2011). Histamine H₄ receptor antagonist reduces dermal inflammation and pruritus in a hapten-induced experimental model. *Eur. J. Pharmacol.* 667: 383-388, ISSN 0014-2999.
- Sweeney SE., & Firestein GS. (2004). Rheumatoid arthritis: regulation of synovial inflammation. *Int. J. Biochem. Cell Biol.* 36: 372-378, ISSN 1357-2725.
- Tanaka M. (2005). Autoimmune abnormality. *Nippon Rinsho* 63(Suppl 1): 80-83 (in Japanese), ISSN 0047-1852.
- Togashi Y., Umeuchi H., Okano K., Ando N., Yoshizawa Y., Honda T., Kawamura K., Endoh T., Utsumi J., Kamei J., Tanaka T., & Nagase H. (2002). Antipruritic activity of the kappa-opioid receptor agonist, TRK-820. *Eur. J. Pharmacol.* 435: 259-264, ISSN 0014-2999.
- Thurmond RL., Gelfand EW., & Dunford PJ. (2008). The role of histamine H₁ and H₄ receptors in allergic inflammation: the search for new antihistamines. *Nat. Rev. Drug. Discov.* 7: 41-53, ISSN 1474-1776.
- Yamaura K., Oda M., Suwa E., Suzuki M., Sato H., & Ueno K. (2009). Expression of histamine H₄ receptor in human epidermal tissues and attenuation of experimental pruritus using H₄ receptor antagonist. *J. Toxicol. Sci.* 34: 427-431, ISSN 0388-1350.

- Yamaura K., Doi R., Suwa E., & Ueno K. (2011). A novel animal model of pruritus induced by successive application of glucocorticoid to mouse skin. *J. Toxicol. Sci.* 36: 395-401, ISSN 0388-1350.
- Yamaura K., Oda M., Suzuki M., & Ueno K. (2011). Lower expression of histamine H(4) receptor in synovial tissues from patients with rheumatoid arthritis compared to those with osteoarthritis. *Rheumatol Int.* Sep 1 (Epub ahead of print), ISSN: 0172-8172
- Yamaura K., Akiyama S., & Ueno K. (2011). Increased expression of the histamine H4 receptor subtype in hypertrophic differentiation of chondrogenic ATDC5 cells. *J Cell Biochem.* In press, ISSN 1097-4644.



Allergic Diseases - Highlights in the Clinic, Mechanisms and Treatment

Edited by Prof. Celso Pereira

ISBN 978-953-51-0227-4

Hard cover, 554 pages

Publisher InTech

Published online 14, March, 2012

Published in print edition March, 2012

The present Edition "Allergic diseases - highlights in the clinic, mechanisms and treatment" aims to present some recent aspects related to one of the most prevalent daily clinical expression disease. The effort of a group of outstanding experts from many countries reflects a set of scientific studies very promising for a better clinical care and also to the treatment and control of the allergy. This book provides a valuable reference text in several topics of the clinical allergy and basic issues related to the immune system response. The inflammatory reaction understanding in allergic disease is clearly evidenced, as well as new strategies for further researches.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Katsunori Yamaura, Masahiko Suzuki, Takao Namiki and Koichi Ueno (2012). Expression of the Histamine H4 Receptor in Human Tissue, Allergic Diseases - Highlights in the Clinic, Mechanisms and Treatment, Prof. Celso Pereira (Ed.), ISBN: 978-953-51-0227-4, InTech, Available from:
<http://www.intechopen.com/books/allergic-diseases-highlights-in-the-clinic-mechanisms-and-treatment/expression-of-the-histamine-h4-receptor-in-human-tissue>

INTECH
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2012 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.