

Molecular Epidemiology of Parasitic Diseases: The Chagas Disease Model

Juan David Ramírez and Felipe Guhl
*Centro de Investigaciones en Microbiología y Parasitología Tropical (CIMPAT),
Universidad de los Andes, Bogotá,
Colombia*

1. Introduction

Parasitic diseases represent one of the most important issues in public health. More than one billion people worldwide are infected by parasites causing different disease scenarios. Parasitic diseases are closely related to geographic, social and economic factors driving the prevalence and incidence of these pathologies (WHO, 2010). These represent a broad group of eukaryotic organisms that may cause severe diseases in animal and human populations. Parasites are the causative agents of pathologies such as Malaria. In 2008, there were 247 million cases of Malaria and nearly one million deaths from the disease, mostly among children living in Africa. In Africa, a child dies of Malaria every 45 seconds; the disease accounts for 20% of all childhood deaths. Leishmaniasis threatens approximately 350 million men, women and children in 88 countries around the world. As many as 12 million people are believed to be currently infected by this disease, with approximately 1–2 million estimated new cases occurring every year. Additionally, an estimated of 10 million people are infected worldwide by Chagas disease (American trypanosomiasis), mostly in Latin America, where Chagas disease is endemic. More than 25 million people are at risk of acquiring this disease. It is estimated that in 2008, Chagas disease killed more than 10,000 people. Schistosomiasis is a chronic, parasitic disease caused by blood flukes (trematode worms) of the genus *Schistosoma*. More than 207 million people are infected with these organisms worldwide, with an estimated 700 million people at risk in 74 endemic countries. Lymphatic filariasis affects more than 1.3 million people in 81 countries. Approximately 65% of those infected live in Southeast Asia, 30% in Africa and the remainder in other tropical areas. Lymphatic filariasis afflicts over 25 million men with genital disease and over 15 million people with lymphoedema. Because the prevalence and intensity of infection are linked to poverty, elimination can contribute to achieving the United Nations Millennium Development Goals. Human African Trypanosomiasis (HAT) affects mostly poor populations living in remote rural areas of Africa. If untreated, it is usually fatal. Travellers also risk becoming infected if they venture through regions where the insect vector (tse tse flies) is common. Generally, the disease is not found in urban areas, although some cases have been reported in suburban areas of Kinshasa, the capital of the Democratic Republic of Congo, and Luanda, the capital city of Angola. In 2004, the number of new reported cases fell to 17,616, which the WHO considered to be due to increased control, estimating the cumulative rate to be between 50,000 and 70,000 cases (WHO, 2010). These trends show the

importance of developing strategies to mitigate the prevalence of these parasitic diseases, and molecular epidemiology arises as a potential tool to understand disease dynamics.

Molecular epidemiology is considered a powerful tool for understanding the genetic variation and evolution of pathogens. The use of the technologies based on molecular biology techniques has allowed the scientific community to reveal disease determinants and the genetic structure of parasites that provoke diseases and cause millions of deaths each year. In recent years, new studies have been conducted with the purpose of elucidating the genetic structure of the etiological agents of these pathologies with the aim of designing strategies that could help to mitigate the associated diseases in human and animal populations. Molecular epidemiology and population genetics have shown to be powerful strategies to understand the genetic structure of parasites, with special emphasis on understanding disease and transmission dynamics. The objective of this chapter is to illustrate for the reader the paramount importance of molecular epidemiology in parasitic diseases, showing some clear examples of parasite disease and transmission dynamics. The focus will be on how molecular epidemiology can be a helpful tool to mitigate disease transmission, prevalence and incidence and can be used as a reliable tool for disease surveillance as well as the need to create synergy between molecular epidemiology and public health programmes to reduce the prevalence of parasitic diseases.

2. Importance and relevance of molecular epidemiology

Molecular epidemiology may be defined as a tool focused on the contribution of potential genetic factors identified by molecular techniques to the aetiology, distribution and prevention of disease across populations (Kilbourne, 1973). It is a field of study that has recently emerged from the integration of molecular biology, epidemiology, biochemistry and public health systems (Figure 1). This approach has been useful in attempting to determine the pathogenesis of certain diseases as well as the genetic variation and genetic structure of pathogens.




Fig. 1. Molecular epidemiology is considered an interdisciplinary science that is a composite of different sciences.

Molecular epidemiology has recently gained paramount importance in the fields of human genetics and in molecular virology. One clear example of this is the use of molecular epidemiology to track viruses that generate severe acute respiratory syndrome (SARS). This allowed researchers to develop strategies with the purpose of tracking the transmissibility and dispersal of this virus, observing that the positive selection pressure associated with human hosts resulted in the emergence of lineages of the virus that became readily transmissible between humans, causing the epidemic outbreak of 2002-2003 (Zhao, 2007). The use of molecular markers also permitted the establishment of prevention and control strategies to mitigate the transmission of these genotypes, thus avoiding increased disease prevalence. Thus, this example shows how traceable pathogens can be and clearly demonstrates the great utility of molecular biology on the basis of molecular epidemiology in obtaining a deeper understanding regarding parasitic diseases.

Molecular epidemiology has been established as a promising science for studying the contribution of potential genetic markers and environmental risk factors of parasitic diseases representing a close synergy between molecular biology and epidemiology. Some of the main objectives of molecular epidemiology focused on the study of parasitic diseases are as follows:

- To enhance our understanding of the pathogenesis of parasitic diseases: Some authors have used the *Toxoplasma gondii* model to describe relationships between nucleoside triphosphate hydrolase (NTPase) isoforms and *Toxoplasma* strain virulence in human toxoplasmosis, reporting that different isoforms are involved in clinical forms of this parasitic disease (Johnson et al., 2003).
- To define genetic susceptibility with genetic markers: The use of human pedigrees to observe patterns of susceptibility to visceral leishmaniasis in Brazil has allowed researchers to develop action plans to mitigate this tropical disease in endemic areas (Jamieson et al., 2007).
- To allow evaluation of subclinical or early disease markers: Prognostic markers are one of the milestone deliverables in molecular epidemiology. In the case of *T. gondii*, a quantitative real-time PCR assay for amniotic fluid has been developed to provide a prognostic marker of foetal infection in pregnant women (Romand et al., 2004)
- To provide new standards for descriptive epidemiology: Some of the problems involved in descriptive epidemiology show how difficult it is to track some kinds of diseases. In particular, the parasites that cause gastrointestinal syndromes fall into this category, such as *Entamoeba histolytica*, *Taenia solium*, *Ascaris lumbricoides*, *Giardia intestinalis*, *Enterobius vermicularis* and others. Microscopic identification becomes tedious and, in some case confusing due to similar morphologies among some parasites. Molecular detection based on PCR assays has provided the field of descriptive epidemiology with a more reliable way to analyse data in population descriptive studies (Singh et al., 2009; Pecson et al., 2006).
- To improve precision in analytical epidemiology: While descriptive epidemiology provides the what, who, when and where; analytical epidemiology attempts to provide the why and how. Few examples are listed related to the detection of emergent genotypes in disease surveillance. A good example is that of HAT, which is caused by two sympatric subspecies (*Trypanosoma brucei rhodesiense* and *T. b. gambiense*); each subspecies is involved in disease severity causing a large number of annual deaths in Africa (Morrison et al., 2011).

As has been shown thus far, molecular epidemiology as applied to the study of parasites is considered an important and relevant tool to investigate these organisms. The important point to focus is on the correct use of the information obtained. Molecular methods are currently available and becoming cheaper every day, and the emergence of new accurate and feasible molecular methods shows promising results for the molecular epidemiology field; however, in molecular epidemiology, the clinical question must be always highlighted to obtain the most reliable results. Thus, the income data become the critical point in the study of parasitic diseases and the basis for obtaining a good outcome that can be translated to meet the main aims of molecular epidemiology studies (Figure 2). This chapter will discuss the molecular methods available to develop molecular epidemiology studies focused on parasites, with some clear examples of how useful and necessary molecular epidemiology is for understanding disease outcomes, transmission dynamics and the current genetic structure of parasitic diseases.




Fig. 2. Flow of information based on the accurate and reliable use of molecular epidemiology focused on the study of parasitic diseases.

3. Molecular biology tools applied in the analysis of parasitic diseases

Molecular biology has made important contributions in the last ten years with respect to understanding the genetics of parasites causing human illness. A broad description of the markers used in molecular epidemiology and their features is presented in Table 1. The first techniques used to track pathogens and disease dispersal were based on biochemical markers, with Multilocus Enzyme Electrophoresis (MLEE) being broadly used to study parasites such as *T. cruzi*, *Leishmania* spp, *E. histolytica* and *T. brucei* (Miles et al., 1977; Pinto et al., 2005; Mathews et al., 1983; Nijokou et al., 2004). This technique is based on differences between loci; according to the obtained banding pattern, it is possible to distinguish among lineages. In the case of *Leishmania*, this technique was used to differentiate species of the genus involved in visceral and cutaneous leishmaniasis (Bañuls et al., 2000; Bañuls et al., 2002; Zhang et al., 2006). A drawback of this technique is the need to culture large quantities of parasites, which are quite difficult to obtain in most parasitic diseases. Subsequently,

molecular biology techniques based on the use of DNA were developed, among which RAPDs (Random Amplified Polymorphic DNA), AFLPs (Amplified Fragment Length Polymorphism), PFGE (Pulse Field Gel Electrophoresis) and RFLPs (Restriction Fragment Length Polymorphism) were the most used techniques within the scientific community. In this sense, analysis of ribosomal markers using RAPDs was of paramount importance in attempting to develop assays to distinguish among morphologically similar amoebas. *E. histolytica* and *E. dispar*, which represent pathogenic and non-pathogenic amoeba species, respectively, have been suggested by many authors to have evolved identically up to the point when a mutation generated a cryptic speciation pattern (Clark et al., 2006). The use of RAPD techniques has also been an important aid in understanding genetic variation within *E. histolytica* isolated from different hosts (Gomes et al., 2000; Prakash et al., 2000). AFLPs and RFLPs are the most recent techniques to be employed to track pathogen dispersal and elucidate their genetic diversity. Fingerprinting based on AFLPs has been useful in differentiating subspecies of *T. brucei*, which is the aetiological agent of sleeping sickness, a pathology that affects more than 8 million people in Africa. These techniques have shown great reproducibility in distinguishing *T. b. rhodesiense* and *T. b. gambiense*, two sympatric species that generate different symptomatologies, demonstrating how molecular epidemiology can assist in understanding disease outcomes in certain pathologies, thus aiding in developing proper treatment and management measures (Agbo et al., 2002; Masiga et al., 2006). RFLPs have been applied to parasites such as *T. gondii* and *Trichinella spiralis*, which are two species for which pigs play an important role in transmission dynamics. *T. gondii* is considered to display a clonal population, but its isolates have been divided into three types that are geographically clustered and, in some cases, are involved in disease outcomes (Wang et al., 1995; Su et al., 2002; Fuentes et al., 2001). These molecular markers have been shown to be important in discriminating species as well as evaluating genetic variability among isolates from the same species. The great advantage of these molecular markers is that they can be used to show the pattern of variation across the whole genome of a pathogen, rather than just a specific region, as will be shown later.

Feature	MLEE ¹	RAPD ² and AFLP ³	RFLP ⁴ and PFGE ⁵	Microarrays	MLMT ⁶	MLST ⁷	qPCR ⁸	Genome Sequencing
Culturing	Yes	No	No	No	No	No	No	No
Analysis of distinct loci	Yes	No	No	Yes	Yes	Yes	Yes	No
Cost	Low	Low	Low	High	Medium	Medium	Medium	High
Labor	High	Low	Low	High	Low	Low	Low	Medium
Informative	Low	Low	Low	High	High	High	High	High
Portability	Low	Medium	Medium	Low	High	High	Medium	Low

¹ Multilocus Enzyme Electrophoresis, ² Random Amplified Polymorphic DNA, ³ Amplified Fragment Length Polymorphisms, ⁴ Restriction Fragment Length Polymorphisms, ⁵ Pulse Field Gel Electrophoresis, ⁶ Multilocus Microsatellite Typing, ⁷ Multilocus Sequence Typing, ⁸ Quantitative Real Time PCR

Table 1. Molecular markers used in molecular epidemiology studies.

The development of Polymerase Chain Reaction (PCR) was of great importance and represents an incredible advance in molecular biology. Since 1990, PCR has been broadly used to study parasites. Modifications of PCR, such as PCR-RFLP, Nested PCR, RT-PCR, AP-PCR (Allele Polymorphic Polymerase Chain Reaction), SHELTA-PCR (Solution Hybridization Enzyme-Linked Assay Polymerase Chain Reaction) and qPCR (Quantitative Real Time Polymerase Chain Reaction), have been applied to study parasites. In these tests, the only thing that a researcher requires is a few aliquots of DNA, which simplifies the analysis. The use of PCR-RFLP is widely reported in discriminating *Leishmania* species. In these studies, the use of Heat Shock proteins with different molecular weights allows discrimination based on PCR amplification of genes that are subsequently digested with restriction endonucleases. According to the band patterns obtained in such analyses, it has been possible to distinguish between the subgenera *Viannia* and *Leishmania* as well as species from the different subgenera in some cases (Volpini et al., 2004; Montalvo et al., 2006). Discrimination among *Leishmania* species or their complexes is necessary in conducting studies on treatment resistance and clinical manifestations associated with leishmaniasis, in which some species cause cutaneous forms, and others cause visceral forms. One of the most important advances in the modification of PCR assays has been the development of quantitative Real-Time PCR (qPCR). This assay involves the quantification of DNA copies in each PCR cycle. The first applications of this technique have been for diagnostic purposes in cases where it has been possible to estimate parasitic loads in infected patients. In the case of *L. infantum*, the species involved in visceral leishmaniasis manifestations, assays have been developed to estimate parasitic loads in biopsies of infected patients and, thus, to estimate the efficacy of treatment using qPCR (Mary et al., 2004; Bretagne et al., 2001; Ranasinghe et al., 2008). Additionally, qPCR using SYBR green and Hybridisation probe chemistry has allowed the development of melting temperature (T_m) analysis according to a dissociation curve. This permits screening for genotypes or species according to the specific temperature of an amplicon and enables observation of single nucleotide polymorphisms, all in the same reaction. Thus, a qPCR Real Time protocol has been proposed to identify *Leishmania* species focused on spliced leader genes and minicircle kDNA regions; according to distinct temperatures, investigators were able to discriminate *Leishmania* species (Wortmann et al., 2005). This approach has also been applied to other parasites, such as *Giardia*, for which qPCR assays were developed to detect *G. lamblia* and to discriminate its genotypes in stool specimens (Guy et al., 2004). Additionally, it has been used to study schistosomiasis, an helminthic disease that affects populations in Africa, Asia and America, with qPCR assays being developed to discriminate species in water where the infective form (cercariae) lives and is transmitted to humans (Lier et al., 2006). qPCR has been shown to be a reliable, feasible, fast and accurate method in molecular epidemiology to discriminate species as well as to determine genotypes within species. This suggests the need to pursue studies involving this method, though in some cases, validation studies are required, and further research is needed. A problem involved in working on parasites is sensitivity because in some parasite diseases, the parasitic loads are quite low and even undetectable in some cases, such that concentration methods must be applied or it may be necessary to analyse the whole sample.

Other molecular markers include microarrays and Southern blot and northern blot techniques. The drawback of microarrays is that they are time consuming, expensive, and in some cases, they do not provide the desired information. In the last decade, the rise of

sequencing procedures has been an important addition to molecular epidemiology investigations. The ability to obtain DNA sequences has allowed researchers to unravel the genetic structure of parasites and to go further in the analyses that can be applied. Thus, molecular phylogenetics and population genetics have provided molecular epidemiology with certain, reliable tools for understanding the genetic structure of parasites. Molecular phylogenetics is the science focused on understanding the evolutionary relationships among groups of organisms based on molecular sequences, and these techniques have been broadly applied to parasites. Hence, molecular phylogenetics has allowed the reconstruction of phylogenetic trees based on maximum parsimony and/or maximum composite likelihood methods with the aim of understanding the evolutionary history of parasites and, in some cases, developing analysis involving loci. Based on the use of DNA sequences and phylogenetic reconstructions, new methods such as MLST (Multilocus Sequence Typing) have arisen. MLST has been broadly used in bacteria and yeast but has only recently been applied to parasites; the drawback of MLST in addressing protozoan parasites associated with working with clonal diploids instead of clonal haploids, such as bacteria. The genetic structure of clonal diploid pathogenic organisms is important in terms of elucidating the drivers of disease prevalence, installation and outcomes as well as the virulence factors and geographical distribution related to the disease. In recent years, the population structure of microorganisms such as *Plasmodium*, *Giardia*, *Entamoeba*, Trypanosomes (*T. brucei*, *T. congolense* and *T. cruzi*), *Candida*, *Leishmania* and *Toxoplasma* has gained paramount relevance due to the discussion of clonal propagation versus sexual recombination (De Meeus et al., 2006; Benett et al., 2010; Grigg and Suzuki, 2003; Morrison et al., 2009; Rougeron et al., 2010; Mzilahowa et al., 2007). There are three hypotheses that describe the genetic structure observed in clonal diploid organisms. In 1987, Harvey and Keymer suggested a panmictic population structure in which sexual recombination is frequent. In 1991, Tibayrenc et al. proposed the clonal theory of parasitic protozoa, suggesting that these organisms display a clonal propagation mode associated with infrequent sexual recombination events. Finally, in 1993, Maynard-Smith et al. proposed an epidemic population structure with a background level of frequent sexual recombination and with occasional clonal expansion of particular genotypes. These hypotheses have been tested using a large number of parasitic protozoa; however, the debate still continues.

The use of new methods for typing and elucidating the genetic variability of parasites like MLST has gained importance because it can be considered to be an improvement of MLEE. MLST makes use of different loci involved in parasite metabolism. Thus, MLST strategies have been developed in *Leishmania* for species identification using five metabolic enzymes that are able to discriminate species and genotypes within complexes (Zemanova et al., 2007). An MLST approach has also recently been described for discriminating among subtypes of *Blastocystis*, which is a protozoan parasite involved in bowel inflammation and acute diarrhoea in immunodeficient patients (Stensvold and Clark, 2011). MLST analyses are not only used to discriminate genotypes of species but also to detect recombination or likely genetic exchanges among parasite populations (diploid clonals). In the case of sexual parasites such as helminthes, these approaches are employed to discriminate among species or to detect genotypes. Another important development in molecular epidemiology has been the use of microsatellite markers in developing MLMT (Multilocus Microsatellite Typing) strategies. Microsatellite markers are defined as tandem repetitions of 1-6 base pair segments of DNA; they are neutral and co-dominant and are useful in developing

population genetics analyses. The variability of microsatellites is due to their higher rate of mutation compared to other neutral regions of DNA. The use of microsatellite markers is widely reported for purposes ranging from species identification to detection of recombination based on population genetics statistics. In *P. vivax*, polymorphic microsatellite markers have been amplified and analysed to unravel the genetic structure of this parasite and to understand its co-evolution with other *Plasmodium* species (Gomes et al., 2003; Imgwon et al., 2006). Population genetics tools present a limitation when working with clonal diploids related to the assumption of Hardy-Weinberg equilibrium and other statistics, such as *Fis* and *Fst*.

In recent years, DNA sequencing has become an important tool in understanding microorganisms, particularly those involved in human pathologies. Sequencing procedures have been improved, and pyrosequencing has become an important method to obtain more feasible and accurate DNA sequences. Pyrosequencing is a method of DNA sequencing based on the "sequencing by synthesis" principle. It differs from Sanger sequencing in that it relies on the detection of pyrophosphate release upon nucleotide incorporation, rather than chain termination with dideoxynucleotides. Genome sequencing methods developed in bacteria have also been applied to sequence whole genomes in parasites. The first parasite genome sequenced was that of *P. falciparum* (Gardner et al., 2002), followed by the genomes of other parasites, such as *Leishmania*, *T. brucei*, *T. cruzi* and *T. gondii* (Ivens et al., 2005; Elsayed et al., 2005; Bontell et al., 2009). New initiatives have been developed to sequence larger genomes, such as those from helminthes including *Ascaris*, *Taenia*, *Schistosoma* and *Echinococcus*. These approaches have allowed scientists to develop projects aimed at annotating parasite genomes for the purpose of detecting possible pharmaceutical markers to develop drugs against these microorganisms. Genome sequencing is becoming cheaper due to the advances made by Illumina, which will allow the scientific community to begin sequencing genomes instead of single genes. The possibility of obtaining this type of metadata permits the application of tools in bioinformatics, metabolomics, immunomics, vaccinomics, proteomics and other field with the purpose of transitioning into the OMICS era. The OMICS era is considered to be associated with the most advanced techniques for understanding the molecular epidemiology of parasitic diseases. These tools will be of paramount importance in developing new drugs against parasites as well as evaluating surveillance disease markers or prognostic disease markers to understand the relatedness between disease outcomes and parasite genetic variability, which is one of the main objectives in molecular epidemiology.

4. Comparative molecular epidemiology: The Chagas disease model

Chagas disease, which is caused by the parasite *T. cruzi*, is a complex zoonosis that is widely distributed throughout the American continent. The infection can be acquired through triatomine faeces, blood transfusion, oral and congenital transmission and laboratory accidents. Chagas disease represents an important public health problem, with estimates by the Pan American Health Organization in 2005 of at least 7.7 million people being infected with *T. cruzi* and another 110 million being at risk (WHO, 2007). Additionally, immigration of infected people from endemic countries is now making Chagas disease a relevant health issue in other regions, including Europe and the United States (Rassi et al., 2009). Chagas disease is comprised of two stages, with the acute phase occurring approximately one week

after the initial infection and approximately 30-40% of infected patients developing the chronic phase of the disease, in which cardiomyopathy is the most frequent and severe clinical manifestation (Rassi et al., 2009). Chagas has lately gained more importance due to recent reports of imported cases in Europe, the United States and Canada (Schmunis and Yadon, 2010).

Obtaining a full understanding of the aetiology and epidemiology of Chagas disease across its distribution has proved elusive and complex and remains the subject of intense investigation to the present day. The difficulty in completely defining the epidemiology of Chagas disease is attributable to several factors. First, Chagas disease is a zoonosis, and a variety of widely distributed mammals serve as reservoirs for *T. cruzi*. Moreover, all mammals are susceptible to *T. cruzi* infection. An additional factor that contributes to the complexity of Chagas disease as a zoonosis is the variety of vectors involved, as they are not simply represented by a range of related species or genera, as is the case for all other known insect vector-associated diseases. Triatomine bugs are a subfamily of insects, and across this relatively broad taxonomic range, there are members from all groups that can harbour *T. cruzi*. However, most transmission is attributable to three main genera: *Rhodnius*, *Panstrongylus*, and *Triatoma*, but this diversity still represents two different tribes of the subfamily (Rhodniini and Triatomini). Furthermore, the insects vary in more than their ancestry, being associated with a diverse range of vertebrate hosts and ecological associations. The third factor that complicates the epidemiology of Chagas disease and accounts for variation in the clinical manifestation of the disease is the subspecific diversity of *T. cruzi* itself. Much work has been conducted over the past 40 years to elucidate the variation of *T. cruzi* across its geographical distribution and associations with hosts and vector species.

The *T. cruzi* parasite comprises a heterogeneous population that displays clonal propagation due to its different cycles of transmission and the possibility of recombination exchanges, which can be found in nature and have previously been reported *in vitro* (Gaunt et al., 2003, Sturm et al., 2003; Westenberger et al., 2002). *T. cruzi* is genetically diverse and is classified into a series of strains or subtypes. This genetic diversity was initially discovered using a panel of isoenzyme markers to investigate differences between parasites involved in domestic and sylvatic cycles in Bahia state in Brazil (Miles et al., 1977). This study represented a breakthrough, revealing that in Bahia, there were substantial genetic differences between the parasites involved in sympatric sylvatic and domestic transmission cycles. These variants were designated zymodemes I and II (ZI and ZII). Soon thereafter, it was revealed that the widespread strain associated with the sylvatic cycle in Brazil (ZI) was the predominant cause of human disease in Venezuela (Miles et al., 1981). These groundbreaking findings paved the way for investigating the aetiology of Chagas disease, allowing host-vector-parasite associations and comparative geographical distributions to be explored, as reviewed by Miles et al. (2009). Subsequently, four additional zymodemes were described from Brazil, Paraguay, and Bolivia. In the following two decades, various authors proceeded to characterise strains of *T. cruzi*, applying other molecular methods as they became available. As a result, further diversity was discovered within the original zymodemes. However, the designations of subtypes in the literature began to become confusing. Recently, a new nomenclature for *T. cruzi* has been adopted that includes six Discrete Taxonomic Units (DTUs) designated *T. cruzi* I (TcI), *T. cruzi* II (TcII), *T. cruzi* III

(TcIII), *T. cruzi* IV (TcIV), *T. cruzi* V (TcV) and *T. cruzi* VI (TcVI) based on different molecular markers and biological features (Zingales et al., 2009). These DTUs are broadly distributed in the American continent in diverse ecotopes (Figure 3). Discrimination of the six DTUs has become an important issue in the molecular epidemiology of *T. cruzi*. There are many reports showing algorithms for the molecular characterisation of these DTUs by performing RAPD, PCR-RFLP, qPCR, MLST, MLMT and DNA sequencing analyses, but to date, there is no consensus protocol for strain typing (Lewis et al., 2009; Rozas et al., 2007; Ramírez et al., 2010; Duffy et al., 2009; Yeo et al., 2011; Llewellyn et al., 2009). One of the most recent and reliable algorithms for *T. cruzi* typing was reported by Ramírez et al., 2010 and has been applied on biological samples (Figure 4).



Fig. 3. Geographical distribution of *T. cruzi* DTUs in the American continent based on Patterson and Guhl, 2010.




Fig. 4. Algorithm for typing *T. cruzi* DTU's based on five molecular markers and also used to genotype biological samples (Ramírez et al., 2010).

In a sense, the findings of Miles et al. (1977) were the tip of the iceberg in unravelling the genetic structure of *T. cruzi*, but at the same time, they hit the nail on the head. The observed predominance of TcI in the human populations in Venezuela and Colombia and TcII, TcV, and TcVI mostly infecting human hosts in Brazil was to prove representative (i.e., it has since been demonstrated that TcI predominates in countries north of the Amazon and TcII-TcVI in Southern Cone countries, but this distribution is not absolute). This is particularly illuminating given that there are distinct clinical differences between patients presenting with Chagas disease in these two geographical regions. Strains appear to differ in terms of both their pathogenicity and response to treatment. TcI and TcII-VI are all associated with cardiac lesions in human infections, but it appears that only TcII, TcV, and TcIV are also associated with digestive tract lesions (Prata, 2001), despite the recent report of digestive tract lesions in Colombia caused by TcI (Mantilla et al., 2010). TcI is generally considered to be less pathogenic with lower parasitemia (Burgos et al., 2007) and more chronic cases being asymptomatic compared to Chagas caused by TcII, TcV, and TcVI in Argentina, Brazil, Chile, Paraguay, and Uruguay. TcI is almost the only form found in human infections north of the Amazon region. Moreover, there is an observed general partitioning of former TcII subtypes between sylvatic and domestic transmission cycles; the human disease is associated with TcII, while TcV is rarely associated with sylvatic hosts (Yeo et al., 2005), and TcIII and TcIV are predominantly sylvatic.

TcI has remained a constant grouping in the nomenclature since it was first described. However, recent studies based on mini-exon gene (SL-IR) sequences have shown

polymorphism on this region, with four genotypes being reported within TcI. These genotypes have also been reported in various regions of South America, where five TcI genotypes have been detected (Figure 5) (Cura et al., 2010; Herrera et al., 2007; Guhl and Ramírez, 2011). Different molecular markers, including a 48 set of microsatellite loci, have also shown the great diversity in TcI (Guhl and Ramírez, 2011; Llewellyn et al., 2009; Spotorno et al., 2008; Ramírez et al., 2011). Primers designed based on TcI sequences confirmed the existence of three genotypes (Ia, Ib and Id) and a new genotype found in the Southern cone countries designated Tcle (Cura et al., 2010; Falla et al., 2009).




Fig. 5. Geographical distribution of *T. cruzi* I genotypes based on SL-IR region (Guhl and Ramírez, 2011)

Genetic variability has been clearly demonstrated in *T. cruzi*, with reports of homogeneous (TcII) and heterogeneous groups considered to be hybrids due to recombination events (TcIII-TcVI) (Gaunt et al., 2003, Sturm et al., 2003; Westenberger et al., 2002). It has been shown that TcIII and TcIV are likely to be a product of recombination of TcI and TcII and

TcIV-TcVI to be a product of recombination of TcII and TcIII/TcIV (Brisse et al., 2003), although this last statement is still controversial. The recent advances in sequencing procedures have allowed three complete *T. cruzi* genomes to be obtained. The first strain fully sequenced was CL Brener (TcVI), which showed a large number of repetitive elements along the core genome (Elsayed et al., 2005). Likewise, the recent sequencing of the Esmeraldo (TcII) and Sylvio X10 (TcI) genomes has shown the relationship between repetitive elements and mucin-like proteins, which are closely associated with parasite cell invasion and survival, showing this area of inquiry to be quite promising with respect to obtaining more information about the genetic structure of *T. cruzi* DTUs (Franzen et al., 2011; Andersson, 2011).

The molecular epidemiology and distribution of *T. cruzi* genotypes may have important implications with respect to characteristics of the disease. However, few correlations have related *T. cruzi* genetic variability and disease outcome, though it has been shown that TcI is more closely related to patients with cardiomyopathy in Colombia and Venezuela, while TcII-TcVI are more associated with patients with digestive syndrome (megaesophagus/megacolon) (Mantilla et al., 2011; Rassi et al., 2009). The distribution of *T. cruzi* genotypes and reservoirs is implicated in the genetic epidemiology of the disease. In the southern part of the American continent, infection of *Canis familiaris* has been found to be related to TcIV, V and VI, whereas infections in the north are associated with genotypes Ia and Ib (Falla et al., 2009; Herrera et al., 2009). Furthermore, a significant number of *D. marsupialis* are infected with the TcId genotype, which suggests an association with the sylvatic transmission cycle. Similar studies in primates have demonstrated that TcI predominantly infects arboreal reservoirs (Cura et al., 2010; Falla et al., 2009). There are several hypotheses regarding the distribution of different genetic groups of *T. cruzi*, suggesting that reservoirs belonging to arboreal ecotopes are preferentially infected with TcI and that terrestrial ecotopes are infected with TcII-TcVI (Yeo et al., 2005). This hypothesis is controversial in light of recent reports demonstrating that the arboreal ecotope reservoirs *Monodelphis brevicaudata*, *Philander frenata* and *Didelphis aurita* are infected with TcIII, TcIV and TcII, respectively (Marcili et al., 2009; Llewellyn et al., 2009b).

However, the associations are not absolute, and in the case of TcI, there is no apparent clustering of particular TcI genotypes with *Didelphis* in comparison to isolates from other arboreal mammals (Llewellyn et al., 2009b). Additionally, with respect to phylogeographical analyses of TcIII, the results indicate that isolates cluster according to geography rather than host association (Marcili et al., 2009; Llewellyn et al., 2009b). This could also be supported by the recent analysis developed in mammals naturally infected with TcI using microsatellite markers revealing the role of mammalian reservoirs in diversifying selection on *T. cruzi* (Llewellyn et al., 2011). Two interesting studies on host responses to different strains have confirmed, by comparative artificial infection, that in the southern USA, two species of opossum (*Monodelphis domestica* and *Didelphis virginiana*) seem to be resistant to TcIV (Roellig et al., 2009; Roellig et al., 2010). This highlights a mechanism for the association of a vertebrate host with one strain over others. The strong association between TcI and *Rhodnius* species can be explained by a similar mechanism: comparative studies on artificial infection of *R. prolixus* with various strains revealed a tendency for this species to be resistant to infection by TcII (Mello et al., 1995). In triatomines, susceptibility or resistance to trypanosome infections seems to be modulated by intestinal symbionts that are vital for

development. *T. cruzi* is considered to be subpathogenic for triatomines, whereas *Trypanosoma rangeli* is a species that commonly infects *Rhodnius* species and causes pathogenicity based on reduction of the number of symbionts (Vallejo et al., 2009). Studies using different species of triatomines, such as *R. pallenscens*, *T. dimidiata*, *R. colombiensis* and *P. geniculatus*, have shown the affinity of TcI for infecting these species in comparison with TcII (Mejia-Jaramillo et al., 2009). At least half of all species of triatomine bugs have been found to be naturally infected with *T. cruzi* (Lent and Wygodzinsky, 1979; Schofield, 1994). Unfortunately, the vast majority of these records do not include specific strain associations. This is clearly an area of potential research. In the context of dispersal triggered by starvation, there is evidence that starvation decreases *T. cruzi* infection in triatomines (Kollien and Schaub, 1998), and in some species, starvation may clear the infection altogether (Phillips et al., 1967; Vargas et al., 1985). This factor could help to explain paradigms such as that observed in Venezuela, where sylvatic and domestic bugs seem to be in panmixia, but TcI shows discrete general clustering of sylvatic and domestic cycles (Fitzpatrick et al., 2009; Llewellyn et al., 2009b). Triatomine bugs directly determine the aetiology of the strains of *T. cruzi* involved in human transmission cycles. This is clear because despite TcI and *Didelphis* being widespread, it is the northern distribution of *Rhodnius* that corresponds with its occurrence in human cycles. Overall, the aspects of epidemiological relevance are that associations between terrestrial ecology, *T. infestans*, terrestrial mammals, and *T. cruzi* strains TcII/TcIV have led to the prominence of TcII, TcV, and TcIV in human infections in the southern cone countries of South America. In the northern cone countries of South America, human American Trypanosomiasis infections seem to stem from TcI associated with arboreal *Rhodnius* and arboreal mammals.

The definition of *T. cruzi* nomenclature must be related to the biological, clinical and pathological characteristics associated with specific populations of *T. cruzi* (Campbell et al., 2004; Zafra et al., 2009). To our knowledge, few correlations reported have been demonstrated to date regarding differences of the host humoral response to specific *T. cruzi* genotypes; however, these findings were flawed because of the low reliability of the diagnostic tests used, leading to a high proportion of false negatives due to variability in the *T. cruzi* strain used for the diagnosis. The implication of TcI in severe forms of myocarditis in cardiac samples from chronic chagasic patients in Argentina and the lack of any specific clinical manifestation related to *T. cruzi* DTUs in Bolivian chagasic patients indicate the pleomorphism of *T. cruzi* (Ramírez et al., 2009; Moncayo and Ortiz, 2006; Burgos et al., 2010; del Puerto et al., 2010). There have been studies reporting detection of *T. cruzi* in blood samples. Direct detection of *T. cruzi* DTUs in the blood of chronic Chagasic patients was carried out by amplification of the 24Sα rDNA divergent domain and the use of mitochondrial house-keeping genes (Zafra et al., 2009). In this study, molecular characterisation of *T. cruzi* DTUs showed that most of the patients were infected with TcI, while some patients were found to be infected with TcII (9.9%). Recently, a new approach to *T. cruzi* DTU detection in chronic Chagasic patients was developed indicating that TcI is the predominant DTU, though TcII was also detected, and it was reported that the genetic characteristics of TcII parasites found in Colombia were similar to those of TcII found in Bolivia and Chile (González et al., 2010). Regarding the genetic variability of the parasite, prognostic markers based on mitochondrial genes where the presence of specific mutations can trigger complications of the chronic phase of the disease in asymptomatic patients have also been demonstrated (dos Santos et al., 2009; Carranza et al., 2009). Despite the observed

genetic variability, it is important to consider the presence of *T. cruzi* clones that can be found in different tissues. Several studies have demonstrated a specific histiotropism of *T. cruzi* in mice showing differences in the pathological, immunological and clinical features that the parasite can elicit in the host (Andrade et al., 2002; Ramírez et al., 2010; Manoel-Caetano et al., 2008). Moreover, some authors have shown that the *T. cruzi* population in a patient's bloodstream could be dissimilar to the parasite population that causes tissue damage (Vago et al., 2000). Differences were found in *T. cruzi* populations in the bloodstreams of patients with chronic Chagasic cardiomyopathy and those of Chagasic patients without cardiomyopathy (Venegas et al., 2009). Microsatellite analyses have also shown multiclonality in heart samples and in the bloodstreams of infected patients, demonstrating that specific populations of *T. cruzi* can probably determine disease outcome (Burgos et al., 2007; Valadares et al., 2008).

Molecular epidemiological studies on *T. cruzi* have attempted to establish the effects of different DTUs in the clinical progression of Chagas disease. Several studies have shown the effect of genetic variability on the host immune response (dos Santos et al., 2009; Melquiades-Rodríguez et al., 2010; Ramírez et al., 2009). It was previously known that cardiopathies in southern cone countries were caused by TcII, TcV and TcVI, but it has recently been demonstrated that TcI can play an important role specifically in severe cardiopathies related to Chagas disease. Studies of cardiac biopsies from Argentinean patients revealed that patients with severe myocarditis were infected with TcI, whereas those with moderate or absent myocarditis were infected with TcII, TcV or TcVI (Burgos et al., 2010). At the same time that the TcI genotype was found in severe myocarditis patients, it was demonstrated that in patients with chronic chagasic cardiopathy, the TcIa genotype was most commonly found in the bloodstream, whereas TcId was most commonly found in cardiac biopsies. These results are consistent with reports from patients in Colombia, where the least and most prevalent TcI genotypes in adult patients with chronic chagasic cardiopathy were TcId and TcIa, respectively (Ramírez et al., 2010). This suggests a possible type of histotropism associated with TcI genotypes as well as the epidemiological importance of this DTU in southern countries, where cardiopathies were previously thought to be caused primarily by TcII, TcV and TcVI. A model of clonal histiotropism has been previously reported showing how a composite of clones is related to disease outcome. Recently reported results from Colombia support this premise, with cardiac biopsies being observed to be infected with TcId, while TcIa is found circulating in the bloodstream (Zafra et al., 2011; Ramírez et al., 2010). This suggests the need to pursue studies to correlate the association between *T. cruzi* genotypes and clinical manifestations of Chagas disease. New studies are also necessary to determine the specific *T. cruzi* populations generating tissue damage in infected patients.

Most of the research performed in Chagas disease is related to understanding the molecular epidemiology of this endemic pathology. Many questions are continually emerging every day in this field based on epidemiological circuits with the aim of better estimating the transmission dynamics of *T. cruzi* in endemic areas. The involvement of *T. cruzi* genetic variability in clinical manifestations is of paramount importance and could resolve the question regarding the high pleomorphism displayed by this clinical entity. New initiatives must be created with interdisciplinary groups with the purpose of unravelling the molecular comparative epidemiology of Chagas disease and attempting to mitigate this pathology in endemic countries.

5. Concluding remarks

In this chapter, many examples regarding the usefulness of molecular epidemiology in parasitic diseases were addressed. These examples illustrated different applications of molecular methods to understand the pathogens that cause human parasitic diseases. It is important to consider the need for synergy between descriptive, analytical and molecular epidemiological methods to develop robust and unbiased data. As a relevant example, we presented the case of *T. cruzi* and described how molecular methods have been useful in defining hypotheses about the parasite's geographical distribution, host associations and the implications of different genotypes for clinical manifestations related to the heart. Despite the studies reported in the literature on molecular epidemiology in parasites, public health systems do not consider the importance of integration between these two areas. We propose the integration of molecular epidemiology and public health systems to mitigate and reduce the prevalence of tropical diseases caused mainly by parasites, and this combination could become a potential tool for disease prevention and control as well as for the development of appropriate programmes for disease surveillance in endemic countries.

6. References

- Agbo E, E., C., Majiwa PAO, Claassen HJ, Te Pas MF (2002) Molecular variation of *Trypanosoma brucei* subspecies as revealed by AFLP fingerprinting. *Parasitology* 124:349-358.
- Andersson B. (2011). The *Trypanosoma cruzi* genome; conserved core genes and extremely variable surface molecule families. *Research in Microbiology* 162:619-625
- Andrade LO, Machado CRS, Chiari E, Pena SDJ, Macedo AM (2002) *Trypanosoma cruzi*: role of host genetic background in the differential tissue distribution of parasite clonal populations. *Experimental Parasitology* 100:269-275
- Bañuls A-L, Dujardin J-C, Guerrini F, De Doncker S, Jacquet D, Arevalo J, Noel S, Le Ray D, Tibayrenc M (2000) Is *Leishmania (Viannia) peruviana* a Distinct Species? A MLEE/RAPD Evolutionary Genetics Answer. *Journal of Eukaryotic Microbiology* 47:197-207.
- Bañuls AL, Hide M, Tibayrenc M (2002) Evolutionary genetics and molecular diagnosis of *Leishmania* species. *Transactions of the Royal Society of Tropical Medicine and Hygiene* 96:S9-S13.
- Bennett RJ (2010) Coming of Age—Sexual Reproduction in *Candida* Species. *PLoS Pathog* 6:e1001155
- Bertram NS (1967) Laboratory studies of *Trypanosoma cruzi* infections in: *Rhodnius prolixus*--larvae and adults in: *Triatoma infestans*, *T. protracta* and *T. maculata*--adults. *J Med Entomol* 4:167-170.
- Bontell I, Hall N, Ashelford K, Dubey J, Boyle J, Lindh J, Smith J (2009) Whole genome sequencing of a natural recombinant *Toxoplasma gondii* strain reveals chromosome sorting and local allelic variants. *Genome Biology* 10:R53.

- Bretagne S, Durand R, Olivi M, Garin J-F, Sulahian A, Rivollet D, Vidaud M, Deniau M (2001) Real-Time PCR as a New Tool for Quantifying *Leishmania infantum* in Liver in Infected Mice. *Clin Diagn Lab Immunol* 8:828-831.
- Brisse S, Henriksson J, Barnabé C, Douzery EJP, Berkvens D, Serrano M, De Carvalho MRC, Buck GA, Dujardin J-C, Tibayrenc M (2003) Evidence for genetic exchange and hybridization in *Trypanosoma cruzi* based on nucleotide sequences and molecular karyotype. *Infection, Genetics and Evolution* 2:173-183.
- Burgos JM, Altchek J, Bisio M, Duffy T, Valadares HMS, Seidenstein ME, Piccinali R, Freitas JM, Levin MJ, Macchi L, Macedo AM, Freilij H, Schijman AG (2007) Direct molecular profiling of minicircle signatures and lineages of *Trypanosoma cruzi* bloodstream populations causing congenital Chagas disease. *International Journal for Parasitology* 37:1319-1327.
- Burgos JM, Diez M, Vigliano C, Bisio M, Risso M, Duffy Ts, Cura C, Brusses B, Favaloro L, Leguizamón MaS, Lucero RH, Laguens R, Levin MJ, Favaloro R, Schijman AG Molecular Identification of *Trypanosoma cruzi* Discrete Typing Units in End-Stage Chronic Chagas Heart Disease and Reactivation after Heart Transplantation. *Clinical Infectious Diseases* 51:485-495.
- Campbell DA WS, Sturm NR. (2004) The determinants of Chagas disease: connecting parasite and host genetics. *Curr Mol Med* 4:549-562.
- Carranza JC, Valadares HMS, D'Ávila DA, Baptista RP, Moreno M, Galvão LMC, Chiari E, Sturm NR, Gontijo ED, Macedo AM, Zingales B (2009) *Trypanosoma cruzi* maxicircle heterogeneity in Chagas disease patients from Brazil. *International Journal for Parasitology* 39:963-973
- Clark CG, Ali IKM, Zaki M, Loftus BJ, Hall N (2006) Unique organisation of tRNA genes in *Entamoeba histolytica*. *Molecular and Biochemical Parasitology* 146:24-29.
- Cura CI, Mejía-Jaramillo AM, Duffy T, Burgos JM, Rodríguez M, Cardinal MV, Kjos S, Gurgel-Gonçalves R, Blanchet D, De Pablos LM, Tomasini N, da Silva A, Russomando G, Cuba CAC, Aznar C, Abate T, Levin MJ, Osuna A, Gürtler RE, Diosque P, Solari A, Triana-Chávez O, Schijman AG *Trypanosoma cruzi* I genotypes in different geographical regions and transmission cycles based on a microsatellite motif of the intergenic spacer of spliced-leader genes. *International Journal for Parasitology* 40:1599-1607.
- da Silva Manoel-Caetano F, Carareto CMA, Borim AA, Miyazaki K, Silva AE (2008) kDNA gene signatures of *Trypanosoma cruzi* in blood and oesophageal mucosa from chronic chagasic patients. *Transactions of the Royal Society of Tropical Medicine and Hygiene* 102:1102-1107.
- del Puerto R, Nishizawa JE, Kikuchi M, Iihoshi N, Roca Y, Avilas C, Gianella A, Lora J, Gutierrez Velarde FU, Renjel LA, Miura S, Higo H, Komiya N, Maemura K, Hirayama K Lineage Analysis of Circulating *Trypanosoma cruzi* Parasites and Their Association with Clinical Forms of Chagas Disease in Bolivia. *PLoS Negl Trop Dis* 4:e687.
- Duffy T, Bisio M, Altchek J, Burgos JM, Diez M, Levin MJ, Favaloro RR, Freilij H, Schijman AG (2009) Accurate Real-Time PCR Strategy for Monitoring Bloodstream Parasitic Loads in Chagas Disease Patients. *PLoS Negl Trop Dis* 3:e419.

- El-Sayed NM, Myler PJ, Bartholomeu DC, Nilsson D, Aggarwal G, Tran A-N, Ghedin E, Worthey EA, Delcher AL, Blandin GI, Westenberger SJ, Caler E, Cerqueira GC, Branche C, Haas B, Anupama A, Arner E, Å...slund L, Attipoe P, Bontempi E, Bringaud Fdr, Burton P, Cadag E, Campbell DA, Carrington M, Crabtree J, Darban H, da Silveira JF, de Jong P, Edwards K, Englund PT, Fazelina G, Feldblyum T, Ferella M, Frasch AC, Gull K, Horn D, Hou L, Huang Y, Kindlund E, Klingbeil M, Kluge S, Koo H, Lacerda D, Levin MJ, Lorenzi H, Louie T, Machado CR, McCulloch R, McKenna A, Mizuno Y, Mottram JC, Nelson S, Ochaya S, Osoegawa K, Pai G, Parsons M, Pentony M, Pettersson U, Pop M, Ramirez JL, Rinta J, Robertson L, Salzberg SL, Sanchez DO, Seyler A, Sharma R, Shetty J, Simpson AJ, Sisk E, Tammi MT, Tarleton R, Teixeira S, Van Aken S, Vogt C, Ward PN, Wickstead B, Wortman J, White O, Fraser CM, Stuart KD, Andersson Br (2005) The Genome Sequence of *Trypanosoma cruzi*, Etiologic Agent of Chagas Disease. *Science* 309:409-415.
- Falla A, Herrera C, Fajardo A, Montilla M, Vallejo GA, Guhl F (2009) Haplotype identification within *Trypanosoma cruzi* I in Colombian isolates from several reservoirs, vectors and humans. *Acta Tropica* 110:15-21.
- Fitzpatrick S, Feliciangeli MD, Sanchez-Martin MJ, Monteiro FA, Miles MA (2008) Molecular Genetics Reveal That Silvatic *Rhodnius prolixus* Do Colonise Rural Houses. *PLoS Negl Trop Dis* 2:e210.
- Franzen O, Ochaya S, Sherwood E, Lewis MD, Llewellyn MS, Miles MA, Andersson Br Shotgun Sequencing Analysis of *Trypanosoma cruzi* I Sylvio X10/1 and Comparison with *T. cruzi* VI CL Brener. *PLoS Negl Trop Dis* 5:e984.
- Fuentes I, Rubio JM, Ramirez C, Alvar J (2001) Genotypic Characterization of *Toxoplasma gondii* Strains Associated with Human Toxoplasmosis in Spain: Direct Analysis from Clinical Samples. *J Clin Microbiol* 39:1566-1570.
- Gardner MJ, Hall N, Fung E, White O, Berriman M, Hyman RW, Carlton JM, Pain A, Nelson KE, Bowman S, Paulsen IT, James K, Eisen JA, Rutherford K, Salzberg SL, Craig A, Kyes S, Chan M-S, Nene V, Shallom SJ, Suh B, Peterson J, Angiuoli S, Perteau M, Allen J, Selengut J, Haft D, Mather MW, Vaidya AB, Martin DMA, Fairlamb AH, Fraunholz MJ, Roos DS, Ralph SA, McFadden GI, Cummings LM, Subramanian GM, Mungall C, Venter JC, Carucci DJ, Hoffman SL, Newbold C, Davis RW, Fraser CM, Barrell B (2002) Genome sequence of the human malaria parasite *Plasmodium falciparum*. *Nature* 419:498-511.
- Gaunt MW, Yeo M, Frame IA, Stothard JR, Carrasco HJ, Taylor MC, Mena SS, Veazey P, Miles GAJ, Acosta N, de Arias AR, Miles MA (2003) Mechanism of genetic exchange in American trypanosomes. *Nature* 421:936-939.
- Gomes MA, Melo MN, Macedo AM, Furst C, Silva EF (2000) RAPD in the analysis of isolates of *Entamoeba histolytica*. *Acta Tropica* 75:71-77 .
- Gomez JC, McNamara DT, Bockarie MJ, Baird JK, Carlton JM, Zimmerman PA (2003) Identification of a polymorphic *Plasmodium vivax* microsatellite marker. *The American Journal of Tropical Medicine and Hygiene* 69:377-379.
- González CI, Ortiz S, Solari A Colombian *Trypanosoma cruzi* major genotypes circulating in patients: Minicircle homologies by cross-hybridization analysis. *International Journal for Parasitology* 40:1685-1692.
- Grigg ME, Suzuki Y (2003) Sexual recombination and clonal evolution of virulence in *Toxoplasma*. *Microbes and Infection* 5:685-690 .

- Guhl F, Ramírez JD (2011) *Trypanosoma cruzi* I diversity: Towards the need of genetic subdivision? *Acta Tropica* 119:1-4.
- Guy RA, Payment P, Krull UJ, Horgen PA (2003) Real-Time PCR for Quantification of *Giardia* and *Cryptosporidium* in Environmental Water Samples and Sewage. *Appl Environ Microbiol* 69:5178-5185 .
- Harvey PH, Keymer AE (1987) Sex among the parasites. *Nature* 330:317-318.
- Herrera C, Bargues MD, Fajardo A, Montilla M, Triana O, Vallejo GA, Guhl F (2007) Identifying four *Trypanosoma cruzi* I isolate haplotypes from different geographic regions in Colombia. *Infection, Genetics and Evolution* 7:535-539.
- Herrera C, Guhl F, Falla A, Fajardo A, Montilla M, Adolfo Vallejo G, Bargues MD (2009) Genetic Variability and Phylogenetic Relationships within *Trypanosoma cruzi* I Isolated in Colombia Based on Miniexon Gene Sequences. *Journal of Parasitology Research* 2009 DOI 10.1155/2009/897364.
- Imwong M, Sudimack D, Pukrittayakamee S, Osorio L, Carlton JM, Day NPJ, White NJ, Anderson TJC (2006) Microsatellite Variation, Repeat Array Length, and Population History of *Plasmodium vivax*. *Molecular Biology and Evolution* 23:1016-1018.
- Ivens AC, Peacock CS, Worthey EA, Murphy L, Aggarwal G, Berriman M, Sisk E, Rajandream M-A, Adlem E, Aert R, Anupama A, Apostolou Z, Attipoe P, Bason N, Bauser C, Beck A, Beverley SM, Bianchetti G, Borzym K, Bothe G, Bruschi CV, Collins M, Cadag E, Ciarloni L, Clayton C, Coulson RMR, Cronin A, Cruz AK, Davies RM, De Gaudenzi J, Dobson DE, Duesterhoeft A, Fazelina G, Fosker N, Frasch AC, Fraser A, Fuchs M, Gabel C, Goble A, Goffeau A, Harris D, Hertz-Fowler C, Hilbert H, Horn D, Huang Y, Klages S, Knights A, Kube M, Larke N, Litvin L, Lord A, Louie T, Marra M, Masuy D, Matthews K, Michaeli S, Mottram JC, Muller-Auer S, Munden H, Nelson S, Norbertczak H, Oliver K, O'Neil S, Pentony M, Pohl TM, Price C, Purnelle Bnd, Quail MA, Rabbinowitsch E, Reinhardt R, Rieger M, Rinta J, Robben J, Robertson L, Ruiz JC, Rutter S, Saunders D, SchÄ¶fer M, Schein J, Schwartz DC, Seeger K, Seyler A, Sharp S, Shin H, Sivam D, Squares R, Squares S, Tosato V, Vogt C, Volckaert G, Wambutt R, Warren T, Wedler H, Woodward J, Zhou S, Zimmermann W, Smith DF, Blackwell JM, Stuart KD, Barrell B, Myler PJ (2005) The Genome of the Kinetoplastid Parasite, *Leishmania major*. *Science* 309:436-442 .
- Jamieson SE, Miller EN, Peacock CS, Fakiola M, Wilson ME, Bales-Holst A, Shaw MA, Silveira F, Shaw JJ, Jeronimo SM, Blackwell JM (2006) Genome-wide scan for visceral leishmaniasis susceptibility genes in Brazil. *Genes Immun* 8:84-90.
- Johnson M, Broady K, Angelici MC, Johnson A (2003) The relationship between nucleoside triphosphate hydrolase (NTPase) isoform and *Toxoplasma* strain virulence in rat and human toxoplasmosis. *Microbes and Infection* 5:797-806.
- Kilbourne ED (1973) The Molecular Epidemiology of Influenza. *Journal of Infectious Diseases* 127:478-487.
- Kollien AH and Schaub AGA (1998) Development of *Trypanosoma cruzi* after starvation and feeding of the vector - a review. *Tokai J Exp Clin Med* 23:335-340.
- Lehane MJ (1994) TRIATOMINAE: BIOLOGY and CONTROL. By C. J. Schofield. *Medical and Veterinary Entomology* 8:218-218 DOI 10.1111/j.1365-2915.1994.tb00501.x.


- Lent and Wygodzinsky (1979) Revision of the Triatominae, Hemiptera Reduviidae and their significance as vectors of Chagas' disease. *Bull Am Mus Nat Hist* 163:123-520.
- Lewis MD, Ma J, Yeo M, Carrasco HnJ, Llewellyn MS, Miles MA (2009) Genotyping of *Trypanosoma cruzi*: Systematic Selection of Assays Allowing Rapid and Accurate Discrimination of All Known Lineages. *The American Journal of Tropical Medicine and Hygiene* 81:1041-1049.
- Lier T, Simonsen GS, Wang T, Lu D, Haukland HH, Vennervald BJ, Hegstad J, Johansen MV (2009) Real-Time Polymerase Chain Reaction for Detection of Low-Intensity *Schistosoma japonicum* Infections in China. *The American Journal of Tropical Medicine and Hygiene* 81:428-432
- Llewellyn MS, Lewis MD, Acosta N, Yeo M, Carrasco HJ, Segovia M, Vargas J, Torrico F, Miles MA, Gaunt MW (2009b) *Trypanosoma cruzi* IIc: Phylogenetic and Phylogeographic Insights from Sequence and Microsatellite Analysis and Potential Impact on Emergent Chagas Disease. *PLoS Negl Trop Dis* 3:e510.
- Llewellyn MS, Miles MA, Carrasco HJ, Lewis MD, Yeo M, Vargas J, Torrico F, Diosque P, Valente V, Valente SA, Gaunt MW (2009) Genome-Scale Multilocus Microsatellite Typing of *Trypanosoma cruzi* Discrete Typing Unit I Reveals Phylogeographic Structure and Specific Genotypes Linked to Human Infection. *PLoS Pathog* 5:e1000410.
- Llewellyn MS, Rivett-Carnac JB, Fitzpatrick S, Lewis MD, Yeo M, Gaunt MW, Miles MA Extraordinary *Trypanosoma cruzi* diversity within single mammalian reservoir hosts implies a mechanism of diversifying selection. *International Journal for Parasitology* 41:609-614.
- Mantilla JC, Zafra GA, Macedo AM, González CI (2010) Mixed infection of *Trypanosoma cruzi* I and II in a Colombian cardiomyopathic patient. *Human Pathology* 41:610-613.
- Marcili A, Lima L, Valente VC, Valente SA, Batista JS, Junqueira ACV, Souza AI, da Rosa JA, Campaner M, Lewis MD, Llewellyn MS, Miles MA, Teixeira MMG (2009) Comparative phylogeography of *Trypanosoma cruzi* TCIIc: New hosts, association with terrestrial ecotopes, and spatial clustering. *Infection, Genetics and Evolution* 9:1265-1274.
- Mary C, Faraut F, Lascombe L, Dumon H (2004) Quantification of *Leishmania infantum* DNA by a Real-Time PCR Assay with High Sensitivity. *J Clin Microbiol* 42:5249-5255.
- Masiga DK, Ndung'u K, Tweedie A, Tait A, Turner CMR (2006) *Trypanosoma evansi*: Genetic variability detected using amplified restriction fragment length polymorphism (AFLP) and random amplified polymorphic DNA (RAPD) analysis of Kenyan isolates. *Experimental Parasitology* 114:147-153.
- Mathews HM, Moss DM, Healy GR, Visvesvara GS (1983) Polyacrylamide gel electrophoresis of isoenzymes from *Entamoeba* species. *J Clin Microbiol* 17:1009-1012.
- Mejía-Jaramillo AM, Peña VH, Triana-Chávez O (2009) *Trypanosoma cruzi*: Biological characterization of lineages I and II supports the predominance of lineage I in Colombia. *Experimental Parasitology* 121:83-91.
- Mello CB, Garcia ES, Ratcliffe NA, Azambuja P (1995) *Trypanosoma cruzi* and *Trypanosoma rangeli*: Interplay with Hemolymph Components of *Rhodnius prolixus*. *Journal of Invertebrate Pathology* 65:261-268.

- Miles MA, Llewellyn MS, Lewis MD, Yeo M, Baleela R, Fitzpatrick S, Gaunt MW, Mauricio IL (2009) The molecular epidemiology and phylogeography of *Trypanosoma cruzi* and parallel research on Leishmania: looking back and to the future. *Parasitology* 136:1509-1528.
- Miles MA, Povoá MM, De Souza AA, Lainson R, Shaw JJ, Ketteridge DS (1981) Chagas's disease in the Amazon Basin: II. The distribution of *Trypanosoma cruzi* zymodemes 1 and 3 in Pará State, north Brazil. *Transactions of the Royal Society of Tropical Medicine and Hygiene* 75:667-674.
- Miles MA, Toye PJ, Oswald SC, Godfrey DG (1977) The identification by isoenzyme patterns of two distinct strain-groups of *Trypanosoma cruzi*, circulating independently in a rural area of Brazil. *Transactions of the Royal Society of Tropical Medicine and Hygiene* 71:217-225.
- Moncayo A, Ortiz Yanine MI (2006) An update on Chagas disease (human American trypanosomiasis). *Annals of Tropical Medicine and Parasitology* 100:663-677.
- Montalvo AM, Fraga J, Aylema Romero J, Monzote L, Montano I, Dujardin JC (2006) PCR-RFLP/Hsp70 para identificar y tipificar *Leishmania* de la región neotropical. *Revista Cubana de Medicina Tropical* 58:0-0.
- Morrison LJ (2011) Parasite-driven pathogenesis in *Trypanosoma brucei* infections. *Parasite Immunology* 33:448-455.
- Morrison LJ, Tait A, McLellan S, Sweeney L, Turner CMR, MacLeod A (2009) A Major Genetic Locus in *Trypanosoma brucei* Is a Determinant of Host Pathology. *PLoS Negl Trop Dis* 3:e557.
- Mzilahowa T, McCall PJ, Hastings IM (2007) "Sexual" Population Structure and Genetics of the Malaria Agent *P. falciparum*. *PLoS ONE* 2:e613.
- Njiokou F, Nkinin SW, Grébaut P, Penchenier L, Barnabé C, Tibayrenc M, Herder S (2004) An isoenzyme survey of *Trypanosoma brucei* s.l. from the Central African subregion: population structure, taxonomic and epidemiological considerations. *Parasitology* 128:645-653.
- Patterson JS, Guhl F (2010) Geographical Distribution of Chagas Disease American Trypanosomiasis. Elsevier, London, pp. 83-114.
- Pecson BM, Barrios JA, Johnson DR, Nelson KL (2006) A Real-Time PCR Method for Quantifying Viable *Ascaris* Eggs Using the First Internally Transcribed Spacer Region of Ribosomal DNA. *Appl Environ Microbiol* 72:7864-7872.
- Pinto M, Rosa JD, Fernandes Z, Graminha M, Mine J, Allegretti S, Delort S, Riedel C, Paes E, Cupolillo E (2005) Isolation and isoenzyme characterization of *Leishmania (Viannia) braziliensis* from a case of human cutaneous leishmaniasis in northeast centre of the state of São Paulo. *Memórias do Instituto Oswaldo Cruz* 100:733-734.
- Prakash A, Chakraborti A, Mahajan RC, Ganguly NK (2000) *Entamoeba histolytica*: Rapid Detection of Indian Isolates by Cysteine Proteinase Gene-Specific Polymerase Chain Reaction. *Experimental Parasitology* 95:285-287.
- Prata A (2001) Clinical and epidemiological aspects of Chagas disease. *The Lancet Infectious Diseases* 1:92-100.
- Quan J-H, Kim TY, Choi I-U, Lee Y-H (2008) Genotyping of a Korean isolate of *Toxoplasma gondii* by multilocus PCR-RFLP and microsatellite analysis. *Korean J Parasitol* 46:105-108.

- Ramírez JD, Duque MC, Guhl F (2011) Phylogenetic reconstruction based on Cytochrome b (Cytb) gene sequences reveals distinct genotypes within Colombian *Trypanosoma cruzi* I populations. *Acta Tropica* 119:61-65.
- Ramírez JD, Guhl F, Rendón LM, Rosas F, Marin-Neto JA, Morillo CA (2010) Chagas Cardiomyopathy Manifestations and *Trypanosoma cruzi* Genotypes Circulating in Chronic Chagasic Patients. *PLoS Negl Trop Dis* 4:e899.
- Ramírez JD, Guhl F, Umezawa ES, Morillo CA, Rosas F, Marin-Neto JA, Restrepo S (2009) Evaluation of Adult Chronic Chagas' Heart Disease Diagnosis by Molecular and Serological Methods. *J Clin Microbiol* 47:3945-3951.
- Ranasinghe S, Rogers ME, Hamilton JGC, Bates PA, Maingon RDC (2008) A real-time PCR assay to estimate *Leishmania chagasi* load in its natural sand fly vector *Lutzomyia longipalpis*. *Transactions of the Royal Society of Tropical Medicine and Hygiene* 102:875-882.
- Rassi A, Marin-Neto JA Chagas disease. *The Lancet* 375:1388-1402.
- Rodrigues CM, Valadares HMS, Francisco AF, Arantes JM, Campos CFa, Teixeira-Carvalho Aa, Martins-Filho OA, Araujo MrSS, Arantes RME, Chiari E, Franco GrR, Machado CR, Pena SrDJ, Faria AMC, Macedo AM (2010) Coinfection with Different *Trypanosoma cruzi* Strains Interferes with the Host Immune Response to Infection. *PLoS Negl Trop Dis* 4:e846.
- Roellig DM, Ellis AE, Yabsley MJ (2009) Genetically different isolates of *Trypanosoma cruzi* elicit different infection dynamics in raccoons (*Procyon lotor*) and Virginia opossums (*Didelphis virginiana*). *International Journal for Parasitology* 39:1603-1610.
- Roellig DM, McMillan K, Ellis AE, Vandenberg JL, Champagne DE, Yabsley MJ (2010) Experimental infection of two South American reservoirs with four distinct strains of *Trypanosoma cruzi*. *Parasitology* 137:959-966 .
- Romand S, Chosson M, Franck J, Wallon M, Kieffer F, Kaiser K, Dumon H, Peyron F, Thulliez P, Picot S (2004) Usefulness of quantitative polymerase chain reaction in amniotic fluid as early prognostic marker of fetal infection with *Toxoplasma gondii*. *American Journal of Obstetrics and Gynecology* 190:797-802 DOI 10.1016/j.ajog.2003.09.039.
- Rougeron V, De Meeüs T, Hide M, Waleckx E, Bermudez H, Arevalo J, Llanos-Cuentas A, Dujardin J-C, De Doncker S, Le Ray D, Ayala FJ, Bañuls A-L (2009) Extreme inbreeding in *Leishmania braziliensis*. *Proceedings of the National Academy of Sciences* 106:10224-10229.
- Rozas M, De Doncker S, Adauí V, Coronado X, Barnabé C, Tibyarenc M, Solari A, Dujardin J-C (2007) Multilocus Polymerase Chain Reaction Restriction Fragment Length Polymorphism Genotyping of *Trypanosoma cruzi* (Chagas Disease): Taxonomic and Clinical Applications. *Journal of Infectious Diseases* 195:1381-1388.
- Santos DM, Talvani A, da Mata Guedes PM, Machado-Coelho GLL, de Lana M, Bahia MT (2009) *Trypanosoma cruzi*: Genetic diversity influences the profile of immunoglobulins during experimental infection. *Experimental Parasitology* 121:8-14.
- Schmunis GA, Yadon ZE Chagas disease: A Latin American health problem becoming a world health problem. *Acta Tropica* 115:14-21.

- Singh U, Ehrenkaufer GM (2009) Recent insights into *Entamoeba* development: Identification of transcriptional networks associated with stage conversion. *International Journal for Parasitology* 39:41-47 .
- Smith JM, Smith NH, O'Rourke M, Spratt BG (1993) How clonal are bacteria? *Proceedings of the National Academy of Sciences* 90:4384-4388.
- Spotorno O AE, Córdova L, Solari I A (2008) Differentiation of *Trypanosoma cruzi* I subgroups through characterization of cytochrome b gene sequences. *Infection, Genetics and Evolution* 8:898-900.
- Stensvold C, Clarck G (2011) Multilocus sequence typing of *Blastocystis* sp. subtype 3. 21st European Congress of Clinical Microbiology and Infectious Diseases (ECCMID), 27th International Congress of Chemotherapy (ICC).
- Sturm NR, Vargas NS, Westenberger SJ, Zingales B, Campbell DA (2003) Evidence for multiple hybrid groups in *Trypanosoma cruzi*. *International Journal for Parasitology* 33:269-279.
- Su C, Zhang X, Dubey JP (2006) Genotyping of *Toxoplasma gondii* by multilocus PCR-RFLP markers: A high resolution and simple method for identification of parasites. *International Journal for Parasitology* 36:841-848 .
- Tibayrenc M, Ayala FJ (1991) Towards a population genetics of microorganisms: The clonal theory of parasitic protozoa. *Parasitology Today* 7:228-232.
- Vago AR, Andrade LO, Leite AA, d'Ávila Reis D, Macedo AM, Adad SJ, Tostes Jr S, Moreira MCV, Filho GB, Pena SDJ (2000) Genetic Characterization of *Trypanosoma cruzi* Directly from Tissues of Patients with Chronic Chagas Disease: Differential Distribution of Genetic Types into Diverse Organs. *The American Journal of Pathology* 156:1805-1809.
- Valadares HMS, Pimenta JR, de Freitas JM, Duffy T, Bartholomeu DC, de Paula Oliveira R, Chiari E, Moreira MdCV, Filho GB, Schijman AG, Franco GR, Machado CR, Pena SDJ, Macedo AM (2008) Genetic profiling of *Trypanosoma cruzi* directly in infected tissues using nested PCR of polymorphic microsatellites. *International Journal for Parasitology* 38:839-850.
- Vallejo GA, Guhl F, Schaub GA *Triatominae-Trypanosoma cruzi/T. rangeli*: Vector-parasite interactions. *Acta Tropica* 110:137-147.
- Vargas LG, Zeledon R (1985) Effect of fasting on *Trypanosoma cruzi* infection in *Triatoma dimidiata* (Hemiptera: Reduviidae). *Journal of Medical Entomology* 22:683-683.
- Venegas J, Coñoepan W, Pichuantes S, Miranda S, Apt W, Arribada A, Zulantay I, Coronado X, Rodriguez J, Reyes E, Solari A, Sanchez G (2009) Differential distribution of *Trypanosoma cruzi* clones in human chronic chagasic cardiopathic and non-cardiopathic individuals. *Acta Tropica* 109:187-193.
- Volpini ÂC, Passos VMA, Oliveira GC, Romanha AJ (2004) PCR-RFLP to identify *Leishmania (Viannia) braziliensis* and *L. (Leishmania) amazonensis* causing American cutaneous leishmaniasis. *Acta Tropica* 90:31-37.
- Westenberger SJ, Barnabé C, Campbell DA, Sturm NR (2005) Two Hybridization Events Define the Population Structure of *Trypanosoma cruzi*. *Genetics* 171:527-543.
- WHO (2007) Report of scientific group in Chagas disease. Buenos Aires, Argentina, April 17-20, 2005. Special Programme for Research and Training in Tropical Diseases (TDR).

- WHO (2010) Millennium Development Goals: progress towards the health-related Millennium Development Goals .
- Wortmann G, Hochberg LP, Arana BA, Rizzo NR, Arana F, Ryan JR (2007) Diagnosis of cutaneous Leishmaniasis in Guatemala using a Real-Time Polymerase Chain Reaction assay and the smartcycler. *The American Journal of Tropical Medicine and Hygiene* 76:906-908.
- Yeo M, Acosta N, Llewellyn M, Sánchez H, Adamson S, Miles GAJ, López E, González N, Patterson JS, Gaunt MW, Arias ARd, Miles MA (2005) Origins of Chagas disease: Didelphis species are natural hosts of *Trypanosoma cruzi* I and armadillos hosts of *Trypanosoma cruzi* II, including hybrids. *International Journal for Parasitology* 35:225-233.
- Yeo M, Mauricio IL, Messenger LA, Lewis MD, Llewellyn MS, Acosta N, Bhattacharyya T, Diosque P, Carrasco HJ, Miles MA Multilocus Sequence Typing (MLST) for Lineage Assignment and High Resolution Diversity Studies in *Trypanosoma cruzi*. *PLoS Negl Trop Dis* 5:e1049.
- Zafra G, Mantilla J, Valadares H, Macedo A, González C (2008) Evidence of *Trypanosoma cruzi* II infection in Colombian chagasic patients. *Parasitology Research* 103:731-734.
- Zafra G, Mantilla JC, Jácome J, Macedo AM, González CI (2011) Direct analysis of genetic variability in *Trypanosoma cruzi* populations from tissues of Colombian chagasic patients. *Human Pathology* 42:1159-1168.
- Zemanová E, Jirku M, Mauricio IL, Horák A, Miles MA, Lukes J (2007) The *Leishmania donovani* complex: Genotypes of five metabolic enzymes (ICD, ME, MPI, G6PDH, and FH), new targets for multilocus sequence typing. *International Journal for Parasitology* 37:149-160.
- Zhang W-W, Miranda-Verastegui C, Arevalo J, Ndao M, Ward B, Cuentas AL, Matlashewski G (2006) Development of a Genetic Assay to Distinguish between *Leishmania viannia* Species on the Basis of Isoenzyme Differences. *Clinical Infectious Diseases* 42:801-809.
- Zhao G-p (2007) SARS molecular epidemiology: a Chinese fairy tale of controlling an emerging zoonotic disease in the genomics era. *Philosophical Transactions of the Royal Society B: Biological Sciences* 362:1063-1081.
- Zingales B, Andrade S, Briones M, Campbell D, Chiari E, Fernandes O, Guhl F, Lages-Silva E, Macedo A, Machado C, Miles M, Romanha A, Sturm N, Tibayrenc M, Schijman A (2009) A new consensus for *Trypanosoma cruzi* intraspecific nomenclature: second revision meeting recommends TcI to TcVI. *Memórias do Instituto Oswaldo Cruz* 104:1051-1054.



Epidemiology - Current Perspectives on Research and Practice

Edited by Prof. Nuno Lunet

ISBN 978-953-51-0382-0

Hard cover, 208 pages

Publisher InTech

Published online 13, March, 2012

Published in print edition March, 2012

This special issue resulted from the invitation made to selected authors to contribute with an overview of a specific subject of their choice, and is based on a collection of papers chosen to exemplify some of the interests, uses and views of the epidemiology across different areas of research and practice. Rather than the comprehensiveness and coherence of a conventional textbook, readers will find a set of independent chapters, each of them of a great interest in their own specialized areas within epidemiology. Taken together, they illustrate the contrast between the attempt to extend the limits of applicability of epidemiological research, and the "regular" scientific activity in this field or an applied epidemiology. Epidemiologists with different levels of expertise and interests will be able to find informative and inspiring readings among the chapters of this book.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Juan David Ramírez and Felipe Guhl (2012). Molecular Epidemiology of Parasitic Diseases: The Chagas Disease Model, *Epidemiology - Current Perspectives on Research and Practice*, Prof. Nuno Lunet (Ed.), ISBN: 978-953-51-0382-0, InTech, Available from: <http://www.intechopen.com/books/epidemiology-current-perspectives-on-research-and-practice/molecular-epidemiology-of-parasitic-diseases-the-chagas-disease-model>

INTECH

open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2012 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.