
Modulating Role of TTR in A β Toxicity, from Health to Disease

Isabel Cardoso, Luis Miguel Santos and
Mobina Alemi

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/63194>

Abstract

Amyloidosis is a generic term that refers to a wide spectrum of diseases that are characterized by the deposition of proteins in different organs, forming insoluble aggregates. Examples include islet amyloid polypeptide (IAPP) associated with diabetes type 2, prion protein (PrP) related with spongiform encephalopathies, (TTR) associated with familial amyloidotic polyneuropathy (FAP), and amyloid-beta (A β) peptide linked to Alzheimer's disease (AD), the most common form of dementia. A β peptide, thought to be the causative agent in AD, is generated upon sequential cleavage of the amyloid precursor protein (APP), by beta- and gamma-secretases, and it is believed that an imbalance between A β production and clearance results in its accumulation in the brain. TTR is a 55 kDa homotetrameric protein synthesized by the liver and choroid plexus of the brain and is involved in the transport of thyroid hormones and retinol. TTR protects against A β toxicity by binding the peptide, thus inhibiting its aggregation. Also, increased A β levels are found in both brain and plasma of AD mice with only one copy of the TTR gene, when compared to animals with two copies of the gene, suggesting a role for TTR in A β clearance. Growing evidence also suggests a wider role for TTR in central nervous system neuroprotection, including in the cases of ischemia, regeneration, and memory.

Keywords: Alzheimer's disease (AD), A β peptide, transthyretin (TTR), neuroprotection, blood-brain barrier (BBB)

1. Amyloidosis

1.1. Amyloidosis definition

Amyloidosis has long been used as a general term referring to a wide spectrum of protein-misfolding diseases [1], which are characterized by the extracellular deposition of those proteins in different organs, consequently forming insoluble aggregates called amyloid, a term popularized by Virchow in 1854 [2]. According to the Nomenclature Committee of the International Society of Amyloidosis (ISA), 31 identified proteins form extracellular amyloid fibrils in humans [3]. Amyloid fibrils are characterized by certain tinctorial properties, independently of the precursor protein forming the deposits, that for a long time were the only diagnosis available. These include apple-green birefringence under polarized light after staining with Congo red and yellow-green fluorescence after staining with thioflavin S and thioflavin T; thioflavin T has also been shown to interact with amyloid in suspension producing a specific fluorescent signal with a new excitation maximum at 450 nm. Ultrastructural studies by transmission electron microscopy (TEM) revealed that the amyloid material is fibrillary appearing as bundles of straight or coiled fibrils, non-branched, 7–10 nm wide and variable in length; in most cases, they seem to be helically twisted. Amyloid fibrils present a high content in β -pleated sheet as demonstrated by x-ray diffraction analysis and extensive antiparallel β -sheet strands with their axes running perpendicularly to the axis of the growing fibril (cross- β pattern).

Amyloid deposits are not entirely composed of the amyloid precursor protein. Several components have been found associated with all amyloid fibrils. These include serum amyloid P component (SAP), sulphonated glycosaminoglycans (GAGs), apolipoproteins E and J, α 1-antichymotrypsin, several basement membrane components such as fibronectin, laminin and collagen type IV, complement proteins, and metal ions.

The extracellular deposition of fibrillary proteins leads to cell damage, organ dysfunction and death, and thus, these proteins are associated with a unique clinical syndrome, as seen in the case of the islet amyloid polypeptide (IAPP) associated with diabetes type 2 [4], prion protein (PrP) associated with the spongiform encephalopathies [5], transthyretin (TTR) associated with familial amyloidotic polyneuropathy [6] (FAP), and amyloid-beta ($A\beta$) peptide associated with Alzheimer's disease [7] (AD), among others.

Amyloid disorders are usually divided into two categories depending on the distribution of the amyloid deposits: localized and systemic amyloidosis. In localized amyloidosis, amyloid is restricted to a single tissue or organ, usually in the surroundings of the cells responsible for the synthesis of the precursor protein; in systemic amyloidosis, the amyloidogenic proteins are usually derived from circulating precursors that are either in excess, abnormal or both. Amyloidosis can also be hereditary or non-hereditary.

This chapter will focus in AD, a form of localized amyloidosis affecting the central nervous system, and the most common form of dementia. In particular, we will discuss the neuroprotective role of TTR in AD, in addition to its amyloidogenic role in FAP, an example of systemic amyloidosis with a special involvement of the peripheral nerve.

2. Overview of AD

AD was firstly described by Alois Alzheimer in 1906 and is characterized by progressive loss of cognitive functions, ultimately leading to death [8]. This condition highly affects not only the life of patients but also the life of their caregivers. Pathologically, the disease is characterized by the presence of extraneuronal amyloid plaques consisting of aggregates of the A β peptide, and neurofibrillary tangles (NFTs) which are intracellular aggregates of abnormally hyperphosphorylated tau protein [9]. A β peptide is generated upon sequential cleavage of the amyloid precursor protein (APP), by beta- and gamma-secretases, and it is believed that an imbalance between A β production and clearance results in its accumulation in the brain.

2.1. From the first description to the confirmation

Alzheimer's disease was first described in the 1907's paper entitled "Über eine eigenartige Erkrankung der Hirnrinde," by Alois Alzheimer [10], in which he reported the behavior of a 51-year-old female patient (Auguste Deter) of the insane asylum of Frankfurt am Main. The patient presented several symptoms that caught Alzheimer's attention, apart from the central nervous system anatomical characteristics. Among others, time and space disorientation, rapid loss of memory, and mood swings were the most prominent symptoms [10]. In relation to pathological features, the observation of something that looked like "thick bundles" [10] of fibrils, later known as senile/amyloid plaques and NFTs [11], transformed AD into a unique condition, distinguishing it from the other neurological conditions known to date.

2.1.1. Symptoms and Diagnosis

Since 1907, clinicians have been trying their best to accurately identify AD-related symptoms and to divide and organize these symptoms in the simplest form. Burns et al. came out in 2002 with three different categories: (1) cognitive deficits that affect memory (amnesia and agnosia), speech (aphasia), and motor behavior; (2) psychiatric symptoms and behavioral disturbances, including depression, anxiety, delusions, and misidentification; (3) difficulties with the daily living activities, such as driving, using the telephone, dealing with money and, later in the disease, all the basic needs (feeding, dressing, toileting) [12–14]. As expected in such a complex condition, a huge symptomatic variation is found in AD patients, although a positive correlation between symptom severity and disease evolution is observed.

Although AD is seen as an elderly disease due to its higher prevalence in the older population (approximately 5.3 million people solely in the US, in 2015) [15], it is also the most frequent form of dementia under the age of 65, with up to 5% of all cases [16]. Of curiosity, every 67 s, one more person is diagnosed with AD, and, by 2050, one new case of AD is expected to develop every 33 s [15]. Due to this disease complexity, diagnosis guidelines had to be established, and for a long time, the main criteria adopted was the one decided at the 1984 consortium, by the National Institute of Neurological and Communicative Disorders and Stroke and the Alzheimer's Disease and Related Disorders Association (NINCDS–ADRDA). These criteria divided AD in three possible diagnosis scenarios, which were possible, probable, or definite AD [17]. More recently, some minor alterations have been proposed in order to

comprise also the stages prior to the clinical observation of symptoms [18, 19], thus prompting three renewed stages: (1) preclinical Alzheimer's disease; (2) mild cognitive impairment (MCI) due to Alzheimer's disease; and (3) dementia due to Alzheimer's disease [20].

Despite all the attentions directed to the establishment of proper diagnostic criteria and guidelines available, the diagnosis of AD is still not an easy task. Actually, a recent meta-analysis showed that the sensitivity and specificity of the clinical diagnosis ranged from 53 to 99% and 55 to 99%, respectively [21]. Although alone it is considered a low value, when combined with other characterizing techniques (as neuroimaging and biomarkers—see Biomarkers section), it is possible to predict/diagnose AD with a high confidence.

Genetically, AD is usually divided in two forms: autosomal dominant familial AD (FAD; predominantly of early-onset—under the age of 65) and sporadic AD (also called and usually associated to the late-onset AD—more than 65 years) [22]. Although extensively used, it is important to point out that this classification is far too simplistic.

Despite all of the effort put into research, the primary event triggering AD remains yet a mystery. Nonetheless, for FAD, several mutations capable of triggering the disease have been identified, especially in three distinct genes: the amyloid precursor protein (APP) [23], the presenilin 1 (PSEN1), and the presenilin 2 (PSEN2) genes [24], in chromosomes 21q, 14q, and 1q, respectively. Although these three genes comprise approximately 55% of all mutations, they are only responsible for less than 1% of all cases of AD (<http://www.molgen.vib-ua.be/ADMutations/>). Contrary to FAD, sporadic AD does not exhibit autosomal-dominant inheritance but up to 60–80% of this form of AD is genetically determined [22].

2.2. The biochemical basis of AD

In spite of its multifactorial etiology, AD is characterized by two specific brain lesions, the amyloid plaques, and the NFTs, considered the hallmarks of AD. Also, associated with these abnormalities, it is often observed severe neural loss and reactive gliosis.

NFTs are filamentous inclusions (intracellular lesions), preferentially observed in pyramidal neurons, which are composed of filamentous aggregates of abnormally hyperphosphorylated microtubule-associated protein tau [25]. Even though NFTs are a hallmark of AD, they are also observed for other neurodegenerative disorders termed tauopathies (e.g., sporadic corticobasal degeneration, palsy, and Pick's disease, progressive supranuclear palsy, Seattle family A, parkinsonism–dementia complex of Guam, and some frontotemporal dementias) [26, 27].

As for amyloid plaques, they can be distinguished in different plaques subtypes, depending on their composition, and being the neuritic and diffuse plaques the two major subtypes in AD. Neuritic plaques are constituted by the 40- and 42-amino acids (aa) A β peptides [28] (A β 40 and A β 42, respectively), surrounded by dystrophic neurites (axons and dendrites), microglia (monocyte- or macrophage-derived cells that reside in the brain), and reactive astrocytes [29]. Diffuse deposits are mainly composed of A β 42 [28] and lack the neuritic and glial components [29], but evolve over time with formation of discrete nodules that eventually become neuritic amyloid plaques [30].

2.3. Amyloid- β precursor protein and A β formation

The amyloid- β precursor protein (APP) is a transmembrane receptor expressed ubiquitously in both neuronal cells and extra-neuronal tissues [31]. In humans, the APP gene is located in the chromosome 21, explaining partially the increased risk for Down syndrome patients to develop AD, and is composed of 18 exons [32]. Three major isoforms are expressed by alternative splicing: APP770 (full length), APP751 (lacking exon 8), and APP695 (lacking exon 7 and exon 8) [23, 31, 33]. APP belongs to a highly conserved family of type 1 transmembrane glycoproteins that extends also to invertebrate species, including the homologous: APL-1 (*Caenorhabditis elegans*), APPL (*Drosophila*), APLP1, and APLP2 (in mammals, besides APP) [34], and appa and appb (zebrafish) [35]. Following translation, APP is trafficked through the endoplasmic reticulum (ER), Golgi and trans-Golgi network (TGN), where it suffers specific endoproteolytic cleavages [33] that will originate several APP metabolites, among them the A β peptide. After reaching the membrane surface, APP can still undergo clathrin-mediated endocytosis and then be recycled to the surface again [36], during which A β can also be produced [37].

The 4 kDa A β peptide was first isolated and sequenced by Glenner and Wong, in 1984 [7] and can be found in the plasma and cerebrospinal fluid (CSF) of healthy humans and other mammals [38]. It was described as a 24 aa peptide but later, sequencing analysis revealed that the peptide could actually comprise 36–43 aa [39], being the two major species A β 40 and A β 42. In healthy individuals, these two forms make up about 90 and 10%, respectively, of the A β peptides that are normally produced by brain cells [40]. Despite the small difference in size and sequence of the various isoforms, they differ greatly in properties; for example, A β 42 is more hydrophobic, thus, more prone to aggregation (compared to the less hydrophobic A β 40). In fact, it readily aggregates *in vitro*, being considered the more amyloidogenic and hence pathogenic species [41].

2.3.1. Towards amyloid or not?

APP processing can originate different metabolites that bear very different physiological functions, depending on the proteolysis pathway adopted: the amyloidogenic or non-amyloidogenic pathway (**Figure 1**). In the non-amyloidogenic pathway, APP is firstly cleaved by the α -secretase, a zinc metalloproteinase of the ADAM family [42], followed by the action of γ -secretase. The latter is a high molecular weight complex of four proteins: presenilin 1 or 2 (PSEN1, PSEN2), nicastrin (NCT) [43, 44], anterior pharynx-defective 1 (APH1), and presenilin enhancer 2 (PEN2) [45]. The cleavage by the α -secretase (at Lys687 of APP770) [46], within the A β domain, abrogates the production of A β , resulting in the release of a large soluble ectodomain of APP (sAPP α , ~100 kDa), leaving behind a 83-residue carboxi-terminal fragment (CTF α , of ~10 kDa) [47]. Then, γ -secretase cuts the CTF α , liberating the extracellular p3 peptide and the 50 aa APP intracellular domain (AICD, of ~6 kDa) [48].

On the other hand, as suggested by its name, the amyloidogenic pathway gives rise to the amyloidogenic A β peptide, and similar to the previous pathway, it consists of two sequential cleavages, first by the β -secretase (beta-site APP-cleaving enzyme 1–BACE-1), and then by γ -secretase. The first protease cleaves APP at Met671 [49], releasing the large soluble ectodomain

sAPP β [33]. The remaining 99 aa CTF β (of ~12 kDa) [50] is then cleaved by the γ -secretase, in the membrane, and originates, as said above, the A β peptide and the AICD [48]. This process generates different A β species, with variable hydrophobic C-termini (related to the γ -secretase cleaving site), that present different propensity to oligomerize [51] and, consequently, to form the amyloid plaques. Noteworthy, AD-linked mutations in the PSEN1 and PSEN2 proteins, particularly important in the case of FAD, influence γ -secretase-mediated processing of APP, and selectively enhance A β 42 production compared to A β 40 [52].




Figure 1. The amyloidogenic and non-amyloidogenic pathways of APP. In the non-amyloidogenic pathway, APP is cleaved by the α -secretase releasing the sAPP α neuroprotective N-terminal fragment, which contains part of the A β sequence. The 83 aa APP fragment (C83) then suffers the action of the γ -secretase, liberating the p3 peptide and the AICD fragment. The amyloidogenic pathway involves the sequential cleavage of APP by the β -secretase which releases the soluble ectodomain sAPP β . The remaining C99 fragment of APP is then cleaved by the γ -secretase, resulting in the formation of the A β peptide. Due to its high propensity to aggregate, A β peptide oligomerizes, accumulates and forms amyloid senile plaques, in turn leading to the described alterations of AD. Current therapeutic approaches in AD include: (1) inhibition of β - and γ -secretases, (2) improving A β clearance, and (3) amelioration of inflammation and synaptic dysregulation.

Although tightly related to AD onset, APP processing is a normal metabolic event and A β is a normal product of cellular metabolism throughout life, circulating as a soluble peptide in biological fluids [53]. Plus, A β deposition can also be found, together with NFTs, in the brain of non-demented elderly people [54].

2.4. Alzheimer's disease hypotheses

AD is one of the human diseases with the highest number of hypothesis formulated trying to explain its pathogenesis, with very different and plausible molecular mechanism to back them up. Within the list, the amyloid cascade hypothesis stands out, together with the so-called "tau

and tangle” hypothesis, strengthened by the fact that they are based in the two hallmarks of AD.

2.4.1. *Amyloid cascade hypothesis*

Since its formalization, in 1992 by Hardy and Higgins, the amyloid cascade hypothesis has had a prominent role in explaining the etiology and pathogenesis of AD. They suggested that amyloid deposition was the primary influence driving AD pathogenesis [55], due to two key observations: the detection of A β as the main constituent of amyloid plaques, and the discovery of mutations in the APP, PSEN1 and PSEN2 genes associated with FAD [56]. This hypothesis stated that a dysregulation in APP processing or A β clearance would provoke an increase in the A β 42/A β 40 ratio, which would promote aggregation, accumulation, and plaque formation. In turn, this would be responsible for the subsequent pathology (including tau aggregation, phosphorylation, neuronal attrition, and clinical dementia) [57]. Due to its inability to entirely explain AD pathogenesis, especially by the lack of correlation between plaque burden and clinical manifestations [58], this hypothesis has been upgraded over the past few years. Scientists started to divert their attention from the effects of the amyloid deposits, to study the other forms (monomers, oligomers, and protofibrils—usually shorter and thinner than mature fibrils) [59] of A β peptide-induced neurotoxicity. Some studies suggest that A β toxicity functions in a plaque-independent manner, indicating that oligomeric intermediates present higher toxicity to the cells [60] and that activation of signaling pathways due to intraneuronal accumulation of A β oligomers is responsible for tau hyperphosphorylation and subsequent deposition [61]. Several explanations have been proposed, with some defending that oligomeric toxicity is related to a greater capacity for diffusion and a larger collective surface area for interacting with neuronal and glial cells [60], while other proposed that it is not related to a specific prefibrillar aggregate (dimer, trimer, and so on) but rather to the propensity that each species has to grow and undergo fibril formation [62].

A more consensual vision about A β is that it possesses a dual role: On one hand, it can be a neurotrophic agent or a neuroprotector against excitotoxicity (by activating the phosphatidylinositol-3-kinase (PI-3K) pathway); on the other hand, an inducer of neuronal degeneration (at high concentrations) in mature neurons [63].

Other interesting and amyloid cascade-opposite hypotheses have been proposed, stating that A β should not be seen as the initiating factor for neurodegeneration in AD, but instead, its deposition is nothing more than a protective mechanism to neuronal insult, in which A β binds and removes harmful substances by blocking them in plaques [64, 65].

2.4.2. *Tau hypothesis*

“Tauists” defend a collection of ideas that maintain the primacy of NFTs formation as the AD-causing event, which Mudher and Lovestone designated as the “tau and tangle hypothesis” [57]. It started to emerge due to solid evidence that amyloid plaques do not account for the complex pathophysiology of AD [66], opposed to the observed highly positive correlation between NFTs and cognitive deficits [67]. It argues that in AD the normal role of tau (micro-

tubules stabilization) is impaired and that NFTs accumulate and occupy much of the neuron, resulting in neuronal death. This was supported by the visualization of the extracellular tangles in the shape of neurons, abundant in the late stages of disease [57]. Also, the discovery of mutations within the tau gene that cause fronto-temporal dementia and parkinsonism linked to chromosome 17 (FTDP-17), demonstrating that tau dysfunction, in the absence of amyloid pathology, was enough to cause neuronal loss and clinical dementia [68], further strengthened this hypothesis. However, tau mutations do not originate amyloid plaques, whereas APP and presenilin gene mutations give rise to amyloid and tau depositions, strongly evidencing that amyloid pathology is upstream of tau pathology [57]. More recently, a more embracing tau hypothesis was proposed, in which a series of damage signals ($A\beta$ oligomers, oxygen-free radicals, iron overload, cholesterol levels in neuronal rafts, low-density lipoprotein (LDL) species and homocysteine, among other) trigger, by innate immunity, the activation of microglial cells with the consequent release of pro-inflammatory cytokines that modify neuronal behavior through anomalous signaling cascades, which finally, promote tau hyperphosphorylation [66]. In turn, tau hyperphosphorylation will contribute to further activation of microglial cells and stimulation of the deleterious cycle, which will lead to progressive neuronal degeneration [66]. The degree of tau phosphorylation in the AD brain is reasonably well correlated with the severity of AD symptoms; however, fetal tau, a much more phosphorylated form of tau than adult tau, does not induce AD-like pathology [69]. In summary, there is no direct evidence for the neurotoxicity of hyperphosphorylated tau (as in the case of $A\beta$ toxicity).

2.4.3. "Other" hypothesis

2.4.3.1. GSK-3 hypothesis

Glycogen synthase kinase-3 (GSK-3), a multi-tasking kinase with major roles in brain signaling, has been recently proposed as a central player in AD pathology. This was supported by observing that the deregulation of this protein is responsible for many of the pathological hallmarks of the disease, in both sporadic and familial AD cases [70]. It was suggested that the hyperactivity of GSK-3 β , the most abundant of two isoforms (GSK-3 α and GSK-3 β) expressed in neurons, is intimately involved with cognitive impairment [71], $A\beta$ production [72], tau hyperphosphorylation [73], acetylcholine synthesis [74], neuronal death [75], and neuroinflammation [76] in AD. Furthermore, regarding $A\beta$ interaction, it was observed that $A\beta$ also regulates GSK-3 β activity [77, 78] making it difficult to establish which event is located upstream. Thus, this hypothesis is seen as an integration and extension of the amyloid cascade hypothesis, still conferring to $A\beta$ a central role in AD pathology. Although GSK-3 modulation appears to be an excellent therapeutic approach, no effective result has been observed in trials, perhaps due to its activity in multiple targets.

2.4.3.2. Oxidative stress/mitochondrial hypothesis

The brain is especially vulnerable to free radical damage as a result of its high oxygen consumption rate, abundant fatty acids content, and the relative low levels of antioxidant enzymes

[79]. The most appealing feature of the oxidative stress hypothesis is its slow and cumulative damaging nature that could, over time, account for the late life onset and slowly progressive nature of AD, and neurodegeneration in general [80]. Also supporting this hypothesis is the suggested unbalanced levels of heavy metals in the brain, among others, iron (Fe), copper (Cu), aluminum, and mercury, which function as catalysts for oxygen free radical generation [80]. There has been high controversy in the measurement of these elements, especially Fe, but most studies reveal an apparent unbalance in AD brains compared to controls. In a recent study, Fe was found significantly increased in patients with severe AD [81] (as previously reported [82]). Nonetheless, some consider that this “accepted” elevation, even if significant in some studies of AD pathology, does not account for brain degeneration, and so, presents itself as a misleading therapeutic target with considerable risks for patients (reviewed in [83]). As for Cu, it has been shown to be decreased in AD brains [81], which at first goes against the oxidation hypothesis. More recently, and also in the presence of some contradictory results [84], copper was found to bind strongly to A β aggregates, inhibiting *in vitro* amyloid fibril formation [85]. In addition, and when bound to the aggregates, copper exhibited a redox role, by degrading hydrogen peroxide [86]. Protein and DNA oxidation (in particular mitochondrial DNA), and lipid peroxidation (which affects the phospholipid-rich membrane) were also found to be increased in AD brains [80].

Another hypothesis intimately related to the brain redox status is the mitochondrial cascade hypothesis [87]. Mitochondria are considered the cell “powerhouses”; however, when in a non-physiological energy production, they can provoke severe damage by increasing the reactive oxygen species (ROS). Curiously, mitochondria are the first target of ROS, suffering DNA oxidation, which may lead to a further increase of ROS production, generating a vicious cycle [88]. The authors of this hypothesis state that sporadic and autosomal dominant AD are not etiologically homogeneous and that mitochondrial dysfunction works as a link for both. Very briefly, in autosomal dominant forms, A β -induced mitochondrial dysfunction leads to the other AD-characteristic histopathologies, while in sporadic AD, mitochondrial malfunction induces the AD pathologies, including processing of APP to A β [89].

2.4.3.3. *The cholinergic hypothesis*

The cholinergic hypothesis was proposed after the observation of a decrease of choline acetyltransferase (ChAT) in AD patients [90]. It states that the loss of cholinergic cells in the septal nuclei and basal forebrain (described in patients with advanced AD [91]) compromises the innervation of the cerebral cortex and related structures, which play an important role in cognitive functions, especially memory [92]. More recent studies have suggested a bidirectional pathway linking A β toxicity in cholinergic dysfunction and the interaction of cholinergic regulatory mechanisms in the processing of APP [93]. Discrediting this hypothesis, later studies showed that the cholinergic degeneration [94] and the decrease in ChAT enzyme activity [95] are not observable in the early stages of disease. This was accompanied by the fact that treatment with acetylcholinesterase inhibitors does not offer long term cure, although it has shown consistent, despite modest, benefits in symptoms improvement [96]. Nonetheless,

some studies point out that compensatory mechanisms could overcome the cholinergic defects, disguising its effects in early stages of AD or mild cognitive impairment [97].

2.4.3.4. Calcium hypothesis

The calcium hypothesis was first introduced by Khachaturian in 1982, stating that cellular mechanisms which maintain the homeostasis of cytosolic Ca^{2+} play a key role in brain aging and that sustained changes in Ca^{2+} homeostasis could provide the final common pathway for age-associated brain changes [98], or in this case, be the proximal cause of neurodegeneration in AD. Out of curiosity, this proposal was purely speculative at the time, only sustained on circumstantial evidence from a handful of studies [98]. It was observed that the persistent elevation of the levels of Ca^{2+} leads to the constant elimination of newly acquired memories, due to a stimulation of long-term depression mechanisms [99]. In fact, calcium signaling dysfunctions occur during the initial phases of the disease, and even before the development of pronounced symptoms [100]. However, whether it is calcium dyshomeostasis that provokes $\text{A}\beta$ production and accumulation [101], or vice-versa [102], remains to be elucidated. Either way, in addition to the $\text{A}\beta$ unbalance, increased levels of Ca^{2+} promoted protein tau hyperphosphorylation [103]. It was also suggested that amyloid oligomers induce membrane permeabilization, leading to increased intracellular Ca^{2+} concentration [104]. Nevertheless, there is some disagreement as to the mechanism by which amyloid oligomers increase intracellular calcium.

2.5. Biomarkers and risk factor

Despite of the extensive Knowledge on the causative gene mutations responsible for familial AD, the sporadic (non-genetic) form of this disease, which results from the diverse interactions between genetic and environmental factors, is still lacking characterization. Thus, researchers are continuously looking for specific molecules that should be altered exclusively in AD, and preferentially in the asymptomatic period. This, combined with the meticulous description of the patient risk factors, may give the opportunity for an early action and increase the success rate of therapeutics.

2.5.1. Biomarkers

By definition, and according to the International Programme on Chemical Safety, biomarker is "any substance, structure, or process that can be measured in the body or its products and influence or predict the incidence of outcome or disease" [105]. The search for early AD biomarkers has been highly targeted over the last years, as investigators believe that the generation of an effective treatment for AD is only possible if the disease is detected at very early stages. Thus, the discovery of biomarkers is of extreme importance for the early diagnosis of AD and even for predicting the conversion of MCI into AD patients.

The search for solid AD biomarkers started with those who seem to be altered in this condition when compared to normality. Several studies showed that the combination of CSF total- and phospho-tau, and CSF and plasma $\text{A}\beta_{42}$ is able to predict with increased sensitivity

and specificity the development of AD in patients with MCI [21, 106, 107], in addition to the already established role of the ApoE- ϵ 4 isoform [108] (major susceptibility gene—see Section 2.5.2. Risk Factors). Studies in FAD-causing mutations carriers showed increased levels of CSF total-tau and plasma A β 42, although having reduced CSF A β 42 levels, at least 10 years before the symptoms establishment [106]. Also, the ratio of A β 42/A β 40 in CSF and plasma was found decreased, respectively, for non-demented mutation carriers [109] and cognitively normal elders (which evolved to MCI or AD) [110]. Other CSF biomarkers, such as BACE-1 [111] and sAPP α / β [112], are also suggested; however, independent studies were not consistent [106], possibly due to technical difficulties in the biomarkers quantification [113]. Our group has also proposed TTR as a biomarker in plasma, demonstrating a negative correlation with AD severity [114], which supported prior observations for CSF levels. In that study, the authors also considered TTR a selective biomarker for AD [115]. Other studies contradict this idea suggesting that TTR potential as biomarker raises some doubt since its levels appear to fluctuate substantially within a single individual over a 2-week interval [116].

In addition to the so-called fluid biomarkers, physicians have at their disposal powerful imaging technology. With the improvement in PET (positron emission tomography) and MRI (magnetic resonance imaging) spectroscopy resolution, neuroimaging has been gaining importance and increasing the confidence in AD diagnosis. Due to the possibility of using specific tracers, such as a derivate of thioflavin T that crosses the blood–brain barrier (BBB) and binds selectively to A β (C11-labeled Pittsburgh Compound B–PiB), it is possible to identify amyloid deposition in the brain in vivo [117]. In 2004, Klunk and colleagues performed the first study of brain amyloid imaging (BAI) using the PiB compound, in which they showed a robust relationship between amyloid deposition and PiB retention [118]. The combination of increased BAI signal, low CSF A β 42, and high CSF p-tau in a subject with dementia is seen as a “definite” diagnosis of AD. Furthermore, BAI and CSF profiles can be used to predict patients with MCI who will progress to frank dementia with high degree of confidence [119]. Despite the value of this compound, the resulting data should be subjected to careful analysis since healthy subjects also present amyloid deposition, hindering the differentiation between symptomatic AD and asymptomatic controls with amyloid plaques [120]. MRI (structural evidence) is also a common technique often used, alone or in combination with CSF tau and A β 42, to predict development of AD [121]. Curiously, a recent study showed that the MRI together with PiB-PET makes the best combination of biomarkers, thus showing the best AD predictive value [119].

2.5.2. Risk factors

2.5.2.1. Environmental factors

As Stephen King wrote in *The Gunslinger*: “Time’s the thief of memory,” and so, the most worldwide accepted (and intuitive) risk factor is aging. In every species, age brings a slowing of brain function [122], thus preventing the brain to properly respond/recover from insults. The increasing of life expectancy, in addition to the increasing of population (attributed to the

postwar “baby boom”), turned aging in a major risk factor. Also gender appears to play a role in disease development, since data show that women are more prone to this disease than men. Although women present higher life expectancy, also in younger study groups (60–80 years), where differences in death rate are insignificant, women present higher incidence of cases [123]. The specific mechanism is unknown; however, several factors have been proposed to influence, such as: age-related sex hormone reduction, risks of other diseases (diabetes, depression, cardiovascular disease), and differences in brain anatomy and metabolism [124].

The cardiovascular risk factors, which appear sometimes as a distinct group of factors, include diabetes mellitus [125], overweight [126], hypertension [127], and high cholesterol levels [128]. Individually or in cooperation, these factors increase the predisposition for cognitive decline. The midlife control of the above cardiovascular factors has been associated with a reduction in white matter lesions in late life [127]. As for cholesterol, results appear to be inconsistent; however, lipid-lowering treatments present benefits against white matter lesions [126].

Contrasting with the previous risk factors, wine consumption, coffee consumption, the use of non-steroidal anti-inflammatory drugs (NSAIDs), and physical activity are associated with reduced risks, thus showing some protective effects [129].

2.5.2.2. Genetic factors

ApoE exists as three isoforms $\epsilon 2$, $\epsilon 3$ and $\epsilon 4$, with $\epsilon 3$ having the highest prevalence. As Corder and colleagues stated, in 1993, ApoE plays an important role in AD, with the risk of developing disease increased in carriers of the ApoE- $\epsilon 4$ allele, such that a double dose of this allele was nearly enough to cause AD by the age of 80 [130]. Despite the broad molecular evidence about ApoE role in AD, its genetic variation is also present in other kinds of neurological disorders, including Parkinson’s disease and multiple sclerosis [22]. In 2009, three novel AD genes were identified, presenting high degree of association: CLU (clusterin or apolipoprotein J), CR1 (complement component (3b/4b) receptor 1), and PICALM (phosphatidylinositol-binding clathrin assembly protein). Down’s syndrome (or trisomy 21) is also considered a genetic risk factor. This condition is the result of a third copy of the chromosome 21, which coincides with the location of the APP gene [131], giving rise to an increased accumulation of A β .

2.6. Drugs and treatments

Due to the complexity of AD, a vast number of targets and pathways may be chosen to intervene. Cholinergic degradation inhibitors, immunotherapy, secretase inhibitors, anti-inflammatory drugs, and tau- and A β -deposition interfering drugs (**Figure 1**) are but a few examples of the many classes of drugs that are being tested at the moment.

The first drugs developed for AD, acetylcholinesterase inhibitors (AChEI), aimed at increasing acetylcholine levels (see “Cholinergic hypothesis”). Currently, FDA has five drugs approved for the “treatment” of AD in the initial stages: 4 AChEI (Donepezil, Rivastigmine, Galantamine, and Tacrine) and 1 NMDA receptor antagonist (Memantine) (<http://www.alzforum.org>). As referred above, they treat and ameliorate the symptoms, but do not cure.

In 2010, Rinne and colleagues showed for the first time target engagement from a disease-modifying drug in humans, using the monoclonal anti-A β antibody bapineuzumab [132, 133]. This study showed a reduction of fibrillar amyloid in the brain of AD individuals, but did not improve cognition and stopped at phase 3 [133]. Crenezumab, another anti-A β , was selected for pre-symptomatic treatment trials of Colombian mutant PSEN1 kindred [133], however, showed extensive cross-reaction with non-A β related proteins [134]. Intravenous immunoglobulins (IVIG) have also been proposed as a potential treatment, based on the hypothesis that IVIG contain naturally occurring antibodies that specifically promote clearance of A β peptides from the brain [135].

β - and γ -secretase modulators [133], A β and tau deposition modulators (e.g., scylloinositol and methylene blue, respectively [39]), and molecules addressing oxidative damage (resveratrol) are also potential drugs under study. Recently, deep brain stimulation (DBS) performed by the group of Dr Lozano in AD patients showed promising results, with improvements and/or slowing in the rate of cognitive decline [136], increased hippocampus volume and glucose metabolism [137]. The authors propose that DBS is able to influence the structure of the brain and that hippocampal atrophy can potentially be slowed, suggesting restorative properties to DBS [137].

A general recommended therapy is a good diet and a healthy lifestyle, in order to control cardiovascular risk factors, decreasing cerebrovascular events. No effective treatment has been found, thus increasing the interest for the early diagnose of AD, to allow a more effective and early stage intervention.

3. Blood–brain barrier and Alzheimer’s disease

The BBB is a profoundly specialized brain endothelial structure of the differentiated neurovascular system. These specialized endothelial cells interacting with astrocytes, microglia, and pericytes, confine components of the circulating blood from neurons. Furthermore, the BBB controls the chemical composition of the neuronal environment which is required for the functioning of neuronal circuits, synaptic transmission, synaptic remodeling, angiogenesis, and neurogenesis. BBB malfunction, through the disruption of the tight junctions (TJs) and alteration of the transport of molecules between blood and brain, brain hypoperfusion, and inflammatory responses, may begin or contribute to the process of different diseases such as AD, Parkinson’s disease, amyotrophic lateral sclerosis, multiple sclerosis. These data support developments of new therapeutic strategies for the neurodegenerative disorders focused at the BBB [138].

3.1. The blood–brain barrier: the earliest findings

In 1885, when scientist Paul Ehrlich injected trypan blue dye into the bloodstream of mice, he noted that the stain colored all of the animal organs, except the brain [139]. In 1913, one of the Ehrlich's students, Edwin Goldmann, performed a follow-up experiment by injecting the same dye into the brain of mice. He observed that injection of trypan blue directly into the CSF

stained all cell types in the brain but failed to penetrate into the periphery [140]. Although aiming at finding new compounds that could attack disease-causing microbes, these experiments suggested a physical barrier between the brain and the blood, becoming the stepping stones on BBB research. Lewandowsky was the first to use the term blood–brain barrier while studying the limited permeation of potassium ferrocyanate into the brain [141]. However, it took until the 1960s for the specific anatomy of the network of brain–blood vessels comprising the BBB to be glimpsed [142]. Using electron microscopy, in 1967, Reese and Karnovsky showed that the BBB is localized at the level of TJs between adjacent brain endothelial cells [143, 144].

It is now known that the brain of mammals is separated from the blood by the BBB, localized to the brain capillaries and pia-subarachnoid membranes, and by the blood–CSF barrier, confined to the choroid plexus [145]. At the junctional complex formed by the TJs and adherens junctions (AJs), brain endothelial cells are connected to each other [142]. Transcellular transport at the BBB occurs through several mechanisms including passive and active transport through different receptors and transporters. Interestingly, endothelial cells, pericytes, and astrocytes present at the BBB also express several enzymes such as cholinesterase, aminopeptidases and endopeptidases that alter endogenous and exogenous molecules, which can negatively influence neuronal function. This produces a metabolic barrier which protects the central nervous system (CNS) [146].

3.2. The neurovascular unit at the BBB; endothelial cells, pericytes, and glial cells

3.2.1. Endothelial cells and pericytes

A cerebral capillary lumen is enclosed by a single endothelial cell. Anatomically, the BBB endothelial cells are distinguished from those at the periphery by enhanced mitochondrial content [147], loss of fenestrations [148], least pinocytotic activity [149], and the presence of TJs [150]. The BBB tightly sealed monolayer of endothelial cells usually prevents the free exchange of solutes between blood and brain [151] except for the lipid-soluble molecules smaller than 400 Da with less than nine hydrogen bonds, which can cross the BBB without any support, via lipid-mediated diffusion [146].

As for pericytes (granular or filamentous), they are connected to the abluminal membrane of the endothelial cells [152]. Pericytes and endothelial cells are ensheathed by the basal lamina, composed of collagen type IV and other extracellular matrix proteins [153]. However, there is not much known about the involvement of pericytes at the BBB, but the addition of pericytes to co-cultures of endothelial cells and astrocytes has been shown to stabilize capillary-like structures [154]. In conditions associated with increased BBB permeability such as hypoxia or traumatic brain injury, pericytes have also been shown to migrate away from brain microvessels [155, 156]. Moreover, pericyte-derived angiopoietin provokes expression of TJs such as occludin in endothelial cells [157], confirming that pericytes are involved in the maintenance of the barrier properties in the cerebral endothelium [142].

3.2.2. *Glial cells*

Astrocytes are another cell type present in the neurovascular unit. These cells are usually located between neurons, pericytes, and capillary endothelium, and communicate with these cells via their several foot processes [138]. It is long believed that astrocytes are crucial in the development of the BBB properties [158]. It has been shown that co-culture of brain endothelial cells with astrocytes can enhance TJs of the brain endothelium [159]. Furthermore, endothelial cultures incubated with astrocyte-conditioned media have shown to improve BBB characteristics *in vitro* by *de novo* protein synthesis of γ GTP in cerebral microvessel endothelial cells, which indicate that the glial cells may induce the cerebral capillary endothelial cells to express differentiated properties which allow the endothelium to function as the blood–brain barrier [160]. Also, astrocytes have been shown to regulate cerebral microvascular permeability [161], via dynamic calcium signaling between astrocytes and the endothelium [162]. This has been explained by Zonta et al., whereby an increase in neuron-induced astrocyte calcium promotes secretion of vasodilatory substances from perivascular astrocyte endfeet, resulting in improved local blood flow [163]. This work constituted a breakthrough in the knowledge of both astrocyte function and regulation of the activity-dependent cerebral blood flow [164].

Finally, another glial cell type present in the BBB is microglia which migrates from the yolk sac into the CNS parenchyma during embryogenesis [165]. Microglia performs critical functions in innate and adaptive brain immune responses. Microglia, when activated, transforms from “ramified” to an “ameboid” and ultimately to a “phagocytic” form. This evolution is correlated with alterations in expression of surface antigens and cytokines release [138]. Studies have shown that perivascular microglial cells are derived from bone marrow [166].

In vivo studies demonstrated that resident microglia cells in the brain parenchyma also communicate with CNS microvessels. This may suggest that microglia plays an important role in regulating BBB features during embryogenesis and diseases [167], besides an indisputable function in CNS development and homeostasis [168]. Furthermore, it has been shown that during embryonic stages of CNS vascularization, stabilization, and fusion of brain endothelial cells are mediated by resident microglial cells [167]. Also, interestingly, it has been shown that specific depletion of microglia results in reduced vessel density in a mouse model of choroidal neo-vascularization [169]. There are studies which suggest that microglial activation may be related to BBB disruption [170] apparently by producing ROS, through NADPH oxidase, which in turn impairs BBB function. Furthermore, TNF- α released from activated microglia has shown to affect BBB integrity, as permeability of endothelial cells co-cultured with microglia, was increased when microglial cells were activated by LPS and it was blocked by a neutralizing antibody against TNF- α , indicating that TNF- α contributes to BBB dysfunction [171]. The post-traumatic inflammatory response is shown to be associated with expression of cytokines such as IL-1 β or metalloproteinases particularly MMP2, 3 and 5 produced mainly by microglia at the lesion site. It has also been demonstrated that these proteins can cause disruption of the basal lamina and/or redistribution and degradation of the TJs complexes [172, 173] resulting in BBB breakdown and neurological disorders after traumatic injury [174].

3.3. Junctional complexes and cytoskeleton linked proteins at the BBB

The presence of junctional complexes is one of the main characteristics of inter-endothelial space of the cerebral microvasculature which include TJs, AJs, and possibly gap junctions. Both TJs and AJs restrict permeability across the endothelium, whereas gap junctions (if present) mediate intercellular communication [175].

3.3.1. Tight junction proteins

Occludin, the first integral membrane protein within the TJ family identified [176], is a 60–65 kDa protein with four transmembrane domains which is highly expressed in the cerebral endothelium [177, 178], whereas it is much less present in non-neural endothelial cells [179]. A construct with a deletion in the N-terminal of occludin showed a considerable effect on the TJ integrity [180]. Deletion of occludin in mice has been shown to cause a complex phenotype and postnatal growth retardation [181]. However, occludin functions are not limited to its role as a TJ protein. For instance, there are studies demonstrating that occludin can regulate epithelial cell differentiation [182] and control cell apoptosis in mouse hepatocytes [183]. Also, it has been shown, in a mouse model of multiple sclerosis, that occludin dephosphorylation leads to noticeable signs of disease which occur just prior to apparent changes in the BBB permeability [184]. In cerebral ischemia, occludin and other TJs were shown to be vulnerable to attack by matrix metalloproteinases [185].

The claudin family, 20–24 kDa proteins, includes more than 20 members that build TJ strands through homophilic interactions [186]. Claudin 3, 5, and 12 play important roles at the BBB [187, 188]. Each claudin regulates the diffusion of a group of molecules of a certain size. For instance, mice neonates with deletion of claudin-5 die due to a size-selective loosening of the BBB for molecules smaller than 800 Da [188]. It is speculated that claudins determine the primary “seal” of the TJ and occludin functions as an additional support structure [142]. It has been demonstrated that overexpression of claudin species can induce development of TJ-like strands, while expression of occludin does not lead to the formation of TJ; rather, occludin only localizes to TJ in cells that have already been transfected with claudins [189].

In addition to the claudin/occluding proteins, junctional adhesion molecules (JAMs) perform an important role in the organization of TJ assembly as it has been reported that these proteins can reduce cell permeability and enhance resistance to macromolecules [190]. JAM-1, a 40-kDa member of the IgG superfamily composed of a single membrane-spanning chain with a large extracellular domain [191], is postulated to mediate the attachment of neighboring cell membranes via homophilic interactions [192]. However, there is not much known about JAMs exact function in the mature BBB, but in the rat cortical cold injury model, a characterized *in vivo* model of BBB breakdown it has been shown that endothelial JAM-1 is significantly reduced which strengthens the idea that JAM-1 contributes to TJs integrity [193].

3.3.2. Adherens junction proteins

The first component of AJs is the vascular endothelial (VE)-cadherin, an endothelial-specific integral membrane protein which is linked to the cytoskeleton via catenins [175] and mediates

cell-cell adhesion via homophilic interactions between the extracellular domains of proteins expressed in adjacent cells [194]. In vitro and in vivo studies have shown that VE-cadherin is required for the cells to regulate vessel maintenance, that is, for the correct organization of the new vessels and for the endothelial integrity in the quiescent vessels [195]. Various mechanisms have been suggested for how VE-cadherin regulates endothelial functions; such as direct activation of signaling molecules with a role in survival and organization of the actin cytoskeleton and regulation of gene transcription cofactors and formation of complexes with growth factor receptors [138].

Even though AJs are required at the BBB to decrease the endothelial permeability [196], it is principally the TJs that present the low paracellular permeability and high electrical resistance [142, 197].

Another component of AJs is Platelet endothelial cell adhesion molecule 1 (PECAM-1), an integral membrane protein of the Ig superfamily with six extracellular domains, a short transmembrane, and a large cytoplasmic domain which is highly expressed in blood and vascular cells especially endothelial cells [198]. PECAM-1 plays a major role in the migration of leukocytes across endothelium [138] and contributes to the steady-state barrier function of endothelial cells. It also functions as a mechano-sensor and stimulates reconstruction of the barrier integrity following perturbations, including the BBB [199, 200]; all physiologic processes that rely on the junctional integrity and signaling [198].

3.3.3. Associated proteins

There are various proteins in the cytoplasm that associate with the transmembrane components of the TJ [142]. Multi-domain scaffolding proteins of membrane-associated guanylate kinase-like homolog family, including zonulae occludentes (ZO) proteins, such as ZO-1 and ZO-2, are characterized for their contributing to the cytoskeletal anchorage for the transmembrane TJ proteins, binding to the claudins, occludin, and actin [201, 202] and controlling the correct distribution of claudins [203].

ZO-1, a 220-kDa protein, is one of the first and best-studied proteins in TJs [204]. It is expressed in epithelial and endothelial cells and even other cells which do not have TJs [205]. It has also been observed to be associated with AJs [206] and gap junction proteins [207]. ZO-1 is located just below the TJ membrane contact points and has been found to be important for both function and stability of TJs. ZO-1 dissociation from the junctional complex is shown to be correlated with increased permeability in the BBB in vitro model [208]. ZO-2, a 160-kDa protein with a high sequence homology to ZO-1 [209], interestingly, has been demonstrated to function redundantly with ZO-1, replacing it and promoting the formation of TJ in the cells lacking ZO-1 [210].

Another component present in the junctional complexes is actin. Although actin has been basically considered structural in function, it is now obvious that the anchorage of AJs and TJs to the actin cytoskeleton is critical for both barrier stability and also for the regulation of cell polarity, cellular movement, fluid sensing, and cell-contact inhibition [211]. Studies using mice lacking the actin-binding protein dystrophin have demonstrated increased brain vascular

permeability due to disorganized α -actin cytoskeleton in endothelial cells and astrocytes [212]. These findings demonstrate that properly arranged actin filaments and their binding to the TJ and/or AJ proteins are critical for normal barrier function. Studies have also shown that HIV-1 gp120 and alcohol are able to alter the cytoskeleton and induce stress fiber actin formation, causing increased permeability of the human BBB endothelium [213]. It has been suggested that alcohol-mediated changes in the brain endothelial cells (BEC) monolayers may increase diffusion of plasma components and viral penetration across the BBB, and therefore, especially at levels attained in heavy drinkers, accelerate HIV-1 penetration into the brain [138].

3.4. Transport at the BBB

Transcellular transport at the BBB happens through several mechanisms. Small lipophilic molecules can access the brain by passive diffusion. Brain CSF bulk flow mediates transport of molecules with different sizes into the CSF at a slow rate [214]. For potentially toxic molecules and metabolic waste products, the CSF works as a sink. These molecules are then eliminated from the CSF back into the blood by active transport or facilitated diffusion across the choroid plexus epithelium, or by vacuolar transport across the epithelial arachnoid granulations [138]. Efflux pumps return many undesired molecules back to the blood to regulate passive transport into the brain [215]. The flow of the plasma oxygen and carbon dioxide across the BBB is diffusive. Therefore, oxygen supply and carbon dioxide elimination are blood-flow dependent, so the gas transport is sufficient as long as cerebral blood flow is within physiological limits [216].

Small polar molecules, such as glucose, amino acids and nucleosides, can pass the BBB by carrier-mediated transport. These carriers can be multi-ligands or specific to only one molecule, such as GLUT1 glucose transporter, the L1 large neutral amino acid transporter, and the CNT2 adenosine transporter which have been cloned from BBB-specific cDNA libraries [146]. The direction of the concentration gradient is usually from blood to brain, regulated by brain demands and the concentration of metabolites in plasma [138]. Ion transporters such as the sodium pump localized on the abluminal membrane are important to sustain the high-concentration gradient for sodium at the BBB, so that sodium-dependent transport can happen. Also, sodium–hydrogen exchanger expressed at the luminal membrane and chloride–bicarbonate exchanger expressed at both sides [217] play significant roles in regulating intracellular pH in the endothelium.

Moreover, large solutes, such as proteins and peptides, are transported across the BBB by receptor-mediated or adsorption-mediated endocytic transport [218, 219]. Analysis of the rat BBB transcriptome has shown that 10–15% of all proteins in the neurovascular unit are transporters which emphasize the critical role of these molecules at the BBB [215, 220]. As a consequence of this controlled transport, the concentration of amino acids and proteins can suffer considerable variations, whereas relatively small differences exist in the concentration of ions between blood and CSF [221].

In the BBB, the ABC transporters for efflux are permeability glycoprotein (P-gp, multidrug resistance protein, ABCB1), the multidrug resistance-associated proteins and breast cancer resistance protein [222, 223] whose major role is to operate as active efflux pumps, transporting

a diverse range of lipid-soluble compounds out of the brain capillary endothelium and the CNS, eliminating potentially neurotoxic endogenous or xenobiotic molecules [216, 224]. P-gp is expressed at the luminal and abluminal membrane, as well as in pericytes and astrocytes [225], and is distributed along the nuclear envelope, in caveolae, cytoplasmic vesicles, Golgi complex, and rough ER [138]. The endothelial cells at the BBB also express several transporters for hormones, some cytokines, and chemokines [226]. Large proteins, such as transferrin, LDL, leptin, immunoglobulin G (IgG), insulin, and insulin-like growth factor also use receptor-mediated transport systems to pass through the BBB [146].

Internalization of ligands and receptors from the plasma membrane is cholesterol sensitive [227] and comprises endocytosis of caveolae, vesicles enriched in caveolin-1 [228]. The caveolar membranes carry several receptors including those for insulin, albumin, receptor for advanced glycation endproducts (RAGE), LDL, HDL [229] and are also closely associated with P-gp. Moreover, caveolin-1 can affect the levels of TJs in endothelial cells of the BBB [230]. Interestingly, study of the ultrastructure of the BBB in young and aged mice during ischemia has demonstrated that permeability is associated with a remarkable increase in endothelial caveolae and vacuoles although TJs were generally intact [231].

3.5. A β clearance at the BBB

Increase in either total A β levels or the relative concentration of both A β 40 and A β 42 (where the former is more concentrated in cerebrovascular plaques and the latter in neuritic plaques) have been implicated in the pathogenesis of both familial and sporadic AD.

There are several identified pathways for the removal of the A β from the brain. A β peptides mainly produced in neurons are degraded by peptidases. Through efflux transporters located in cerebral vessels, A β flows out from brain parenchyma into the plasma. A β is also removed through perivascular pathways into the cervical lymph nodes as A β within ISF diffuses in the extracellular spaces of the brain parenchyma entering basement membranes of capillaries, passing into the tunica media of arteries, and draining out of the brain. A β can also be taken up by different cells in the brain [232]. A few AD cases are familial AD, associated with genetic mutations which promote an increase in the production of A β [233]. On the other hand, the cause of the sporadic AD, the majority of the AD cases, is considered to be the impaired clearance of A β from the brain [234, 235]. In this viewpoint, AD is associated with cerebrovascular disorder, which drives the accumulation of A β at the blood vessels (cerebral amyloid angiopathy, CAA) and in the brain parenchyma, extracellularly [138, 236], and intraneuronal lesions—NTFs [237].

In the healthy brain, A β concentration is accurately regulated by its rate of production, its enzymatic degradation [238], its rapid clearance across the BBB through LRP1 [239, 240], and influx back into the brain by RAGE [241]. These receptors are multi-ligand cell surface receptors that, in addition to A β , mediate the clearance of a large number of proteins. P-gp, belonging to the superfamily of ATP-binding cassette (ABC) transporters, is also involved in effluxing A β out of the brain. While LRP1 and P-gp appear to mediate the efflux of A β from the brain to the periphery, RAGE has been strongly implicated in A β influx back into the CNS. With increasing age and also in AD pathology, the expression of the A β efflux transporters is

decreased and the A β influx transporter expression is increased at the BBB, adding to the amyloid burden in the brain and its gradual neurotoxic oligomerization [242]. Thus, continuous A β elimination by transport across the BBB and/or metabolism is essential to prevent its potentially neurotoxic accumulations in the brain [234]. Studies have demonstrated several transport proteins such as α 2-macroglobulin, TTR, apolipoprotein E (apoE), and apolipoprotein J (apoJ), which bind to A β and control its clearance, metabolism, and aggregation [243]. It has been shown that apoJ can increase the BBB clearance of A β 42 [244], while apoE disrupts the clearance of free A β across the mouse BBB, in an isoform-specific manner (apoE4>apoE3 or apoE2), by driving A β transport from LRP1 to VLDLR which internalizes A β -apoE complex at a slower rate than LRP1 [245]. Another transport pathway is the bulk flow of the ISF into the CSF through the perivascular Virchow-Robin arterial spaces, which is followed by drainage into the plasma across the arachnoid villi [243].

3.5.1. A β transport by LRP1

LRP1, the major efflux transporter for A β across the BBB [239] and a member of the LDL receptor family, acts as both a multifunctional scavenger and a signaling receptor. LRP1 is synthesized as a precursor molecule (600 kDa) in the ER. Then in the Golgi network, a cleavage generates an 85 kDa transmembrane beta-subunit (containing two intracellular NPxY motifs) that remains non-covalently connected to the extracellular 515 kDa alpha-subunit (containing 4 ligand-binding domains for more than 30 ligands) [246].

Transcytosis of A β across the BBB starts with its binding to LRP1 at the abluminal side of the cerebral endothelium [239, 240]. However, this has been controversial due to the studies that failed to demonstrate a role for LRP-mediated transcytosis, but rather showed a role for LRP receptors in endocytosis and degradation of A β [247]. Anyway, the significant function of LRP1 in AD is not only portrayed by LRP1-mediated endocytosis of A β but also by data showing that the cytoplasmic domain of LRP1 has been involved in APP processing [248]. Cleavage of the extracellular domain of LRP1 by beta-secretase (BACE1) releases soluble LRP1 (sLRP1) in plasma [249]. Reduced expression of LRP1 has been described during aging in animal models and in AD individuals [240, 249]. In astrocytes, LRP1 also mediates degradation of amyloid deposits via apoE [250].

3.5.2. A β transport by RAGE

RAGE, a multiligand receptor in the immunoglobulin superfamily, which can bind to various ligands including A β and advanced glycation end products (AGE proteins) [251], is the most influential influx transporter for A β across the BBB [241]. Interestingly, and unlike many receptors (including LRP1), RAGE expression is triggered by the accumulation of RAGE ligands, meaning that the levels of RAGE expression are determined by the levels of its ligands. In the healthy brain, RAGE is expressed at minimal levels at the BBB, except at the endothelium of bigger microvessels of the brain. However, when RAGE ligands increase in the AD brain, RAGE expression rises in the affected cerebral vessels, neurons or microglia [251]. This mechanism worsens the cellular dysfunction due to RAGE-A β interactions. Circulating A β can enter the brain by a special receptor-mediated transport mechanism that is dependent on

RAGE expression on the luminal surface of brain vessels [252]. Following A β binding to RAGE at the luminal membrane of the BBB, transcytosis of circulating A β across the BBB into the brain parenchyma and its binding to neurons occurs. Moreover, activation of NF- κ B in the endothelial cell leads to proinflammatory cytokines secretion and cerebral blood flow suppression [241]. A β -RAGE interaction not only generates oxidative damage to RAGE-expressing neurons, which results in neuronal degeneration, but also activates microglial cells, indirectly leading to inflammation [251]. Therefore, repression of A β -RAGE interaction in the BBB can inhibit A β influx, oxidant stress, and cytokine production. The inhibitors of A β /RAGE interaction have been shown to improve the BBB function and the cerebral blood flow responses to the brain activation, and to reduce neuroinflammation. Some RAGE/A β blockers are currently being tested in AD patients [138]. While RAGE is involved in the influx of A β into the brain, the soluble isoform of RAGE (sRAGE) has been detected in the plasma. It seems that sRAGE competes with cell-surface RAGE for ligand binding, thus increasing the elimination of circulating A β [145].

3.6. BBB dysfunction in AD

Failure in the BBB function is an outstanding event in the development and progression of several CNS diseases including multiple sclerosis [253], ischemia [254], Parkinson's disease, and AD [255]. While in some of the diseases increased BBB permeability is an outcome and consequence of the pathology (such as ischemic stroke), in other cases, BBB failure may be a causative event for the disease (such as multiple sclerosis). Furthermore, BBB dysfunction can be mild with temporary opening of TJs, or it can be a chronic breakdown [256], with changes in transporters and enzymes happening at the same time [216].

Although cerebrovascular abnormalities have been noted in AD, the starting point between BBB failure and AD pathology is not clear yet [142, 257]. Nevertheless, BBB homeostasis is altered in the initial stages of AD leading to the production of proinflammatory cytokines and suppressors of the cerebral blood flow by endothelial cells; then, amyloid deposits are observed in cerebral capillaries and vessels in the later stages of AD [258]. A large number of alterations in the structure and function of the BBB were shown to occur in AD. For instance: decreased LRP1 expression in human brain microvasculature [239], increased LRP1 oxidative damage [259], impaired microvascular P-gp [260], increased expression of RAGE in the cerebral vessels, neurons and microglia [251]. Moreover, it has been shown to occur: decreased glucose consumption by the brain, thus predicting a decay in cognitive function [261], which has been described by the reduction in the GLUT-1 transporters expression (protein levels) in AD hippocampus microvessels [262] but not a decrease in GLUT-1 mRNA levels [263]. Other alterations have been described, such as: decreased endothelial mitochondrial density, increased endothelial vacuolization, accumulation of collagen and perlecan in the basement capillary membrane [264], and increased pinocytotic vesicles [265]. Also, it has been reported a reduced number and smaller diameter of capillaries in CNS implying the diminished overall surface for LRP1-mediated transcytosis of A β across the BBB, and also a decreased length of brain capillaries [266] which lowers transport of energy substrates and nutrients across the BBB, and reduces the clearance of neurotoxins from the brain [138]; Overexpression and

accumulation of occludin in frontal cortex and basal ganglia of AD brains [267] have also been described as well as lower expression of genes such as the homeobox gene MEOX2 (or GAX), a regulator of vascular differentiation [268]. Finally, activated microglia [269] and astrocytes, the resident brain immune cells present in neurovascular unit of the BBB, have been observed surrounding A β plaques in AD brains, releasing inflammatory cytokines, such as IL-1 and IL-6, TNF- α , and transforming growth factor- β [174].

4. Periphery and Alzheimer's disease

4.1. A β levels at the periphery

Although studies in rodents have shown an increase in plasma A β levels, data in human AD patients have been contradictory; while some demonstrate increased levels of circulating A β [270], others report decreased [271] or unchanged [272] levels. Nevertheless, the significant role of circulating A β in AD pathology cannot be neglected. Interestingly, A β is also produced outside the brain in considerable amounts by the platelets, skeletal muscle, vascular walls, kidney, heart, liver, and by other non-neural tissues [273–275]. These pools may also provide a dynamic exchange of A β between the brain and periphery. However, A β peptides in the periphery cannot form filamentous structures, probably due to the presence of multiple circulating molecules that bind A β and thereby change its free-plasma levels [276].

4.1.1. Plasma proteins involved in peripheral sink and clearance of A β

Continuous removal of A β , not only from the brain but also from blood and from the entire organism, is essential for preventing its accumulation in the brain. A β in the plasma is bound to a number of proteins such as albumin [277], apoE and apoJ [278], TTR and a soluble form of LRP1 (sLRP1). In healthy human plasma, sLRP1 is a major endogenous brain A β sinker that sequesters 70–90% of plasma A β . sLRP1 has been demonstrated to bind the majority of circulating A β preventing RAGE-dependent influx of A β back to the brain, while improving its systemic clearance [279]. These data confirm the peripheral sink hypothesis, which imply systemic clearance of A β via binding proteins in serum and preventing its uptake through RAGE. Accordingly, this explains how some therapeutics, such as peripheral administration of sLRP1 or antibodies against A β [280], decrease A β levels in the brain. Moreover, AD patients show decreased plasma sLRP1 levels and increased oxidative damage to sLRP1, reducing its binding capacity for A β . This, in turn, increases the free A β fraction in plasma [279], that is, accessible for RAGE-dependent influx back to the brain [281].

Higher sRAGE levels are linked to reduced risk of developing several disorders such as cardiovascular disease and AD. Levels of sRAGE are decreased in AD and in vascular dementia, which may confer a target for therapeutic purposes [282]. It has been demonstrated that systemic administration of a truncated form of RAGE decreases A β load in a transgenic mouse model [241]. Moreover, sRAGE has an inverse relationship with cholesterol, presenting another modulatory impact on A β metabolism due to the role of cholesterol as a mediator of inflammation and APP processing [145].

4.1.2. Alterations in other plasma proteins in AD

Few proteins are synthesized solely by the brain or are present in higher concentrations in CSF compared with the blood. In conditions of the BBB failure, these CSF markers can appear or be increased in the plasma [283]. Hence, estimating levels of CSF proteins in the plasma may be a reliable method to control BBB integrity. For instance, S-100 is primarily produced in the brain by astrocytes and when the BBB is disturbed it is quickly released from the brain and appears in the blood [284].

4.2. Liver and A β elimination

At the periphery, the liver play important roles not only in the storage and in the release of nutrients and proteins but also in the neutralization and elimination of a variety of toxic substances such as A β from the plasma.

The receptor for LDL (LDLr), LRP1, and megalin/LRP2 play important roles in endocytosis of lipoproteins and systemic lipid homeostasis. Moreover, LRP1 mediates the clearance of a multitude of extracellular ligands such as A β and regulates diverse signaling processes such as growth factor signaling, inflammatory signaling pathways, apoptosis, and phagocytosis in liver.

In the liver, LRP1, which can be blocked by the receptor-associated protein (RAP), has been shown to be the predominant transporter that mediates systemic clearance of A β [285]. LRP1 localized to hepatic cells binds to and systemically clears circulating A β . Reduced hepatic LRP1 levels are associated with decreased peripheral A β clearance in aged rats. In aged squirrel monkeys, systemic clearance of A β is also reduced and associated with increased A β levels in the brain. In addition to the liver, sLRP1-A β complexes and free A β are removed through the kidneys [279]. Liver was also demonstrated to be the major source of A β and to be able to regulate brain A β levels [286].

The half-life of circulating A β is short, in the range of minutes [287, 288]. This also suggests that rapid systemic clearance of A β prevents reuptake by RAGE after efflux.

It has been shown that some proteins such as insulin [289] and TTR [290] increase LRP1 levels in hepatic plasma membrane, and in turn enhance peripheral A β clearance. Insulin-degrading enzyme, a zinc metalloendopeptidase that hydrolyzes numerous peptides, including A β [238], insulin [291] and the AICD, has been purified from several mammalian tissues including liver, brain and blood cells [292]. Furthermore, experiments in rats demonstrated that after 3.5 min post-infusion of radiolabelled A β into the lateral ventricle, 40% of the injected radioactivity was already in the blood and urine, and internalized by the liver and the kidneys, indicating not only a quick clearance mechanism but also the involvement of systemic organs in the elimination and catabolism of A β [293]. Therefore, the capacity of the liver to internalize, catabolize, and eliminate large doses of A β , may explain not only the low plasma A β levels but also its small variation noted with age and disease stages.

In some cases, both parenchymal and non-parenchymal liver cells take up proteins, whereas in other cases this is done mostly by hepatocytes, as happens for TTR, apoA-I, SAP and A β . In

vivo and in vitro experiments showed that hepatocytes are the main cells involved in A β uptake (about 90%) and in its catabolism [288].

5. Transthyretin

Transthyretin (TTR), formerly called prealbumin due to its electrophoretic characteristics, located just in front of the albumin band, is a plasma protein secreted mainly by the liver and choroid plexus (CP) [294]. The name “transthyretin” discloses its dual physiological role as a carrier for thyroid hormones [295] and retinol, the latter through the binding to retinol-binding protein (RBP) [296]. TTR was first described in the CSF [297] and shortly after in the plasma [298].

Although the involvement of TTR in the transport of thyroid hormones and RBP, as well as in FAP, is very well established, its involvement in neuroprotection is part of a very recent knowledge and constantly evolving.

5.1. TTR gene structure and expression

TTR is codified by a single copy gene localized in the long arm of chromosome 18 [299]. The entire nucleotide sequence including the 5' (transcription initiating site) and the 3' (untranslated region) flanking regions was determined [300, 301] and attributed to the region 18q11.2–q12.1 [302]. The full gene is 7.6 kilobase (kb) long comprising 4 exons and 3 introns [300, 301]. Exon 1 contains 95 basepairs (bp), including 26 bp 5' untranslated, and codes for a 20 aa residue leader peptide and aa 1–3 of the mature protein; exon 2 (131 bp), 3 (136 bp), and 4 (253 bp) hold the coding sequences for aa residues 4–47, 48–92 and 93–127, respectively. The introns (A, B and C) are 934, 2090 and 3308 bp long, respectively. Introns A and C contain two open reading frames (orf) of unknown significance [301].

The TTR mRNA spans ~0.7 kb and contains a 5'-untranslated region (26–27 nucleotides), a coding region (441 nucleotides), and a 3'-untranslated region (145–148 nucleotides) preceding the poly(A) tail [294, 303]. Human [299], rat [304, 305] and mouse [306] coding regions exhibit a considerable degree of homology (~85%), suggesting a phylogenetically preserved modulating role in gene expression.

TTR is predominantly synthesized by the liver where more than 90% of the protein is produced. The remaining is produced by the CP and the retina. TTR is detected in the fetal blood very early during development, as soon as eight weeks after conception [307]. TTR plasma concentration is age dependent, and in healthy newborns, it is about half of that in adults [308, 309]. TTR values vary from 20 to 40 mg/dL. In spite of the low TTR levels in CSF (~2 mg/dL), the CP is presented as the major site of TTR expression, expressed as a ratio of TTR/mass of tissue, corresponding to a ~30-fold higher than that found in plasma [310]. TTR represents 20% of the total CSF proteins [310].

With respect to the regulation of TTR expression, several studies showed that liver and CP TTR expression are increased in response to sex hormones, as demonstrated in mice [311,

312]. In rats, hydrocortisone and psychosocial stress are also inducers of TTR in the CP [313]. Others studies indicate that TTR can be expressed by brain cells, for instance in response to the heat shock factor 1 (HSF1) and to the AICD fragment of APP, as we will discuss further ahead in the context of TTR protection in AD.

5.2. TTR protein structure

The TTR mRNA codifies for the TTR-monomer; the polypeptide of 147 aa residues whose N-terminal region is a hydrophobic signal peptide of 20 aa residues. The TTR monomer is subjected to a cleaving process, during its migration through the ER, giving rise to the native TTR monomer after breaking of the signal peptide [294]. Assembly of four identical subunits (13,745 Da) occurs yielding the mature tetrameric protein, with a molecular mass of 54,980 Da [314].




Figure 2. The human TTR tetrameric structure complexed with two molecules of RBP (light green), which in turn is bound to retinol, and with T4 (binding in the central channel).

The tridimensional structure of TTR was made available at 1.8 Å resolution by X-ray diffraction studies on the crystallized protein [315]. Each monomer contains two β -sheets formed by strands DAGH and CBEF. All, except strands A and G, display an antiparallel orientation, and are arranged in a topology similar to the classic Greek key barrel. The strands are 7–8 residues long, except strand D, which, is only 3 residues in length. Only about 5% of the aa residues are located in one segment of α -helix comprising residues 75–83, at the end of strand E. Two monomers associate forming the dimer by interactions between chains F and H of each monomer. Fixing the chains of one monomer as A–H and the ones of the other monomer as A'–H', the arrangement within a dimer is DAGHH'G'A'D' and CBEFF'E'B'C'. The tetramer consists of two dimers with connecting edges occurring between the AB loop of one dimer with the H strand of the other dimer (**Figure 2**). The quaternary structure of TTR has the shape of a globular protein with an overall size of 70 Å \times 55 Å \times 50 Å [315]. The two dimers are slightly rotated in relation to each other along the y axis.

5.3. Physiology and Metabolism

5.3.1. Transport of thyroxine (T_4)

Thyroid hormones are transported in blood circulation and delivered to the target tissues, with an incredible high amount of the hormones (99%) bound to serum proteins. In human plasma, TTR, thyroxine-binding protein (TBG) and albumin are responsible for the delivery of T_4 into the target tissues. Although TBG is much less concentrated in the plasma than TTR, it presents the highest affinity constant for T_4 ($K_a = 1 \times 10^{10} \text{ M}^{-1}$) and transports about 70% of the plasma T_4 . TTR has an intermediate affinity for T_4 ($K_a = 7 \times 10^7 \text{ M}^{-1}$) and transports about 15% of the hormone, and finally, albumin presents the lowest binding affinity ($K_a = 7 \times 10^5 \text{ M}^{-1}$) [316].

The four monomers of the TTR tetramer, demarcate an open channel through the molecule where two binding sites for thyroid hormones are located (**Figure 2**). These two binding sites present negative cooperativity [317] implying that when the first thyroid hormone molecule occupies the first site, the affinity for the second molecule is highly reduced.

T_4 transport into the brain is of particular interest and has raised much controversy [318]. Although TTR is the major T_4 -binding protein in the CSF, studies done in *ttr* knock-out mice ($TTR^{-/-}$) [319] revealed that TTR is not essential for thyroxine to reach the brain and other tissues [320]. In addition, measurement of several parameters of thyroid hormone function indicates that these mice are euthyroid despite strongly reduced total T_4 plasma levels [321]. No other protein was found to replace TTR in the transport of T_4 in the CSF, and T_4 levels were normal in the cortex, cerebellum, and hippocampus while strongly reduced in the CP of the $TTR^{-/-}$ mice [322].

Therefore, it was suggested that TTR might be a reservoir for T_4 both in the plasma and in the CP and CSF, which might become important under conditions of increased hormone demand.

5.3.2. Transport of vitamin A

TTR is also responsible for retinol transport through binding to retinol-binding protein (RBP, **Figure 2**). RBP is a 21 kDa monomeric protein comprising 182 aa residues [323]. The conformational structure of RBP bound to retinol was determined by x-ray crystallography [324]. In most instances, RBP is transported in the bloodstream in the form of a saturated holo-RBP protein equimolarly attached to TTR [325]. Although TTR has four binding sites for RBP [326, 327], under physiological conditions only one RBP molecule is bound to TTR due to the RBP limiting concentration.

5.3.3. TTR metabolism

The biological half-life of TTR is about 2–3 days in humans [328, 329], 22–23 h in monkeys [330] and 10–13 h in rats [331]. The major sites of TTR degradation in the rat are liver (36–38%), followed by the muscle (12–15%) and skin (8–10%) [332].

The normal physiology of TTR is not completely understood, in particular, its cellular uptake. Nevertheless, several observations suggest that TTR internalization is receptor-mediated, both

in human hepatoma cells (HepG2) [333] and in chicken oocytes [334]. Megalin, a receptor implicated in the renal re-uptake of plasma proteins carriers of lipophilic compounds, was shown to play a role in renal uptake of TTR [335]. Furthermore, different TTR mutations presented different levels of cell association and degradation, suggesting that the structure of TTR is important for megalin recognition. TTR internalization was further explored by studying TTR uptake using hepatomas and primary hepatocytes [336]. This work showed direct evidence for TTR internalization by a specific receptor, forming a ~90 kDa complex. TTR internalization was inhibited by RBP (70% decrease) and T₄ (20% decrease) and TTR mutants revealed differences in uptake, indicating again that the recognition by receptor is structure dependent. Internalization was also inhibited by lipoproteins and RAP, a ligand for all members of the low-density lipoprotein receptor family (LDLr). All together, these results suggest a common pathway for TTR and lipoprotein metabolism and the existence of a RAP-sensitive receptor for TTR internalization. TTR is also internalized by several other cells types in cell culture, such as astrocytoma cells [337], cardiomyocytes [338, 339], neurons [340, 341], endothelial cells and others. More recently, TTR was also shown to be uptaken by sensory neurons, an event also mediated by megalin [340].

5.3.4. Proteolytic activity

TTR was also found to act as a cryptic protease and the first substrate described was apoA-I [342, 343]. TTR is also able to cleave lipidated apoA-I (mainly in the lipid poor pre β -HDL subpopulation) which can be relevant in the lipoprotein metabolism [343]. Moreover, apoA-I cleaved by TTR presents less ability to promote cholesterol efflux [343] and shows increased amyloidogenicity and propensity to form aggregates. Liz et al. also described that TTR was able to cleave amidated neuropeptide Y (NPY) and that its proteolytic activity affects axonal growth, leading to the conclusion that TTR has natural substrates in the nervous system [344]. Newly, the same authors described TTR as a metallopeptidase [345], and this result was supported by another study that showed the involvement of a carboxylate and an ammonium group, possibly from a lysine side chain, in the TTR hydrolytic activity [346]. Costa et al. showed for the first time that TTR can cleave A β peptide in vitro [347] and implications in A β clearance in AD will be addressed further ahead.

5.4. TTR as a cause of disease

TTR is associated with the most prevalent type of hereditary systemic amyloidosis. The pathologic conditions include FAP and familial amyloidotic cardiomyopathy (FAC). A non-hereditary condition is also related to TTR, the systemic senile amyloidosis (SSA), and affects about 25% of people over 80 years of age. In SAA, the deposits occur in the heart and are composed of wild-type (WT) TTR.

FAP is related to a peculiar form of hereditary autosomal dominant polyneuropathy. Corino de Andrade first described the disease in 1952 [6] in the Portuguese population mainly from the northern part of the country. Characterized by systemic deposition of amyloid and with a special involvement of the peripheral nerves, the age of onset of the disease is usually between 20 and 35 years of age, with a fast progression to death within 10–15 years.

Clinically, FAP is characterized by early impairment of temperature and pain sensation in the feet, and autonomic dysfunction leading to paresis, malabsorption, and emaciation. Painless injury to the feet complicated by ulcers, cellulitis, osteomyelitis, and Charcots joints may also occur [348]. Motor involvement occurs with disease development causing wasting and weakness, and there is a progressive loss of reflexes. The amyloid deposits can occur in any part of the peripheral nervous system, including the nerve trunks, plexus, and sensory and autonomic ganglia. The other organ frequently involved in FAP, which is the heart. Clinically, the cardiomyopathy may present as an arrhythmia, heart block, or heart failure.

The first report relating immunologically TTR as the main protein in FAP fibrils was in 1978 by Costa and colleagues [349]. In 1984, the Val30Met mutation was identified in the protein isolated from Portuguese FAP patients. This variant was shown to be a biochemical marker for FAP [350] that resulted from a point mutation in the exon 2 of the TTR gene [351]. Since the identification of the Val30Met variant, many others aa substitutions were identified in the TTR protein and now over 100 mutations are described (<http://amyloidosismutations.com/mut-attr.php>); these variants are associated with different clinical phenotypes and a considerable number of non-pathogenic TTR mutations have been identified, including a T119M variant described to be protective.

TTR tetramer dissociation is believed to be on the basis of a series of events leading to TTR amyloid formation. In fact, the amyloidogenic potential of the TTR variants relates inversely with its tetrameric stability, and it is thought that upon dissociation of the tetramer, non-native TTR monomers are formed which can assemble, forming amyloidogenic intermediate species, such as oligomers and aggregates. Similarly to other amyloidogenic proteins, it is now believed that cellular toxicity is derived from the initial intermediate species occurring in the initial stages of FAP [352]. TTR stabilization has been proposed as a key step for the inhibition of TTR fibril formation and has been the basis for FAP therapeutic strategies. Such stabilization can be achieved through the use of small compounds sharing molecular structural similarities with T₄ and binding in the T₄ central-binding channel. Most of such compounds belong to the class of NSAIDs as it is the case of diflunisal and tafamidis, currently being administered to FAP patients. For instance, in FAP patients, diflunisal was shown to stabilize TTR, increasing its serum concentration [353], and to reduce the rate of progression of neurological impairment and to preserve quality of life [354].

6. TTR and neuroprotection

Several lines of evidence indicate that TTR possesses neuroprotective properties in multiple contexts. Studies with TTR^{-/-} mice revealed that these animals show reduced signs of depressive-like behavior, probably due to the modulation of noradrenergic system by the increase of noradrenaline in the limbic forebrain [355]. Additionally, increased levels of NPY, known as an antidepressant neurotransmitter [356], were reported in dorsal root ganglia (DRG), sciatic nerve, spinal cord, hippocampus, cortex, and CSF of TTR^{-/-} mice [357], supporting the importance of TTR in the modulation of depressive behavior. Furthermore,

Sousa and co-workers also described that TTR^{-/-} mice present memory impairment compared with wild-type (TTR^{+/+}) animals, indicating that the absence of TTR accelerates cognitive deficits usually associated with aging [358].

In addition, TTR was associated with nerve regeneration. Fleming et al. revealed, for the first time, that TTR acts as an enhancer of nerve regeneration, following the observation that TTR^{-/-} mice have decreased ability to regenerate from a sciatic crushed nerve [359]. Later, the same authors showed that the absence of TTR leads to impaired retrograde transport and decreased axonal growth, and also that the effect of TTR in neurite outgrowth and nerve regeneration is mediated by megalin-dependent internalization [340].

It was also established a relationship between TTR and ischemia, one of the major causes of brain injuries in world. Santos and co-workers proposed that in a compromised heat-shock response, CSF TTR contributes to control neuronal cell death, edema, and inflammation, influencing the survival endangered neurons [360].

More recently, a new neuroprotective role in the CNS was attributed to TTR, as a transcription inducer of insulin-like growth factor receptor I (IGF-IR), known as a protective receptor against apoptosis [361]. Vieira and colleagues described, for the first time, that TTR induces increased levels of IGF-IR, showing that TTR triggers IGF-IR nuclear translocation in cultured neurons [341].

6.1. TTR protection in Alzheimer's Disease

There are several reports suggesting a relevant protective role of TTR in AD. However, the precise mechanism(s) is not entirely understood.

In 1993, Wisniewski described that A β 40 fibril formation was inhibited upon incubation with human CSF [362], which was explained by the sequestration of A β by extracellular proteins circulating in CSF such as apoE and apoJ [362–364]. The earliest description of TTR protection in AD was presented by Schwarzman and colleagues in 1994 when they observed that, in CSF and contrarily to the expectations, apoE was not the major protein binding to A β , but TTR [365], proposing the sequestration hypothesis as a possible explanation for the peptide aggregation and consequent progression of AD. This hypothesis suggested that certain extracellular proteins sequester normally produced A β , thereby preventing amyloid formation and its toxicity. Amyloid formation would occur when sequestration failed [365, 366], which could be related either with an A β overproduction, a reduction in the levels of the sequestering proteins, inability of those proteins to interact with the peptide, deficient clearance mechanisms, or a combination of all the events above stated.

Supporting a protective role for TTR in AD are its decreased levels observed both in the CSF [367] and in the plasma [114, 368] of AD patients as compared to age-matched subjects. Serot and coworkers suggested that the decrease in TTR levels in the CSF of AD brains was possibly related with an epithelial atrophy in the CP [367]. Interestingly, very recently, studies in an AD transgenic mouse model reported CP dysfunction and revealed a specific increase only of the A β 42 isoform in epithelial cytosol and in stroma surrounding choroidal capillaries, accompanied by a thickening of the epithelial basal membrane, greater collagen-IV deposition

around capillaries in CP that probably restrain solute exchanges, and attenuated expression of epithelial aquaporin-1 and TTR protein compared to non-transgenic mice [369].

Modulation of A β aggregation and toxicity was also investigated in brain vascular smooth muscle cells, isolated from dogs and AD patients, containing intracytoplasmic granules of A β produced in the presence of apoE. In this model, TTR was able to rescue the cells from this accumulation and the positive thioflavin S staining, initially observed, was no longer detected [370].

Although not fully consensual, several authors reported the presence of TTR in amyloid plaques both in AD patients [371–373] and in AD transgenic models [372], strengthening a role for TTR in A β deposition and in AD. TTR was also identified as a survival gene, and its differential overexpression in mice hippocampus was suggested to be responsible for the lack of neurodegeneration observed in the Tg2576 transgenic mice overexpressing the mutant form of human amyloid precursor protein with the Swedish mutation (APP_{sw}) [374]. Up-regulation of TTR and other survival genes was induced by the sAPP α , a neuroprotective fragment resulting from APP processing by α -secretase [373]. TTR up-regulation was also reported in situations of exposure of AD transgenic mice to an “enriched environment,” also resulting in pronounced reductions in cerebral A β levels and amyloid deposits, compared to animals raised under “standard housing” conditions [375]. Later on, AD transgenic models with genetic reduction of TTR and/or overexpression of human TTR [372] further showed the ability of TTR to modulate A β aggregation and toxicity. While the overexpression of human TTR ameliorated AD features in APP transgenic mice [372], the ablation of the mouse TTR gene resulted in accelerated amyloid deposition and increased A β brain levels [372, 376, 377]. In vitro studies further demonstrated that a direct interaction between TTR and A β abrogated the noxious properties of A β oligomers [378].

Animal models, in particular mice models, also provided evidence for a gender-associated modulation of brain A β levels [377] as elevated brain levels of A β 42 were observed in AD female mice with only one copy of TTR when compared to female with the two copies of the TTR gene, while no significant differences were observed in males. Additionally, this work also indicated that reduced levels of brain testosterone and 17 β -estradiol in female mice with TTR genetic reduction might underlie their increased AD-like neuropathology [377]. Interestingly, estradiol was found to be decreased in female AD patients when compared to healthy age and gender-matched controls [114], which in conjunction with TTR regulation by sex hormones, as already described, can account for TTR lowering in AD and the prevalence of this disease in women. In fact, plasma TTR levels were found decreased in AD women, as compared to healthy age- and gender-matched controls, whereas plasma TTR levels in AD men were not significantly different from their respective controls [114], further confirming the gender modulation by TTR.

6.1.1. TTR and A β interaction

TTR and A β binding was initially demonstrated by adding radiolabelled A β 1-28 synthetic peptide to human CSF and subsequent analysis by SDS-PAGE [365]. Later on, the TTR/A β

complex was also demonstrated in plasma [379], although the details of the interaction as well as the effects in A β fibrillogenesis and toxicity were not known.

In 2008, Costa and colleagues characterized the TTR/A β interaction by competition-binding assays using synthetic A β 42, which revealed that the WT TTR binds to different A β peptide species: soluble (with a K_d of 28 ± 5 nM), oligomers, and fibrils [378]. Other studies showed that TTR drastically decreased the rate of aggregation without affecting the fraction of A β in the aggregate pool and an estimated apparent K_S of 2300 M^{-1} was calculated [380]. These data support a hypothesis, wherein TTR preferentially binds to aggregated rather than monomeric A β and arrests further growth of the aggregates. Recent work indicates that the intensity of TTR binding to A β peptide is highest for partially aggregated materials and decreased for freshly prepared or heavily aggregated A β , suggesting that TTR binds selectively to soluble toxic A β aggregates [381]. Although Schwarzman and colleagues had shown in 1994 that TTR is capable of inhibiting A β 1-28 aggregation [365], Costa and co-workers showed by transmission electron microscopy (TEM) analysis that TTR is capable of interfering with A β fibrillization, both at inhibiting its aggregation and at disrupting pre-formed A β fibrils [378]. Thus, new and innovative studies are necessary to clarify the details of this interaction.

Another point of controversy refers to the TTR species involved in A β binding. Some studies support that the TTR monomer rather than the tetramer binds more strongly to A β [382], and it is even suggested that while the TTR monomer arrests A β aggregate growth, the tetramer modestly enhances the peptide aggregation [382]. Another study performed using diverse natural TTR mutants showed that different TTR variants bind differentially to A β in the following manner: T119M>WT>V30M³Y78F>L55P TTR [378], indicating the lower the amyloidogenic potential of TTR, the stronger the affinity toward the peptide. Since the amyloidogenic potential correlates inversely with TTR tetrameric stability, authors concluded that the TTR tetramer is the species binding to A β peptide. Previous work had already shown that amyloidogenic TTR mutants such as L55P and E42G, the only ones able to form TTR amyloid fibrils at pH 6.8 amongst the forty-seven variants tested, were unable to bind A β [383].

Given the above considerations, it is therefore conceivable that mutations of the TTR gene could alter the TTR/A β sequestration properties. However, the screening study in AD patients found no correlation between TTR variants and AD [384], and therefore other factors, namely conformational changes resulting from aging, should be affecting TTR levels and its binding properties towards A β .

Computer-assisted modelling was developed to determine the possible key residues participating in the interaction and the data suggested that residues 38–42, Asparagine 62 (E62) and E66 of each TTR monomer had a central role in the interaction [365]. Later studies have confirmed that only the residues 38–42 of TTR were important for the interaction [366]. More recently, Du and Murphy, identified the A strand, in the inner β -sheet of TTR, as well as the EF helix, as regions of TTR that are involved with A β [382] association. New data from the same group now indicates the involvement of the G strand of TTR with the particular involvement of L82A and L110A, suggesting that A β binding to TTR is mediated through these bulky hydrophobic leucines [385].

6.1.2. Effects of TTR stabilization in AD

The decrease in TTR levels in the context of AD is found early in disease development as indicated by the lower levels also found in plasma TTR in MCI patients [114]. TTR levels continue to decrease as disease progresses correlating negatively with disease severity, both in CSF [115] and plasma [114], and with senile plaque burden [386]. Similar results were found in AD transgenic mice as TTR was decreased as early as 3 months of age [377], well before amyloid deposition. Mice, however, seem to be able to compensate and female showed restored TTR levels at the age of 10 months [377].

The reason for TTR decrease is not known, but its tetrameric stability seems to play an important role. An unstable TTR can result in accelerated clearance, accounting for the lower levels observed. Further, such instability can also affect the A β sequestration properties of TTR. Supporting this stability hypothesis is the observation that plasma TTR from AD patients presents impaired ability to carry T₄. This also supports that it is the TTR tetramer that binds A β , since T₄ binding to TTR implies that the tetramer is assembled. Very interestingly, TTR genetic stabilization, that is, the presence of the T119M allele, was associated to decreased cerebrovascular disease and increased life expectancy [387]. In the context of AD, and arguing in favor of the stability hypothesis is the observation that the TTR/A β interaction can be improved in vitro in the presence of small chemical molecules known to bind to the T₄-binding channel and to stabilize its tetrameric fold [388]. Importantly, in vivo studies using one of such stabilizer—iododiflunisal, known to be a very potent TTR stabilizer [389] and shown to improve TTR/A β interaction [388]—administrated to AD transgenic mice resulted in amelioration of AD features such as cognitive function and A β brain deposition [390]. In addition, plasma levels of A β 42 were decreased upon iododiflunisal administration. These results opened the possibility for the use of TTR stabilizers in AD therapeutic drug development.

6.1.3. Mechanisms of TTR protection in AD

A β sequestration by TTR and other extracellular proteins was the first hypothesis proposed to explain why CSF is able to inhibit A β amyloid formation, implying that, when sequestered, A β cannot aggregate to form amyloid [362, 365]. However, the precise mechanism leading to final A β removal is not yet elucidated.

Following the identification of the TTR proteolytic activity, it was also described that A β peptide is cleaved by TTR, in vitro, with consequent generation of non-amyloidogenic fragments or fragments with amyloidogenic potential inferior to the full-length peptide [347]. Since clearance of A β from the brain can occur via proteolytic degradation of the peptide by several enzymes, such as neprilysin (NEP), insulin-degrading enzyme (IDE), Endothelin-converting enzyme (ECE), angiotensin-converting enzyme (ACE), uPA/tPA-plasmin system, cathepsin D, and matrix metalloendopeptidase 9 [391, 392], lower levels of TTR or inhibition of its proteolytic activity would result in less A β peptide eliminated by the cells, and therefore in its accumulation and amyloid formation. Interestingly, several sites of A β cleavage by TTR are common to several of the proteases mentioned [347]. Further, NEP is known to cleave not only monomeric but also oligomeric forms of A β localized intra and extracellularly, as determined in vitro and in vivo [393, 394]; similarly, TTR was shown to degrade

both monomers and aggregates of A β in vitro. Nevertheless, A β degradation by TTR was not yet shown in vivo. It is also not known if binding/sequestration of A β and its degradation by TTR are part of the same mechanism or are independent processes.

In addition to the peptidolytic removal of A β , clearance of the peptide from the brain also occurs via active transport at the BBB and BCSFB, as already discussed. The receptors for A β at the BBB bind A β directly, or bind to one of its carrier proteins, and transport it across the endothelial cell. The first hint pointing to the involvement of TTR in A β transport and clearance came from the analysis of brain and plasma levels of the peptide in mice with different TTR genetic backgrounds. Results showed that AD transgenic mice with just one copy of TTR had lower brain and plasma A β levels, as compared to animals with two TTR gene copies, raising the hypothesis that TTR might be involved both in A β brain efflux and in its peripheral removal at the liver. Very recently, it has been showed that TTR promotes A β internalization and efflux in hCMEC/D3 cells, a BBB cellular model widely used. Importantly, TTR stimulated brain-to-blood A β permeability in hCMEC/D3 which in turn can be explained because TTR itself can only cross the BBB only in the brain-to-blood direction [290, 332]. Thus, TTR can transport A β from, but not into the brain, acting as a neuroprotective molecule.

The presence of TTR in brain areas other than its site of synthesis and secretion—CP and CSF, respectively—has been already shown. In situations of injury, such as ischemia, TTR was detected at the local of infarct and shown to derive from CSF TTR [360]. However, other studies demonstrated TTR synthesis by cortical [395] or hippocampal neurons both in vitro [396], and in vivo [397] showing that TTR expression in the brain can be regulated [396]. For instance, Kerridge and colleagues showed that TTR expressed in SH-SY5Y neuroblastoma cell line is up-regulated by the AICD fragment of amyloid precursor protein (APP), specifically derived from the APP695 isoform [396]. Induced accumulation of functional AICD resulted in TTR up-regulation with concomitant A β decreased levels. Wang and colleagues reported that TTR expression in SH-SY5Y cells, primary hippocampal neurons, and the hippocampus of APP23 mice is significantly enhanced by HSF1 [397]. In any case, TTR is available in the brain and might participate in brain A β efflux by promoting BBB permeability to the peptide. It is also possible that TTR contributes to A β clearance from the brain through the BCSFB.

TTR was also able to increase A β internalization by hepatocytes prompting TTR as an A β transporter both in the brain and at periphery. Previous work showed that TTR is internalized by hepatocytes using a RAP-sensitive receptor, which together with the knowledge that as follows: (1) LRP1 is the main A β receptor both at the BBB and at the liver, (2) LRP1 is preferentially expressed at the basolateral membrane of the endothelial cells of the BBB, and (3) TTR can only cross the BBB in the brain-to-blood direction, indicates this receptor is involved in TTR-assisted A β transport. So far, it has been shown that mice with TTR genetic ablation present decrease levels of brain and liver LRP1 and that TTR added to hCMEC/D3 cells results in increased LRP1 expression. These findings open new perspectives for TTR/LRP-related therapeutic interventions in AD. However, a direct interaction between LRP1 and TTR is yet to be demonstrated. TTR has also been suggested to act in a chaperone-like manner by binding toxic or pretoxic A β aggregates in both the intracellular and extracellular environment [372].




Figure 3. Schematic representation of the proposed mechanisms underlying TTR Protection in AD. (A) In a healthy individual, TTR binds A β peptide in the brain at the CSF promoting its degradation and elimination through the epithelial cells of the CP. At the BBB, A β effluxes through LRP1 to the blood, a process in which TTR also participates. At the periphery, TTR and other A β bindable substances create a peripheral sink avoiding the return of A β to the brain, and transport the peptide to the liver, where it will be internalized by liver LRP1 for final degradation. (B) In AD patients, TTR decreased tetrameric stability for yet unknown reasons, and decreased expression due to decreased sex hormones, result in lower protein levels in the brain and plasma. In turn, this contributes to failure in A β sequestration, efflux at the BBB, and peripheral transport to the liver. Impaired A β transport at the BBB and internalization/degradation by the liver is aggravated by LRP1 decreased expression, modulated by TTR.

Therefore, more studies are necessary to unravel the mechanism(s) underlying TTR protection in AD, and to clarify how the hypothesis presented so far fit together (**Figure 3**).

Acknowledgements

This work was funded by national funds through FCT—Fundação para a Ciência e a Tecnologia/MEC—Ministério da Educação e Ciência and when applicable co-funded by FEDER funds within the partnership agreement PT2020 related with the research unit number 4293, and by FEDER funds through the Operational Competitiveness Programme—COMPETE and by National Funds through FCT—Fundação para a Ciência e a Tecnologia under the projects FCOMP-01-0124-FEDER-037277 (PEst-C/SAU/LA0002/2013).

Author details

Isabel Cardoso^{1,2*}, Luis Miguel Santos^{1,2} and Mobina Alemi^{1,2,3}

*Address all correspondence to: icardoso@ibmc.up.pt

1 IBMC-Instituto de Biologia Molecular e Celular, Porto, Portugal

2 i3S-Instituto de Investigação e Inovação em Saúde, Universidade do Porto, Porto, Portugal

3 Faculdade de Medicina, Universidade do Porto, Porto, Portugal

References

- [1] Westermark GT, Fändrich M, Westermark P. AA amyloidosis: pathogenesis and targeted therapy. *Annu Rev Pathol* [Internet]. 2015;10:321–44 [cited 2015 Dec 10]. <http://www.ncbi.nlm.nih.gov/pubmed/25387054>
- [2] Virchow R. Zur Cellulose—Frage. *Arch für Pathol Anat und Physiol und für Klin Med* [Internet]. 1854;6(3):416–26 [cited 2016 Jan 5]. <http://link.springer.com/10.1007/BF02116546>
- [3] Sipe JD, Benson MD, Buxbaum JN, Ikeda S, Merlini G, Saraiva MJM, et al. Nomenclature 2014: amyloid fibril proteins and clinical classification of the amyloidosis. *Amyloid* [Internet]. 2014;21(4):221–4 [cited 2015 May 28]. <http://www.ncbi.nlm.nih.gov/pubmed/25263598>
- [4] Westermark P. Quantitative studies on amyloid in the islets of Langerhans. *Ups J Med Sci* [Internet]. 1972;77(2):91–4 [cited 2015 Oct 28]. <http://www.ncbi.nlm.nih.gov/pubmed/4116019>
- [5] Prusiner SB. Novel proteinaceous infectious particles cause scrapie. *Science* [Internet]. 1982;216(4542):136–44 [cited 2015 Aug 2]. <http://www.ncbi.nlm.nih.gov/pubmed/6801762>
- [6] Andrade C. A peculiar form of peripheral neuropathy; familiar atypical generalized amyloidosis with special involvement of the peripheral nerves. *Brain* [Internet]. 1952;75(3):408–27 [cited 2016 Jan 13]. <http://www.ncbi.nlm.nih.gov/pubmed/12978172>
- [7] Glenner GG, Wong CW. Alzheimer's disease: initial report of the purification and characterization of a novel cerebrovascular amyloid protein. *Biochem Biophys Res Commun* [Internet]. 1984;120(3):885–90 [cited 2015 Jan 26]. <http://www.ncbi.nlm.nih.gov/pubmed/6375662>
- [8] Stelzmann R a., Schnitzlein HN, Murtagh FR. An English translation of Alzheimer's 1907 paper, "uber eine eigenartige erkankung der hirnrinde." *Clin Anat*. 1995;8(6):429–31.
- [9] Pimplikar SW. Reassessing the amyloid cascade hypothesis of Alzheimer's disease. *Int J Biochem Cell Biol* [Internet]. 2009;41(6):1261–8. <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2680505&tool=pmcentrez&rendertype=abstract>
- [10] Alzheimer A, Stelzmann RA, Schnitzlein HN, Murtagh FR. An English translation of Alzheimer's 1907 paper, "Über eine eigenartige Erkankung der Hirnrinde". *Clin Anat* [Internet]. 1995;8(6):429–31 [cited 2015 Mar 10]. <http://www.ncbi.nlm.nih.gov/pubmed/8713166>
- [11] Kumar A, Singh A. A review on Alzheimer's disease pathophysiology and its management: an update. *Pharmacol Rep* [Internet]. 2015;67(2):195–203 [cited 2015 Jan 9]. <http://www.ncbi.nlm.nih.gov/pubmed/25712639>

- [12] Burns A, Byrne EJ, Maurer K. Alzheimer's disease. *Lancet* (London, England) [Internet] (Elsevier). 2002;360(9327):163–5 [cited 2016 Jan 6]. <http://www.thelancet.com/article/S0140673602094205/fulltext>
- [13] van der Linde RM, Dening T, Matthews FE, Brayne C. Grouping of behavioural and psychological symptoms of dementia. *Int J Geriatr Psychiatry* [Internet]. 2014;29(6):562–8 [cited 2015 Oct 20]. <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=4255309&tool=pmcentrez&rendertype=abstract>
- [14] Petrovic M, Hurt C, Collins D, Burns A, Camus V, Liperoti R, et al. Clustering of behavioural and psychological symptoms in dementia (BPSD): a European Alzheimer's disease consortium (EADC) study. *Acta Clin Belg* [Internet]. 2016;62(6):426–32 [cited 2016 Jan 6]. <http://www.ncbi.nlm.nih.gov/pubmed/18351187>
- [15] 2015 Alzheimer's disease facts and figures. *Alzheimers Dement* [Internet]. 2015;11(3):332–84 [cited 2015 Jun 2]. <http://www.ncbi.nlm.nih.gov/pubmed/25984581>
- [16] Reitz C, Mayeux R. Alzheimer disease: epidemiology, diagnostic criteria, risk factors and biomarkers. *Biochem Pharmacol* [Internet] (NIH Public Access) 2014;88(4):640–51 [cited 2014 Jul 10]. <http://pmc/articles/PMC3992261/?report=abstract>
- [17] McKhann G, Drachman D, Folstein M, Katzman R, Price D, Stadlan EM. Clinical diagnosis of Alzheimer's disease: report of the NINCDS-ADRDA Work Group* under the auspices of Department of Health and Human Services Task Force on Alzheimer's Disease. *Neurology* [Internet]. 1984;34(7):939–939 [cited 2015 Jun 7]. <http://www.neurology.org/content/34/7/939>
- [18] McKhann GM, Knopman DS, Chertkow H, Hyman BT, Jack CR, Kawas CH, et al. The diagnosis of dementia due to Alzheimer's disease: recommendations from the National Institute on Aging-Alzheimer's Association workgroups on diagnostic guidelines for Alzheimer's disease. *Alzheimers Dement* [Internet] (Elsevier). 2011;7(3):263–9 [cited 2014 Jul 9]. <http://www.alzheimersanddementia.com/article/S1552526011001014/fulltext>
- [19] Hyman BT, Phelps CH, Beach TG, Bigio EH, Cairns NJ, Carrillo MC, et al. National Institute on Aging-Alzheimer's Association guidelines for the neuropathologic assessment of Alzheimer's disease. *Alzheimers Dement* [Internet]. 2012;8(1):1–13 [cited 2015 Aug 6]. <http://www.sciencedirect.com/science/article/pii/S1552526011029803>
- [20] 2013 Alzheimer's disease facts and figures. *Alzheimers Dement* [Internet]. 2013;9(2):208–45 [cited 2015 Sep 2]. <http://www.sciencedirect.com/science/article/pii/S1552526013000769>
- [21] Gaugler JE, Kane RL, Johnston J a, Sarsour K. Sensitivity and specificity of diagnostic accuracy in Alzheimer's disease: a synthesis of existing evidence. *Am J Alzheimer's Dis Other Dement*. 2013;28(4):337–47.

- [22] Bertram L, Lill CM, Tanzi RE. The genetics of Alzheimer disease: back to the future. *Neuron* [Internet]. 2010;68(2):270–81 [cited 2015 Jun 14]. <http://www.sciencedirect.com/science/article/pii/S0896627310008378>
- [23] Goate A, Chartier-Harlin MC, Mullan M, Brown J, Crawford F, Fidani L, et al. Segregation of a missense mutation in the amyloid precursor protein gene with familial Alzheimer's disease. *Nature* [Internet]. 1991;349(6311):704–6 [cited 2014 Dec 28]. <http://www.ncbi.nlm.nih.gov/pubmed/1671712>
- [24] Scheuner D, Eckman C, Jensen M, Song X, Citron M, Suzuki N, et al. Secreted amyloid beta-protein similar to that in the senile plaques of Alzheimer's disease is increased in vivo by the presenilin 1 and 2 and APP mutations linked to familial Alzheimer's disease. *Nat Med* [Internet]. 1996;2(8):864–70 [cited 2015 Aug 4]. <http://www.ncbi.nlm.nih.gov/pubmed/8705854>
- [25] Grundke-Iqbal I, Iqbal K, Quinlan M, Tung YC, Zaidi MS, Wisniewski HM. Microtubule-associated protein tau. A component of Alzheimer paired helical filaments. *J Biol Chem* [Internet]. 1986;261(13):6084–9 [cited 2016 Jan 8]. <http://www.ncbi.nlm.nih.gov/pubmed/3084478>
- [26] Spillantini MG, Goedert M. Tau protein pathology in neurodegenerative diseases. *Trends Neurosci* [Internet]. 1998;21(10):428–33 [cited 2016 Jan 8]. <http://www.sciencedirect.com/science/article/pii/S016622369801337X>
- [27] Lee VM, Goedert M, Trojanowski JQ. Neurodegenerative tauopathies. *Annu Rev Neurosci* [Internet]. 2011;24:1121–59 (Annual Reviews 4139 El Camino Way, P.O. Box 10139, Palo Alto, CA 94303-0139, USA; 2001 Jan 28 [cited 2015 Nov 22]). http://www.annualreviews.org/doi/full/10.1146/annurev.neuro.24.1.1121?url_ver=Z39.88-2003&rft_id=ori%3Arid%3Aacrossref.org&rft_dat=cr_pub%3Dpubmed&
- [28] Iwatsubo T, Odaka A, Suzuki N, Mizusawa H, Nukina N, Ihara Y. Visualization of A beta 42(43) and A beta 40 in senile plaques with end-specific A beta monoclonals: evidence that an initially deposited species is A beta 42(43). *Neuron* [Internet]. 1994;13(1):45–53 [cited 2015 Nov 18]. <http://www.ncbi.nlm.nih.gov/pubmed/8043280>
- [29] Selkoe DJ. Alzheimer's disease: genes, proteins, and therapy. *Physiol Rev* [Internet]. 2001;81(2):741–66 [cited 2015 Dec 12]. <http://www.ncbi.nlm.nih.gov/pubmed/11274343>
- [30] Dickson DW. The pathogenesis of senile plaques. *J Neuropathol Exp Neurol* [Internet]. 1997;56(4):321–39 [cited 2015 Nov 29]. <http://www.ncbi.nlm.nih.gov/pubmed/9100663>
- [31] Nalivaeva NN, Turner AJ. The amyloid precursor protein: a biochemical enigma in brain development, function and disease. *FEBS Lett* [Internet]. 2013;587(13):2046–54 [cited 2016 Jan 10]. <http://www.sciencedirect.com/science/article/pii/S0014579313003529>

- [32] Yoshikai S, Sasaki H, Doh-ura K, Furuya H, Sakaki Y. Genomic organization of the human amyloid beta-protein precursor gene. *Gene* [Internet]. 1990;87(2):257–63 [cited 2015 Dec 31]. <http://www.ncbi.nlm.nih.gov/pubmed/2110105>
- [33] Zhang Y, Thompson R, Zhang H, Xu H. APP processing in Alzheimer's disease. *Mol Brain* [Internet]. 2011;4:3 [cited 2015 Nov 18]. <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3022812&tool=pmcentrez&rendertype=abstract>
- [34] Zhou Z, Chan CH, Ma Q, Xu X, Xiao Z, Tan E. The roles of amyloid precursor protein (APP) in neurogenesis: implications to pathogenesis and therapy of Alzheimer disease. *Cell Adh Migr*. 2016;5(4):280–92.
- [35] Musa A, Lehrach H, Russo VA. Distinct expression patterns of two zebrafish homologues of the human APP gene during embryonic development. *Dev Genes Evol* [Internet]. 2001;211(11):563–7 [cited 2015 Nov 18]. <http://www.ncbi.nlm.nih.gov/pubmed/11862463>
- [36] Xiao Q, Gil S-C, Yan P, Wang Y, Han S, Gonzales E, et al. Role of phosphatidylinositol clathrin assembly lymphoid-myeloid leukemia (PICALM) in intracellular amyloid precursor protein (APP) processing and amyloid plaque pathogenesis. *J Biol Chem* [Internet]. 2012;287(25):21279–89 [cited 2016 Jan 10]. <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3375549&tool=pmcentrez&rendertype=abstract>
- [37] Cirrito JR, Kang J-E, Lee J, Stewart FR, Verges DK, Silverio LM, et al. Endocytosis is required for synaptic activity-dependent release of amyloid-beta in vivo. *Neuron* [Internet]. 2008;58(1):42–51 [cited 2015 Nov 1]. <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2390913&tool=pmcentrez&rendertype=abstract>
- [38] Seubert P, Vigo-Pelfrey C, Esch F, Lee M, Dovey H, Davis D, et al. Isolation and quantification of soluble Alzheimer's beta-peptide from biological fluids. *Nature* [Internet]. 1992;359(6393):325–7 [cited 2015 Dec 4]. <http://www.ncbi.nlm.nih.gov/pubmed/1406936>
- [39] Querfurth HW, LaFerla FM. Alzheimer's disease. *N Engl J Med* [Internet]. 2010;362(4):329–44 [cited 2014 Nov 18]. <http://www.ncbi.nlm.nih.gov/pubmed/20107219>
- [40] Selkoe DJ. Alzheimer disease: mechanistic understanding predicts novel therapies. *Ann Intern Med* [Internet]. 2004;140(8):627–38 [cited 2016 Jan 10]. <http://www.ncbi.nlm.nih.gov/pubmed/15096334>
- [41] Jarrett JT, Berger EP, Lansbury PT. The carboxy terminus of the beta amyloid protein is critical for the seeding of amyloid formation: implications for the pathogenesis of Alzheimer's disease. *Biochemistry* [Internet]. 1993;32(18):4693–7 [cited 2015 Oct 5]. <http://www.ncbi.nlm.nih.gov/pubmed/8490014>
- [42] Rossner S. New players in old amyloid precursor protein-processing pathways. *Int J Dev Neurosci* [Internet]. 2004;22(7):467–74 [cited 2016 Jan 11]. <http://www.sciencedirect.com/science/article/pii/S0736574804000899>

- [43] Yu G, Nishimura M, Arawaka S, Levitan D, Zhang L, Tandon A, et al. Nicastrin modulates presenilin-mediated notch/glp-1 signal transduction and betaAPP processing. *Nature* [Internet]. 2000;407(6800):48–54 [cited 2015 Dec 9]. <http://www.ncbi.nlm.nih.gov/pubmed/10993067>
- [44] Esler WP, Kimberly WT, Ostaszewski BL, Ye W, Diehl TS, Selkoe DJ, et al. Activity-dependent isolation of the presenilin–secretase complex reveals nicastrin and a substrate. *Proc Natl Acad Sci* [Internet]. 2002;99(5):2720–5 [cited 2016 Jan 11]. <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=122414&tool=pmcentrez&rendertype=abstract>
- [45] Francis R, McGrath G, Zhang J, Ruddy DA, Sym M, Apfeld J, et al. aph-1 and pen-2 are required for Notch pathway signaling, gamma-secretase cleavage of betaAPP, and presenilin protein accumulation. *Dev Cell* [Internet]. 2002;3(1):85–97 [cited 2016 Jan 11]. <http://www.ncbi.nlm.nih.gov/pubmed/12110170>
- [46] Helm K von der, Korant BD, Cheronis JCD. Proteases as targets for therapy, Issue 146 [Internet]. Springer Science & Business Media; 2000; p. 410 [cited 2016 Jan 11]. https://books.google.com/books?id=a5_eckV79UEC&pgis=1
- [47] Esposito LA. Measuring APP carboxy-terminal fragments. *Methods Mol Biol* [Internet]. 2011;670:71–84 [cited 2016 Jan 11]. <http://www.ncbi.nlm.nih.gov/pubmed/20967584>
- [48] Multhaup G, Huber O, Buée L, Galas M-C. Amyloid precursor protein (APP) metabolites APP intracellular fragment (AICD), A β 42, and Tau in nuclear roles. *J Biol Chem* [Internet]. 2015;290(39):23515–22 [cited 2016 Jan 11]. <http://www.jbc.org/content/290/39/23515.long>
- [49] Makarova A, Williams SE, Strickland DK. Proteases and lipoprotein receptors in Alzheimer’s disease. *Cell Biochem Biophys* [Internet]. 2004;41(1):139–78 [cited 2016 Jan 11]. <http://www.ncbi.nlm.nih.gov/pubmed/15371644>
- [50] Vingtdeux V, Hamdane M, Gompel M, Bégard S, Drobecq H, Ghestem A, et al. Phosphorylation of amyloid precursor carboxy-terminal fragments enhances their processing by a gamma-secretase-dependent mechanism. *Neurobiol Dis* [Internet]. 2005;20(2):625–37 [cited 2016 Jan 11]. <http://www.ncbi.nlm.nih.gov/pubmed/15936948>
- [51] Haass C, Selkoe DJ. Soluble protein oligomers in neurodegeneration: lessons from the Alzheimer’s amyloid beta-peptide. *Nat Rev Mol Cell Biol* [Internet]. Nature Publishing Group; 2007;8(2):101–12 [cited 2014 Jul 9]. doi:10.1038/nrm2101
- [52] Thinakaran G. The role of presenilins in Alzheimer’s disease. *J Clin Invest* [Internet]. 1999;104(10):1321–7 [cited 2016 Jan 11]. <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=409849&tool=pmcentrez&rendertype=abstract>
- [53] Shoji M, Golde TE, Ghiso J, Cheung TT, Estus S, Shaffer LM, et al. Production of the Alzheimer amyloid β protein by normal proteolytic processing. *Science* (80). 1992;258(5079):126–9.

- [54] Bennett DA, Schneider JA, Arvanitakis Z, Kelly JF, Aggarwal NT, Shah RC, et al. Neuropathology of older persons without cognitive impairment from two community-based studies. *Neurology*. 2006;66(12):1837–44.
- [55] Hardy JA, Higgins GA. Alzheimer's disease: the amyloid cascade hypothesis. *Science*. 1992;256(5054):184–5.
- [56] Reitz C. Alzheimer's disease and the amyloid cascade hypothesis: a critical review. *Int J Alzheimers Dis* [Internet]. 2012;2012:369808. <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3313573&tool=pmcentrez&rendertype=abstract>
- [57] Mudher A, Lovestone S. Alzheimer's disease - Do tauists and baptists finally shake hands? *Trends Neurosci*. 2002;25(1):22–6.
- [58] Chételat G. Alzheimer disease: β -independent processes-rethinking preclinical AD. *Nat Rev Neurol* [Internet]. 2013;9(3):123–4. <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3935395&tool=pmcentrez&rendertype=abstract>
- [59] Klein WL, Krafft G a., Finch CE. Targeting small A β oligomers: the solution to an Alzheimer's disease conundrum? *Trends Neurosci*. 2001;24(4):219–24.
- [60] Gilbert BJ. The role of amyloid β in the pathogenesis of Alzheimer's disease. *J Clin Pathol* [Internet]. 2013;66(5):362–6 [cited 2016 Jan 16]. <http://www.ncbi.nlm.nih.gov/pubmed/23526599>
- [61] Jin M, Shepardson N, Yang T, Chen G, Walsh D, Selkoe DJ. Soluble amyloid beta-protein dimers isolated from Alzheimer cortex directly induce Tau hyperphosphorylation and neuritic degeneration. *Proc Natl Acad Sci USA* [Internet]. 2011;108(14):5819–24. <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3078381&tool=pmcentrez&rendertype=abstract>
- [62] Jan A, Adolfsson O, Allaman I, Buccarello AL, Magistretti PJ, Pfeifer A, et al. A β 42 neurotoxicity is mediated by ongoing nucleated polymerization process rather than by discrete A β 42 species. *J Biol Chem*. 2011;286(10):8585–96.
- [63] Puzzo D, Arancio O. Amyloid- β peptide: Dr. Jekyll or Mr. Hyde? *J Alzheimers Dis* [Internet]. 2013;33 Suppl 1:S111–20 [cited 2016 Jan 30]. <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3696497&tool=pmcentrez&rendertype=abstract>
- [64] Robinson SR, Bishop GM. A β as a bioflocculant: implications for the amyloid hypothesis of Alzheimer's disease. *Neurobiol Aging* [Internet] (Elsevier). 2002;23(6):1051–72 [cited 2016 Jan 30]. <http://www.neurobiologyofaging.org/article/S0197458001003426/fulltext>
- [65] Lee H, Zhu X, Castellani RJ, Nunomura A, Perry G, Smith MA. Amyloid-beta in Alzheimer disease: the null versus the alternate hypotheses. *J Pharmacol Exp Ther* [Internet]. 2007;321(3):823–9. <http://www.ncbi.nlm.nih.gov/pubmed/17229880>

- [66] Maccioni RB, Farías G, Morales I, Navarrete L. The revitalized tau hypothesis on Alzheimer's disease. *Arch Med Res* [Internet]. 2010;41(3):226–31. <http://www.science-direct.com/science/article/pii/S0188440910000500>
- [67] Nagy Z, Esiri MM, Jobst KA, Morris JH, King EM, McDonald B, et al. Relative roles of plaques and tangles in the dementia of Alzheimer's disease: correlations using three sets of neuropathological criteria. *Dementia*. 1995;6(1):21–31.
- [68] Goedert M, Jakes R. Mutations causing neurodegenerative tauopathies. *Biochim Biophys Acta Mol Basis Dis*. 2005;1739(2):240–50.
- [69] Hashiguchi M, Hashiguchi T. Kinase-Kinase interaction and modulation of Tau phosphorylation. *Int Rev Cell Mol Biol*. 2013;300:121–60.
- [70] Hooper C, Killick R, Lovestone S. The GSK3 hypothesis of Alzheimer's disease. *J Neurochem*. 2008;104(6):1433–9.
- [71] Hernández F, Borrell J, Guaza C, Avila J, Lucas JJ. Spatial learning deficit in transgenic mice that conditionally over-express GSK-3 β in the brain but do not form tau filaments. *J Neurochem*. 2002;83:1529–33.
- [72] DaRocha-Souto B, Coma M, Pérez-Nievas BG, Scotton TC, Siao M, Sánchez-Ferrer P, et al. Activation of glycogen synthase kinase-3 beta mediates β -amyloid induced neuritic damage in Alzheimer's disease. *Neurobiol Dis* [Internet]. 2012;45(1):425–37. doi: 10.1016/j.nbd.2011.09.002
- [73] Hernández F, Gómez de Barreda E, Fuster-Matanzo A, Lucas JJ, Avila J. GSK3: a possible link between beta amyloid peptide and tau protein. *Exp Neurol* [Internet]. 2010;223(2):322–5. doi: 10.1016/j.expneurol.2009.09.011
- [74] Hoshi M, Takashima a, Noguchi K, Murayama M, Sato M, Kondo S, et al. Regulation of mitochondrial pyruvate dehydrogenase activity by tau protein kinase I/glycogen synthase kinase 3beta in brain. *Proc Natl Acad Sci USA*. 1996;93(7):2719–23.
- [75] Hetman M, Cavanaugh JE, Kimelman D, Xia Z. Role of glycogen synthase kinase-3beta in neuronal apoptosis induced by trophic withdrawal. *J Neurosci*. 2000;20(7):2567–74.
- [76] Lucas JJ, Hernández F, Gómez-Ramos P, Morán MA, Hen R, Avila J. Decreased nuclear beta-catenin, tau hyperphosphorylation and neurodegeneration in GSK-3beta conditional transgenic mice. *EMBO J* [Internet]. 2001;20(1–2):27–39. <http://www.pubmed-central.nih.gov/articlerender.fcgi?artid=140191&tool=pmcentrez&rendertype=abstract>
- [77] Takashima A, Noguchi K, Michel G, Mercken M, Hoshi M, Ishiguro K, et al. Exposure of rat hippocampal neurons to amyloid β peptide (25-35) induces the inactivation of phosphatidylinositol-3 kinase and the activation of tau protein kinase I/glycogen synthase kinase-3 β . *Neurosci Lett*. 1996;203:33–6.

- [78] Lei P, Ayton S, Bush AI, Adlard P a. GSK-3 in Neurodegenerative Diseases. *Int J Alzheimers Dis*. 2011;2011:189246.
- [79] Padurariu M, Ciobica A, Lefter R, Serban IL, Stefanescu C, Chirita R. The oxidative stress hypothesis in Alzheimer's disease. *Psychiatr Danub* [Internet]. 2013;25(4):401–9. <http://www.ncbi.nlm.nih.gov/pubmed/24247053>
- [80] Markesbery WR. Oxidative stress hypothesis in Alzheimer's disease. *Free Radic Biol Med*. 1997;23(1):134–47.
- [81] Graham SF, Nasaruddin M Bin, Carey M, Holscher C, McGuinness B, Kehoe PG, et al. Age-associated changes of brain copper, iron, and zinc in Alzheimer's disease and dementia with Lewy bodies. *J Alzheimers Dis* [Internet]. 2014;42(4):1407–13. <http://content.iospress.com/articles/journal-of-alzheimers-disease/jad140684>
- [82] Deibel MA, Ehmann WD, Markesbery WR. Copper, iron, and zinc imbalances in severely degenerated brain regions in Alzheimer's disease: possible relation to oxidative stress. *J Neurol Sci*. 1996;143(606):137–42.
- [83] Schrag M, Mueller C, Oyoyo U, Smith M a., Kirsch WM. Iron, zinc and copper in the Alzheimer's disease brain: a quantitative meta-analysis. Some insight on the influence of citation bias on scientific opinion. *Prog Neurobiol* [Internet]. 2011;94(3):296–306. doi:10.1016/j.pneurobio.2011.05.001
- [84] Tiiman A, Palumaa P, Tõugu V. The missing link in the amyloid cascade of Alzheimer's disease-Metal ions. *Neurochem Int* [Internet]. 2013;62(4):367–78. doi:10.1016/j.neuint.2013.01.023
- [85] Mold M, Ouro-Gnao L, Wieckowski BM, Exley C. Copper prevents amyloid- β (1-42) from forming amyloid fibrils under near-physiological conditions in vitro. *Sci Rep*. 2013;3(i):1256.
- [86] Mayes J, Tinker-Mill C, Kolosov O, Zhang H, Tabner BJ, Allsop D. Amyloid fibrils in alzheimer disease are not inert when bound to copper ions but can degrade hydrogen peroxide and generate reactive oxygen species. *J Biol Chem*. 2014;289(17):12052–62.
- [87] Swerdlow RH, Khan SM. A "mitochondrial cascade hypothesis" for sporadic Alzheimer's disease. *Med Hypotheses*. 2004;63(1):8–20.
- [88] Grimm A, Friedland K, Eckert A. Mitochondrial dysfunction: the missing link between aging and sporadic Alzheimer's disease. *Biogerontology* [Internet]. 2015 [cited 2015 Oct 16]. <http://www.ncbi.nlm.nih.gov/pubmed/26468143>
- [89] Russell H, Swerdlow SMK. The Alzheimer's Disease mitochondrial cascade hypothesis: an update. *Exp Neurol*. 2009;218(2):308–15.
- [90] Perry E, Perry R, Blessed G, Tomlinson B. Necropsy evidence of central cholinergic deficits in senile dementia. *Lancet* [Internet]. 1977;309(8004):189 [cited 2016 Jan 15]. <http://www.sciencedirect.com/science/article/pii/S0140673677917809>

- [91] Pinto T, Lanctôt KL, Herrmann N. Revisiting the cholinergic hypothesis of behavioral and psychological symptoms in dementia of the Alzheimer's type. *Ageing Res Rev* [Internet]. 2011;10(4):404–12. doi:10.1016/j.arr.2011.01.003
- [92] Coyle JT, Price DL, DeLong MR. Alzheimer's disease: a disorder of cortical cholinergic innervation. *Science*. 1983;219(4589):1184–90.
- [93] Pákási M, Kálmán J. Interactions between the amyloid and cholinergic mechanisms in Alzheimer's disease. *Neurochem Int*. 2008;53(3):103–11.
- [94] Gilmor ML, Erickson JD, Varoqui H, Hersh LB, Bennett DA, Cochran EJ, et al. Preservation of nucleus basalis neurons containing choline acetyltransferase and the vesicular acetylcholine transporter in the elderly with mild cognitive impairment and early Alzheimer's disease. *J Comp Neurol* [Internet]. 1999;411(4):693–704. <http://www.ncbi.nlm.nih.gov/pubmed/10421878>
- [95] DeKosky ST, Ikonomic MD, Styren SD, Beckett L, Wisniewski S, Bennett DA, et al. Upregulation of choline acetyltransferase activity in hippocampus and frontal cortex of elderly subjects with mild cognitive impairment. *Ann Neurol* [Internet]. 2002;51(2):145–55. <http://www.ncbi.nlm.nih.gov/pubmed/11835370>
- [96] Di Santo SG, Prinelli F, Adorni F, Caltagirone C, Musicco M. A meta-analysis of the efficacy of donepezil, rivastigmine, galantamine, and memantine in relation to severity of Alzheimer's disease. *J Alzheimer's Dis*. 2013;35(2):349–61.
- [97] Craig L a., Hong NS, McDonald RJ. Revisiting the cholinergic hypothesis in the development of Alzheimer's disease. *Neurosci Biobehav Rev* [Internet]. 2011;35(6):1397–409. doi:10.1016/j.neubiorev.2011.03.001
- [98] Khachaturian ZS. Calcium, membranes, aging, and Alzheimer's disease. Introduction and overview. *Ann N Y Acad Sci* [Internet]. 1989;568:1–4 [cited 2016 Jan 16]. <http://www.ncbi.nlm.nih.gov/pubmed/2629579>
- [99] Berridge MJ. Calcium signalling and Alzheimer's disease. *Neurochem Res*. 2011;36(7):1149–56.
- [100] LaFerla FM. Calcium dyshomeostasis and intracellular signalling in Alzheimer's disease. *Nat Rev Neurosci*. 2002;3(11):862–72.
- [101] Pierrot N, Ghisdal P, Caumont A-S, Octave J-N. Intraneuronal amyloid-beta1-42 production triggered by sustained increase of cytosolic calcium concentration induces neuronal death. *J Neurochem*. 2004;88(5):1140–50.
- [102] Mattson MP, Cheng B, Davis D, Bryant K, Lieberburg I, Rydel RE. beta-Amyloid peptides destabilize calcium homeostasis and render human cortical neurons vulnerable to excitotoxicity. *J Neurosci*. 1992;12(2):376–89.

- [103] Pierrot N, Santos SF, Feyt C, Morel M, Brion JP, Octave JN. Calcium-mediated transient phosphorylation of tau and amyloid precursor protein followed by intraneuronal amyloid- β accumulation. *J Biol Chem*. 2006;281(52):39907–14.
- [104] Glabe CG. Common mechanisms of amyloid oligomer pathogenesis in degenerative disease. *Neurobiol Aging*. 2006;27(4):570–5.
- [105] Strimbu K, Tavel JA. What are biomarkers? *Curr Opin HIV AIDS* [Internet]. 2010;5(6):463–6. <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3078627&tool=pmcentrez&rendertype=abstract>
- [106] Rosén C, Hansson O, Blennow K, Zetterberg H. Fluid biomarkers in Alzheimer's disease—current concepts. *Mol Neurodegener* [Internet]. 2013;8(1):20. <http://www.molecularneurodegeneration.com/content/8/1/20>
- [107] Hansson O, Zetterberg H, Buchhave P, Londos E, Blennow K, Minthon L. Association between CSF biomarkers and incipient Alzheimer's disease in patients with mild cognitive impairment: a follow-up study. *Lancet Neurol*. 2006;5(3):228–34.
- [108] Andreasson U, Lautner R, Schott JM, Mattsson N, Hansson O, Herukka S-K, et al. CSF biomarkers for Alzheimer's pathology and the effect size of APOE ϵ 4. *Mol Psychiatry* [Internet]. 2014;19(2):148–9 [cited 2016 Jan 17]. <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3903112&tool=pmcentrez&rendertype=abstract>
- [109] Ringman JM, Younkin SG, Pratico D, Seltzer W, Cole GM, Geschwind DH, et al. Biochemical markers in persons with preclinical familial Alzheimer disease. *Neurology*. 2008;71(2):85–92.
- [110] Graff-Radford NR, Crook JE, Lucas J, Boeve BF, Knopman DS, Ivnik RJ, et al. Association of low plasma A β 42/A β 40 ratios with increased imminent risk for mild cognitive impairment and Alzheimer disease. *Arch Neurol* [Internet]. 2007;64(3):354–62. <http://archneur.jamanetwork.com/article.aspx?articleid=793567>
- [111] Verheijen JH, Huisman LGM, van Lent N, Neumann U, Paganetti P, Hack CE, et al. Detection of a soluble form of BACE-1 in human cerebrospinal fluid by a sensitive activity assay. *Clin Chem* [Internet]. 2006;52(6):1168–74. <http://www.ncbi.nlm.nih.gov/pubmed/16614000>
- [112] Perneczky R, Tsolakidou a., Arnold a., Diehl-Schmid J, Grimmer T, Förstl H, et al. CSF soluble amyloid precursor proteins in the diagnosis of incipient Alzheimer disease. *Neurology*. 2011;77(1):35–8.
- [113] Lopez-Font I, Cuchillo-Ibañez I, Sogorb-Esteve A, García-Ayllón M-S, Sáez-Valero J. Transmembrane amyloid-related proteins in CSF as potential biomarkers for Alzheimer's disease. *Front Neurol* [Internet]. 2015;6(June):125. <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=4451586&tool=pmcentrez&rendertype=abstract>
- [114] Ribeiro CA, Santana I, Oliveira C, Baldeiras I, Moreira J, Saraiva MJ, et al. Transthyretin decrease in plasma of MCI and AD patients: investigation of mechanisms for disease

- modulation. *Curr Alzheimer Res* [Internet]. 2012;9(8):881–9. <http://www.ncbi.nlm.nih.gov/pubmed/22698061>
- [115] Gloeckner SF, Meyne F, Wagner F, Heinemann U, Krasnianski A, Meissner B, et al. Quantitative analysis of transthyretin, tau and amyloid-beta in patients with dementia. *J Alzheimers Dis* [Internet]. 2008;14(1):17–25 [cited 2016 Jan 5]. <http://www.ncbi.nlm.nih.gov/pubmed/18525124>
- [116] Fagan AM, Perrin RJ. Upcoming candidate cerebrospinal fluid biomarkers of Alzheimer's disease. *Biomark Med* [Internet]. 2012;6(4):455–76. <http://www.ncbi.nlm.nih.gov/pubmed/22917147>
- [117] Petersen RC, Jack Jr. CR. Imaging and biomarkers in early Alzheimer's disease and mild cognitive impairment. *Clin Pharmacol Ther* [Internet]. 2009;86(4):438–41. <http://www.nature.com/clpt/journal/v86/n4/pdf/clpt2009166a.pdf>
- [118] Klunk WE, Engler H, Nordberg A, Wang Y, Blomqvist G, Holt DP, et al. Imaging brain amyloid in Alzheimer's disease with Pittsburgh compound-B. *Ann Neurol*. 2004;55(3):306–19.
- [119] Trzepacz PT, Yu P, Sun J, Schuh K, Case M, Witte MM, et al. Comparison of neuroimaging modalities for the prediction of conversion from mild cognitive impairment to alzheimer's dementia. *Neurobiol Aging*. 2014;35(1):143–51.
- [120] Castellani RJ, Rolston RK, Smith MA. Alzheimer disease. *Dis Mon* [Internet]. 2010;56(9):484–546 [cited 2015 Dec 11]. <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2941917&tool=pmcentrez&rendertype=abstract>
- [121] Brys M, Glodzik L, Mosconi L, Switalski R, De Santi S, Pirraglia E, et al. Magnetic resonance imaging improves cerebrospinal fluid biomarkers in the early detection of Alzheimer's disease. *J Alzheimer's Dis*. 2009;16(2):351–62.
- [122] Herrup K. Reimagining Alzheimer's disease—an age-based hypothesis. *J Neurosci* [Internet]. 2010;30(50):16755–62. <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3004746&tool=pmcentrez&rendertype=abstract>
- [123] Viña J, Lloret A. Why women have more Alzheimer's disease than men: gender and mitochondrial toxicity of amyloid-beta peptide. *J Alzheimers Dis* [Internet]. 2010;20 Suppl 2(S2):S527–33. <http://content.iospress.com/articles/journal-of-alzheimers-disease/jad100501?resultNumber=0&totalResults=105&start=0&q=Why+women+have+more+Alzheimer%27s+disease+than+men%3A+gender+and+mitochondrial+toxicity+of+amyloid-beta+peptid&resultsPageSize=10&rows=1>
- [124] Li R, Singh M. Sex differences in cognitive impairment and Alzheimer's disease. *Front Neuroendocrinol* [Internet]. 2014;35(3):385–403. doi:10.1016/j.yfrne.2014.01.002
- [125] Arvanitakis Z, Wilson RS, Bienias JL, Evans D a, Bennett D a. Diabetes mellitus and risk of Alzheimer disease and decline in cognitive function. *Arch Neurol*. 2004;61(5):661–6.

- [126] Vuorinen M, Solomon A, Rovio S, Nieminen L, Kåreholt I, Tuomilehto J, et al. Changes in vascular risk factors from midlife to late life and white matter lesions: a 20-year follow-up study. *Dement Geriatr Cogn Disord*. 2011;31(2):119–25.
- [127] Imtiaz B, Tolppanen A-M, Kivipelto M, Soininen H. Future directions in Alzheimer's disease from risk factors to prevention. *Biochem Pharmacol* [Internet]. 2014;88(4):661–70. <http://www.ncbi.nlm.nih.gov/pubmed/24418410>
- [128] Shepardson NE, Shankar GM, Selkoe DJ. Cholesterol level and statin use in Alzheimer disease: II. Review of human trials and recommendations. *Arch Neurol* [Internet]. 2011;68(11):1385–92. <http://eutils.ncbi.nlm.nih.gov/entrez/eutils/elink.fcgi?dbfrom=pubmed&id=22084122&retmode=ref&cmd=prlinks\ npapers3://publication/doi/10.1001/archneurol.2011.242>
- [129] Lindsay J, Laurin D, Verreault R, Hébert R, Helliwell B, Hill GB, et al. Risk factors for Alzheimer's disease: a prospective analysis from the Canadian Study of Health and Aging. *Am J Epidemiol*. 2002;156(5):445–53.
- [130] Corder EH, Saunders a M, Strittmatter WJ, Schmechel DE, Gaskell PC, Small GW, et al. Gene dose of apolipoprotein E type 4 allele and the risk of Alzheimer's disease in late onset families. *Science*. 1993;261(5123):921–3.
- [131] Tanzi RE, Bertram L. Twenty years of the Alzheimer's disease amyloid hypothesis: a genetic perspective. *Cell*. 2005;120(4):545–55.
- [132] Rinne JO, Brooks DJ, Rossor MN, Fox NC, Bullock R, Klunk WE, et al. 11C-PiB PET assessment of change in fibrillar amyloid-beta load in patients with Alzheimer's disease treated with bapineuzumab: a phase 2, double-blind, placebo-controlled, ascending-dose study. *Lancet Neurol* [Internet]. 2010;9(4):363–72 [cited 2015 Aug 12]. <http://www.ncbi.nlm.nih.gov/pubmed/20189881>
- [133] Gandy S, DeKosky ST. Toward the treatment and prevention of Alzheimer's disease: rational strategies and recent progress. *Annu Rev Med* [Internet]. 2013;64:367–83. <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3625402&tool=pmcentrez&rendertype=abstract>
- [134] Watt AD, Crespi GAN, Down RA, Ascher DB, Gunn A, Perez KA, et al. Do current therapeutic anti-A β antibodies for Alzheimer's disease engage the target? *Acta Neuropathol*. 2014;127(6):803–10.
- [135] Gong B, Pan Y, Zhao W, Knable L, Vempati P, Begum S, et al. IVIG immunotherapy protects against synaptic dysfunction in Alzheimer's disease through complement anaphylatoxin C5a-mediated AMPA-CREB-C/EBP signaling pathway. *Mol Immunol*. 2013;56(4):619–29.
- [136] Laxton AW, Tang-Wai DF, McAndrews MP, Zumsteg D, Wennberg R, Keren R, et al. A phase I trial of deep brain stimulation of memory circuits in Alzheimer's disease.

- Ann Neurol [Internet]. 2010;68(4):521–34 [cited 2016 Jan 18]. <http://www.ncbi.nlm.nih.gov/pubmed/20687206>
- [137] Laxton AW, Lozano AM. Deep brain stimulation for the treatment of alzheimer disease and dementias. World Neurosurg [Internet]. 2013;80(3–4):S28.e1–S28.e8. doi: 10.1016/j.wneu.2012.06.028
- [138] Zlokovic BV. The blood-brain barrier in health and chronic neurodegenerative disorders. Neuron [Internet]. 2008;57(2):178–201 [cited 2015 Jul 15]. <http://www.ncbi.nlm.nih.gov/pubmed/18215617>
- [139] Ehrlich P. Das sauerstoffbedürfnis des organismus. Eine Farbenanalytische Stud. 1885
- [140] Goldmann E. Vitalfärbung am zentralnervensystem. Abh Preuss Akd Wiss Phys Math. 1913;1(1):1–13.
- [141] Lewandowsky M. Zur lehre von der cerebrospinalflüssigkeit. Z Klin Med. 1900;40:480–94.
- [142] Hawkins BT, Davis TP. The blood-brain barrier/neurovascular unit in health and disease. Pharmacol Rev [Internet]. 2005;57(2):173–85. <http://www.ncbi.nlm.nih.gov/pubmed/15914466>
- [143] Brightman MW, Reese TS. Junctions between intimately apposed cell membranes in the vertebrate brain. J Cell Biol. 1969;40(3):648–77.
- [144] Reese TS, Karnovsky MJ. Fine structural localization of a blood-brain barrier to exogenous peroxidase. J Cell Biol [Internet]. 1967;34(1):207–17. <http://www.pubmed-central.nih.gov/articlerender.fcgi?artid=2107213&tool=pmcentrez&rendertype=abstract>
- [145] Bates KA, Verdile G, Li Q-X, Ames D, Hudson P, Masters CL, et al. Clearance mechanisms of Alzheimer’s amyloid-beta peptide: implications for therapeutic design and diagnostic tests. Mol Psychiatry [Internet]. Nature Publishing Group; 2009;14(5):469–86 [cited 2015 Dec 28]. doi: 10.1038/mp.2008.96
- [146] Pardridge WM. Molecular biology of the blood-brain barrier. Mol Biotechnol [Internet]. 2005;30(1):57–70. <http://www.ncbi.nlm.nih.gov/pubmed/15805577>
- [147] Oldendorf WH, Cornford ME, Brown WJ. The large apparent work capability of the blood-brain barrier: a study of the mitochondrial content of capillary endothelial cells in brain and other tissues of the rat. Ann Neurol [Internet]. 1977;1(5):409–17 [cited 2015 Dec 1]. <http://www.ncbi.nlm.nih.gov/pubmed/617259>
- [148] Fenstermacher J, Gross P, Sposito N, Acuff V, Pettersen S, Gruber K. Structural and functional variations in capillary systems within the brain. Ann N Y Acad Sci. 1988;529:21–30.
- [149] Sedlakova R, Shivers RR, Del Maestro RF. Ultrastructure of the blood-brain barrier in the rabbit. J Submicrosc Cytol Pathol. 1999;31:149–61.

- [150] Kniessel U, Wolburg H. Tight junctions of the blood-brain barrier. *Cell Mol Neurobiol*. 2000;20(1):57–76.
- [151] Ohtsuki S, Terasaki T. Contribution of carrier-mediated transport systems to the blood-brain barrier as a supporting and protecting interface for the brain; importance for CNS drug discovery and development. *Pharm Res*. 2007;24(9):1745–58.
- [152] Tagami M, Nara Y, Kubota A, Fujino H, Yamori Y. Ultrastructural changes in cerebral pericytes and astrocytes of stroke-prone spontaneously hypertensive rats. *Stroke*. 1990;21:1064–71.
- [153] Farkas E, Luiten PGM. Cerebral microvascular pathology in aging and Alzheimer's disease. Vol. 64, *Progress in Neurobiology*. 2001. 575-611 p.
- [154] Ramsauer M, Krause D, Dermietzel R. Angiogenesis of the blood-brain barrier in vitro and the function of cerebral pericytes. *FASEB J*. 2002;16(10):1274–6.
- [155] Gonul E, Duz B, Kahraman S, Kayali H, Kubar A, Timurkaynak E. Early pericyte response to brain hypoxia in cats: an ultrastructural study. *Microvasc Res* [Internet]. 2002;64(1):116–9. <http://www.ncbi.nlm.nih.gov/pubmed/12074637>
- [156] Dore-Duffy P, Owen C, Balabanov R, Murphy S, Beaumont T, Rafols JA. Pericyte migration from the vascular wall in response to traumatic brain injury. *Microvasc Res* [Internet]. 2000;60(1):55–69. <http://www.ncbi.nlm.nih.gov/pubmed/10873515>
- [157] Hori S, Ohtsuki S, Hosoya K, Nakashima E, Terasaki T. A pericyte-derived angiopoietin-1 multimeric complex induces occludin gene expression in brain capillary endothelial cells through Tie-2 activation in vitro. *J Neurochem* [Internet]. 2004;89(2):503–13. <http://www.ncbi.nlm.nih.gov/pubmed/15056293>
- [158] Davson H, Oldendorf WH. Symposium on membrane transport. Transport in the central nervous system. *Proc R Soc Med* [Internet]. 1967;60(4):326–9 [cited 2016 Jan 12]. <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=1901728&tool=pmcentrez&rendertype=abstract>
- [159] Tao-Cheng JH, Nagy Z, Brightman MW. Tight junctions of brain endothelium in vitro are enhanced by astroglia. *J Neurosci* [Internet]. 1987;7(10):3293–9 [cited 2016 Jan 12]. <http://www.ncbi.nlm.nih.gov/pubmed/3668629>
- [160] Maxwell K, Berliner JA, Cancilla PA. Induction of gamma-glutamyl transpeptidase in cultured cerebral endothelial cells by a product released by astrocytes. *Brain Res* [Internet]. 1987;410(2):309–14 [cited 2016 Jan 12]. <http://www.ncbi.nlm.nih.gov/pubmed/2885071>
- [161] Ballabh P, Braun A, Nedergaard M. The blood-brain barrier: an overview: structure, regulation, and clinical implications. *Neurobiol Dis* [Internet]. 2004;16(1):1–13 [cited 2014 Jul 11]. <http://www.ncbi.nlm.nih.gov/pubmed/15207256>

- [162] Braet K, Paemeleire K, D'Herde K, Sanderson MJ, Leybaert L. Astrocyte-endothelial cell calcium signals conveyed by two signalling pathways. *Eur J Neurosci*. 2001;13(1):79–91.
- [163] Zonta M, Angulo MC, Gobbo S, Rosengarten B, Hossmann K-A, Pozzan T, et al. Neuron-to-astrocyte signaling is central to the dynamic control of brain microcirculation. *Nat Neurosci* [Internet]. 2003;6(1):43–50 [cited 2016 Jan 12]. <http://www.ncbi.nlm.nih.gov/pubmed/12469126>
- [164] Anderson CM, Nedergaard M. Astrocyte-mediated control of cerebral microcirculation. *Trends Neurosci* [Internet]. 2003;26(7):340–4 [cited 2016 Jan 12] (author reply 344–5). <http://www.ncbi.nlm.nih.gov/pubmed/12850427>
- [165] Alliot F, Godin I, Pessac B. Microglia derive from progenitors, originating from the yolk sac, and which proliferate in the brain. *Brain Res Dev Brain Res* [Internet]. 1999;117(2):145–52 [cited 2015 Dec 20]. <http://www.ncbi.nlm.nih.gov/pubmed/10567732>
- [166] Hickey WF, Kimura H. Perivascular microglial cells of the CNS are bone marrow-derived and present antigen in vivo. *Science* [Internet]. 1988;239(4837):290–2 [cited 2015 Dec 20]. <http://www.ncbi.nlm.nih.gov/pubmed/3276004>
- [167] Fantin A, Vieira JM, Gestri G, Denti L, Schwarz Q, Prykhodzhiy S, et al. Tissue macrophages act as cellular chaperones for vascular anastomosis downstream of VEGF-mediated endothelial tip cell induction. *Blood* [Internet]. 2010;116(5):829–40 [cited 2015 Dec 8]. <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2938310&tool=pmcentrez&rendertype=abstract>
- [168] Nayak D, Roth TL, McGavern DB. Microglia development and function. *Annu Rev Immunol* [Internet]. 2014;32:367–402 [cited 2015 Nov 24]. <http://www.ncbi.nlm.nih.gov/pubmed/24471431>
- [169] Espinosa-Heidmann DG, Suner IJ, Hernandez EP, Monroy D, Csaky KG, Cousins SW. Macrophage depletion diminishes lesion size and severity in experimental choroidal neovascularization. *Invest Ophthalmol Vis Sci* [Internet]. 2003;44(8):3586–92 [cited 2016 Jan 13]. <http://www.ncbi.nlm.nih.gov/pubmed/12882811>
- [170] Sumi N, Nishioku T, Takata F, Matsumoto J, Watanabe T, Shuto H, et al. Lipopolysaccharide-activated microglia induce dysfunction of the blood-brain barrier in rat microvascular endothelial cells co-cultured with microglia. *Cell Mol Neurobiol* [Internet]. 2010;30(2):247–53 [cited 2016 Jan 13]. <http://www.ncbi.nlm.nih.gov/pubmed/19728078>
- [171] Nishioku T, Matsumoto J, Dohgu S, Sumi N, Miyao K, Takata F, et al. Tumor necrosis factor-alpha mediates the blood-brain barrier dysfunction induced by activated microglia in mouse brain microvascular endothelial cells. *J Pharmacol Sci* [Internet]. 2010;112(2):251–4 [cited 2016 Jan 13]. <http://www.ncbi.nlm.nih.gov/pubmed/20118615>
- [172] Obermeier B, Daneman R, Ransohoff RM. Development, maintenance and disruption of the blood-brain barrier. *Nat Med* [Internet]. 2013;19(12):1584–96 [cited 2014 Jul 10].

<http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=4080800&tool=pmcentrez&rendertype=abstract>

- [173] Bolton SJ, Anthony DC, Perry VH. Loss of the tight junction proteins occludin and zonula occludens-1 from cerebral vascular endothelium during neutrophil-induced blood-brain barrier breakdown in vivo. *Neuroscience* [Internet]. 1998;86(4):1245–57 [cited 2016 Jan 13]. <http://www.ncbi.nlm.nih.gov/pubmed/9697130>
- [174] da Fonseca ACC, Matias D, Garcia C, Amaral R, Geraldo LH, Freitas C, et al. The impact of microglial activation on blood-brain barrier in brain diseases. *Front Cell Neurosci* [Internet]. 2014;8:362 [cited 2016 Jan 13]. <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=4217497&tool=pmcentrez&rendertype=abstract>
- [175] Bazzoni G, Dejana E. Endothelial cell-to-cell junctions: molecular organization and role in vascular homeostasis. *Physiol Rev* [Internet]. 2004;84(3):869–901 [cited 2015 Aug 6]. <http://www.ncbi.nlm.nih.gov/pubmed/15269339>
- [176] Furuse M, Hirase T, Itoh M, Nagafuchi A, Yonemura S, Tsukita S. Occludin: a novel integral membrane protein localizing at tight junctions. *J Cell Biol* [Internet]. 1993;123(6 Pt 2):1777–88 [cited 2015 Mar 19]. <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2290891&tool=pmcentrez&rendertype=abstract>
- [177] Lippoldt A, Kniesel U, Liebner S, Kalbacher H, Kirsch T, Wolburg H, et al. Structural alterations of tight junctions are associated with loss of polarity in stroke-prone spontaneously hypertensive rat blood-brain barrier endothelial cells. *Brain Res* [Internet]. 2000;885(2):251–61 [cited 2016 Jan 13]. <http://www.ncbi.nlm.nih.gov/pubmed/11102579>
- [178] Hawkins BT, Abbruscato TJ, Egleton RD, Brown RC, Huber JD, Campos CR, et al. Nicotine increases in vivo blood-brain barrier permeability and alters cerebral microvascular tight junction protein distribution. *Brain Res* [Internet]. 2004;1027(1–2):48–58 [cited 2015 Dec 22]. <http://www.ncbi.nlm.nih.gov/pubmed/15494156>
- [179] Hirase T, Staddon JM, Saitou M, Ando-Akatsuka Y, Itoh M, Furuse M, et al. Occludin as a possible determinant of tight junction permeability in endothelial cells. *J Cell Sci* [Internet]. 1997;110(Pt 1):1603–13 [cited 2016 Jan 13]. <http://www.ncbi.nlm.nih.gov/pubmed/9247194>
- [180] Bamforth SD, Kniesel U, Wolburg H, Engelhardt B, Risau W. A dominant mutant of occludin disrupts tight junction structure and function. *J Cell Sci* [Internet]. 1999;112(Pt 1):1879–88. <http://www.ncbi.nlm.nih.gov/pubmed/10341207>
- [181] Saitou M, Furuse M, Sasaki H, Schulzke JD, Fromm M, Takano H, et al. Complex phenotype of mice lacking occludin, a component of tight junction strands. *Mol Biol Cell*. 2000;11(12):4131–42.

- [182] Schulzke JD, Gitter a. H, Mankertz J, Spiegel S, Seidler U, Amasheh S, et al. Epithelial transport and barrier function in occludin-deficient mice. *Biochim Biophys Acta Biomembr.* 2005;1669(1):34–42.
- [183] Murata M, Kojima T, Yamamoto T, Go M, Takano KI, Osanai M, et al. Down-regulation of survival signaling through MAPK and Akt in occludin-deficient mouse hepatocytes in vitro. *Exp Cell Res.* 2005;310:140–51.
- [184] Morgan L, Shah B, Rivers LE, Barden L, Groom AJ, Chung R, et al. Inflammation and dephosphorylation of the tight junction protein occludin in an experimental model of multiple sclerosis. *Neuroscience [Internet].* 2007;147(3):664–73. <http://www.sciencedirect.com/science/article/pii/S030645220700468X>
- [185] Rosenberg GA, Yang Y. Vasogenic edema due to tight junction disruption by matrix metalloproteinases in cerebral ischemia. *Neurosurg Focus [Internet].* 2007;22(5):E4 [cited 2016 Jan 14]. <http://www.ncbi.nlm.nih.gov/pubmed/17613235>
- [186] Piontek J, Winkler L, Wolburg H, Müller SL, Zuleger N, Piehl C, et al. Formation of tight junction: determinants of homophilic interaction between classic claudins. *FASEB J.* 2008;22(1):146–58.
- [187] Rolf Dermietzel, David C. Spray MN, editor. *Blood-brain barriers: from ontogeny to artificial interfaces*, vol 1. Wiley Online Library. Wiley-VCH Verlag GmbH & Co. KGaA; 2007. 741 p.
- [188] Nitta T, Hata M, Gotoh S, Seo Y, Sasaki H, Hashimoto N, et al. Size-selective loosening of the blood-brain barrier in claudin-5-deficient mice. *J Cell Biol [Internet].* 2003;161(3): 653–60. <http://jcb.rupress.org/content/161/3/653.long>
- [189] Kubota K, Furuse M, Sasaki H, Sonoda N, Fujita K, Nagafuchi A, et al. Ca(2+)-independent cell-adhesion activity of claudins, a family of integral membrane proteins localized at tight junctions. *Curr Biol [Internet].* 1999;9(18):1035–8 [cited 2016 Jan 14]. <http://www.ncbi.nlm.nih.gov/pubmed/10508613>
- [190] Jia W, Martin TA, Zhang G, Jiang WG. Junctional adhesion molecules in cerebral endothelial tight junction and brain metastasis. *Anticancer Res [Internet].* 2013;33(6): 2353–9 [cited 2016 Jan 14]. <http://www.ncbi.nlm.nih.gov/pubmed/23749882>
- [191] Martín-Padura I, Lostaglio S, Schneemann M, Williams L, Romano M, Fruscella P, et al. Junctional adhesion molecule, a novel member of the immunoglobulin superfamily that distributes at intercellular junctions and modulates monocyte transmigration. *J Cell Biol [Internet].* 1998;142(1):117–27 [cited 2016 Jan 14]. <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2133024&tool=pmcentrez&rendertype=abstract>
- [192] Dejana E, Lampugnani MG, Martinez-Estrada O, Bazzoni G. The molecular organization of endothelial junctions and their functional role in vascular morphogenesis and permeability. *Int J Dev Biol [Internet].* 2000;44(6):743–8 [cited 2016 Jan 14]. <http://www.ncbi.nlm.nih.gov/pubmed/11061439>

- [193] Yeung D, Manias JL, Stewart DJ, Nag S. Decreased junctional adhesion molecule-A expression during blood-brain barrier breakdown. *Acta Neuropathol* [Internet]. 2008;115(6):635–42 [cited 2016 Jan 14]. <http://www.ncbi.nlm.nih.gov/pubmed/18357461>
- [194] Vincent PA, Xiao K, Buckley KM, Kowalczyk AP. VE-cadherin: adhesion at arm's length. *Am J Physiol Cell Physiol* [Internet]. 2004;286(5):C987–97 [cited 2016 Jan 14]. <http://www.ncbi.nlm.nih.gov/pubmed/15075197>
- [195] Lampugnani MG, Dejana E. Adherens junctions in endothelial cells regulate vessel maintenance and angiogenesis. *Thromb Res* [Internet]. 2007;120 Suppl :S1–6 [cited 2016 Jan 14]. <http://www.ncbi.nlm.nih.gov/pubmed/18023702>
- [196] Abbruscato TJ, Davis TP. Protein expression of brain endothelial cell E-cadherin after hypoxia/aglycemia: influence of astrocyte contact. *Brain Res* [Internet]. 1999;842(2):277–86 [cited 2016 Jan 14]. <http://www.ncbi.nlm.nih.gov/pubmed/10526124>
- [197] Romero IA, Radewicz K, Jubin E, Michel CC, Greenwood J, Couraud P-O, et al. Changes in cytoskeletal and tight junctional proteins correlate with decreased permeability induced by dexamethasone in cultured rat brain endothelial cells. *Neurosci Lett* [Internet]. 2003;344(2):112–6 [cited 2015 Dec 8]. <http://www.ncbi.nlm.nih.gov/pubmed/12782340>
- [198] Privratsky JR, Newman PJ. PECAM-1: regulator of endothelial junctional integrity. *Cell Tissue Res* [Internet]. 2014;355(3):607–19 [cited 2016 Jan 14]. <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3975704&tool=pmcentrez&rendertype=abstract>
- [199] Graesser D, Solowiej A, Bruckner M, Osterweil E, Juedes A, Davis S, et al. Altered vascular permeability and early onset of experimental autoimmune encephalomyelitis in PECAM-1-deficient mice. *J Clin Invest* [Internet]. 2002;109(3):383–92 [cited 2016 Jan 14]. <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=150854&tool=pmcentrez&rendertype=abstract>
- [200] Tietz S, Engelhardt B. Brain barriers: crosstalk between complex tight junctions and adherens junctions. *J Cell Biol* [Internet]. 2015;209(4):493–506 [cited 2016 Jan 14]. <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=4442813&tool=pmcentrez&rendertype=abstract>
- [201] Fanning AS, Jameson BJ, Jesaitis LA, Anderson JM. The tight junction protein ZO-1 establishes a link between the transmembrane protein occludin and the actin cytoskeleton. *J Biol Chem* [Internet]. 1998;273(45):29745–53 [cited 2016 Jan 15]. <http://www.ncbi.nlm.nih.gov/pubmed/9792688>
- [202] Van Itallie CM, Anderson JM. Architecture of tight junctions and principles of molecular composition. *Semin Cell Dev Biol* [Internet]. 2014;36:157–65 [cited 2016 Jan 6]. <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=4254347&tool=pmcentrez&rendertype=abstract>
- [203] Kirk J, Plumb J, Mirakhur M, McQuaid S. Tight junctional abnormality in multiple sclerosis white matter affects all calibres of vessel and is associated with blood-brain

- barrier leakage and active demyelination. *J Pathol* [Internet]. 2003;201(2):319–27 [cited 2016 Jan 15]. <http://www.ncbi.nlm.nih.gov/pubmed/14517850>
- [204] Stevenson BR, Siliciano JD, Mooseker MS, Goodenough DA. Identification of ZO-1: a high molecular weight polypeptide associated with the tight junction (zonula occludens) in a variety of epithelia. *J Cell Biol* [Internet]. 1986;103(3):755–66 [cited 2016 Jan 15]. <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2114282&tool=pmcentrez&rendertype=abstract>
- [205] Howarth AG, Hughes MR, Stevenson BR. Detection of the tight junction-associated protein ZO-1 in astrocytes and other nonepithelial cell types. *Am J Physiol* [Internet]. 1992;262(2 Pt 1):C461–9 [cited 2016 Jan 15]. <http://www.ncbi.nlm.nih.gov/pubmed/1539634>
- [206] Itoh M, Morita K, Tsukita S. Characterization of ZO-2 as a MAGUK family member associated with tight as well as adherens junctions with a binding affinity to occludin and alpha catenin. *J Biol Chem* [Internet]. 1999;274(9):5981–6 [cited 2016 Jan 15]. <http://www.ncbi.nlm.nih.gov/pubmed/10026224>
- [207] Toyofuku T, Yabuki M, Otsu K, Kuzuya T, Hori M, Tada M. Direct association of the gap junction protein connexin-43 with ZO-1 in cardiac myocytes. *J Biol Chem* [Internet]. 1998;273(21):12725–31 [cited 2016 Jan 15]. <http://www.ncbi.nlm.nih.gov/pubmed/9582296>
- [208] Abbruscato TJ, Lopez SP, Mark KS, Hawkins BT, Davis TP. Nicotine and cotinine modulate cerebral microvascular permeability and protein expression of ZO-1 through nicotinic acetylcholine receptors expressed on brain endothelial cells. *J Pharm Sci* [Internet]. 2002;91(12):2525–38 [cited 2016 Jan 15]. <http://www.ncbi.nlm.nih.gov/pubmed/12434396>
- [209] Gumbiner B, Lowenkopf T, Apatira D. Identification of a 160-kDa polypeptide that binds to the tight junction protein ZO-1. *Proc Natl Acad Sci USA* [Internet]. 1991;88(8):3460–4 [cited 2016 Jan 15]. <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=51467&tool=pmcentrez&rendertype=abstract>
- [210] Umeda K, Matsui T, Nakayama M, Furuse K, Sasaki H, Furuse M, et al. Establishment and characterization of cultured epithelial cells lacking expression of ZO-1. *J Biol Chem* [Internet]. 2004;279(43):44785–94 [cited 2016 Jan 15]. <http://www.ncbi.nlm.nih.gov/pubmed/15292177>
- [211] Dejana E, Tournier-Lasserre E, Weinstein BM. The control of vascular integrity by endothelial cell junctions: molecular basis and pathological implications. *Dev Cell* [Internet]. 2009;16(2):209–21 [cited 2015 Sep 8]. <http://www.ncbi.nlm.nih.gov/pubmed/19217423>
- [212] Nico B, Frigeri A, Nicchia GP, Corsi P, Ribatti D, Quondamatteo F, et al. Severe alterations of endothelial and glial cells in the blood-brain barrier of dystrophic mdx

- mice. *Glia* [Internet]. 2003;42(3):235–51 [cited 2016 Jan 16]. <http://www.ncbi.nlm.nih.gov/pubmed/12673830>
- [213] Shiu C, Barbier E, Di Cello F, Choi HJ, Stins M. HIV-1 gp120 as well as alcohol affect blood-brain barrier permeability and stress fiber formation: involvement of reactive oxygen species. *Alcohol Clin Exp Res* [Internet]. 2007;31(1):130–7 [cited 2016 Jan 16]. <http://www.ncbi.nlm.nih.gov/pubmed/17207111>
- [214] Davson H. Review lecture. The blood-brain barrier. *J Physiol* [Internet]. 1976;255(1):1–28 [cited 2016 Jan 17]. <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=1309232&tool=pmcentrez&rendertype=abstract>
- [215] Wong AD, Ye M, Levy AF, Rothstein JD, Bergles DE, Searson PC. The blood-brain barrier: an engineering perspective. *Front Neuroeng* [Internet]. 2013;6:7 [cited 2016 Jan 16]. <http://journal.frontiersin.org/article/10.3389/fneng.2013.00007/abstract>
- [216] Abbott NJ, Patabendige AAK, Dolman DEM, Yusof SR, Begley DJ. Structure and function of the blood-brain barrier. *Neurobiol Dis* [Internet]. 2010;37(1):13–25 [cited 2014 Jul 15]. <http://www.ncbi.nlm.nih.gov/pubmed/19664713>
- [217] Taylor CJ, Nicola PA, Wang S, Barrand MA, Hladky SB. Transporters involved in regulation of intracellular pH in primary cultured rat brain endothelial cells. *J Physiol* [Internet]. 2006;576(Pt 3):769–85 [cited 2016 Jan 17]. <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=1890423&tool=pmcentrez&rendertype=abstract>
- [218] Giacomini KM, Huang S-M, Tweedie DJ, Benet LZ, Brouwer KLR, Chu X, et al. Membrane transporters in drug development. *Nat Rev Drug Discov* [Internet]. 2010;9(3):215–36 [cited 2014 Jul 11]. <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3326076&tool=pmcentrez&rendertype=abstract>
- [219] Ueno M. Mechanisms of the penetration of blood-borne substances into the brain. *Curr Neuropharmacol* [Internet]. 2009;7(2):142–9 [cited 2016 Jan 16]. <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2730006&tool=pmcentrez&rendertype=abstract>
- [220] Enerson BE, Drewes LR. The rat blood-brain barrier transcriptome. *J Cereb Blood Flow Metab*. 2006;26(7):959–73.
- [221] Abbott NJ, Rönnbäck L, Hansson E. Astrocyte-endothelial interactions at the blood-brain barrier. *Nat Rev Neurosci* [Internet]. 2006;7(1):41–53 [cited 2014 Jul 9]. <http://www.ncbi.nlm.nih.gov/pubmed/16371949>
- [222] Dauchy S, Dutheil F, Weaver RJ, Chassoux F, Daumas-Duport C, Couraud P-O, et al. ABC transporters, cytochromes P450 and their main transcription factors: expression at the human blood-brain barrier. *J Neurochem* [Internet]. 2008;107(6):1518–28 [cited 2016 Jan 17]. <http://www.ncbi.nlm.nih.gov/pubmed/19094056>

- [223] Begley DJ. ABC transporters and the blood-brain barrier. *Curr Pharm Des* [Internet]. 2004;10(12):1295–312 [cited 2016 Jan 17]. <http://www.ncbi.nlm.nih.gov/pubmed/15134482>
- [224] Dallas S, Miller DS, Bendayan R. Multidrug resistance-associated proteins: expression and function in the central nervous system. *Pharmacol Rev* [Internet]. 2006;58(2):140–61 [cited 2016 Jan 17]. <http://www.ncbi.nlm.nih.gov/pubmed/16714484>
- [225] Bendayan R, Ronaldson PT, Gingras D, Bendayan M. In situ localization of P-glycoprotein (ABCB1) in human and rat brain. *J Histochem Cytochem* [Internet]. 2006;54(10):1159–67 [cited 2016 Jan 12]. <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3957801&tool=pmcentrez&rendertype=abstract>
- [226] Banks WA. The CNS as a target for peptides and peptide-based drugs. *Expert Opin Drug Deliv* [Internet]. 2006;3(6):707–12 [cited 2016 Jan 17]. <http://www.ncbi.nlm.nih.gov/pubmed/17076593>
- [227] Lajoie P, Nabi IR. Regulation of raft-dependent endocytosis. *J Cell Mol Med* [Internet]. 2016;11(4):644–53 [cited 2016 Jan 17]. <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3823247&tool=pmcentrez&rendertype=abstract>
- [228] Parton RG, Richards AA. Lipid rafts and caveolae as portals for endocytosis: new insights and common mechanisms. *Traffic* [Internet]. 2003;4(11):724–38 [cited 2016 Jan 17]. <http://www.ncbi.nlm.nih.gov/pubmed/14617356>
- [229] Wolburg H. Blood-Brain Barriers. In: Dermietzel R, Spray DC, Nedergaard M, editors. Weinheim, Germany: Wiley-VCH Verlag GmbH & Co. KGaA; 2006 [cited 2016 Jan 17]. doi: 10.1002/9783527611225
- [230] Song L, Ge S, Pachter JS. Caveolin-1 regulates expression of junction-associated proteins in brain microvascular endothelial cells. *Blood* [Internet]. 2007;109(4):1515–23 [cited 2016 Jan 17]. <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=1794065&tool=pmcentrez&rendertype=abstract>
- [231] Nahirney PC, Reeson P, Brown CE. Ultrastructural analysis of blood-brain barrier breakdown in the peri-infarct zone in young adult and aged mice. *J Cereb Blood Flow Metab* [Internet]. 2015 [cited 2016 Jan 1]. <http://www.ncbi.nlm.nih.gov/pubmed/26661190>
- [232] Ueno M, Chiba Y, Matsumoto K, Nakagawa T, Miyanaka H. Clearance of beta-amyloid in the brain. *Curr Med Chem* [Internet]. 2014;21(35):4085–90 [cited 2016 Jan 17]. <http://www.ncbi.nlm.nih.gov/pubmed/25312211>
- [233] Citron M, Oltersdorf T, Haass C, McConlogue L, Hung AY, Seubert P, et al. Mutation of the beta-amyloid precursor protein in familial Alzheimer's disease increases beta-protein production. *Nature* [Internet]. 1992;360(6405):672–4 [cited 2015 Dec 29]. <http://www.ncbi.nlm.nih.gov/pubmed/1465129>

- [234] Tanzi RE, Moir RD, Wagner SL. Clearance of Alzheimer's Aβ peptide: the many roads to perdition. *Neuron* [Internet]. 2004;43(5):605–8 [cited 2016 Jan 17]. <http://www.ncbi.nlm.nih.gov/pubmed/15339642>
- [235] Zlokovic B V, Yamada S, Holtzman D, Ghiso J, Frangione B. Clearance of amyloid beta-peptide from brain: transport or metabolism? *Nat Med* [Internet]. 2000;6(7):718–9 [cited 2016 Jan 17]. <http://www.ncbi.nlm.nih.gov/pubmed/10888892>
- [236] Hardy J. Amyloid double trouble. *Nat Genet* [Internet]. 2006;38(1):11–2 [cited 2016 Jan 17]. <http://www.ncbi.nlm.nih.gov/pubmed/16380721>
- [237] Ballatore C, Lee VM-Y, Trojanowski JQ. Tau-mediated neurodegeneration in Alzheimer's disease and related disorders. *Nat Rev Neurosci* [Internet]. 2007;8(9):663–72 [cited 2014 Jul 21]. doi: 10.1038/nrn2194
- [238] Selkoe DJ. Clearing the brain's amyloid cobwebs. *Neuron* [Internet]. 2001;32(2):177–80 [cited 2016 Jan 17]. <http://www.ncbi.nlm.nih.gov/pubmed/11683988>
- [239] Shibata M, Yamada S, Kumar SR, Calero M, Bading J, Frangione B, et al. Clearance of Alzheimer's amyloid-ss(1-40) peptide from brain by LDL receptor-related protein-1 at the blood-brain barrier. *J Clin Invest* [Internet]. 2000;106(12):1489–99 [cited 2015 Dec 14]. <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=387254&tool=pmcentrez&rendertype=abstract>
- [240] Deane R, Wu Z, Sagare A, Davis J, Du Yan S, Hamm K, et al. LRP/amyloid beta-peptide interaction mediates differential brain efflux of Aβ isoforms. *Neuron* [Internet]. 2004;43(3):333–44 [cited 2015 Nov 26]. <http://www.ncbi.nlm.nih.gov/pubmed/15294142>
- [241] Deane R, Du Yan S, Subramanyam RK, LaRue B, Jovanovic S, Hogg E, et al. RAGE mediates amyloid-beta peptide transport across the blood-brain barrier and accumulation in brain. *Nat Med* [Internet]. 2003;9(7):907–13 [cited 2015 Sep 21]. <http://www.ncbi.nlm.nih.gov/pubmed/12808450>
- [242] Pascale CL, Miller MC, Chiu C, Boylan M, Caralopoulos IN, Gonzalez L, et al. Amyloid-beta transporter expression at the blood-CSF barrier is age-dependent. *Fluids Barriers CNS* [Internet]. 2011;8:21 [cited 2015 Dec 5]. <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3162580&tool=pmcentrez&rendertype=abstract>
- [243] Deane R, Bell RD, Sagare A, Zlokovic B V. Clearance of amyloid-beta peptide across the blood-brain barrier: implication for therapies in Alzheimer's disease. *CNS Neurol Disord Drug Targets* [Internet]. 2009;8(1):16–30 [cited 2015 Oct 5]. <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2872930&tool=pmcentrez&rendertype=abstract>
- [244] Bell RD, Sagare AP, Friedman AE, Bedi GS, Holtzman DM, Deane R, et al. Transport pathways for clearance of human Alzheimer's amyloid β-peptide and apolipoproteins E and J in the mouse central nervous system. *J Cereb Blood Flow Metab* [Internet].

- 2007;27(5):909–18 [cited 2015 Oct 22]. <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2853021&tool=pmcentrez&rendertype=abstract>
- [245] Deane R, Sagare A, Hamm K, Parisi M, Lane S, Finn MB, et al. apoE isoform-specific disruption of amyloid β peptide clearance from mouse brain. *J Clin Invest* [Internet]. 2008;118(12):4002–13 [cited 2015 Oct 17]. <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2582453&tool=pmcentrez&rendertype=abstract>
- [246] Herz J, Strickland DK. LRP: a multifunctional scavenger and signaling receptor. *J Clin Invest* [Internet]. 2001;108(6):779–84 [cited 2015 Nov 12]. <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=200939&tool=pmcentrez&rendertype=abstract>
- [247] Nazer B, Hong S, Selkoe DJ. LRP promotes endocytosis and degradation, but not transcytosis, of the amyloid-beta peptide in a blood-brain barrier in vitro model. *Neurobiol Dis* [Internet]. 2008;30(1):94–102 [cited 2016 Jan 17]. <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2376120&tool=pmcentrez&rendertype=abstract>
- [248] Pflanzner T, Janko MC, André-Dohmen B, Reuss S, Weggen S, Roebroek AJM, et al. LRP1 mediates bidirectional transcytosis of amyloid- β across the blood-brain barrier. *Neurobiol Aging* [Internet]. 2011;32(12):2323.e1–11 [cited 2016 Jan 17]. <http://www.ncbi.nlm.nih.gov/pubmed/20630619>
- [249] von Arnim CAF, Kinoshita A, Peltan ID, Tangredi MM, Herl L, Lee BM, et al. The low density lipoprotein receptor-related protein (LRP) is a novel beta-secretase (BACE1) substrate. *J Biol Chem* [Internet]. 2005;280(18):17777–85 [cited 2015 Dec 15]. <http://www.ncbi.nlm.nih.gov/pubmed/15749709>
- [250] Koistinaho M, Lin S, Wu X, Esterman M, Koger D, Hanson J, et al. Apolipoprotein E promotes astrocyte colocalization and degradation of deposited amyloid-beta peptides. *Nat Med* [Internet]. 2004;10(7):719–26 [cited 2016 Jan 18]. <http://www.ncbi.nlm.nih.gov/pubmed/15195085>
- [251] Yan S Du, Chen X, Fu J, Chen M, Zhu H, Roher A, et al. RAGE and amyloid- β peptide neurotoxicity in Alzheimer's disease. *Nature* [Internet]. 1996;382(6593):685–91 [cited 2016 Jan 14]. <http://www.ncbi.nlm.nih.gov/pubmed/8751438>
- [252] Zlokovic B V, Ghiso J, Mackic JB, McComb JG, Weiss MH, Frangione B. Blood-brain barrier transport of circulating Alzheimer's amyloid beta. *Biochem Biophys Res Commun* [Internet]. 1993;197(3):1034–40 [cited 2016 Jan 18]. <http://www.ncbi.nlm.nih.gov/pubmed/8280117>
- [253] Correale J, Villa A. The blood-brain-barrier in multiple sclerosis: functional roles and therapeutic targeting. *Autoimmunity* [Internet]. 2007;40(2):148–60 [cited 2016 Jan 13]. <http://www.ncbi.nlm.nih.gov/pubmed/17453713>

- [254] Kaur C, Ling EA. Blood brain barrier in hypoxic-ischemic conditions. *Curr Neurovasc Res* [Internet]. 2008;5(1):71–81 [cited 2016 Jan 18]. <http://www.ncbi.nlm.nih.gov/pubmed/18289024>
- [255] Desai BS, Monahan AJ, Carvey PM, Hendey B. Blood-brain barrier pathology in Alzheimer's and Parkinson's disease: implications for drug therapy. *Cell Transplant* [Internet]. 2007;16(3):285–99 [cited 2016 Jan 18]. <http://www.ncbi.nlm.nih.gov/pubmed/17503739>
- [256] Förster C. Tight junctions and the modulation of barrier function in disease. *Histochem Cell Biol* [Internet]. 2008;130(1):55–70 [cited 2015 Nov 30]. <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2413111&tool=pmcentrez&rendertype=abstract>
- [257] Wardlaw JM, Sandercock PAG, Dennis MS, Starr J. Is breakdown of the blood-brain barrier responsible for lacunar stroke, leukoaraiosis, and dementia? *Stroke* [Internet]. 2003;34(3):806–12 [cited 2015 Dec 5]. <http://www.ncbi.nlm.nih.gov/pubmed/12624314>
- [258] Zlokovic B V. Clearing amyloid through the blood-brain barrier. *J Neurochem* [Internet]. 2004;89(4):807–11 [cited 2016 Jan 14]. <http://www.ncbi.nlm.nih.gov/pubmed/15140180>
- [259] Owen JB, Sultana R, Aluise CD, Erickson MA, Price TO, Bu G, et al. Oxidative modification to LDL receptor-related protein 1 in hippocampus from subjects with Alzheimer disease: implications for A β accumulation in AD brain. *Free Radic Biol Med* [Internet]. 2010;49(11):1798–803 [cited 2016 Jan 18]. <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2970765&tool=pmcentrez&rendertype=abstract>
- [260] Cirrito JR, Deane R, Fagan AM, Spinner ML, Parsadanian M, Finn MB, et al. P-glycoprotein deficiency at the blood-brain barrier increases amyloid-beta deposition in an Alzheimer disease mouse model. *J Clin Invest* [Internet]. 2005;115(11):3285–90 [cited 2016 Jan 18]. <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=1257538&tool=pmcentrez&rendertype=abstract>
- [261] Landau SM, Harvey D, Madison CM, Koeppe RA, Reiman EM, Foster NL, et al. Associations between cognitive, functional, and FDG-PET measures of decline in AD and MCI. *Neurobiol Aging* [Internet]. 2011;32(7):1207–18 [cited 2016 Jan 18]. <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2891865&tool=pmcentrez&rendertype=abstract>
- [262] Harik SI. Changes in the glucose transporter of brain capillaries. *Can J Physiol Pharmacol* [Internet]. 1992;70 Suppl:S113–7 [cited 2016 Jan 18]. <http://www.ncbi.nlm.nih.gov/pubmed/1295661>
- [263] Mooradian AD, Chung HC, Shah GN. GLUT-1 expression in the cerebra of patients with Alzheimer's disease. *Neurobiol Aging* [Internet]. 2016;18(5):469–74 [cited 2016 Jan 18]. <http://www.ncbi.nlm.nih.gov/pubmed/9390772>
- [264] Sagare AP, Bell RD, Zlokovic BV. Neurovascular dysfunction and faulty amyloid β -peptide clearance in Alzheimer disease. *Cold Spring Harb Perspect Med*. 2012 Oct

- 1;2(10). pii: a011452. doi: 10.1101/cshperspect.a011452. Review. PubMed PMID: 23028132; PubMed Central PMCID: PMC3475405.
- [265] Claudio L. Ultrastructural features of the blood-brain barrier in biopsy tissue from Alzheimer's disease patients. *Acta Neuropathol* [Internet]. 1996;91(1):6–14 [cited 2016 Feb 5]. <http://www.ncbi.nlm.nih.gov/pubmed/8773140>
- [266] Bailey TL, Rivara CB, Rocher AB, Hof PR. The nature and effects of cortical microvascular pathology in aging and Alzheimer's disease. *Neurol Res* [Internet]. 2004;26(5):573–8 [cited 2016 Jan 8]. <http://www.ncbi.nlm.nih.gov/pubmed/15265277>
- [267] Romanitan MO, Popescu BO, Winblad B, Bajenaru OA, Bogdanovic N. Occludin is overexpressed in Alzheimer's disease and vascular dementia. *J Cell Mol Med* [Internet]. 2016;11(3):569–79 [cited 2016 Jan 18]. <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3922362&tool=pmcentrez&rendertype=abstract>
- [268] Wu Z, Guo H, Chow N, Sallstrom J, Bell RD, Deane R, et al. Role of the MEOX2 homeobox gene in neurovascular dysfunction in Alzheimer disease. *Nat Med* [Internet]. 2005;11(9):959–65 [cited 2016 Jan 18]. doi: 10.1038/nm1287
- [269] Rogers J, Lubner-Narod J, Styren SD, Civin WH. Expression of immune system-associated antigens by cells of the human central nervous system: relationship to the pathology of Alzheimer's disease. *Neurobiol Aging*. Jan 1988; ;9(4):339–49.
- [270] Mehta PD, Pirttilä T, Mehta SP, Sersen EA, Aisen PS, Wisniewski HM. Plasma and cerebrospinal fluid levels of amyloid beta proteins 1-40 and 1-42 in Alzheimer disease. *Arch Neurol* [Internet]. 2000;57(1):100–5 [cited 2016 Jan 18]. <http://www.ncbi.nlm.nih.gov/pubmed/10634455>
- [271] Pesaresi M, Lovati C, Bertora P, Mailland E, Galimberti D, Scarpini E, et al. Plasma levels of beta-amyloid (1-42) in Alzheimer's disease and mild cognitive impairment. *Neurobiol Aging* [Internet]. 2006;27(6):904–5 [cited 2016 Jan 18]. <http://www.ncbi.nlm.nih.gov/pubmed/16638622>
- [272] Hansson O, Zetterberg H, Vanmechelen E, Vanderstichele H, Andreasson U, Londos E, et al. Evaluation of plasma A β (40) and A β (42) as predictors of conversion to Alzheimer's disease in patients with mild cognitive impairment. *Neurobiol Aging* [Internet]. 2010;31(3):357–67 [cited 2016 Jan 18]. <http://www.ncbi.nlm.nih.gov/pubmed/18486992>
- [273] Roher AE, Esh CL, Kokjohn TA, Castaño EM, Van Vickle GD, Kalback WM, et al. Amyloid beta peptides in human plasma and tissues and their significance for Alzheimer's disease. *Alzheimers Dement* [Internet]. 2009;5(1):18–29 [cited 2016 Jan 18]. <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2663406&tool=pmcentrez&rendertype=abstract>

- [274] Li QX, Whyte S, Tanner JE, Evin G, Beyreuther K, Masters CL. Secretion of Alzheimer's disease Abeta amyloid peptide by activated human platelets. *Lab Invest* [Internet]. 1998;78(4):461–9 [cited 2016 Jan 18]. <http://www.ncbi.nlm.nih.gov/pubmed/9564890>
- [275] Kuo YM, Kokjohn TA, Watson MD, Woods AS, Cotter RJ, Sue LI, et al. Elevated abeta42 in skeletal muscle of Alzheimer disease patients suggests peripheral alterations of AbetaPP metabolism. *Am J Pathol* [Internet]. 2000;156(3):797–805 [cited 2016 Jan 18]. <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=1876838&tool=pmcentrez&rendertype=abstract>
- [276] Kuo YM, Kokjohn TA, Kalback W, Luehrs D, Galasko DR, Chevallier N, et al. Amyloid-beta peptides interact with plasma proteins and erythrocytes: implications for their quantitation in plasma. *Biochem Biophys Res Commun* [Internet]. 2000;268(3):750–6 [cited 2016 Jan 18]. <http://www.ncbi.nlm.nih.gov/pubmed/10679277>
- [277] Biere AL, Ostaszewski B, Stimson ER, Hyman BT, Maggio JE, Selkoe DJ. Amyloid beta-peptide is transported on lipoproteins and albumin in human plasma. *J Biol Chem* [Internet]. 1996;271(51):32916–22 [cited 2016 Jan 18]. <http://www.ncbi.nlm.nih.gov/pubmed/8955133>
- [278] Strittmatter WJ, Weisgraber KH, Huang DY, Dong LM, Salvesen GS, Pericak-Vance M, et al. Binding of human apolipoprotein E to synthetic amyloid beta peptide: isoform-specific effects and implications for late-onset Alzheimer disease. *Proc Natl Acad Sci U S A* [Internet]. 1993;90(17):8098–102 [cited 2016 Jan 18]. <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=47295&tool=pmcentrez&rendertype=abstract>
- [279] Sagare A, Deane R, Bell RD, Johnson B, Hamm K, Pendu R, et al. Clearance of amyloid-beta by circulating lipoprotein receptors. *Nat Med* [Internet]. 2007;13(9):1029–31 [cited 2015 Oct 1]. <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2936449&tool=pmcentrez&rendertype=abstract>
- [280] Pan W, Solomon B, Maness LM, Kastin AJ. Antibodies to beta-amyloid decrease the blood-to-brain transfer of beta-amyloid peptide. *Exp Biol Med (Maywood)* [Internet]. 2002;227(8):609–15 [cited 2016 Jan 18]. <http://www.ncbi.nlm.nih.gov/pubmed/12192102>
- [281] Erickson MA, Banks WA. Blood-brain barrier dysfunction as a cause and consequence of Alzheimer's disease. *J Cereb Blood Flow Metab* [Internet]. 2013;33(10):1500–13 [cited 2016 Jan 16]. <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3790938&tool=pmcentrez&rendertype=abstract>
- [282] Geroldi D, Falcone C, Emanuele E. Soluble receptor for advanced glycation end products: from disease marker to potential therapeutic target. *Curr Med Chem* [Internet]. 2006;13(17):1971–8 [cited 2016 Jan 18]. <http://www.ncbi.nlm.nih.gov/pubmed/16842191>
- [283] Marchi N, Fazio V, Cucullo L, Kight K, Masaryk T, Barnett G, et al. Serum transthyretin monomer as a possible marker of blood-to-CSF barrier disruption. *J Neurosci* [Internet]. 2003;23(5):1949–55 [cited 2016 Jan 18]. <http://www.ncbi.nlm.nih.gov/pubmed/12629200>

- [284] Kapural M, Krizanac-Bengez L, Barnett G, Perl J, Masaryk T, Apollo D, et al. Serum S-100beta as a possible marker of blood-brain barrier disruption. *Brain Res* [Internet]. 2002;940(1-2):102-4 [cited 2016 Jan 19]. <http://www.ncbi.nlm.nih.gov/pubmed/12020881>
- [285] Tamaki C, Ohtsuki S, Iwatsubo T, Hashimoto T, Yamada K, Yabuki C, et al. Major involvement of low-density lipoprotein receptor-related protein 1 in the clearance of plasma free amyloid beta-peptide by the liver. *Pharm Res* [Internet]. 2006;23(7):1407-16 [cited 2015 Dec 15]. <http://www.ncbi.nlm.nih.gov/pubmed/16779710>
- [286] Sagare AP, Deane R, Zlokovic B V. Low-density lipoprotein receptor-related protein 1: a physiological A β homeostatic mechanism with multiple therapeutic opportunities. *Pharmacol Ther* [Internet]. 2012;136(1):94-105. doi: 10.1016/j.pharmthera.2012.07.008
- [287] Kandimalla KK, Curran GL, Holasek SS, Gilles EJ, Wengenack TM, Poduslo JF. Pharmacokinetic analysis of the blood-brain barrier transport of 125I-amyloid beta protein 40 in wild-type and Alzheimer's disease transgenic mice (APP,PS1) and its implications for amyloid plaque formation. *J Pharmacol Exp Ther*. 2005;313(3):1370-8.
- [288] Ghiso J, Shayo M, Calero M, Ng D, Tomidokoro Y, Gandy S, et al. Systemic catabolism of Alzheimer's A β 40 and A β 42. *J Biol Chem* [Internet]. 2004;279(44):45897-908 [cited 2016 Jan 19]. <http://www.ncbi.nlm.nih.gov/pubmed/15322125>
- [289] Tamaki C, Ohtsuki S, Terasaki T. Insulin facilitates the hepatic clearance of plasma amyloid beta-peptide (1-40) by intracellular translocation of low-density lipoprotein receptor-related protein 1 (LRP-1) to the plasma membrane in hepatocytes. *Mol Pharmacol* [Internet]. 2007;72(4):850-5 [cited 2016 Jan 19]. <http://www.ncbi.nlm.nih.gov/pubmed/17609417>
- [290] Alemi M, Gaiteiro C, Ribeiro CA, Santos LM, Gomes JR, Oliveira SM, et al. Transthyretin participates in beta-amyloid transport from the brain to the liver- involvement of the low-density lipoprotein receptor-related protein 1? *Sci Rep* [Internet]. 2016;6:20164 [cited 2016 Feb 4]. <http://www.ncbi.nlm.nih.gov/pubmed/26837706>
- [291] Duckworth WC, Bennett RG, Hamel FG. Insulin degradation: progress and potential. *Endocr Rev* [Internet]. 1998;19(5):608-24 [cited 2016 Jan 19]. <http://www.ncbi.nlm.nih.gov/pubmed/9793760>
- [292] Wang D-S, Dickson DW, Malter JS. beta-Amyloid degradation and Alzheimer's disease. *J Biomed Biotechnol* [Internet]. 2006;2006(3):58406 [cited 2016 Jan 19]. <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=1559921&tool=pmcentrez&rendertype=abstract>
- [293] Ghersi-Egea JF, Gorevic PD, Ghiso J, Frangione B, Patlak CS, Fenstermacher JD. Fate of cerebrospinal fluid-borne amyloid beta-peptide: rapid clearance into blood and appreciable accumulation by cerebral arteries. *J Neurochem* [Internet]. 1996;67(2):880-3 [cited 2016 Jan 19]. <http://www.ncbi.nlm.nih.gov/pubmed/8764620>

- [294] Soprano DR, Herbert J, Soprano KJ, Schon EA, Goodman DS. Demonstration of transthyretin mRNA in the brain and other extrahepatic tissues in the rat. *J Biol Chem* [Internet]. 1985;260(21):11793–8 [cited 2015 Dec 30]. <http://www.ncbi.nlm.nih.gov/pubmed/4044580>
- [295] Woeber KA, Ingbar SH. The contribution of thyroxine-binding prealbumin to the binding of thyroxine in human serum, as assessed by immunoabsorption. *J Clin Invest*. 1968;47(7):1710–21.
- [296] Goodman D. Retinoids and retinol-binding proteins. *Harvey Lect*. 1987;Series 81:111–32.
- [297] Kabat EA, Moore DH, Landow H. An electrophoretic study of the protein components in cerebrospinal fluid and their relationship to the serum proteins. *J Clin Invest* [Internet]. 1942;21(5):571–7 [cited 2015 Dec 30]. <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=435175&tool=pmcentrez&rendertype=abstract>
- [298] Seibert FB NJ. Electrophoretic study of the blood response in tuberculosis. *J Biol Chem*. 1942;143:29–38.
- [299] Wallace MR, Naylor SL, Kluge-Beckerman B, Long GL, McDonald L, Shows TB, et al. Localization of the human prealbumin gene to chromosome 18. *Biochem Biophys Res Commun* [Internet]. 1985;129(3):753–8. <http://www.ncbi.nlm.nih.gov/pubmed/2990465>
- [300] Sasaki H, Yoshioka N, Takagi Y, Sakaki Y. Structure of the chromosomal gene for human serum prealbumin. *Gene* [Internet]. 1985;37(1–3):191–7 [cited 2015 Dec 30]. <http://www.ncbi.nlm.nih.gov/pubmed/4054629>
- [301] Tsuzuki T, Mita S, Maeda S, Araki S, Shimada K. Structure of the human prealbumin gene. *J Biol Chem*. 1985;260(22):12224–7.
- [302] Whitehead AS, Skinner M, Bruns GA, Costello W, Edge MD, Cohen AS, et al. Cloning of human prealbumin complementary DNA. Localization of the gene to chromosome 18 and detection of a variant prealbumin allele in a family with familial amyloid polyneuropathy. *Mol Biol Med* [Internet]. 1984;2(6):411–23 [cited 2015 Dec 30]. <http://www.ncbi.nlm.nih.gov/pubmed/6100724>
- [303] Mita S, Maeda S, Shimada K, Araki S. Cloning and sequence analysis of cDNA for human prealbumin. *Biochem Biophys Res Commun* [Internet]. 1984;124(2):558–64. <http://www.sciencedirect.com/science/article/pii/0006291X84915900>
- [304] Sundelin J, Melhus H, Das S, Eriksson U, Lind P, Trägårdh L, et al. The primary structure of rabbit and rat prealbumin and a comparison with the tertiary structure of human prealbumin [Internet]. *J Biol Chem*. 1985;260:6481–7. http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=3922975 \n<http://www.jbc.org/cgi/reprint/260/10/6481>

- [305] Dickson PW, Howlett GJ, Schreiber G. Rat transthyretin (prealbumin). Molecular cloning, nucleotide sequence, and gene expression in liver and brain. *J Biol Chem* [Internet]. 1985;260(13):8214–9. <http://www.ncbi.nlm.nih.gov/pubmed/3839240>
- [306] Costa RH, Lai E, Darnell JE. Transcriptional control of the mouse prealbumin (transthyretin) gene: both promoter sequences and a distinct enhancer are cell specific. *Mol Cell Biol*. 1986;6(12):4697–708.
- [307] Andreoli M, Robbins J. Serum proteins and thyroxine-protein interaction in early human fetuses. *J Clin Invest* [Internet]. 1962;41:1070–7 [cited 2015 Dec 30]. <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=291012&tool=pmcentrez&rendertype=abstract>
- [308] Stabilini R, Vergani C, Agostoni A, Agostoni RP. Influence of age and sex on prealbumin levels. *Clin Chim Acta* [Internet]. 1968;20(2):358–9 [cited 2015 Dec 30]. <http://www.ncbi.nlm.nih.gov/pubmed/4968450>
- [309] Vahlquist A, Rask L, Peterson PA, Berg T. The concentrations of retinol-binding protein, prealbumin, and transferrin in the sera of newly delivered mothers and children of various ages. *Scand J Clin Lab Invest* [Internet]. 1975;35(6):569–75 [cited 2015 Dec 30]. <http://www.ncbi.nlm.nih.gov/pubmed/1239075>
- [310] Weisner B, Roethig HJ. The concentration of prealbumin in cerebrospinal fluid (CSF), indicator of CSF circulation disorders. *Eur Neurol* [Internet]. 1983;22(2):96–105 [cited 2016 Jan 12]. <http://www.ncbi.nlm.nih.gov/pubmed/6840150>
- [311] Gonçalves I, Alves CH, Quintela T, Baltazar G, Socorro S, Saraiva MJ, et al. Transthyretin is up-regulated by sex hormones in mice liver. *Mol Cell Biochem* [Internet]. 2008;317(1–2):137–42 [cited 2016 Feb 2]. <http://www.ncbi.nlm.nih.gov/pubmed/18568387>
- [312] Quintela T, Gonçalves I, Martinho A, Alves CH, Saraiva MJ, Rocha P, et al. Progesterone enhances transthyretin expression in the rat choroid plexus in vitro and in vivo via progesterone receptor. *J Mol Neurosci*. 2011;44(3):152–8.
- [313] Martinho A, Gonçalves I, Costa M, Santos CR. Stress and glucocorticoids increase transthyretin expression in rat choroid plexus via mineralocorticoid and glucocorticoid receptors. *J Mol Neurosci*. 2012;48(1):1–13.
- [314] Kanda Y, Goodman DS, Canfield RE, Morgan FJ. The amino acid sequence of human plasma prealbumin. *J Biol Chem* [Internet]. 1974;249(21):6796–805 [cited 2016 Jan 12]. <http://www.ncbi.nlm.nih.gov/pubmed/4607556>
- [315] Blake CC, Geisow MJ, Oatley SJ, Rérat B, Rérat C. Structure of prealbumin: secondary, tertiary and quaternary interactions determined by Fourier refinement at 1.8 Å. *J Mol Biol* [Internet]. 1978;121(3):339–56 [cited 2015 Dec 31]. <http://www.ncbi.nlm.nih.gov/pubmed/671542>
- [316] Loun B, Hage DS. Characterization of thyroxine-albumin binding using high-performance affinity chromatography. I. Interactions at the warfarin and indole sites of

- albumin. *J Chromatogr* [Internet]. 1992;579(2):225–35 [cited 2016 Jan 12]. <http://www.ncbi.nlm.nih.gov/pubmed/1429970>
- [317] Ferguson RN, Edelhoach H, Saroff HA, Robbins J, Cahnmann HJ. Negative cooperativity in the binding of thyroxine to human serum prealbumin. Preparation of tritium-labeled 8-anilino-1-naphthalenesulfonic acid. *Biochemistry*. 1975;14(2):282–9.
- [318] Richardson SJ, Wijayagunaratne RC, D'Souza DG, Darras VM, Van Herck SLJ. Transport of thyroid hormones via the choroid plexus into the brain: The roles of transthyretin and thyroid hormone transmembrane transporters. Vol. 9, *Frontiers in Neuroscience*. 2015.
- [319] Manral P, Reixach N. Amyloidogenic and non-amyloidogenic transthyretin variants interact differently with human cardiomyocytes: insights into early events of non-fibrillar tissue damage. *Biosci Rep*. 2015 Jan 14;35(1). pii: e00172. doi: 10.1042/BSR20140155. PubMed PMID: 25395306; PubMed Central PMCID: PMC4293901.
- [320] Palha JA, Hays MT, Morreale de Escobar G, Episkopou V, Gottesman ME, Saraiva MJ. Transthyretin is not essential for thyroxine to reach the brain and other tissues in transthyretin-null mice. *Am J Physiol* [Internet]. 1997;272(3 Pt 1):E485–93 [cited 2016 Jan 13]. <http://www.ncbi.nlm.nih.gov/pubmed/9124556>
- [321] Palha JA, Episkopou V, Maeda S, Shimada K, Gottesman ME, Saraiva MJ. Thyroid hormone metabolism in a transthyretin-null mouse strain. *J Biol Chem* [Internet]. 1994;269(52):33135–9 [cited 2016 Jan 13]. <http://www.ncbi.nlm.nih.gov/pubmed/7806543>
- [322] Palha JA, Fernandes R, de Escobar GM, Episkopou V, Gottesman M, Saraiva MJ. Transthyretin regulates thyroid hormone levels in the choroid plexus, but not in the brain parenchyma: study in a transthyretin-null mouse model. *Endocrinology* [Internet]. 2000;141(9):3267–72 [cited 2016 Jan 13]. <http://www.ncbi.nlm.nih.gov/pubmed/10965897>
- [323] Rask L, Anundi H, Peterson PA. The primary structure of the human retinol-binding protein. *FEBS Lett* [Internet]. 1979;104(1):55–8 [cited 2016 Jan 13]. <http://www.ncbi.nlm.nih.gov/pubmed/573217>
- [324] Newcomer ME, Jones TA, Aqvist J, Sundelin J, Eriksson U, Rask L, et al. The three-dimensional structure of retinol-binding protein. *EMBO J* [Internet]. 1984;3(7):1451–4 [cited 2016 Jan 13]. <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=557543&tool=pmcentrez&rendertype=abstract>
- [325] Blaner WS. Retinol-binding protein: the serum transport protein for vitamin A. *Endocr Rev* [Internet]. 1989;10(3):308–16 [cited 2015 Dec 12]. <http://www.ncbi.nlm.nih.gov/pubmed/2550213>
- [326] Kopelman M, Cogan U, Mokady S, Shinitzky M. The interaction between retinol-binding proteins and prealbumins studied by fluorescence polarization. *Biochim*

- Biophys Acta [Internet]. 1976;439(2):449–60 [cited 2016 Jan 13]. <http://www.ncbi.nlm.nih.gov/pubmed/986177>
- [327] van Jaarsveld PP, Edelhoek H, Goodman DS, Robbins J. The interaction of human plasma retinol-binding protein and prealbumin. *J Biol Chem* [Internet]. 1973;248(13):4698–705 [cited 2016 Jan 13]. <http://www.ncbi.nlm.nih.gov/pubmed/4718739>
- [328] Vahlquist A, Peterson PA, Wibell L. Metabolism of the vitamin A transporting protein complex. I. Turnover studies in normal persons and in patients with chronic renal failure. *Eur J Clin Invest* [Internet]. 1973;3(4):352–62 [cited 2016 Jan 13]. <http://www.ncbi.nlm.nih.gov/pubmed/4760057>
- [329] Socolow EL, Woeber KA, Purdy RH, Holloway MT, Ingbar SH. Preparation of I-131-labeled human serum prealbumin and its metabolism in normal and sick patients. *J Clin Invest* [Internet]. 1965;44(10):1600–9 [cited 2016 Jan 13]. <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=292644&tool=pmcentrez&rendertype=abstract>
- [330] Vahlquist A, Peterson PA. Comparative studies on the vitamin A transporting protein complex in human and cynomolgus plasma. *Biochemistry* [Internet]. 1972;11(24):4526–32 [cited 2016 Jan 13]. <http://www.ncbi.nlm.nih.gov/pubmed/4631638>
- [331] Peterson PA, Nilsson SF, Ostberg L, Rask L, Vahlquist A. Aspects of the metabolism of retinol-binding protein and retinol. *Vitam Horm* [Internet]. 1974;32:181–214 [cited 2016 Jan 13]. <http://www.ncbi.nlm.nih.gov/pubmed/4617399>
- [332] Makover A, Moriwaki H, Ramakrishnan R, Saraiva MJ, Blaner WS, Goodman DS. Plasma transthyretin. Tissue sites of degradation and turnover in the rat. *J Biol Chem*. 1988;263:8598–603.
- [333] Receptor-mediated uptake and internalization of transthyretin. PubMed—NCBI [Internet]. 2016 [cited 2016 Jan 14]. <http://www.ncbi.nlm.nih.gov/pubmed/2153133>
- [334] Vieira A V, Sanders EJ, Schneider WJ. Transport of serum transthyretin into chicken oocytes. A receptor-mediated mechanism. *J Biol Chem* [Internet]. 1995;270(7):2952–6 [cited 2016 Jan 14]. <http://www.ncbi.nlm.nih.gov/pubmed/7852374>
- [335] Sousa MM, Norden AG, Jacobsen C, Willnow TE, Christensen EI, Thakker R V, et al. Evidence for the role of megalin in renal uptake of transthyretin. *J Biol Chem* [Internet]. 2000;275(49):38176–81 [cited 2016 Jan 14]. <http://www.ncbi.nlm.nih.gov/pubmed/10982792>
- [336] Sousa MM, Saraiva MJ. Internalization of transthyretin. Evidence of a novel yet unidentified receptor-associated protein (RAP)-sensitive receptor. *J Biol Chem* [Internet]. 2001;276(17):14420–5. <http://www.ncbi.nlm.nih.gov/pubmed/11278770>
- [337] Divino CM, Schussler GC. Transthyretin receptors on human astrocytoma cells. *J Clin Endocrinol Metab* [Internet]. 1990;71(5):1265–8 [cited 2016 Jan 14]. <http://www.ncbi.nlm.nih.gov/pubmed/2172276>

- [338] Bourgault S, Choi S, Buxbaum JN, Kelly JW, Price JL, Reixach N. Mechanisms of transthyretin cardiomyocyte toxicity inhibition by resveratrol analogs. *Biochem Biophys Res Commun* [Internet]. 2011;410(4):707–13 [cited 2016 Jan 14]. <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3145458&tool=pmcentrez&rendertype=abstract>
- [339] Manral P, Reixach N. Amyloidogenic and non-amyloidogenic transthyretin variants interact differently with human cardiomyocytes: insights into early events of non-fibrillar tissue damage. *Biosci Rep* [Internet]. 2015;35(1) [cited 2016 Jan 14]. <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=4293901&tool=pmcentrez&rendertype=abstract>
- [340] Fleming CE, Mar FM, Franquinho F, Saraiva MJ, Sousa MM. Transthyretin internalization by sensory neurons is megalin mediated and necessary for its neurotogenic activity. *J Neurosci* [Internet]. 2009;29(10):3220–32 [cited 2016 Jan 14]. <http://www.ncbi.nlm.nih.gov/pubmed/19279259>
- [341] Vieira M, Gomes JR, Saraiva MJ. Transthyretin Induces Insulin-like Growth Factor I Nuclear Translocation Regulating Its Levels in the Hippocampus. *Mol Neurobiol* [Internet]. 2015;51(3):1468–79. <http://link.springer.com/10.1007/s12035-014-8824-4>
- [342] Liz MA, Faro CJ, Saraiva MJ, Sousa MM. Transthyretin, a new cryptic protease. *J Biol Chem* [Internet]. 2004;279(20):21431–8 [cited 2016 Jan 13]. <http://www.ncbi.nlm.nih.gov/pubmed/15033978>
- [343] Liz MA, Gomes CM, Saraiva MJ, Sousa MM. ApoA-I cleaved by transthyretin has reduced ability to promote cholesterol efflux and increased amyloidogenicity. *J Lipid Res* [Internet]. 2007;48(11):2385–95 [cited 2016 Jan 13]. <http://www.ncbi.nlm.nih.gov/pubmed/17693625>
- [344] Liz MA, Fleming CE, Nunes AF, Almeida MR, Mar FM, Choe Y, et al. Substrate specificity of transthyretin: identification of natural substrates in the nervous system. *Biochem J* [Internet]. 2009;419(2):467–74 [cited 2016 Jan 13]. <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=4153561&tool=pmcentrez&rendertype=abstract>
- [345] Liz MA, Leite SC, Juliano L, Saraiva MJ, Damas AM, Bur D, et al. Transthyretin is a metallopeptidase with an inducible active site. *Biochem J* [Internet]. 2012;443(3):769–78 [cited 2016 Jan 13]. <http://www.ncbi.nlm.nih.gov/pubmed/22332999>
- [346] Gouvea IE, Kondo MY, Assis DM, Alves FM, Liz MA, Juliano MA, et al. Studies on the peptidase activity of transthyretin (TTR). *Biochimie* [Internet]. 2013;95(2):215–23 [cited 2016 Jan 13]. <http://www.ncbi.nlm.nih.gov/pubmed/23000319>
- [347] Costa R, Ferreira-da-Silva F, Saraiva MJ, Cardoso I. Transthyretin protects against A-beta peptide toxicity by proteolytic cleavage of the peptide: a mechanism sensitive to the Kunitz protease inhibitor. *PLoS One* [Internet]. 2008;3(8):e2899 [cited 2015 Dec 9]. <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2483353&tool=pmcentrez&rendertype=abstract>

- [348] Reilly MM, King RH. Familial amyloid polyneuropathy. *Brain Pathol* [Internet]. 1993;3(2):165–76 [cited 2016 Jan 14]. <http://www.ncbi.nlm.nih.gov/pubmed/8293178>
- [349] Costa PP, Figueira AS, Bravo FR. Amyloid fibril protein related to prealbumin in familial amyloidotic polyneuropathy. *Proc Natl Acad Sci USA* [Internet]. 1978;75(9):4499–503 [cited 2016 Jan 14]. <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=336143&tool=pmcentrez&rendertype=abstract>
- [350] Biochemical marker in familial amyloidotic polyneuropathy, Portuguese type. Family studies on the transthyretin (prealbumin)-methionine-30 variant. PubMed—NCBI [Internet]. 2016 [cited 2016 Jan 14]. <http://www.ncbi.nlm.nih.gov/pubmed/?term=sar-aiva+mj+1985>
- [351] Maeda S, Mita S, Araki S, Shimada K. Structure and expression of the mutant prealbumin gene associated with familial amyloidotic polyneuropathy. *Mol Biol Med* [Internet]. 1986;3(4):329–38 [cited 2016 Jan 14]. <http://www.ncbi.nlm.nih.gov/pubmed/3022108>
- [352] Sousa MM, Cardoso I, Fernandes R, Guimarães A, Saraiva MJ. Deposition of transthyretin in early stages of familial amyloidotic polyneuropathy: evidence for toxicity of nonfibrillar aggregates. *Am J Pathol* [Internet]. 2001;159(6):1993–2000 [cited 2016 Feb 2]. <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=1850610&tool=pmcentrez&rendertype=abstract>
- [353] Berk JL, Suhr OB, Obici L, Sekijima Y, Zeldenrust SR, Yamashita T, et al. Repurposing diflunisal for familial amyloid polyneuropathy: a randomized clinical trial. *JAMA*. 2013;310(24):2658–67.
- [354] Sekijima Y, Tojo K, Morita H, Koyama J, Ikeda S. Safety and efficacy of long-term diflunisal administration in hereditary transthyretin (ATTR) amyloidosis. *Amyloid*. 2015;22(2):79–83.
- [355] Sousa JC, Grandela C, Fernández-Ruiz J, de Miguel R, de Sousa L, Magalhães AI, et al. Transthyretin is involved in depression-like behaviour and exploratory activity. *J Neurochem* [Internet]. 2004;88(5):1052–8 [cited 2016 Jan 14]. <http://www.ncbi.nlm.nih.gov/pubmed/15009661>
- [356] Heilig M. The NPY system in stress, anxiety and depression. *Neuropeptides* [Internet]. 2004;38(4):213–24 [cited 2016 Jan 14]. <http://www.ncbi.nlm.nih.gov/pubmed/15337373>
- [357] Nunes AF, Saraiva MJ, Sousa MM. Transthyretin knockouts are a new mouse model for increased neuropeptide Y. *FASEB J* [Internet]. 2006;20(1):166–8 [cited 2016 Jan 14]. <http://www.ncbi.nlm.nih.gov/pubmed/16263939>
- [358] Sousa JC, Marques F, Dias-Ferreira E, Cerqueira JJ, Sousa N, Palha JA. Transthyretin influences spatial reference memory. *Neurobiol Learn Mem* [Internet]. 2007;88(3):381–5. <http://www.ncbi.nlm.nih.gov/pubmed/17698379>

- [359] Fleming CE, Saraiva MJ, Sousa MM. Transthyretin enhances nerve regeneration. *J Neurochem*. 2007;103(2):831–9.
- [360] Santos SD, Lambertsen KL, Clausen BH, Akinc A, Alvarez R, Finsen B, et al. CSF transthyretin neuroprotection in a mouse model of brain ischemia. *J Neurochem* [Internet]. 2010;115(6):1434–44. <http://www.ncbi.nlm.nih.gov/pubmed/21044072>
- [361] Kooijman R. Regulation of apoptosis by insulin-like growth factor (IGF)-I. *Cytokine Growth Factor Rev* [Internet]. 2006;17(4):305–23 [cited 2016 Jan 14]. <http://www.ncbi.nlm.nih.gov/pubmed/16621671>
- [362] Wisniewski T, Castano E, Ghiso J, Frangione B. Cerebrospinal fluid inhibits Alzheimer's amyloid fibril formation in vitro. *Ann Neurol* [Internet]. 1993;34(4):631–3 [cited 2016 Jan 4]. <http://www.ncbi.nlm.nih.gov/pubmed/8215255>
- [363] Ghiso J, Matsubara E, Koudinov A, Choi-Miura NH, Tomita M, Wisniewski T, et al. The cerebrospinal-fluid soluble form of Alzheimer's amyloid beta is complexed to SP-40,40 (apolipoprotein J), an inhibitor of the complement membrane-attack complex. *Biochem J* [Internet]. 1993;293(Pt 1):27–30 [cited 2016 Jan 4]. <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=1134315&tool=pmcentrez&rendertype=abstract>
- [364] Goldgaber D, Schwarzman AI, Bhasin R, Gregori L, Schmechel D, Saunders AM, et al. Sequestration of amyloid beta-peptide. *Ann N Y Acad Sci* [Internet]. 1993;695:139–43 [cited 2016 Jan 4]. <http://www.ncbi.nlm.nih.gov/pubmed/8239272>
- [365] Schwarzman AL, Gregori L, Vitek MP, Lyubski S, Strittmatter WJ, Enghilde JJ, et al. Transthyretin sequesters amyloid beta protein and prevents amyloid formation. *Proc Natl Acad Sci USA* [Internet]. 1994;91(18):8368–72 [cited 2015 Dec 6]. <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=44607&tool=pmcentrez&rendertype=abstract>
- [366] Schwarzman AL, Goldgaber D. Interaction of transthyretin with amyloid beta-protein: binding and inhibition of amyloid formation. *Ciba Found Symp* [Internet]. 1996;199:146–60 [cited 2015 Nov 30]; discussion 160–4. <http://www.ncbi.nlm.nih.gov/pubmed/8915609>
- [367] Serot JM, Christmann D, Dubost T, Couturier M. Cerebrospinal fluid transthyretin: aging and late onset Alzheimer's disease. *J Neurol Neurosurg Psychiatry*. 1997;63(4):506–8.
- [368] Han S-H, Jung ES, Sohn J-H, Hong HJS, Hong HJS, Kim JW, et al. Human serum transthyretin levels correlate inversely with Alzheimer's disease. *J Alzheimers Dis* [Internet]. 2011;25(1):77–84. <http://www.ncbi.nlm.nih.gov/pubmed/21335655>
- [369] González-Marrero I, Giménez-Llort L, Johanson CE, Carmona-Calero EM, Castañeyra-Ruiz L, Brito-Armas JM, et al. Choroid plexus dysfunction impairs beta-amyloid clearance in a triple transgenic mouse model of Alzheimer's disease. *Front Cell*

- Neurosci [Internet]. 2015;9:17. <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=4319477&tool=pmcentrez&rendertype=abstract>
- [370] Mazur-Kolecka B, Frackowiak J, Wiśniewski HM. Apolipoproteins E3 and E4 induce, and transthyretin prevents accumulation of the Alzheimer's beta-amyloid peptide in cultured vascular smooth muscle cells. *Brain Res* [Internet]. 1995;698(1-2):217-22 [cited 2016 Jan 4]. <http://www.ncbi.nlm.nih.gov/pubmed/8581485>
- [371] Shirahama T, Skinner M, Westermarck P, Rubinow A, Cohen AS, Brun A, et al. Senile cerebral amyloid. Prealbumin as a common constituent in the neuritic plaque, in the neurofibrillary tangle, and in the microangiopathic lesion. *Am J Pathol*. 1982;107(1):41-50.
- [372] Buxbaum JN, Ye Z, Reixach N, Friske L, Levy C, Das P, et al. Transthyretin protects Alzheimer's mice from the behavioral and biochemical effects of Abeta toxicity. *Proc Natl Acad Sci USA* [Internet]. 2008;105(7):2681-6 [cited 2015 Dec 7]. <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2268196&tool=pmcentrez&rendertype=abstract>
- [373] Stein TD, Anders NJ, DeCarli C, Chan SL, Mattson MP, Johnson JA. Neutralization of transthyretin reverses the neuroprotective effects of secreted amyloid precursor protein (APP) in APPSW mice resulting in tau phosphorylation and loss of hippocampal neurons: support for the amyloid hypothesis. *J Neurosci* [Internet]. 2004;24(35):7707-17 [cited 2016 Jan 4]. <http://www.ncbi.nlm.nih.gov/pubmed/15342738>
- [374] Stein TD, Johnson JA. Lack of neurodegeneration in transgenic mice overexpressing mutant amyloid precursor protein is associated with increased levels of transthyretin and the activation of cell survival pathways. *J Neurosci* [Internet]. 2002;22(17):7380-8 [cited 2016 Jan 4]. <http://www.ncbi.nlm.nih.gov/pubmed/12196559>
- [375] Lazarov O, Robinson J, Tang Y-P, Hairston IS, Korade-Mirnic Z, Lee VM-Y, et al. Environmental enrichment reduces A β levels and amyloid deposition in transgenic mice. *Cell* [Internet]. 2005;120(5):701-13 [cited 2015 Oct 7]. <http://www.ncbi.nlm.nih.gov/pubmed/15766532>
- [376] Choi SH, Leight SN, Lee VM-Y, Li T, Wong PC, Johnson J a, et al. Accelerated Abeta deposition in APP^{swe}/PS1 Δ E9 mice with hemizygous deletions of TTR (transthyretin). *J Neurosci*. 2007;27(26):7006-10.
- [377] Oliveira SM, Ribeiro CA, Cardoso I, Saraiva MJ. Gender-dependent transthyretin modulation of brain amyloid- β Levels: evidence from a mouse model of alzheimer's disease. *J Alzheimer's Dis*. 2011;27(2):429-39.
- [378] Costa R, Gonçalves A, Saraiva MJ, Cardoso I. Transthyretin binding to A-Beta peptide –impact on a-beta fibrillogenesis and toxicity. *FEBS Lett*. 2008;582(6):936-42.

- [379] Carro E, Trejo JL, Gomez-Isla T, LeRoith D, Torres-Aleman I. Serum insulin-like growth factor I regulates brain amyloid-beta levels. *Nat Med* [Internet]. 2002;8(12):1390–7 [cited 2015 Dec 29]. <http://www.ncbi.nlm.nih.gov/pubmed/12415260>
- [380] Liu L, Murphy RM. Kinetics of inhibition of beta-amyloid aggregation by transthyretin. *Biochemistry* [Internet]. 2006;45(51):15702–9 [cited 2016 Jan 4]. <http://www.ncbi.nlm.nih.gov/pubmed/17176092>
- [381] Yang DT, Joshi G, Cho PY, Johnson JA, Murphy RM. Transthyretin as both a sensor and a scavenger of β -amyloid oligomers. *Biochemistry* [Internet]. 2013;52(17):2849–61 [cited 2016 Jan 5]. <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3658121&tool=pmcentrez&rendertype=abstract>
- [382] Du J, Murphy RM. Characterization of the interaction of β -amyloid with transthyretin monomers and tetramers. *Biochemistry* [Internet]. 2010;49(38):8276–89 [cited 2016 Jan 5]. <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2943652&tool=pmcentrez&rendertype=abstract>
- [383] Schwarzman AL, Tsiper M, Wenthe H, Wang A, Vitek MP, Vasiliev V, et al. Amyloidogenic and anti-amyloidogenic properties of recombinant transthyretin variants. *Amyloid* [Internet]. 2004;11(1):1–9 [cited 2015 Dec 23]. <http://www.ncbi.nlm.nih.gov/pubmed/15185492>
- [384] Palha JA, Moreira P, Wisniewski T, Frangione B, Saraiva MJ. Transthyretin gene in Alzheimer's disease patients. *Neurosci Lett* [Internet]. 1996;204(3):212–4 [cited 2016 Jan 5]. <http://www.ncbi.nlm.nih.gov/pubmed/8938268>
- [385] Du J, Cho PY, Yang DT, Murphy RM. Identification of beta-amyloid-binding sites on transthyretin. *Protein Eng Des Sel* [Internet]. 2012;25(7):337–45 [cited 2016 Feb 1]. <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3530273&tool=pmcentrez&rendertype=abstract>
- [386] Merched A, Serot JM, Visvikis S, Aguillon D, Faure G, Siest G. Apolipoprotein E, transthyretin and actin in the CSF of Alzheimer's patients: relation with the senile plaques and cytoskeleton biochemistry. *FEBS Lett* [Internet]. 1998;425(2):225–8 [cited 2016 Jan 5]. <http://www.ncbi.nlm.nih.gov/pubmed/9559653>
- [387] Hornstrup LS, Frikke-Schmidt R, Nordestgaard BG, Tybjaerg-Hansen A. Genetic stabilization of transthyretin, cerebrovascular disease, and life expectancy. *Arterioscler Thromb Vasc Biol* [Internet]. 2013;33(6):1441–7 [cited 2015 Oct 27]. <http://www.ncbi.nlm.nih.gov/pubmed/23580146>
- [388] Ribeiro CA, Saraiva MJ, Cardoso I. Stability of the transthyretin molecule as a key factor in the interaction with a-beta peptide—relevance in Alzheimer's Disease. *PLoS One* [Internet]. 2012;7(9):e45368. <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3444465&tool=pmcentrez&rendertype=abstract>

- [389] Cardoso I, Almeida MR, Ferreira N, Arsequell G, Valencia G, Saraiva MJ. Comparative in vitro and ex vivo activities of selected inhibitors of transthyretin aggregation: relevance in drug design. *Biochem J*. 2007;408(1):131–8.
- [390] Ribeiro C a., Oliveira SM, Guido LF, Magalhães A, Valencia G, Arsequell G, et al. Transthyretin stabilization by iododiflunisal promotes amyloid- β peptide clearance, decreases its deposition, and ameliorates cognitive deficits in an Alzheimer's disease mouse model. *J Alzheimer's Dis*. 2014;39:357–70.
- [391] Eckman EA, Reed DK, Eckman CB. Degradation of the Alzheimer's amyloid beta peptide by endothelin-converting enzyme. *J Biol Chem [Internet]*. 2001;276(27):24540–8 [cited 2015 Dec 27]. <http://www.ncbi.nlm.nih.gov/pubmed/11337485>
- [392] Proteolytic mechanisms in amyloid-beta metabolism: therapeutic implications for Alzheimer's disease. *PubMed—NCBI [Internet]*. 2016 [cited 2016 Jan 6]. <http://www.ncbi.nlm.nih.gov/pubmed/16153892>
- [393] Malito E, Hulse RE, Tang W-J. Amyloid beta-degrading cryptidases: insulin degrading enzyme, presequence peptidase, and neprilysin. *Cell Mol Life Sci [Internet]*. 2008;65(16):2574–85 [cited 2016 Jan 14]. <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2756532&tool=pmcentrez&rendertype=abstract>
- [394] Wang Y-J, Zhou H-D, Zhou X-F. Clearance of amyloid-beta in Alzheimer's disease: progress, problems and perspectives. *Drug Discov Today [Internet]*. 2006;11(19–20):931–8 [cited 2016 Jan 14]. <http://www.ncbi.nlm.nih.gov/pubmed/16997144>
- [395] Li X, Masliah E, Reixach N, Buxbaum JN. Neuronal production of transthyretin in human and murine Alzheimer's disease: is it protective? *J Neurosci [Internet]*. 2011;31(35):12483–90 [cited 2015 Nov 16]. <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3172869&tool=pmcentrez&rendertype=abstract>
- [396] Kerridge C, Belyaev ND, Nalivaeva NN, Turner AJ. The A β -clearance protein transthyretin, like neprilysin, is epigenetically regulated by the amyloid precursor protein intracellular domain. *J Neurochem*. 2014;130(3):419–31.
- [397] Wang X, Cattaneo F, Ryno L, Hulleman J, Reixach N, Buxbaum JN. The systemic amyloid precursor transthyretin (TTR) behaves as a neuronal stress protein regulated by HSF1 in SH-SY5Y human neuroblastoma cells and APP23 Alzheimer's disease model mice. *J Neurosci [Internet]*. 2014;34(21):7253–65. <http://www.jneurosci.org/content/34/21/7253.short>

