
Subcritical Water Extraction and Its Prospects for Aflatoxins Extraction in Biological Materials

Sefater Gbashi, Ntakadzeni Edwin Madala,
Oluwafemi Ayodeji Adebo, Lizelle Piater,
Judith Zanele Phoku and Patrick Berka Njobeh

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/intechopen.68706>

Abstract

Aflatoxins (AFs) are well-known mycotoxins and contaminants of various agricultural commodities globally that are linked to a wide range of adverse health and economic complications. Because of their incessant proliferation and deleterious consequences, it has become mandatory to routinely monitor the levels of these toxins in agricultural products before they go into the market. Essentially, effective analysis is an important component of AFs control, and extraction is a necessary step for their analysis, irrespective of the protocol adopted. Conventional methods for AF extraction are expensive, the processes involved are tedious and utilize large quantities of organic solvents that are environmentally unfriendly. This has necessitated the quest for alternatives that are 'green', cost-effective and easy to perform. In this regard, subcritical water extraction (SWE) is a viable alternative that has proven to be effective in the extraction of other bio-active compounds. This chapter presents a critical appraisal of the principles and dynamics of SWE, and its current applications as a viable tool in the extraction of AFs from various biological matrices. Although further research needs to be performed to enhance its applicability, the adoption of SWE in the extraction of AFs seems very promising and needs to be properly exploited.

Keywords: subcritical water extraction, aflatoxins, temperature, biological materials

1. Introduction

The proliferated contamination of agricultural commodities by AFs has become a serious global concern because of their severe impact on health and the economy [1, 2]. This group of mycotoxins is food contaminants produced by filamentous toxigenic fungal species [3, 4] principally those members within the *Aspergillus* genera. Relative to their contamination of food and feed commodities, approximately 4.5 billion people in the world are at the risk of being chronically exposed to mycotoxins, in particular, AFs [5, 6]. Several reports have implicated AFs as very poisonous human and environmental pollutants [2, 7–10]. In fact, one of the AFs, aflatoxin B₁ (AFB₁), has been recognized as the most potent naturally occurring carcinogen known to man [2]. As a result of their widespread proliferation and associated deleterious effects, there is a growing concern over their intake via consumption of contaminated food and feed by humans and animals alike. This has led to more stringent guidelines and regulatory limits of these toxins, especially with the globalization of the food supply chain, and consequently, necessitating routine surveillance of the prevalence as well as levels of AFs in food and feed [11, 12]. As such, the need for more sensitive and robust analytical methods for the determination of AFs is eminent [12], particularly one that is carried out following the greener route.

Extraction is an important step in AFs analysis. It is inevitable irrespective of the protocol adopted. Although different methods exist for extracting AFs from food and feed such as solvent extraction, solid-phase extraction, and immuno-affinity column (IAC) extraction, there are anxieties over human and environmental health regarding safety in their applications [13, 14]. Conventional techniques also involve labor-intensive and time-consuming procedures [15], requiring relatively large volumes of organic solvents, which are expensive and hazardous [13, 16]. Bearing in mind these concerns associated with extraction of AFs, the design of a greener route that is efficient, cheap, fast and relatively easy to address these challenges is significant. Subcritical water extraction (SWE) seems promising in this regard. Better results, recoveries, and effectiveness have been reported for SWE as compared to other traditional methods for extracting different bioactive compounds [17–19]. In this chapter, a comprehensive review on the implications of AFs and issues with their analysis is presented. The need and potential applicability of SWE in AF analysis are highlighted. Lastly, herein, we demonstrate the basic principle of SWE, underscoring its advantages and disadvantages, and wrapping up the chapter with a discussion on how SWE can be suitable in extracting AFs from biological matrices for routine analysis.

2. Aflatoxins

2.1. Definition and concept of aflatoxins

Aflatoxins are the most perilous and troublesome group of mycotoxins to humans and animals that are generally produced by toxigenic strains of fungi, notably *Aspergillus flavus*, *A. parasiticus* and *A. niger* [2, 20, 21]. At least 14 different types of AFs are known to exist in nature, however, the major ones of economic and health significance are aflatoxin B₁ (AFB₁),

B₂ (AFB₂), G₁ (AFG₁), G₂ (AFG₂), M₁ (AFM₁) and M₂ (AFM₂). AFM₁ and M₂ are hydroxylated metabolites of AFB₁ and B₂, respectively, bio-transformed by the liver and found in milk, urine, and other body fluids, being less harmful than their precursor toxins [22, 23]. Among the AF group, AFB₁ is considered the most toxic. This one has been established as the most notorious naturally occurring carcinogen [2, 24, 25]. For that reason, it has been classified as a Group 1A human carcinogen [26]. Cereals such as maize are common crops that are contaminated by AFs. Additionally, crops such as oilseeds, including peanuts, different kinds of spices, figs and other dried fruit, are also familiar but most susceptible substrates.

2.2. Physicochemical properties of aflatoxins

Aflatoxins are a group of closely related difuranocoumarin derivatives, with similar structures as they constitute a unique group of naturally occurring heterocyclic compounds that are highly oxygenated [22]. AFs can be broadly classified into two groups based on their chemical structure namely, difurocoumarocyclopentenone series (AFB₁, B₂, M₁, M₂, and other derivatives) and difurocoumarolactone series (AFG₁, G₂, and others) [22]. Their chemistry constitutes highly substituted coumarins containing a fused dihydrofurofuran moiety. The AFBs (*i.e.*, members of the blue fluorescent series) generally feature a fusion of a cyclopentenone ring to the lactone ring of the coumarin moiety, while the AFGs possess a fused lactone ring [27]. AFB₁ and G₁ contain an unsaturated bond on the terminal furan ring at the 8, 9 position. Epoxidation at this position has shown to be essential for their carcinogenicity [28]. The intensity of fluorescence (light) emission differs greatly among the four compounds. This property plays a significant role in their quantification by fluorescence techniques [29]. AFs are also very stable chemical compounds and notoriously difficult to eradicate in food commodities [30, 31]. They are chemically stable during processing and storage, even when heated at quite elevated temperatures such as those achieved during the production of breakfast cereals or baking of bread [31, 32]. This necessitates the avoidance of conditions that favor their production, which is not always feasible in practice [31, 33].

2.3. Exposure and health implications of aflatoxins

The presence of AFs in foods and feeds is problematic as it induces vicious health repercussions in humans and animals when exposed to them. Poisoning from AFs has been reported in different parts of the world, and victims include humans, animals and other non-human primates [34]. Common exposure routes include ingestion of AF contaminated foods and feeds; however, aerosol, parental (placental and breastfeeding) and dermal routes have also been reported [9, 35], but it can be supposed that ingestion is the main source of AF exposure among humans and animals. This group of poisons enters the blood stream and lymphatic system with the liver as targeted organ and damage macrophage systems inhibit protein synthesis and increase sensitivity to opportunistic infections [36]. Exposure to AFs can be chronic or acute, and symptoms and degree of illness depend on the type of AF, concentration, and duration of exposure, as well as species, age, sex, and health status of the exposed individual [37].

Aflatoxicosis refers to poisoning and associated illness resulting from AF exposure [38, 39]. There are numerous cases of aflatoxicosis reported in the literature. In Ibadan, Nigeria, the

death of some children who consumed mould-infested *Kulikuli*, was suggested to be due to aflatoxicosis [36]. An outbreak of hepatitis in 1974 in India that killed 100 people and caused ailment in hundreds of others was as a result of AFs via consumption heavily contaminated maize [40]. Incidence of liver cancer and aggravated cases of over 40% of diseases in developing nations including kwashiorkor, growth stunting, and HIV are directly or indirectly associated with dietary AF exposure [5, 27, 41]. One of the most epic episodes of aflatoxicosis reported in human history occurred in rural Kenya, of which 317 cases of illness and 125 deaths were reported [42]. The cause of this outbreak was deciphered to be consumption of maize products heavily contaminated with AFs (several folds above the Kenyan regulatory limit of 20 µg/kg). An outbreak of canine aflatoxicosis occurred in South Africa in 2011 leaving over 220 dogs dead and several others seriously affected after consuming pet food contaminated with high levels of AFs [43]. It is, however, very problematic that aflatoxicosis often remains unrecognized by health workers for an extended period of time, except when a large number of people are affected [9].

2.4. Economic implications of aflatoxins

2.4.1. *Economic losses due to aflatoxins contamination of food and feed*

The economic significance of AFs is globally illustrious both in the developed and in the underdeveloped nations. In the United States, losses due to AF contamination of maize are estimated at up to 1.68 billion US dollars annually [44, 45]. Globalization of trade has added to the cost and complexity of the situation. For example, adopting the EU standard limit of 4 µg/kg for AFs in peanuts was estimated to cost about 450 million dollars in annual losses on exports [45, 46]. Although estimates on the economic impact of AFs are scarce in the developing countries, based on the literature reports of high levels of mycotoxins found in agricultural commodities in these countries, it is probable that losses consistently far exceed those reported in the United States [45]. To give an instance, in Southeast Asia, the impact of AFs is calculated to a level of 900 million US dollars, of which 500 million are costs directly related to human health effects [45]. In the coastal and eastern regions of Kenya, 2.3 million bags of maize worth over Ksh 3.2 billion (roughly 30 million US dollars) were declared unfit for human consumption by the Ministry of Public Health and Sanitation in 2010 due to high levels of AF contamination [47]. The change in policy by the European Union (EU) is expected to reduce imports on cereals, dried fruits and oilseeds (mainly nuts) by 64%, costing some nine African countries including Egypt, Nigeria and South Africa about 670 million US dollars in trade per year [48]. These economic impacts add to the complexity of the AF malice in developing countries, as they may be compelled to export their best quality produce and sadly retain the poorer commodities for domestic use [47].

2.4.2. *Research and surveillance costs of aflatoxins in food and feed*

The economic impact of AFs is felt across the entire food and feed supply chains, that is, “from farm to fork.” Costs associated with AF management, that is, from sampling and related research expenses, surveillance, mitigation to litigation are also very significant [45, 49]. A

study conducted in West Africa estimated annual costs averaging 466 million US dollars from testing, regulatory enforcement, to other quality control measures [50]. In 2000, the USDA's Agricultural Research Service (ARS) instituted a mycotoxin research program worth 17.7 million US dollars primarily geared towards prevention of fungal contamination and toxin production in crops [49]. On average, total value of commercially available test kits for AFs on the market is approximately 10 million US dollars annually, whereas the cost for analysis of AFs alone is placed at 30–50 million US dollars on annual basis [49].

2.5. Regulation of aflatoxins

As elucidated earlier in Section 2.4, AFs constitute a major concern to human health and national economies around the world. Due to the fact that AFs are ubiquitous contaminants and potent carcinogens even at low concentrations, they require stringent regulation to ensure food safety and human health. Different countries have established various limits for AFs in agricultural commodities marketed within their jurisdiction based upon their own perception of risk assessment. At present, over 100 countries have regulations in limiting AFs and other mycotoxins in the food and feed industry [51]. AFs are the most regulated mycotoxins, and 61 countries have regulatory limits of AFB₁ in foodstuffs ranging from 1 to 20 µg/kg, 76 countries have limits up to 35 µg/kg for total AFs in foodstuffs, whereas 21 countries have limits of up to 50 µg/kg for total AFs in animal feeds [52]. In South Africa, regulations exist for total AFs in peanuts intended for further processing (15 µg/kg), in ready-to-eat foodstuffs for humans (10 µg/kg of which AFB₁ is not more than 5 µg/kg), and AFM₁ in dairy milk (0.05 µg/L) [53, 54]. The Joint Expert Committee on Food Additives (JECFA) of the Food and Agriculture Organization (FAO) and World Health Organization (WHO) serving as a scientific advisory body to CODEX Alimentarius Commission recommended that levels of AFs in food should be kept As Low As Reasonably Achievable (ALARA) [52]. FAO [52] has published a compendia summarizing worldwide regulations for AFs and other mycotoxins. Many other similar synopses on limits and regulations for AFs are available in literature and could be consulted for additional information [55–57].

2.6. Analysis of aflatoxins in food and feed

Due to the severe effects that AFs elicit in animal and man, several countries and politico-economic unions have placed high priority on the safety of agricultural commodities marketed and consumed within their jurisdiction [51, 52, 58]. Particularly with the globalization of trade, much efforts have been put into mitigation and control of the prevalence of this toxin group in food and feed [59–62]; however, it is apparent that the complete elimination of AFs from foods is an unattainable objective [37]. This has led to various interventions put in place to manage and minimize risk exposure to them [63, 64]. Adequate risk management has been identified as a critical frontline defense in the overall control of AFs in food and feed supplies [36, 63, 65–67]. Any good food safety management program for naturally occurring toxicants [such as Hazard Analysis and Critical Control Point (HACCP)] assumes a holistic approach, involving various phases such as determination of exposure levels, establishment of analytical capabilities, setting and ensuring compliance with regulatory limits, and establishment of surveillance programs [36, 66].

Such critical approaches provide for routine and detailed analysis of every step throughout the food supply chain (from farm to fork) [36, 63]. This positions analysis at the epicenter of AFs management and risk control, which is a global priority [11, 68]. Bearing in mind that decisions relating to regulatory issues or commercial arbitration need to be based on well-defined methods of analysis [66], it is thus vital to ensure that methods for AF analysis are sensitive, efficient and validated against standard guidelines [69, 70]. In AF analysis, the role played by extraction and sample preparation, in general, cannot be overemphasized. Extraction is an inevitable step in AFs analysis no matter the analytical method employed. It has been estimated that up to 70% and perhaps even more of the effort and time that goes into sample analysis comprises the extraction and sample preparation process [71]. Proper design of the extraction process facilitates rapid, efficient and quality analytical results [71].

2.6.1. Extraction of aflatoxins

Many efforts have been geared towards developing suitable methods to quantitatively extract and detect AFs in agricultural commodities. For any bioanalytical chemists, the goal is to develop methods with improved sensitivity and selectivity, while at the same time maintaining the credibility of the results, as well as reduce cost and time [72].

2.6.1.1. Conventional extraction methods for aflatoxins in food and feed

Different methods have been used for AFs extraction in food and feed. Of these methods, solvent extraction is one of the oldest but still most frequently used method [73]. This method separates analytes based on their relative solubility in two different liquids that are immiscible [74]. One or more solutes contained in a feed solution are transferred to another immiscible solvent, often by rigorously mixing the two immiscible phases, then allowing the two phases to separate [74, 75]. The enriched solvent is called the extract [76]. Common solvents used for solvent extraction include methanol, acetonitrile, chloroform, ethyl acetate, isooctane, ethanol and dichloromethane [4, 73, 77]. The most commonly used solvent extraction approach for AFs is the multi-mycotoxin extraction method of Patterson and Roberts [78]. This method utilizes different organic solvents and reagents such as acetonitrile, isooctane, potassium chloride, dichloromethane and sulfuric acid. It has been widely favored because it selectively extracts several mycotoxins in a single extraction. However, the application of solvent extraction has been greatly limited because it enables the consumption of large quantities of organic solvents, which pose hazards to the environment [16, 79]. Furthermore, solvent extraction often involves long extraction times and laborious procedures with the process extending up to 24 h or more [4, 78]. Moreover, solvents of the required purity tend to be expensive and there are often additional costs with proper disposal of wastes after use [74, 79].

Solid phase extraction (SPE) is another very commonly used extraction method for AFs. It involves the separation of analytes between a liquid mobile phase and a stationary phase contained in a cartridge. Typical materials used at the solid adsorbent phase include ethyl (C2), octyl (C8), octadecyl (C18), cyanopropyl (CN), aminopropyl (NH₂), and an ion exchange phase [80]. Non-specific SPE materials are commonly still employed in AF analysis, which is often used for the extraction of more than one mycotoxin [73]. The use of more analyte-specific stationary

phases such as immunoaffinity (IA) materials that contain specific antibodies that bind to the analyte of interest is also gaining much attention [70, 81]. Although SPE techniques are relatively simple, have higher specificity and require little quantities of solvents, they are also very expensive and the antibodies are not available for some mycotoxins and products [80].

2.6.1.2. Other methods for aflatoxin extraction

Aside from the extraction methods discussed above, several other methods have been investigated for the extraction of AFs, some of which include quick, easy, cheap, effective, rugged, and safe (QuEChERS) [82, 83], supercritical fluid extraction (SFE) [84], ultrasonic extraction [85], and many others reviewed in the literature [12, 70, 73, 86, 87]. However, as elucidated previously in Section 2.6.1.1, these techniques are fundamentally limited by the use of large volumes of organic solvents, some of which are well known to be toxic and considered as environmental hazards, issues of low recovery, long and laborious procedures and high costs involved amongst others [72]. Moreover, novel advancements in spectrometric analysis of bioactive compounds (e.g., “omics”) are pushing the limits of conventional techniques of extraction [72, 88].

Further to this, the adoption of an extraction method strongly depends on the analytical objectives; hence, for AF analysis, methods are required to meet established benchmark standards of the survey, monitory work, legislation and research [89]. It is in line with this that we propose the adoption of SWE as an alternative to conventional extraction methods for AFs, particularly with respect to improved recovery and selectivity, reduced organic solvent consumption and extraction time, at a lower cost. SWE has been in the spotlight as an efficacious and highly promising alternative to traditional techniques of extraction, whose successful applications in the biochemical, pharmaceutical and chemical engineering fields have been well documented in the literature [14, 90–96]. A brief description of this method of extraction is presented in the proceeding sections of this chapter.

3. Subcritical water extraction

3.1. Concept and principle of subcritical water extraction

The term subcritical water refers to liquid water between the boiling point temperature and critical point temperature of water (100–374°C) (**Figure 1**) [14]. Pressure is applied to keep the water in liquid state. Subcritical water extraction (SWE) is a green, cheap and easy-to-adopt extraction technique that utilizes water within its subcritical region as the extraction solvent [14]. The phenomenon behind the extractability of subcritical water is based on the fact that when the temperature of water is raised and the pressure kept sufficient to maintain it in its liquid state (e.g., 250°C and 50 bar), the dielectric constant of water decreases and the hydrogen bond and other intermolecular forces of water weakens, which greatly enhancing its extractability [14, 97].

At atmospheric temperature and pressure (25°C at 1 bar), water has one of the highest dielectric constants amongst non-metallic liquids ($\epsilon=80$) [98]. However, when the temperature and pressure of water are raised to 250°C and 50 bar, respectively, the dielectric constant falls ($\epsilon=27$), which is around the range of non-polar solvents such as methanol ($\epsilon=33$), acetone




Figure 1. Phase diagram of water as a function of temperature and pressure [14].

($\epsilon=20.7$), ethanol ($\epsilon=24$), and acetonitrile ($\epsilon=37$) [13, 97]. As a result of the drop in the dielectric constant of subcritical water, its surface tension and viscosity decrease, while its diffusivity increases [13, 14]. As such, water behaves like an organic solvent, dissolving a wide range of low and medium polarity analytes [14]. Interestingly, the extractability and selectivity of subcritical water can be easily maneuvered to extract a range of analytes by simply varying the temperature conditions of the water [90]. Another theoretical explanation on the extractability of subcritical water basis this ability on the fact that, as the temperature of water increases, the average kinetic energy of the molecules of the mixture also increases. This thus disrupts the bonds that exist within and between the molecules, as such, increasing extraction rate.

3.2. Instrumentation of subcritical water extraction

A typical setup of a laboratory scale SWE unit comprises a source of water, temperature retention coil, a solvent pump, an oven and extraction cell, a backpressure valve and a condenser connected to the outlet (Figure 2). The grounded sample to be extracted is placed inside the extraction cell, which is located inside the oven. The oven, which usually has an automatic thermostat mechanism, is set to the desired temperature, the backpressure valve is partially locked to maintain the desired pressure and water is pumped at a preset flow rate through the retention coil into the extraction cell. The extraction takes place in the extraction cell as the subcritical water flows through it and mixes with the sample. The hot water extract flows through the condenser and is collected at the outlet [14].




Figure 2. Simple laboratory setup of a PHWE unit [95].

3.3. Operational modes of pressurized hot water extraction

Extraction with subcritical water is performed in two common modes, static mode and flow-through (dynamic) mode. Extraction in the static mode involves retaining the sample in the extraction cell with subcritical water for short periods after which the fluid is allowed to flow out purging the extraction cell and extract collected. It is important to optimize the retention periods to allow for an equilibrium to be reached between the solvent and analyte. The disadvantage of operating in this mode is that within a short time the analyte fluid equilibrium is reached, and hence, no further extraction of the analyte occurs no matter how long the samples are retained in the extraction. On the other hand, extraction in the dynamic mode allows for a continuous flow of fresh fluid through the extraction cell, which reduces or eliminates analyte-fluid equilibrium in a single operation when properly optimized. As such, recovery efficiency is higher in the dynamic mode, although, fluid consumption could be more, resulting in lower energy efficiency compared to the static mode [13, 14]. In a study by Yang *et al.* [99], it was observed that extraction in dynamic mode resulted in the higher recovery of lignin and hemicelluloses from maize stover cellulose than the static mode.

3.4. Factors affecting subcritical water extraction

A number of factors such as temperature, flow rate, pressure, particle size, co-solvents and surfactants affect the performance of SWE. Some of these factors are further described below.

3.4.1. Temperature

The extraction efficiency of SWE is strongly affected by changes in temperature [100]. Generally, extraction efficiency increases with increase in temperature. A higher recovery of total antioxidants was achieved from grape pomace by increasing the extraction temperature [101]. Despite the increase in efficiency by increase in temperature, excess temperatures can result in degradation of thermolabile analytes, hence the need for optimization [100, 102]. The

recovery of carvacrol and thymol from *Zataria multiflora* between 100 and 175°C indicated that recoveries increased steadily with the increase in temperature up to 150°C, then a degradation phenomenon followed with a noticeable burning smell [103].

3.4.2. Pressure

The effect of pressure on the extraction efficiency SWE has been described as insignificant [104, 105]. In a study by Shalmashi *et al.* [106], changes in pressure, that is, 20, 30 and 40 bar during SWE did not show any significant effect on the recovery caffeine from tea waste. This is because water is fairly incompressible at temperatures below 300°C, which implies that pressure has very little influence on the physicochemical properties of water, as long as it can maintain in a liquid state [107, 108]. Nevertheless, increased pressure can compromise matrix tissue membranes and force the extraction fluid deep into matrix pores where water at lower pressure may not normally reach [109].

3.4.3. Cosolvents and modifiers

Cosolvents and solvent modifiers are often used to enhance the extractability of SWE. Cosolvents are secondary solvents (usually organic solvents) that are added to subcritical water to enhance its solvation power [95, 110]. The incorporation of methanol during SWE was observed to significantly ($p < 0.05$) increase yield of flavonoids and di-acylated cinnamic acids from *Bidens pilosa* [95, 111]. Solvent modifiers such as salts and other reagents can alter important physicochemical properties of water such as polarity, surface tension, and hydrogen bonding strength, which results in an enhanced extractability [13, 112]. Modifiers can also interact directly with the sample matrix, reducing the activation energy required for analyte desorption and diffusion [14, 113]. Elsewhere [112], it was observed that the solubility of atrazine can be doubled when urea was added to subcritical water, and when ethanol was used, the solubility increased by over 10-folds.

In addition to the above-described factors, other factors that influence the extractability of SWE include solvent flow rate, physicochemical and functional characteristics of the sample matrix and analyte, matrix particle size and geometry of extraction cell [13, 14, 114].

3.5. Advantages and disadvantages of subcritical water extraction

3.5.1. Advantages of subcritical water extraction

The major advantage of SWE is that it is a green (i.e., environmentally friendly) extraction method. The extractant is water, which is non-toxic, non-flammable and renewable. Moreover, water is readily available and cheap, and extraction with it does not generate harmful by-products [90, 115]. In comparison with traditional extraction methods, SWE is less time-consuming and much easier to perform with very few extraction steps, as such, human errors are greatly minimized. When put side-by-side with supercritical fluid extraction (SFE), SWE edges on the basis of being a simple technology, hence, requiring much lower maintenance and engineering cost for equipment [14, 19]. During extraction with subcritical water, the fluid can be maneuvered to selectively extract a range of analytes with different polarities by mere adjusting the temperature of the water, whereas SFE extracts only nonpolar or light-weight

compounds [90, 112]. Further to this, SWE is very compatible with various analytical instrumentations because water is colorless and may not interfere with sorts of photodetection such as UV detection or flame ionization detection [14, 116].

3.5.2. Disadvantages of subcritical water extraction

A major setback of SWE is the thermal degradation of some thermolabile analytes at elevated temperatures [117, 118]. When the temperature and pressure of water are extremely high (*i.e.*, above 374°C and 221 bar), there is also the risk that water can become very reactive and could oxidize or catalyze the hydrolysis of some compounds [13]. However, optimization by means of the adoption of a cosolvent or modifier could ameliorate or eliminate these issues [95].

3.6. Application of subcritical water extraction

In the last decade, SWE has been widely investigated for the extraction of various nutritional constituents, organic pollutants, and pharmacoactive compounds from vegetal tissues, food products, soil residues and other ecological biomasses [13, 14, 100, 119]. Free fatty acids and other oils were extracted from spent bleaching earth using SWE [120]. Likewise, it was possible to recover important metabolites from *Moringa oleifera* leaves using SWE [117]. The similar extraction method was used for the recovery of proteins, carbohydrates, and lignans from flaxseed meal [121], catechins and proanthocyanidins from grape seeds [122], flavonoids from aspen knotwood [123] and antioxidants from microalga *Spirulina platensis* [124]. The use of SWE in various applications in different scientific disciplines has been reviewed [90, 100, 107].

3.7. Prospects of subcritical water extraction of aflatoxins

In a recent study [125], we developed and validated an SWE method for the extraction of AFB₁ from maize and subsequently, analysis on high-performance liquid chromatography followed. Results obtained from that study revealed that SWE is suitable for the effective extraction of AFB₁ from maize matrix, with recovery rates ranging from 37 to 128%. Subsequent validation of the optimized method showed acceptable values for accuracy or recovery rate (116%), linearity (%RSD 0.93) and repeatability (%RSD 1.63). It has been stated earlier in Section 2.5 that more countries are enforcing stringent regulations limiting AFs in food and feed, which is increasing demand for their analysis. The efficiency, simplicity, safety and low-cost implications of using SWE are very attractive and compelling in this regard. In comparison with conventional solvent extraction techniques, SWE is very easy to use and requires less time and money [18].

It is known that AFs occur in a diverse manner and can be found deeply deposited inside the food matrices, and as such, their extraction usually requires a process that allows the solvent to penetrate all areas of the matrix to reach hidden toxins trapped in matrix pores [126]. The high pressures involved in SWE seem very suitable in meeting this requirement. Although issues with thermal degradation of some analytes have been a major limiting factor of SWE, it is interesting to know that AFs and most other mycotoxins are relatively thermally stable [31, 62]. Moreover, optimization using cosolvents has been found effective in ameliorating this setback [95, 127]. Accordingly, in our recent study [125] described in the beginning of this section (Section 3.7), we observed a clear positive enhancement on the recovery of AFB₁ by means of a

cosolvent (methanol). Based on these observations and other consulted literature reports, it is evident that SWE is a viable alternative to conventional extraction methods for AFs [14, 125].

4. Conclusion

From the literature documents reviewed herein, it has been established that AFs are very potent natural toxins that constitute a significant nuisance to human and animal health as well as the economy. One way to amply combat the prevalence of these toxic substances is by frequent monitoring of their occurrence levels at various critical points along the food supply chain. To this effect, various national and international regulations have been established and are being enforced. Efficient analytical capabilities provide adequate insights on the prevalence of these toxins, which constitute a basis to monitor and where necessary readress such interventions. This has positioned analysis as a critical element in AF management and control. Extraction is an important step during AF analysis, and hence, improvement in extraction has been a priority in aflatoxicology research. There is a continual quest for efficient extraction methods that are fast, safe and deliver suitable results at reduced cost. SWE meets all these requirements and could make for efficient routine analysis of AFs and other important fungal metabolites in foods and feeds. These observations could stimulate interest and further propel the adoption of SWE in many other applications even beyond the mycotoxicology domain, as well as its scale-up for subsequent industrial applications.

Acknowledgements

This piece of work was financially supported via the Global Excellence and Stature Fellowship of the University of Johannesburg granted to the main author. This chapter was supported in part via the NRF Center of Excellence (CoE) in Food Security co-hosted by the University of Pretoria and the University of the Western Cape in South Africa.

Author details

Sefater Gbashi^{1*}, Ntakadzeni Edwin Madala², Oluwafemi Ayodeji Adebo¹, Lizelle Piater², Judith Zanele Phoku³ and Patrick Berka Njobeh¹

*Address all correspondence to: sefatergbashi@gmail.com

1 Department of Biotechnology and Food Technology, Faculty of Science, University of Johannesburg, South Africa

2 Department of Biochemistry, Faculty of Science, University of Johannesburg, South Africa

3 Water and Health Research Centre, Faculty of Health Sciences, University of Johannesburg, South Africa

References

- [1] Lizárraga-Paulín EG, Moreno-Martínez E, Miranda-Castro SP. Aflatoxins and their impact on human and animal health: An emerging problem. In: Guevara-Gonzalez RG, editor. *Aflatoxins – Biochemistry and Molecular Biology*. Rijeka, Croatia: InTech; 2011. pp. 255-282
- [2] Makun HA, Dutton MF, Njobeh PB, Gbodi TA, Ogbadu GH. Aflatoxin contamination in foods and feeds: A special focus on Africa. In: Ayman AE, editor. *Trends in Vital Food and Control Engineering*. Rijeka, Croatia: InTech; 2012. pp. 187-234
- [3] Njobeh BP, Dutton FM, Makun HA. Mycotoxins and human health: Significance, prevention and control. In: Mishra AK, Tiwari A, Mishra SB, editors. *Smart Biomolecules in Medicine*. India: VBRI Press; 2010. pp. 132-177
- [4] Chilaka CA, Kock S De, Phoku JZ, Mwanza M, Augustina M, Dutton MF. Fungal and mycotoxin contamination of South African commercial maize. *Journal of Food, Agriculture and Environment*. 2012;**10**(2):296-303
- [5] Turner PC, Collinson AC, Cheung YB, Gong Y, Hall AJ, Prentice AM, et al. Aflatoxin exposure in utero causes growth faltering in Gambian infants. *International Journal of Epidemiology*. 2007;**36**:1119-1125
- [6] Bryden WL. Mycotoxins in the food chain: Human health. *Asia Pacific Journal of Clinical Nutrition*. 2007;**16**(1):95-101
- [7] Zain ME. Impact of mycotoxins on humans and animals. *Journal of Saudi Chemical Society*. 2011;**15**(2):129-144
- [8] Akande KE, Abubakar MM, Adegbola TA, Bogoro SE. Nutritional and health implications of mycotoxins in animal feeds: A review. *Pakistan Journal of Nutrition*. 2006;**5**(5):398-403
- [9] Peraica M, Radic B, Lucic A, Pavlovic M. Toxic effects of mycotoxins in humans. *The Bulletin of the World Health Organization*. 1999;**77**(9):754-766
- [10] Shephard GS. Impact of mycotoxins on human health in developing countries. *Food Additives & Contaminants*. 2008;**25**(2):146-151
- [11] Krska R, Schubert-ullrich P, Molinelli A, Sulyok M, MacDonald S, Crews C. Mycotoxin analysis: An update. *Food Additives & Contaminants*. 2008;**25**(2):152-163
- [12] Rahmani A, Jinap S, Soleimany F. Qualitative and quantitative analysis of mycotoxins. *Comprehensive Reviews in Food Science and Food Safety*. 2009;**8**:202-251
- [13] Teo CC, Tan SN, Yong J, Hew CS, Ong ES. Pressurized hot water extraction (PHWE). *Journal of Chromatography A*. 2010;**1217**:2484-2494
- [14] Gbashi S, Adebo OA, Piater L, Madala NE, Njobeh PB. Subcritical water extraction in biological materials. *Separation and Purification Reviews*. 2016;**46**:21-34

- [15] Choi MPK, Chan KKC, Leung HW, Huie CW. Pressurized liquid extraction of active ingredients (ginsenosides) from medicinal plants using non-ionic surfactant solutions. *Journal of Chromatography A*. 2003;**983**:153-162
- [16] Santana MC, Ferrera ZS, Padrón MET, Rodríguez JJS. Methodologies for the extraction of phenolic compounds from environmental samples: New approaches. *Molecules*. 2009;**14**:298-320
- [17] Hawthorne SB, Grabanski CB, Martin E, Miller DJ. Comparisons of Soxhlet extraction, pressurized liquid extraction, supercritical fluid extraction and subcritical water extraction for environmental solids: Recovery, selectivity and effects on sample matrix. *Journal of Chromatography A*. 2000;**892**(1-2):421-33
- [18] Kubátová A, Miller DJ, Hawthorne SB. Comparison of subcritical water and organic solvents for extracting kava lactones from kava root. *Journal of Chromatography A*. 2001;**923**(1-2):187-194
- [19] Bart JCJ. Additives in Polymers: Industrial Analysis and Applications. West Sussex, England: John Wiley and Sons Ltd, 2005. p. 100
- [20] Yang J, Li J, Jiang Y, Duan X, Qu H, Yang B, et al. Natural occurrence, analysis, and prevention of mycotoxins in fruits and their processed product. *Critical Reviews in Food Science and Nutrition*. 2014;**54**(1):64-83
- [21] Mircea C, Poiata A, Tuchilus C, Agoroae L, Butnaru E, Stanescu U. Aflatoxigenic fungi isolated from medicinal herbs. *Toxicology Letters*. 2008;**180**:32-246
- [22] Bbosa GS, Kitya D, Lubega A, Ogwal-Okeng J, Anokbonggo WW, Kyegombe DB. Review of the biological and health effects of aflatoxins on body organs and body systems. In: Razzaghi-Abyaneh M, editor. *Aflatoxins – Recent advances and future prospects*. Rijeka, Croatia: InTech; 2013. pp. 239-265
- [23] Prado G, de Oliveira MS, Lima AS, Moreira APA. Occurrence of aflatoxin M₁ in parmesan cheese consumed in Minas Gerais, Brazil. *Ciência e Agrotecnologia*. 2008;**32**(6):1906-1911
- [24] IARC. IARC Monographs on the Evaluation of Carcinogenic Risks to Humans. Some Traditional Herbal Medicines, Some Mycotoxins, Naphthalene and Styrene. Vol. 82. Lyon, France: IARC; 2002
- [25] D'Mello J. Mycotoxins in Cereal Grains, Nuts and Other Plant Products: Food Safety: Contaminants and Toxins. Wallingford, England: CAB International; 2003. pp. 65-90
- [26] Park JW, Kim EK, Kim YB. Estimation of the daily exposure of Koreans to aflatoxin B₁ through food consumption. *Food Additives & Contaminants*. 2004;**21**:70-75
- [27] Kensler TW, Roebuck BD, Wogan GN, Groopman JD. Aflatoxin: A 50-year odyssey of mechanistic and translational toxicology. *Toxicological Sciences*. 2011;**120**(SI):S28-S48
- [28] Groopman JD, Kensler TW. Role of metabolism and viruses in aflatoxin-induced liver cancer. *Toxicology and Applied Pharmacology*. 2005;**206**:131-137

- [29] Dhanasekaran D, Shanmugapriya S, Thajuddin N, Panneerselvam A. Aflatoxins and aflatoxicosis in human and animals. In: Guevara-Gonzalez RG, editor. *Aflatoxins – Biochemistry and molecular Biology*. 1st ed. Rijeka, Croatia: InTech; 2011. pp. 221-254
- [30] Bullerman LB, Schroeder LL, Park KY. Formation and control of mycotoxins in food. *Journal of Food Protection*. 1984;**47**:637-646
- [31] Turner NW, Subrahmanyam S, Piletsky SA. Analytical methods for determination of mycotoxins: A review. *Analytica Chimica Acta*. 2009;**632**:168-180
- [32] Kabak B. The fate of mycotoxins during thermal food processing. *Journal of the Science of Food and Agriculture*. 2009;**89**(4):549-554
- [33] Fernández-Cruz ML, Mansilla ML, Tadeo JL. Mycotoxins in fruits and their processed products: Analysis, occurrence and health implications. *Journal of Advanced Research*. 2010;**1**:113-122
- [34] Reddy L. Chemoprotective action of natural products on cultured human epithelial cells exposed to aflatoxin B₁. Durban Institute of Technology, Durban, South Africa; 2005
- [35] Boonen J, Malysheva S V, Taevernier L, Diana Di Mavungu J, De Saeger S, De Spiegeleer B. Human skin penetration of selected model mycotoxins. *Toxicology*. 2012;**301**(1-3):21-32
- [36] Atanda O, Makun HA, Ogara IM, Edema M, Idahor KO, Eshiett ME, et al. Fungal and mycotoxin contamination of Nigerian foods and feeds. In: Makun HA, editor. *Mycotoxin and Food Safety in Developing Countries*. Rijeka, Croatia: InTech; 2013. pp. 1-38
- [37] Bennett J, Klich M. Mycotoxins. *Clinical Microbiology Reviews*. 2003;**16**(3):497-516
- [38] Pettersson H. Controlling mycotoxins in animal feed. In: Magan N, Olsen M, editors. *Mycotoxins in Food: Detection and Control*. Cambridge, England: Woodhead Publishing Ltd. 2004; pp. 269-273
- [39] Williams JH, Phillips TD, Jolly PE, Stiles JK, Jolly CM, Aggarwal D. Human aflatoxicosis in developing countries: A review of toxicology, exposure, potential health consequences, and interventions 1, 2, 3. *The American Journal of Clinical Nutrition*. 2004;**80**:106-122
- [40] Montville TJ, Matthews KR. *Food Microbiology: An Introduction*. 2nd ed. Washington DC, USA: ASM Press. 2008; p. 428
- [41] Jolly PE, Shuaib FM, Jiang Y, Preko P, Baidoo J, Stiles JK, et al. Association of high viral load and abnormal liver function with high aflatoxin B₁-albumin adduct levels in HIV-positive Ghanaians: Preliminary observations. *Food Additives & Contaminants. Part A, Chemistry, Analysis, Control, Exposure & Risk Assessment*. 2011;**28**(9):1224-1234
- [42] Lewis L, Onsongo M, Njapau H, Schurz-Rogers H, Lubber G, Kieszak S, et al. Aflatoxin contamination of commercial maize products during an outbreak of acute aflatoxicosis in eastern and central Kenya. *Environmental Health Perspectives*. 2005;**113**(12):1763-1767
- [43] Arnot LF, Duncan NM, Coetzer H, Botha CJ. An outbreak of canine aflatoxicosis in Gauteng Province, South Africa. *Journal of the South African Veterinary Association*. 2012;**83**(1):1-4

- [44] Mitchell NJ, Bowers E, Hurburgh C, Wu F. Potential economic losses to the US corn industry from aflatoxin contamination. *Food Additives & Contaminants. Part A, Chemistry, Analysis, Control, Exposure & Risk Assessment*. 2016;**33**(3):540-550
- [45] Schmale DG, Munkvold GP. Mycotoxins in crops: A threat to human and domestic animal health. *Plant Health Instructor Journal*. 2009;**3**:340-353
- [46] Wilson SJ, Otsuki T. What price precaution? European harmonization of aflatoxin regulations and African groundnut exports. *European Review of Agricultural Economics*. 2001;**28**(3):263-284
- [47] Marechera G, Ndwiga J. Farmer perceptions of aflatoxin management strategies in lower Eastern Kenya. *Journal of Agricultural Extension and Rural Development*. 2014;**6**(12):382-392
- [48] Bankole SA, Adebajo A. Mycotoxins in food in West Africa: Current situation and possibilities of controlling it. *African Journal of Biotechnology*. 2003;**2**(9):254-263
- [49] Robens J, Cardwell K. The costs of mycotoxin management to the USA: Management of aflatoxins in the United States. *Journal of Toxicology Toxin Reviews*. 2003;**22**(2-3):139-152
- [50] CAST. Mycotoxins: Risks in plant, animal, and human systems. Council for Agricultural Science and Technology (CAST). Task Force Report No. 139. Ames, Iowa, USA; 2003
- [51] Van Egmond HP, Schothorst RC, Jonker MA. Regulations relating to mycotoxins in food. *Analytical and Bioanalytical Chemistry*. 2007;**389**:147-157
- [52] FAO. Worldwide Regulations for Mycotoxins in Food and Feed in 2003. Food and Agriculture Organization of the United Nations (FAO) Food Quality and Standards Service. Food and Nutrition Paper 81. Rome, Italy: FAO; 2004
- [53] MRC. Guidelines on mycotoxin control in South African foodstuffs: From the application of the Hazard Analysis and Critical Control Point (HACCP) system to new national mycotoxin regulations. Medical Research Council Policy Brief. 2009:1-12
- [54] Government Gazette. Regulations Governing Tolerances for Fungus-produced Toxins in Foodstuffs. Government Gazette 32409 South Africa: Section 15(1) of the Foodstuffs, Cosmetics and Disinfectants Act, 1972. 2009; p. 6-7
- [55] Anukula N, Vangnaib K. Significance of regulation limits in mycotoxin contamination in Asia and risk management programs at the national level. *Journal of Food and Drug Analysis*. 2013;**21**(3):227-241
- [56] Weidenbörner M. *Encyclopedia of Food Mycotoxins*. 1st ed. Berlin Heidelberg, Germany: Springer-Verlag Berlin Heidelberg; 2001
- [57] Guzmán-de-Peña D, Peña-Cabriaes JJ. Regulatory considerations of aflatoxin contamination of food in Mexico. *Revista Latinoamericana de Microbiología*. 2005;**47**(3-4):160-164
- [58] Creppy EE. Update of survey, regulation and toxic effects of mycotoxins in Europe. *Toxicology Letters*. 2002;**127**(1-3):19-28

- [59] Suttajit M. Prevention and Control of Mycotoxins [Internet]. Mycotoxin prevention and control in food grains. FAO Agriculture and Consumer Protection; 1989. Available from: <http://www.fao.org/docrep/X5036E/x5036E0q.htm> [Accessed: February 15, 2017]
- [60] Bhat R, Rai R V, Karim AA. Mycotoxins in food and feed: Present status and future concerns. *Comprehensive Reviews in Food Science and Food Safety*. 2010;**9**(1):57-81
- [61] Karlovsky P, Suman M, Berthiller F, De Meester J, Eisenbrand G, Perrin I, et al. Impact of food processing and detoxification treatments on mycotoxin contamination. *Mycotoxin Research*. 2016;**32**(4):179-205
- [62] Bullerman LB, Bianchini A. Stability of mycotoxins during food processing. *International Journal of Food Microbiology*. 2007;**119**:140-146
- [63] Aldred D, Magan N, Olsen M. The use of HACCP in the control of mycotoxins: The case of cereals. In: Magan N, Olsen M, editors. *Mycotoxins in Food: Detection and Control*. Cambridge, England: Woodhead Publishing Ltd.; 2004. pp. 139-173
- [64] Lopez-Garcia R, Park DL, Phillips TD. Integrated mycotoxin management systems. *Recent Patents on Food, Nutrition & Agriculture*. 1999;**23**:38-48
- [65] Toregeani-Mendes KA, Arroiteia CC, Kemmelmeier C, Dalpasquale VA, Bando É, Alves AF, et al. Application of hazard analysis critical control points system for the control of aflatoxins in the Brazilian groundnut-based food industry. *International Journal of Food Science & Technology*. 2011;**46**(12):2611-2618
- [66] Gil L, Ruiz P, Font G, Manyes L. An overview of the applications of hazards analysis and critical control point (HACCP) system to mycotoxins. *Critical Reviews in Toxicology*. 2016;**33**:50-55
- [67] FAO/IAEA. Manual on the application of the HACCP system in mycotoxin prevention and control. Joint Food and Agriculture Organization of the United Nations (FAO), and the International Agency for Atomic Energy (IAEA) Training and Reference Centre for Food and Pesticide Control, Rome, Italy. 2001; FAO Food and Nutrition Paper **73**:1-104
- [68] Patel P. Mycotoxin analysis: Current and emerging technologies. In: Magan N, Olsen M, editors. *Mycotoxins in Food: Detection and Control*. Cambridge, England: Woodhead Publishing Limited; 2004. pp. 88-110
- [69] Pittet A. Modern methods and trends in mycotoxin analysis. *Mitteilungen aus Leb und Hyg*. 2005;**96**:424-444
- [70] Arroyo-Manzanares N, Huertas-Pérez JF, García-Campaña AM, Gámiz-Gracia L. Mycotoxin analysis: New proposals for sample treatment. *Advances in Chemistry*. 2014;**2014**:1-12
- [71] Rezaee M, Khalilian F, Pourjavid MR, Seidi S, Chisvert A, Abdel-Rehim M. Extraction and sample preparation. *International Journal of Analytical Chemistry*. 2015;**2015**:1-2
- [72] Augusto F, Hantao LW, Mogollo NGS, Braga SCGN. New materials and trends in sorbents for solid-phase extraction. *Trends in Analytical Chemistry*. 2013;**43**:14-23

- [73] Cigić IK, Prosen H. An overview of conventional and emerging analytical methods for the determination of mycotoxins. *International Journal of Molecular Sciences*. 2009;**10**(1):62-115
- [74] Sapkale GN, Patil SM, Surwase US, Bhatbhage PK. Supercritical fluid extraction. *International Journal of Chemical Sciences*. 2010;**8**(2):729-743
- [75] Zaiz T, Lanez H, Kechida B. Aspen hysys simulation and comparison between organic solvents (sulfolane and DMSO) used for benzene extraction. *International Journal of Chemical and Petroleum Sciences*. 2013;**2**(1):10-19
- [76] Gbashi S. Pressurized hot water extraction (PHWE) and chemometric fingerprinting of phytochemicals from *Bidens pilosa* [thesis]. South Africa: University of Johannesburg, South Africa; 2016
- [77] Zabe N, Ayalew K, Powers S. Ethanol extraction method for a rapid test for aflatoxin in corn. *Food Additives & Contaminants*. 2008;**1001**:297-305
- [78] Patterson DSP, Roberts BA. Mycotoxins in animal feedstuffs: Sensitive thin layer chromatographic detection of aflatoxin, ochratoxin A, sterigmatocystin, zearalenone and T-2 toxin. *Journal Association of Official Analytical Chemists*. 1979;**62**:1265-1267
- [79] Tabaraki R, Nateghi A. Optimization of ultrasonic-assisted extraction of natural antioxidants from rice bran using response surface methodology. *Ultrasonics Sonochemistry*. 2011;**18**(6):1279-1286
- [80] Limsuwan S. Development, evaluation and application of methods for mycotoxin analysis [thesis]. Germany. Georg-August-University Göttingen; 2011
- [81] Gam L, Tham S, Latiff A. Immunoaffinity extraction and tandem mass spectrometric analysis of human chorionic gonadotropin in doping analysis. *Journal of Chromatography B*. 2003;**792**(2):187-196
- [82] Bardsley J, Oliver M. Determination of multiple mycotoxins in grain using a QuEChERS sample preparation approach and LC-MS/MS detection. *Appl Note* 21121. 2015;Thermo Fisher Scientific, Runcorn, UK:1-7
- [83] Sirhan AY, Tan GH, Al-Shunnaq A, Abdulra'uf L, Wong RCS. QuEChERS-HPLC method for aflatoxin detection of domestic and imported food in Jordan. *Journal of Liquid Chromatography & Related Technologies*. 2014;**37**(3):321-342
- [84] Brown JD. Development of a method for the simultaneous detection of mycotoxins in corn using supercritical fluid extraction (SFE), high performance liquid chromatography (HPLC), and electrospray-atmospheric pressure ionization/mass spectrometry (ES-API/MS) for extraction, separation, and identification [thesis]. USA: The University of Iowa; 1999
- [85] Kong WJ, Liu SY, Qiu F, Xiao XH, Yang MH. Simultaneous multi-mycotoxin determination in nutmeg by ultrasound-assisted solid-liquid extraction and immunoaffinity column clean-up coupled with liquid chromatography and on-line post-column photochemical derivatization-fluorescence detection. *Analyst*. 2013;**138**(9):2729-2739

- [86] Turner NW, Bramhmbhatt H, Szabo-Vezse M, Poma A, Coker R, Piletsky SA. Analytical methods for determination of mycotoxins: An update (2009-2014). *Analytica Chimica Acta*. 2015;**901**:12-33
- [87] Anfossi L, Baggiani C, Giovannoli C, Giraudi G. Mycotoxins in food and feed: Extraction, analysis and emerging technologies for rapid and on-field detection. *Recent Patents Food, Nutrition & Agriculture*. 2010;**2**(2):140-153
- [88] Zhang A, Sun H, Wang P, Han Y, Wang X. Modern analytical techniques in metabolomics analysis. *Analyst*. 2012;**137**(2):293-300
- [89] Hussain I. Aflatoxin measurement and analysis. In: Torres-Pacheco I, editor. *Aflatoxins – Detection, Measurement and Control*. 1st ed. Rijeka, Croatia: InTech; 2011. pp. 129-146
- [90] Liang X, Fan Q. Application of sub-critical water extraction in pharmaceutical industry. *Journal of Chemical Engineering and Materials Science*. 2013;**1**:1-6
- [91] Wilson ID, Davis P, Ruane RJ. Supercritical fluid chromatography and extraction of pharmaceuticals. In: Dean JR, editor. *Applications of Supercritical Fluids in Industrial Analysis*. Dordrecht, Netherlands: Springer Netherlands; 1993. p. 74-103
- [92] Ozel MZ, Gogus F, Lewis AC. Subcritical water extraction of essential oils from *Thymbra spicata*. *Food Chemistry*. 2003;**82**(3):381-386
- [93] Gupta A, Naraniwal M, Kothari V. Modern extraction methods for preparation of bioactive plant extracts. *International Journal of Applied and Natural Sciences*. 2012;**1**(1):8-26
- [94] Ibañez E, Kubátová A, Señoráns FJ, Cavero S, Reglero G, Hawthorne SB. Subcritical water extraction of antioxidant compounds from rosemary plants. *Journal of Agricultural and Food Chemistry*. 2003;**51**(2):375-382
- [95] Gbashi S, Njobeh P, Steenkamp P, Tutu H, Madala N. The effect of temperature and methanol-water mixture on pressurized hot water extraction (PHWE) of anti-HIV analogues from *Bidens pilosa*. *Chemistry Central Journal*. 2016;**10**(37):1-12
- [96] Shaddel R, Maskooki A, Haddad-Khodaparast MH, Azadmard-Damirchi S, Mohamadi M, Fathi-Achachlouei B. Optimization of extraction process of bioactive compounds from Bene hull using subcritical water. *Food Science and Biotechnology*. 2014;**23**(5):1459-1468
- [97] Alupului A, Călinescu I, Lavric V. Microwave extraction of active principles from medicinal plants. *UPB Scientific Bulletin, Series B: Chemistry and Materials Science*. 2012;**74**(2):129-142
- [98] Cabane B, Vuilleumier R. The physics of liquid water. *Comptes Rendus Geoscience*. 2005;**337**:1-23
- [99] Yang B, Wyman CE. Effect of xylan and lignin removal by batch and flow through pretreatment on the enzymatic digestibility of corn stover Cellulose. *Biotechnology and Bioengineering*. 2004;**86**(1):88-95

- [100] Asl AH, Khajenoori M. Subcritical water extraction. In: Nakajima H, editor. Mass Transfer – Advances in Sustainable Energy and Environment Oriented Numerical Modeling. Rijeka, Croatia: InTech; 2013. pp. 459-487
- [101] Vergara-Salinas JR, Bulnes P, Zúñiga MC, Pérez-Jiménez J, Torres JL, Mateos-Martín ML, et al. Effect of pressurized hot water extraction on antioxidants from grape pomace before and after enological fermentation. Journal of Agricultural and Food Chemistry. 2013;**61**(28):6929-6936
- [102] Khoza BS, Dubery IA, Byth-Illing HA, Steenkamp PA, Chimuka L, Madala NE. Optimization of pressurized hot water extraction of flavonoids from *Momordica foetida* using UHPLC-qTOF-MS and multivariate chemometric approaches. Food Analytical Methods. 2015;**2015**:1-10
- [103] Khajenoori M, Asl AH, Hormozi F, Eikani MH, Bidgoli HN. Subcritical water extraction of essential oils from *Zataria multiflora* Boiss. Journal of Food Process Engineering. 2009;**32**(6):804-816
- [104] Hawthorne SB, Yang Y, Miller DJ. Extraction of organic pollutants from environmental solids with sub- and supercritical water. Analytical Chemistry. 1994;**66**:2912-2920
- [105] Deng CH, Li N, Zhang XM. Rapid determination of essential oil in "*Acorus tatarinowii* Chott" by pressurized hot water extraction followed by solid-phase microextraction and gas chromatography-mass spectrometry. Journal of Chromatography A. 2004;**1059**(1):149-155
- [106] Shalmashi A, Abedi M, Golmohammad F, Eikani MH. Isolation of caffeine from tea waste using subcritical water extraction. Journal of Food Process Engineering. 2010;**33**(4):701-711
- [107] Ramos L, Kristenson EM, Brinkman UAT. Current use of pressurized liquid extraction and subcritical water extraction in environmental analysis. Journal of Chromatography A. 2002;**975**:3-29
- [108] Carr AG, Mammucari R, Foster NR. A review of subcritical water as a solvent and its utilization for the processing of hydrophobic organic compounds. Chemical Engineering Journal. 2011;**172**(1):1-17
- [109] Ong ES, Cheong JSH, Goh D. Pressurized hot water extraction of bioactive or marker compounds in botanicals and medicinal plant materials. Journal of Chromatography A. 2006;**1112**(1-2):92-102
- [110] Williams HD, Trevaskis NL, Charman SA, Shanker RM, Charman WN, Pouton CW, et al. Strategies to address low drug solubility in discovery and development. Pharmacological Reviews. 2013;**65**:315-449
- [111] Gbashi S, Njobeh P, Steenkamp P, Madala N. Pressurized hot water extraction and chemometric fingerprinting of flavonoids from *Bidens pilosa* by UPLC-tandem mass spectrometry. CYTA - J Food. 2016;**15**(2):171-180.

- [112] Curren MSS, King JW. Solubility of Triazine pesticides in pure and modified subcritical water. *Analytical Chemistry*. 2001;**73**:740-745
- [113] Plaza M, Turner C. Pressurized hot water extraction of bioactives. *TrAC Trends in Analytical Chemistry*. 2015;**71**:39-54
- [114] Shitu A, Izhar S, Tahir TM. Sub-critical water as a green solvent for production of valuable materials from agricultural waste biomass: A review of recent work. *Global Journal of Environmental Science and Management*. 2015;**1**(3):255-264
- [115] Chemat F, Vian M, Cravotto G. Green extraction of natural products: Concept and principles. *International Journal of Molecular Sciences*. 2012;**13**:8615-8627
- [116] Khoza BS, Gbashi S, Steenkamp PA, Njobeh PB, Madala NE. Identification of hydroxylcinnamoyl tartaric acid esters in *Bidens pilosa* by UPLC-tandem mass spectrometry. *South African Journal of Botany*. 2016;**2016**:96-100
- [117] Khoza BS, Chimuka L, Mukwevho E, Mukwevho E, Steenkamp PA, Madala NE. The effect of temperature on pressurized hot water extraction of pharmacologically-important metabolites as analyzed by UPLC-qTOF-MS and PCA. *Evidence-Based Complementary and Alternative Medicine*. 2014;**2014**:1-9. (Article ID 914759)
- [118] Moreno E, Reza J, Trejo A. Extraction of polycyclic aromatic hydrocarbons from soil using water under subcritical conditions. *Polycyclic Aromatic Compounds*. 2007;**27**(4):239-260
- [119] Duy L, Sephton MA, Watson JS. Subcritical water extraction of organic matter from sedimentary rocks. *Analytica Chimica Acta*. 2015;**879**:48-57
- [120] Fattah RA, Mostafa NA, Mahmoud MS, Abdelmoez W. Recovery of oil and free fatty acids from spent bleaching earth using sub-critical water technology supported with kinetic and thermodynamic study. *Advances in Bioscience and Biotechnology*. 2014;**5**:261-272
- [121] Ho CHL, Cacace JE, Mazza G. Extraction of lignans, proteins and carbohydrates from flaxseed meal with pressurized low polarity water. *Leb Technology*. 2007;**40**:1637-1647
- [122] García-Marino M, Rivas-Gonzalo J, Ibáñez E, García-Marino C. Recovery of catechins and proanthocyanidins from winery by-products using subcritical water extraction. *Analytica Chimica Acta*. 2006;**563**(1-2):44-50
- [123] Hartonen K, Parshintsev J, Sandberg K, Bergelin E, Nisula L, Riekkola ML. Isolation of flavonoids from aspen knotwood by pressurized hot water extraction and comparison with other extraction techniques. *Talanta*. 2007;**74**(1):32-38
- [124] Herrero M, Ibanez E, Senorans J, Cifuentes A. Pressurized liquid extracts from *Spirulina platensis* microalga: Determination of their antioxidant activity and preliminary analysis by micellar electrokinetic chromatography. *Journal of Chromatography A*. 2004;**1047**:195-203

- [125] Gbashi S, Madala N, Adebo OA, Njobeh PB. Development and validation of a pressurized hot water extraction method for the extraction of aflatoxin B₁ in maize. In: 7th Euro Global Summit on Clinical Microbiology and Mycotoxins February 27-28, 2017 Amsterdam, Netherlands. Amsterdam, Netherlands; 2017. p. Full text data not published.
- [126] Scudamore KA. Mycotoxins. In: Gilbert J, Senyuva H, editors. Bioactive Compounds in Foods. West Sussex, United Kingdom: Blackwell Publishing Ltd; 2008. pp. 134-162
- [127] Curren MSS, King JW. Ethanol-modified subcritical water extraction combined with solid-phase microextraction for determining Atrazine in beef kidney. Journal of Agricultural and Food Chemistry. 2001;**49**:2175-2180