

Plant Metabolites in Plant Defense Against Pathogens

*Xóchitl S. Ramírez-Gómez, Sandra N. Jiménez-García,
Vicente Beltrán Campos and Ma. Lourdes García Campos*

Abstract

Medicinal plants are widely used worldwide to treat various diseases. Its widespread use is due in part to the cultural acceptance of traditional medicine in different regions of the world, as well as its effectiveness in treating various diseases. Many of its active substances or secondary metabolites are formed to a response of various situations that generate stress in their habitat, such as sudden changes in environmental temperature, humidity, rain, drought, and infections by phytopathogens (fungi, bacteria, viruses, nematodes, protozoa). The production of these secondary metabolites is a mechanism of defense of plants. In this context, the objective of this chapter is to study the secondary metabolites of medicinal plants that could have a promising application in the control of different phytopathogens in crops of agricultural and economic interest.

Keywords: medicinal plants, phytopathogens, secondary metabolites, pesticides, biotic and abiotic elicitors

1. Introduction

Phytopathogens generally attack plants during their growth, causing alterations in their cellular metabolism and/or interfering with the absorption of nutrients [1]. The crops of cereals, vegetables, and fruits are affected by these organisms during harvest and postharvest [2]. However, one of the main control measures to eradicate phytopathogens is the use of pesticides. Although they are effective, easy to access, and easy to use, they have several disadvantages, generate resistance, and are considered toxic substances, not only for bacteria, fungi, viruses, protozoa, and nematodes but also for the humans, animals, and the environment [3, 4]. In this context, the pesticides can induce acute and chronic toxicity, to persist in the environment and pollute soil and water. So, they are easily incorporated into the food chain, bioaccumulation, and biomagnification [5]. Regarding their toxicity mechanisms, it has been described that they can act as endocrine disruptors and as reactive species that generate oxidative stress in the cell [6–9].

On the other hand, the study of medicinal plants as possible natural sources of obtaining active compound (secondary metabolites) against phytopathogens has gained increasing interest in recent years, due to several aspects, mainly that they are obtained from a natural source through the production or synthesis of secondary metabolites considered as nontoxic such as phenols, flavonoids, terpenes, alkaloids, etc. [10–13]. Another advantage is that phytopathogens still do not develop resistance to the antifungal, antimicrobial, and nematocide effect of the phytochemical

compounds produced by some medicinal plants. When carrying out an exhaustive search in the literature, it was found that the potential use of the secondary metabolites obtained from medicinal plant extracts is fungicide [14–16]. Most of the research in this area focuses on evaluating the effects of these active compounds on fungi such as *Fusarium*, maybe because it is one of the main phytopathogens that cause economic losses mainly in cereal crops and health problems by their aflatoxins [17, 18]. This chapter shows an overview of the recent research on this topic, emphasizing the effect of biotic and abiotic elicitors on the secondary metabolite production, as well as a brief description of the scientific name of the plant, metabolites with antifungal and antibacterial effect, and their limitations and perspectives of its use in the biological control of phytopathogens.

2. Pesticides in the control of phytopathogens

In the market, there are a variety of pesticides that are used alone or in combination to eradicate, control, or prevent pests [4]. Pesticides can be classified according to the chemical group to which they belong, to their selectivity toward a certain phytopathogen, its mechanism of action, and its use or application. However, the most widely used for their effectiveness and a broad spectrum of activity against various pests and diseases in plants are insecticides, herbicides, and fungicides [4, 19].

Pesticides used in agriculture mainly contaminate the soil by direct application and water by leaching, and it is very easy for them to be present either in trace quantities or high in food and to enter the food chain, which facilitates its accumulation and biomagnification [5, 20]. In general pesticides are considered dangerous substances for living beings since they can produce acute or chronic toxicity; however the magnitude of the poisoning depends on several aspects to be considered such as the physicochemical characteristics of the pesticide, the concentration, the exposure time, the route of entry to organisms, their toxicodynamics and toxicokinetics (absorption, distribution, half-life, metabolism, and elimination), as well as the use of mixtures of different pesticides, the components of their formulation, and the general state of health of the individual [21, 22]. All these aspects influence that pesticides represent a risk or danger for those who use them in the fields of cultivation, as well as for those who consume foods that contain substances in trace quantities in prolonged consumption.

Regarding its toxicity, it has been described that pesticides act as endocrine disruptors and generators of free radicals and enzymatic inhibitors [8, 9]. Unfortunately, the cellular targets to which most of these pesticides are directed coincide with cellular targets that are also present in man, such as the case of the mechanisms of action of organophosphorus insecticides, which inhibit the activity of acetylcholinesterase enzyme present in different insects; unfortunately man and other mammals also have acetylcholinesterase, so their toxicity is not selective toward the pests that they wish to control, but they also affect man, and depending on the magnitude of the poisoning, they can cause death [19–22]. However, until today an ideal pesticide does not exist, and the correct use of herbicides, fungicides, insecticides, etc. has many benefits to control plagues and increase the yield of the crops [19].

3. Secondary metabolites of medicinal plants as biological control of phytopathogens

There are several methods of biological control against phytopathogens. The use of extracts of medicinal plants to eradicate diseases in crops caused mainly by

viruses, bacteria, and fungi is one of them [23]. The above makes sense if we analyze the fact that plants have mechanisms to protect themselves from both biotic and abiotic stress agents. That is, if the phytopathogens (biotic agents) are attacking the plants, why not think what the plant does to defend itself?

In this context, it is interesting to analyze the secondary metabolism of plants know which phytochemical substances are produced and what biological activity they present.

3.1 The bioactive potential of secondary metabolites derived from the medicinal plant

Plants are formed by a primary metabolism that is responsible for the physiological processes and development of the plant, such as lipids, carbohydrates, and proteins [23]. The secondary metabolism is not essential in the basic processes of plants. However, these bioactive compounds play an important role in the defense of plants, and these secondary metabolites can be classified as phenolic compounds, carotenoids, terpenes, alkaloids, and sulfur compounds, among others, as shown in Table 1 [24].

Phenolic compounds are aromatic substances formed during the passage of the shikimic acid pathway or mainly the mevalonic pathway. These can be divided into insoluble compounds such as condensed tannins, lignins, and hydroxamic acids bound to the cell walls, and soluble compounds are phenolic acids, flavonoids, and kinases [25]. Carotenoids are lipophilic molecules and are found in plants giving orange tones. The importance of these compounds is the intervention they have in photosynthesis, and they also protect the photosynthetic apparatus from excess

Classification	Types	Example
Terpenes	Monoterpene	Geraniol
	Sesquiterpenes	Humulene
	Diterpenes	Cafestol
	Sesterpenes	Geranylarsol
	Triterpenes	Squalene
	Sesquaterpenes	Ferrugicadiol
	Tetraterpenes	Lycopenes
	Polyterpenes	Gutta-percha
Phenolics	Coumarin	Hydroxycoumarins
	Furano-coumarins	Psoralin
	Lignin	Resveratrol
	Flavonoids	Quercitin
	Isoflavonoids	Genistein
	Tanins	Tanins acid
N Containing Compounds	Alkaloids	Cocaine
	Cyanogenic glucosides	Dhurrin
	Non-Protein amino acids	Canavanin
S Containing compounds	Glutathione	
	Glucosinolate	β -D-Glucopyrinose
	Thionins	
	Defensins	
	Allinin	

Table 1.
Types of plant secondary metabolites.

energy [25]. The carotenoid contents in plants are affected by various factors, such as plant development, stress conditions, postharvest conditions, or cooking treatments, but the interest of these compounds has been increasing due to their potential antioxidant activity [26]. Terpenes are lipid-soluble compounds that include one- or more five-carbon isoprene units, which are synthesized by all organisms through two pathways, mevalonate and deoxy-D-xylulose [27]. Terpenoids are classified according to the number of isoprene units they contain; terpenes and terpenoids are basic constituents of many types of plant essential oils [28]. Alkaloids are bioactive compounds that generally contain nitrogen derived from an amino acid of great importance because it has physiological and medicinal properties, for example, caffeine, nicotine, morphine, atropine, and quinine [29].

Now well, all these compounds mentioned above help the plants to develop complex defense systems against different types of stress for the survival or the systematic forces in their metabolism for resistance against pests and diseases. Stress provoked in the plant involves several signaling response pathways for pathogens and insects, and some of these response pathways are induced by the microorganisms themselves. Also, the plants have specific recognition and signaling systems allowing them to detect the pathogens and initiate an effective defense response [30, 31]. The defence system broadest have the plants against pathogens are the phenolic compounds (phenylpropanoids and flavonoids). These substances have different mechanisms of action they can dissociate the ions of the phenolic hydroxyl and forming phenolates, ionic and hydrogen bonds with peptides and proteins causing a high astringency and protein denaturation. In the other hand, they interfere with the pathogen's cell signalling compounds and affect their physiological activities through enzymatic inhibition, DNA alkylation and altering their reproductive system [31]. The compounds with allelopathic effects affect positively or negatively on the ecosystem's structure to remove or eliminate microorganisms from the plants. Some phenolic compounds are allelochemicals that have been shown to have an activity as antibiotics, antifungals, and antipredator [31]. Phenolic acids, such as benzoic, hydroxybenzoic, vanillic, and caffeic, have antimicrobial and antifungal properties produced by the inhibition of enzymes. Caffeic, chlorogenic, sinapic, ferulic, and p-coumaric acids have antioxidant activity by the inhibition of oxidation of lipids and the elimination of reactive oxygen species. These effects are important to the plant defense [32].

3.2 Improving production of plant secondary metabolites through biotic and abiotic stresses

Classification of secondary metabolites related to the defense of plants is commonly used in the form of synthesis and accumulation of phytochemicals with interaction effect of the pathogenic plant against plant insect, virus, fungi, and antibacterial compounds. For example, phytoalexins are produced very quickly after infection of a pathogen producing toxicity to an ambiguous environment of fungi or bacteria [33, 34].

Phenylpropanoids and flavonoids have hydroxyl groups that contain phenolic compounds, which dissociate into phenolate ions, and the phenolic hydroxyl groups form ionic bonds and hydrogen bonds with peptides and protons, producing a high astringency and denaturation that thus show an antifungal effect acting together with cellular signaling compounds and physiological activities or acting on the parts of the pathogen, reproductive system, enzymatic inhibition, etc. [35]. The properties of the proteins change with any change in protein conformation, for example, by changing the three-dimensional structure forming covalent bonds with SH, OH or free amino groups there is inactivation or protein function loss. When polyphenols of the plants bind to some proteins of phytopathogens are less toxic for them

but can protect the plant of abiotic elicitors [36]. On the other hand, phytoalexins are induced against the attack of microbes and insects activated by β -glucosidase by the release of biocidal aglycones [37]. In the same way act the benzoxazinoids (BX), these phytochemical compounds are produced and released by tissue damage and hydrolysis by β -glucosidase and act as insect repellents too [38].

At present, several biotechnological strategies have been used to increase the productivity of secondary metabolites, using different inducers of secondary metabolites such as at the cellular, organic, and plant levels, as well as the most effective methods to improve the synthesis of these secondary metabolites in endemic and medicinal plants [39]. These secondary metabolites accumulate in plants when they are prone to various stress types, inducers, or signal molecules. Thus, there are different modulating factors of secondary metabolites, as well as microbial, physical, or chemical effects such as abiotic or biotic elicitors, inducing the biosynthesis of specific compound that plays an important role in the adaptations of plants to stress conditions, and these phenomena cause a greater synthesis and accumulation of secondary metabolites [40]. In **Table 2** the authors focus on the abiotic elicitors that are substances of biological origin such as proteins and carbohydrates that are initiator compounds or coupling responses at the cellular level activating several enzymes or signaling canals. There are also microorganisms and chemical compounds with elicitor effect that stress the plant and produce the expression of a greater amount of metabolites or new metabolites which cause physiological changes in the plant against pathogens. As shown in **Table 2**, glycoprotein-type proteins produce phytoalexins that have been used to identify ion channels in cell membranes and thus transfer signals by external stimuli, as demonstrated by Alami [41] where the *Plantanus x acerifolia* cultures were applied to an inducer of *Ceratocystis fimbriata* f. sp. These, in turn, induced the synthesis of phytoalexins (hydroxycoumarin, scopoletin, and umbelliferone), and upon isolating the glycoprotein produced the synthesis of coumarin by 80%. On the other hand, oligogalacturonic acids are found in the cell wall of the plant inducing the biosynthesis of phytoalexins, whereas chitin is found in the cell wall of fungi, generating signaling factors in plants such as *Hypericum perforatum* production stress in the plant and increasing the production of phenolic compounds for their defense against pathogens [39, 42]. Rhizobacteria function as modelers of secondary metabolites with pharmacological activity. Rhizobacteria colonize the rhizospheres of the plants and improve the growth of the plant, being localized in the bark or root nodules acting as inducers of the enzymes that participate in the metabolic pathways of bioactive compounds and jasmonic acid biosynthesis; these act as signal transducers [43, 44]. Other signal inducers are the mycorrhizal fungi that help the plant to absorb more water and show defense against other pathogens such as fungi, bacteria, or parasites that affect the roots of the plant. These mycorrhizal fungi produce secondary metabolites such as phenolic compounds and alkaloids, among others [45–48]. Elicitors such as salicylic acid, jasmonic acid, hydrogen peroxide, chitosan, etc. act as plant hormones in the expression of genes interacting as target signaling causing a physiological response in the plant which increases the production of phenolic compounds, vitamin C, carotenoids, or defense stimuli against pathogens; there are also synergistic effects between salicylic acid and jasmonic acid providing resistance against pathogens by the induction of the octadecanoic acid pathway [49–52, 53].

On the other hand, **Table 3** shows some research that has the influence of different abiotic elicitors that are considered substance and that are not of biological origin such as salt, drought, light or heavy metals, and temperature, among others. **Table 3** shows different perspectives of research on medicinal or aromatic plants in hydroponic crops, outdoors, and the application of elicitors in different stages of growth or postharvest. For example, heavy metals such as Al^{3+} , Cr^{3+} , Co^{2+} , Ni^{2+} ,

	Classification	Elicitor/ Species	Compounds	Plant Species	References
	Proteins	Glycoprotein	Coumarin	<i>Plantanus acerifolia</i>	[41]
	Carbohydrates	Chitosan	Phytoalexin	<i>Nicotiana tabacum</i> <i>Eschscholzia californica</i>	[54]
		Ologogalacturonic acid	Saponin	<i>Panax ginseng</i>	[55]
		Ologogalacturonic acid	Trans-resveratrol Viniferins	<i>Vitis vinifera</i>	[56]
		Chitin Pectin	Hipericin Pseudohypericin	<i>Hipericum perforatum</i>	[39]
		Chitin	Phenilpropanoid Naphtodianthrone	<i>Hipericum perforatum</i>	[42]
		Chitosan	Curcumin	<i>Curcuma longa</i> L.	[40]
		Chitosan	Withaferin	<i>Wiyhania somnifera</i>	[57]
	Plant growth promoting rhizobacteria (PGPR)	<i>Pseudomonas putida</i> <i>Pseudomonas fluorescens</i>	Cis-Thujone Camphor 1,8-cineole	<i>Salvia officinalis</i>	[58]
		<i>Pseudomonas fluorescens</i> <i>Bacillus subtilis</i> <i>Azospirillum brasilense</i> <i>Bacillus solanum</i>	γ- terpinene Trans-sabinene hydrate Cis- sabinene hydrate Thymol	<i>Origanum majoricum</i>	[59]
		<i>Glomus aggregatum</i> <i>Bacillus coagulans</i> <i>Trichoderma harzianum</i>	Phenols Tannins Flavonoids Saponins Alkaloids	<i>Solanum viarum</i>	[43]
		<i>Bacillus polymyxa</i> <i>Pseudomonas putida</i> <i>Azotobacter chroococcum</i> <i>Glomus intraradices</i>	Stevioside	<i>Stevia rebaudiana</i>	[42]
		<i>Bacillus subtilis</i>	Phenolic compounds (gallic, cinnamic, ferulic acid)	<i>Ocimum basilicum</i>	[60]
		<i>Pseudomonas flurescens</i> <i>Azopirillum brasilense</i>	Monoterpenes Phenolic compounds	<i>Tagetes minuta</i>	[61]
	Fungus	<i>Fusarium oxysporum</i> <i>Botrytis cinerea</i>	Phenylpropanoid Naphtodianthrone	<i>Hypericum perforatum</i>	[42]
		<i>Phytophthora megaspema</i> <i>Rhizopus arrhizus</i>	Alkaloids (tropane)	<i>Datura stramonium</i>	[45]
		<i>Aspergillus niger</i>	Rosmarinic acid	<i>Ocimum basilium</i>	[46]
		<i>Rhizostonia solani</i>	Sesquiterpenes	<i>Hyoscyamus muticus</i>	[47]
	Phytohormones	Absciscic acid Gibberellin	Phenolic acids Tanshinones	<i>Salvia miltiorrhiza</i>	[48, 62]
		Ethylene	Anthocyanin	<i>Fragaria ananassa</i>	[63]
		Gibberellic acid	Caffeic acid derivatives	<i>Echinacea pupurea</i>	[64]
	Elicitors	Salicylic acid	Stilbene Tanshinones Monoterpene Gymnemic acid	<i>Vitis vinifera</i> <i>Salvia miltiorrhiza</i> <i>Houttuynia cordata</i> <i>Gymnema sylvestre</i>	[49-52]
		Methyl salicilate	Withaferin A	<i>Withania somnifera</i>	[57]
		Methyl Jasmonate	Anthocynin Stilbene Trans-resveratrol Rosmarinic acid Saponin Soyasaponin	<i>Vitis vinifera</i> <i>Mentha piperita</i> <i>Glycyrrhiza globa</i>	[56, 53, 54]
		Jasmonic acid	Plumgagin	<i>Plumbago indica</i> <i>Plumbago rosea</i>	[53]

Table 2.
Effect of biotic elicitor on the production of various secondary metabolites in plants.

Classification	Elicitor/ Species	Compounds	Plant Species	References
Heavy metals	Al ³⁺ , Cr ³⁺ , Co ²⁺ , Ni ²⁺ , Cu ²⁺ , Zn ²⁺ , Cd ²⁺	Sesquiterpenoid Lubimin 3-hydroxylubimin	<i>Datura stramonium</i>	[70, 71]
	Ag ⁺	Atropine	<i>Salvia castanea</i> <i>Datura metel</i>	[72]
	Cd ²⁺ , Co ²⁺ , Ag ⁺	Resveratrol	<i>Vitis vinifera</i>	[73]
	Ca ²⁺ , Mg ²⁺ , Mn ²⁺ , n ²⁺ , Cu ²⁺ , Fe ²⁺ , Co ²⁺	Betalain	<i>Beta vulgaris</i>	[74]
Temperature	High temperature	Hypericin Hyperforin Ginsenoside	<i>Hypericum perforatum</i> <i>Panax quinquefolius</i>	[65, 75]
	Low temperature	Melatonin Anthocyanin	<i>Rhodiola crenulata</i> <i>Melastoma malabathricum</i>	[66, 78]
Light	UV-B light	Vinblastine Vincristine	<i>Catharanthus roseus</i>	[66]
	UV-C irradiation	Flavonoid Stilbene	<i>Catharanthus roseus</i>	[78]
	Light radiation	Anthocyanins	<i>Melastoma malabathric</i>	[77]
Salinity	Salinity	Sorbitol Jasmonic acid Flavonoids Anthocyanin GABA Phenylpropeno	<i>Lycopersicon esculentum</i> <i>Hordeum vulgare</i> <i>Brivillea ilicifolia</i> <i>G. Arenaria</i> <i>Sesamum indicum</i>	[67-79]
Drought	Drought stress	Rosmarinic Ursolic Oleanolic acid	<i>Prunella vulgaris</i>	[80]
	Water Stress	Salvianolic acid	<i>Salvia miltiorrhiza</i>	[68]
	Drought	Steviol glycosides	<i>Stevia rebaudiana</i>	[69]
	Water Osmotic imbalance	Hypericin	<i>Hypericum perforatum</i>	[81]

Table 3.
Effect of abiotic elicitor on the production of various secondary metabolites in plants.

Cu²⁺, Zn²⁺, and Cd²⁺, among others, are considered high toxicity compounds depending on the concentrations applied in the sprinkler system or because they are used as biocontrol since they alter the production of metabolites in plants. Similarly, Zobayed [65] demonstrated that the temperature in high concentrations in *Panax quinquefolius* improves the senescence of the leaves and produces a greater quantity of bioactive compounds in the root of the plant. So the investigations using high or low temperatures demonstrate the production of secondary metabolites, but the temperatures that have been investigated the most are the low producing physiological changes in the plant, increasing the lignification by the production of suberin in the cell wall and the metabolites such as sorbitol, raffinose, proline, melatonin, anthocyanins, etc. However, light by means of ultraviolet radiations generates the production of essential oils and phenolic compounds and decreases the production of toxic compounds in some plants [66]. On the other hand, salinity and drought produce death leading to cellular dehydration or osmotic stress and in certain concentrations can reduce the growth or development of plants but alter many physiological and metabolic processes that stimulate the production of polyphenolic compounds, anthocyanins, terpenes, and alkaloids, among others. Salinity can be produced in plants by ionic or osmotic means and drought by environmental or intentional changes due to water deficit which are always accompanied by temperature or solar radiation [67–69]. Then we can say that the biotic and abiotic factors are modular secondary metabolites influencing the metabolic level and the

production of secondary metabolites. Therefore, the current research focuses on the use of elicitors, for the regulation of metabolic pathways, and target signaling in genes that influence the overproduction of secondary metabolites using various applications but taking care of the production performance of fruits, vegetables, or different plants.

Recent studies focused on evaluating the secondary metabolites of medicinal plants that are active against phytopathogens show that the potential use that these compounds can have in the future is for the control of phytopathogenic fungi, mainly against different species of *Fusarium* [14–16]. In this regard, the most active compounds have been found mainly in the essential oil obtained from the aerial parts of various medicinal plants, which suggests that the bioactive compounds are liposoluble; this may explain why they are active mainly against fungi, because the cell wall of these specimens are composed mainly of ergosterol, the active liposoluble compounds present in the essential oil to easily cross the cell wall of the fungus and in the interior act on their cell target, or they can alter the permeability of the wall of the fungus [82]. It can cause rupture and lysis of the fungal cell; however, it is necessary to study the toxicodynamics of these substances in order for them to know how to act in the fungi cell. On the other hand, the antifungal activity has been evaluated *in vitro*, by the agar diffusion and microdilution method; in general terms the range of the evaluated IC₅₀ varies in a range that goes from 0.0035 to 8 mg/ml of the extract. It is important to mention that one of the main limitations of these studies is that this activity has only been evaluated at the laboratory level [83–85]. **Table 4** shows different types of extracts made with medicinal plants, and their biological activity reported *in vitro* tests at the laboratory level.

Finally, in the realization of a retrospective of the secondary metabolite modulating factors in our workgroup, Garcia-Mier [95] demonstrated that the use of mixtures of elicitors such as jasmonic acid, hydrogen peroxide, and chitosan in different concentrations applied in various stages of plant development of the sweet bell red pepper and in different stages of ripening of the fruit has a positive effect on the increase of polyphenolic and carotenoid compounds, where the results showed that the maturation stage of 95% produces a greater quantity of bioactive compounds. On the other hand, Vargas-Hernández [96] demonstrated that the foliar application of hydrogen peroxide in *Capsicum chinense* Jacq. has an effect on the antimicrobial activity, where the different concentrations of hydrogen peroxide potentiated the production of secondary metabolites such as flavonoids, capsaicin, and dihydrocapsaicin, where these metabolites had an effect on microorganisms such as *Staphylococcus aureus*, *Escherichia coli*, *Streptococcus mutant*, *Salmonella thompson*, *Listeria monocytogenes*, *Streptococcus faecalis*, and *Candida albicans*, and the results showed that the application of hydrogen peroxide increases the inhibitory effect against pathogenic microorganisms, showing greater activity against *S. aureus*, *S. Thompson*, and *C. albicans* in the jaguar variety, while the variety Chichen-Itza was more potent against *E. faecalis* and *E. coli*. Also, Zunun-Pérez [97] evaluated the effect of modulating factors of secondary metabolites by spray application that is performed in *Capsicum annum* L. in weekly applications and 1 day before collection with elicitors such as hydrogen peroxide, salicylic acid, and oligosaccharide of xyloglucan on capsate concentration and the expression of genes such as phenylalanine ammonia-lyase, aminotransferase, capsaicin synthase, and β -keto acyl synthase where the results showed that hydrogen peroxide in weekly applications significantly increases capsate concentrations and gene expression and the yields of the production of the plant are not affected by the application of these elicitors.

Medicinal Plant	Type of Extract	Phytochemistry Compounds	Biological Activity Against	Reference
<i>Acacia farnesiana</i>	Hydroalcoholic extract	Tannins Flavonoids Saponins Alkaloids Triterpenes Quinones	<i>Pyricularia grisea</i> <i>Phytophthora parasitica</i> var. <i>nicotianae</i>	[86]
<i>Artemisia herba alba</i>	Aqueous extract	Tannins Flavonoids Saponins Steroids Alkaloids	<i>Fusarium graminearum</i> <i>Fusarium sporotrichioides</i>	[87]
<i>Asphodelus tenuifolius</i>	Aqueous extract	Tannins Flavonoids Steroids Alkaloids	<i>Fusarium graminearum</i> <i>Fusarium sporotrichioides</i>	[87]
<i>Bauhinia galpinii</i>	Methanol:Dichloromethane extract	Phenols Flavonoids	<i>Aspergillus parasiticus</i> <i>Aspergillus flavus</i>	[88]
<i>Breonadia salicina</i>	Hexane and Methanol extracts	No data	<i>Penicillium janthinellum</i> <i>Trichoderma harzianum</i> <i>Fusarium oxysporum</i>	[89]
<i>Bucida buceras</i>	Hexane, Dichloromethane, Acetone and Methanol extracts	No data	<i>Penicillium janthinellum</i> <i>Penicillium expansum</i> <i>Trichoderma harzianum</i> <i>Fusarium oxysporum</i>	[89]
<i>Carpobrotus eludis</i>	Methanol:Dichloromethane extract	Phenols Flavonoids	<i>Aspergillus parasiticus</i>	[88]
<i>Cotula cinerea</i>	Aqueous extract	Tannins Flavonoids Saponins Alkaloids	<i>Fusarium graminearum</i> <i>Fusarium sporotrichioides</i>	[87]
<i>Euphorbia guyoniana</i>	Aqueous extract	Tannins Flavonoids Saponins Steroids Alkaloids Anthocyanins	<i>Fusarium graminearum</i> <i>Fusarium sporotrichioides</i>	[87]
<i>Harpephyllum caffrum</i>	Acetone and Methanol extracts	No data	<i>Penicillium janthinellu</i> , <i>Trichoderma harzianum</i> <i>Fusarium oxysporum</i>	[88, 89]
	Methanol:Dichloromethane extract	Phenols Flavonoids	<i>Aspergillus parasiticus</i> <i>Aspergillus ochraceous</i>	
<i>Maesa lanceolata</i>	Methanol:Dichloromethane extract	Phenols Flavonoids	<i>Aspergillus parasiticus</i>	[88]
<i>Milletia grandis</i> (E. Mey)	Aqueous, Methanol:Dichloromethane extracts	Phenols Flavonoids	<i>Aspergillus ochraceous</i> <i>Fusarium graminearum</i> <i>Fusarium oxysporum</i>	[88]
<i>Morinda citrifolia</i> L.	Essential oil	Methyl octanoate, Octanoic acid, Ethyl octanoate, Isopentyl hexanoate, 3- Methyl-2- butenyl hexanoate, 3-Methylbutyl octanoate, Methylbutyl-2- enyl octanoate	<i>Exserohilum turcicum</i>	[90]
<i>Olinia ventosa</i>	Hexane, Dichloromethane, Acetone and Methanol extracts	No data	<i>Trichoderma harzianum</i>	[89]
<i>Parthenium hysterophorus</i>	Hydroalcoholic extract	Tannins Flavonoids Saponins Amino acids Triterpenes Phenols	<i>Pyricularia grisea</i> <i>Phytophthora parasitica</i> var. <i>nicotianae</i> , <i>Phytophthora parasitica</i> <i>Fusarium oxysporum</i> <i>Stemphylium solani</i> Weber	[86]
<i>Pluchea carlinensi</i>	Hydroalcoholic extract	Phenols Tannins Flavonoids Saponins Steroids Alkaloids Quinone	<i>Pyricularia grisea</i> <i>Phytophthora parasitica</i> var. <i>nicotianae</i> , <i>Phytophthora parasitica</i> <i>Fusarium oxysporum</i> <i>Stemphylium solani</i> Weber	[86]

<i>Ricinus communis</i>	Methanol:Dichloromethane extract	Phenols Flavonoids	<i>Aspergillus parasiticus</i> <i>Fusarium verticillioides</i>	[88]
<i>Salvia africana- lutea</i> L.	Methanol:Dichloromethane extract	Propanoic acid, Dodecane Phosphoric acid, Glycerol, Succinic acid, Malic acid Pentadecane, Rythronic acid, Xylitol, Ribitol, 2-keto-1-gluconic acid, 1,3-dibromobicyclon, D-fructose, Fructose oxime, D-glucose, D-mannose, D-galactose, Sedoheptulose, o-methyloxime, Galactonic acid, Hexadecanoic acid Myo-inositol, Mannitol Caffeic acid, Octadecanoic acid, Alpha-D-glucopyranoside, Octacosane	<i>Fusarium verticillioides</i> <i>Fusarium proliferatum</i>	[91]
<i>Solanum panduriforme</i>	Methanol:Dichloromethane extract	Phenols Flavonoids	<i>Aspergillus parasiticus</i> <i>Aspergillus ochraceous</i> <i>Fusarium graminearum</i> <i>Fusarium oxysporum</i>	[88]
<i>Solidago canadensis</i> L.	Essential oil	Germacrene D, limonene, α -pinene, β -elemene, and bornyl acetate. Camphene, α -caryophyllene, sesquibinene bicyclosesquiphellandrene, β -cadinene, β -caryophyllene, β -pinene, β -sabinene Hydrocarbon monoterpenes, Oxygenated monoterpenes, Sesquiterpene hydrocarbons	<i>Monilinia fruticola</i> <i>Botrytis cinerea</i> <i>Aspergillus niger</i> <i>Penicillium expansum</i> <i>Bacillus megaterium</i> <i>Clavibacter michiganensis</i> <i>Xanthomonas campestris</i> <i>Pseudomonas fluorescens</i> <i>Pseudomonas syringae</i> pv. <i>phaseolicola</i>	[92]
<i>Thymus capitatus</i>	Essential oil	Thymol Carvacrol	<i>Aspergillus niger</i> <i>Aspergillus oryzae</i> <i>Penicillium digitatum</i> <i>Fusarium solani</i>	[93]
<i>Vangueria infausta</i>	Dichloromethane and Acetone extracts	No data	<i>Aspergillus parasiticus</i> <i>Trichoderma harzianum</i>	[89]
<i>Viola odorata</i> L.	Methanol:Water extract	Cyclotides: cy02, cy03, cy013, and cy019	<i>Fusarium oxysporum</i> <i>Fusarium graminearum</i> <i>Fusarium culmorum</i> <i>Mycosphaerella fragariae</i> <i>Botrytis cinerea</i> <i>Pseudomonas. syringae</i> pv. <i>syringae</i> <i>Pectobacterium atrosepticum</i> <i>Dickeya dadantii</i>	[94]
<i>Waburgia salutaris</i>	Methanol:Dichloromethane extract	Phenols Flavonoids	<i>Aspergillus parasiticus</i> <i>Aspergillus ochraceous</i> <i>Fusarium verticillioides</i> <i>Fusarium oxysporum</i>	[88]
<i>Xylothea kraussiana</i>	Dichloromethane and Acetone extracts	No data	<i>Trichoderma harzianum</i> <i>Fusarium oxysporum</i>	[89]
<i>Ziziphus mucronata</i>	Methanol:Dichloromethane extract	Phenols Flavonoids	<i>Fusarium graminearum</i>	[88]

Table 4.
Secondary metabolites of medicinal plants with biological activity against phytopathogens.

4. Conclusions

The phytochemicals that produce medicinal plants derived from their secondary metabolism represent a safe and effective alternative to control various phytopathogens that affect various crops of agricultural products of economic and nutritional

interest. There are different challenges in the use of biopesticides obtained from medicinal plants, such as evaluating the costs of obtaining these compounds on a large scale or exploring the possibility of them being obtained through chemical synthesis to increase yield and reduce costs. On the other hand, the various studies that exist on the effectiveness of these compounds are only at the laboratory level, which is why it is still necessary to explore and evaluate their effectiveness at the greenhouse and field levels.

Acknowledgements

The authors would like to acknowledge the University of Guanajuato for the grant of this publication.

Conflict of interest

The authors declare no conflict of interest.

Author details

Xóchitl S. Ramírez-Gómez^{1*}, Sandra N. Jiménez-García², Vicente Beltrán Campos¹ and Ma. Lourdes García Campos¹

¹ Department of Clinical Nursing, Division of Health Sciences and Engineering, University of Guanajuato, Celaya, Guanajuato, Mexico

² Department of Nursing and Obstetrics, Division of Health Sciences and Engineering, University of Guanajuato, Celaya, Guanajuato, Mexico

*Address all correspondence to: xosofira2002@yahoo.com.mx

IntechOpen

© 2019 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. 

References

- [1] Jones JDG, Dangel JL. The plant immune system. *Nature*. 2006;**444**: 323-329. DOI: 10.1038/nature05286
- [2] FHIA. Post-Harvest Deterioration of Fresh Fruits and Vegetables by Fungi and Bacteria [Internet]. 2007. Available from: http://www.fhia.org.hk/download/fhia_informa/fhiainfdic2007.pdf [Accessed: January 4, 2019]
- [3] Gene SM, Hoekstra PF, Hannam C, White M, Truman C, Hanson ML, et al. The role of vegetated buffers in agriculture and their regulation across Canada and the United States. *Journal of Environmental Management*. 2019;**243**: 12-21. DOI: 10.1016/j.jenvman.2019.05.003
- [4] Sun C, Chen L, Zhai L, Liu H, Jiang Y, Wang K, et al. National assessment of spatiotemporal loss in agricultural pesticides and related potential exposure risks to water quality in China. *Science of the Total Environment*. 2019;**677**:98-107. DOI: 10.1016/j.scitotenv.2019.04.34
- [5] Katagi T. Bioconcentration, bioaccumulation, and metabolism of pesticides in aquatic organisms. *Reviews of Environmental Contamination and Toxicology*. 2010;**204**:1-132. DOI: 10.1007/978-1-4419-1440-8_1
- [6] Pelkonen O, Bennekou SH, Crivellente F, Terron A, Hernandez AF. Integration of epidemiological findings with mechanistic evidence in regulatory pesticide risk assessment: EFSA experiences. *Archives of Toxicology*. 2019;1-10. DOI: 10.1007/s00204-019-02467-w
- [7] Panseri S, Chiesa L, Ghisleni G, Marano G, Boracchi P, Ranghieri V, et al. Persistent organic pollutants in fish: Biomonitoring and cocktail effect with implications for food safety. *Food Additives & Contaminants. Part A, Chemistry, Analysis, Control, Exposure & Risk Assessment*. 2019;**36**:601-611. DOI: 10.1080/19440049.2019.1579926
- [8] Hao Y, Zhang H, Zhang P, Yu S, Ma D, Li L, et al. Chlorothalonil inhibits mouse ovarian development through endocrine disruption. *Toxicology Letters*. 2019;**303**:38-47. DOI: 10.1016/j.toxlet.2018.12.011
- [9] Abolhassani M, Asadikaram G, Paydar P, Fallah H, Aghaee-Afshar M, Moazed V, et al. Organochlorine and organophosphorus pesticides may induce colorectal cancer; a case-control study. *Ecotoxicology and Environmental Safety*. 2019;**178**: 168-177. DOI: 10.1016/j.ecoenv.2019.04.030
- [10] Żaczek M, Weber-Dąbrowska B, Górski A. Phages in the global fruit and vegetable industry. *Journal of Applied Microbiology*. 2015;**118**:537-556. DOI: 10.1111/jam.12700
- [11] Pino-Otín MR, Ballesteros D, Navarro E, González-Coloma A, Val J, Mainar AM. Ecotoxicity of a novel biopesticide from *Artemisia absinthium* on non-target aquatic organisms. *Chemosphere*. 2019;**216**:131-146. DOI: 10.1016/j.chemosphere.2018.09.071
- [12] Pino-Otín MR, Val J, Ballesteros D, Navarro E, Sánchez E, González-Coloma A, et al. Ecotoxicity of a new biopesticide produced by *Lavandula luisieri* on non-target soil organisms from different trophic levels. *Science of the Total Environment*. 2019;**671**: 83-93. DOI: 10.1016/j.scitotenv.2019.03.293
- [13] Yamada T, Hamada M, Floreancig P, Nakabachi A. Diaphorin, a polyketide synthesized by an intracellular symbiont of the Asian citrus psyllid, is potentially harmful for biological control agents. *PLoS One*. 2019;**14**:e0216319. DOI: 10.1371/journal.pone.0216319

- [14] Mafakheri H, Mirghazanfari SM. Antifungal activity of the essential oils of some medicinal plants against human and plant fungal pathogens. *Cellular and Molecular Biology*. 2018;**64**:13-19
- [15] Isaac GS, Abu-Tahon MA. *In vitro* antifungal activity of medicinal plant extract against *Fusarium oxysporum* F. sp. *lycopersici* race 3 the causal agent of tomato wilt. *Acta Biologica Hungarica*. 2014;**65**:107-118. DOI: 10.1556/ABiol.65.2014.1.10
- [16] Askarne L, Talibi I, Boubaker H, Boudyach EH, Msanda F, Saadi B, et al. Use of Moroccan medicinal plant extracts as botanical fungicide against citrus blue mould. *Letters in Applied Microbiology*. 2013;**56**:37-43. DOI: 10.1111/lam.12012
- [17] Mohamed EZ. Impact of mycotoxins on humans and animals. *Journal of Saudi Chemical Society*. 2011;**15**: 129-144. DOI: 10.1016/j.jscs.2010.06.006
- [18] Molina A, Chavarría G, Alfaro-Cascante M, Leiva A, Granados-Chinchilla F. Mycotoxins at the start of the food chain in Costa Rica: Analysis of six *Fusarium* toxins and ochratoxin a between 2013 and 2017 in animal feed and aflatoxin M1 in dairy products. *Toxins*. 2019;**11**:E312. DOI: 10.3390/toxins11060312
- [19] Thiour-Mauprivez C, Martin-Laurent F, Calvayrac C, Barthelmebs L. Effects of herbicide on non-target microorganisms: Towards a new class of biomarkers? *Science of the Total Environment*. 2019;**684**:314-325. DOI: 10.1016/j.scitotenv.2019.05.230
- [20] Tahir HM, Basheer T, Ali S, Yaqoob R, Naseem S, Khan SY. Effect of pesticides on biological control potential of *Neoscona theisi* (Araneae: Araneidae). *Journal of Insect Science*. 2019;**19**:pii:17. DOI: 10.1093/jisesa/iez024
- [21] Brunton LL, Chabner BA, Knollman B. Goodman & Gilman's the Pharmacological Basis of Therapeutics. 12th ed. Vol. 17-39. United States: McGraw-Hill; 2012. pp. 123-143
- [22] Katzung BG, Masters SB, Trevor AJ. Basic & Clinical Pharmacology. 12th ed. United States: McGraw-Hill; 2013. pp. 37-68
- [23] Jimenez-Garcia SN, Guevara-Gonzalez RG, Miranda-Lopez R, Feregrino-Perez AA, Torres-Pacheco I, Vazquez-Cruz MA. Functional properties and quality characteristics of bioactive compounds in berries: Biochemistry, biotechnology, and genomics. *Food Research International*. 2013;**54**:1195-1207. DOI: 10.1016/j.foodres.2012.11.004
- [24] Jimenez-Garcia SN, Vazquez-Cruz MA, Guevara-Gonzalez RG, Torres-Pacheco I, Cruz-Hernandez A, Feregrino-Perez AA. Current approaches for enhanced expression of secondary metabolites as bioactive compounds in plants for agronomic and human health purposes—A review. *Polish Journal of Food and Nutrition Sciences*. 2013;**63**:67-78. DOI: 10.2478/v10222-012-0072-6
- [25] Khalid M, Rahman S-u, Bilal M. Role of flavonoids in plant interactions with the environment and against human pathogens—A review. *Journal of Integrative Agriculture*. 2019;**18**:211-230. DOI: 10.1016/s2095-3119(19)62555-4
- [26] Zhao C, Liu Y, Lai S, Cao H, Guan Y, San Cheang W, et al. Effects of domestic cooking process on the chemical and biological properties of dietary phytochemicals. *Trends in Food Science and Technology*. 2019;**01**:55-66. DOI: 10.1016/j.tifs.2019.01.004
- [27] Sahebi M, Hanafi MM, van Wijnen AJ, Akmar ASN, Azizi P, Idris AS, et al. Profiling secondary metabolites of plant defence mechanisms and oil palm in

- response to *Ganoderma boninense* attack. International Biodeterioration and Biodegradation. 2017;**122**:151-164. DOI: 10.1016/j.ibiod.2017.04.016
- [28] Valduga AT, Gonçalves IL, Magri E, Delalibera FJR. Chemistry, pharmacology and new trends in traditional functional and medicinal beverages. Food Research International. 2019;**120**:478-503. DOI: 10.1016/j.foodres.2018.10.091
- [29] Chakraborty P. Herbal genomics as tools for dissecting new metabolic pathways of unexplored medicinal plants and drug discovery. Biochime Open. 2018;**6**:9-16. DOI: 10.1016/j.biopen.2017.12.003
- [30] Jamwal K, Bhattacharya S, Puri S. Plant growth regulator mediated consequences of secondary metabolites in medicinal plants. Journal of Applied Research on Medicinal and Aromatic Plants. 2018;**9**:26-38. DOI: 10.1016/j.jarmap.2017.12.003
- [31] Frérot B, Leppik E, Groot AT, Unbehend M, Holopainen JK. Chemical signatures in plant-insect interactions. In: Sauvion N, Thiéry D, Calatayud PA, editors. Insect-Plant Interact a Crop Prot Perspect. 1st ed. London, United Kingdom: Academic Press; 2017. pp. 139-177. Available from: <http://www.sciencedirect.com/science/article/pii/S0065229616300982>
- [32] McCutcheon SC, Jørgensen SE. Phytoremediation. In: Jørgensen SE, Fath BD, editors. Encyclopedia of Ecology. 1st ed. Oxford: Academic Press; 2008. pp. 2751-2766. Available from: <http://www.sciencedirect.com/science/article/pii/B9780080454054000690>
- [33] Hounsborne N, Hounsborne B, Tomos D, Edwards-Jones G. Plant metabolites and nutritional quality of vegetables. Journal of Food Science. 2008;**73**:48-65. DOI: 10.1111/j.1750-3841.2008.00716.x
- [34] VanEtten HD, Sandrock RW, Wasmann CC, Soby SD, McCluskey K, Wang P. Detoxification of phytoanticipins and phytoalexins by phytopathogenic fungi. Canadian Journal of Botany. 1995;**73**:518-525. DOI: 10.1139/b95-291
- [35] Morrissey J, Lou GM. Iron uptake and transport in plants: The good, the bad, and the ionome. Chemical Reviews. 2009;**109**:4553-4567. DOI: 10.1021/cr900112r
- [36] Zaynab M, Fatima M, Abbas S, Sharif Y, Umair M, Zafar MH, et al. Role of secondary metabolites in plant defense against pathogens. Microbial Pathogenesis. 2018;**124**:198-202. DOI: 10.1016/j.micpath.2018.08.034
- [37] Seppänen SK, Syrjälä L, Von Weissenberg K, Teeri TH, Paajanen L, Pappinen A. Antifungal activity of stilbenes in *in vitro* bioassays and in transgenic *Populus* expressing a gene encoding pinosylvin synthase. Plant Cell Reports. 2004;**22**:584-593. DOI: 10.1007/s00299-003-0728-0
- [38] Del Cueto J, Møller BL, Dicenta F, Sánchez-Pérez R. β -Glucosidase activity in almond seeds. Plant Physiology and Biochemistry. 2018;**126**:163-172. DOI: 10.1016/j.plaphy.2017.12.028
- [39] Gadzovska Simic S, Tusevski O, Maury S, Delaunay A, Joseph C, Hagège D. Effects of polysaccharide elicitors on secondary metabolite production and antioxidant response in *Hypericum perforatum* L. shoot cultures. Scientific World Journal. 2014;**2014**:609649. DOI: 10.1155/2014/609649
- [40] Sathiyabama M, Bernstein N, Anusuya S. Chitosan elicitation for increased curcumin production and stimulation of defence response in turmeric (*Curcuma longa* L.). Industrial Crops and Products. 2016;**89**:87-94. DOI: 10.1016/j.indcrop.2016.05.007
- [41] Alami I, Mari S, Clérivet A. A glycoprotein from *Ceratocystis fimbriata*

- f. sp. *platani* triggers phytoalexin synthesis in *Platanus* × *acerifolia* cell-suspension cultures. *Phytochemistry*. 1998;**48**:771-776. DOI: 10.1016/S0031-9422(97)00892-3
- [42] Gadzovska Simic S, Tusevski O, Maury S, Delaunay A, Lainé E, Joseph C, et al. Polysaccharide elicitors enhance phenylpropanoid and naphthodianthrone production in cell suspension cultures of *Hypericum perforatum*. *Plant Cell, Tissue and Organ Culture*. 2015;**122**: 649-663. DOI: 10.1007/s11240-015-0798-z
- [43] Hemashenpagam N, Selvaraj T. Effect of arbuscular mycorrhizal (AM) fungus and plant growth promoting rhizomicroorganisms (PGPR's) on medicinal plant *Solanum viarum* seedlings. *Journal of Environmental Biology*. 2011;**32**:579-583
- [44] Vafadar F, Amooaghaie R, Otrushy M. Effects of plant-growth-promoting rhizobacteria and arbuscular mycorrhizal fungus on plant growth, stevioside, NPK, and chlorophyll content of *Stevia rebaudiana*. *Journal of Plant Interactions*. 2013;**9**:128-136. DOI: 10.1080/17429145.2013.779035
- [45] Kurosaki F, Yamashita A, Arisawa M. Involvement of GTP-binding protein in the induction of phytoalexin biosynthesis in cultured carrot cells. *Plant Science*. 2001;**161**:273-278. DOI: 10.1016/S0168-9452(01)00407-1
- [46] Bais HP, Walker TS, Schweizer HP, Vivanco JM. Root specific elicitation and antimicrobial activity of rosmarinic acid in hairy root cultures of *Ocimum basilicum*. *Plant Physiology and Biochemistry*. 2002;**40**:983-995. DOI: 10.1016/S0981-9428(02)01460-2
- [47] Singh G, Gavrieli J, Oakey JS, Curtis WR. Interaction of methyl jasmonate, wounding and fungal elicitation during sesquiterpene induction in *Hyoscyamus muticus* in root cultures. *Plant Cell Reports*. 1998;**17**:391-395. DOI: 10.1007/s002990050412
- [48] Yang D, Ma P, Liang X, Wei Z, Liang Z, Liu Y, et al. PEG and ABA trigger methyl jasmonate accumulation to induce the MEP pathway and increase tanshinone production in *Salvia miltiorrhiza* hairy roots. *Physiologia Plantarum*. 2012;**146**:173-183. DOI: 10.1111/j.1399-3054.2012.01603.x
- [49] Xu A, Zhan JC, Huang WD. Effects of ultraviolet C, methyl jasmonate and salicylic acid, alone or in combination, on stilbene biosynthesis in cell suspension cultures of *Vitis vinifera* L. cv. Cabernet Sauvignon. *Plant Cell, Tissue and Organ Culture*. 2015; **122**:197-211. DOI: 10.1007/s11240-015-0761-z
- [50] Hao X, Shi M, Cui L, Xu C, Zhang Y, Kai G. Effects of methyl jasmonate and salicylic acid on tanshinone production and biosynthetic gene expression in transgenic *Salvia miltiorrhiza* hairy roots. *Biotechnology and Applied Biochemistry*. 2015;**62**:24-31. DOI: 10.1002/bab.1236
- [51] Xu YW, Lv SS, Zaho D, Chen JW, Yang WT, Wu W. Effects of salicylic acid on monoterpene production and antioxidant systems in *Houttuynia cordata*. *African Journal of Biotechnology*. 2012;**11**:1364-1372. DOI: 10.5897/AJB11.1524
- [52] Chodisetti B, Rao K, Gandi S, Giri A. Gymnemic acid enhancement in the suspension cultures of *Gymnema sylvestre* by using the signaling molecules —Methyl jasmonate and salicylic acid. *In Vitro Cellular & Developmental Biology—Plant*. 2015;**51**:88-92. DOI: 10.1007/s11627-014-9655-8
- [53] Krzyzanowska J, Czubacka A, Pecio L, Przybys M, Doroszewska T, Stochmal A, et al. The effects of jasmonic acid and methyl jasmonate on rosmarinic acid production in *Mentha piperita* cell

- suspension cultures. *Plant Cell, Tissue and Organ Culture*. 2012;**108**:73-81. DOI: 10.1007/s11240-011-0014-8
- [54] Hayashi H, Huang P, Inoue K. Up-regulation of soyasaponin biosynthesis by methyl jasmonate in cultured cells of *Glycyrrhiza glabra*. *Plant & Cell Physiology*. 2003;**44**:404-411. DOI: 10.1093/pcp/pcg054
- [55] Hu X, Neill S, Cai W, Tang Z. Hydrogen peroxide and jasmonic acid mediate oligogalacturonic acid-induced saponin accumulation in suspension-cultured cells of *Panax ginseng*. *Physiologia Plantarum*. 2003;**118**: 414-421. DOI: 10.1034/j.1399-3054.2003.00124.x
- [56] Taurino M, Ingrosso I, D'amico L, De Domenico S, Nicoletti I, Corradini D, et al. Jasmonates elicit different sets of stilbenes in *Vitis vinifera* cv. Negramaro cell cultures. *Springerplus*. 2015;**4**:49. DOI: 10.1186/s40064-015-0831-z
- [57] Gorelick J, Rosenberg R, Smotrich A, Hanuš L, Bernstein N. Hypoglycemic activity of withanolides and elicited *Withania somnifera*. *Phytochemistry*. 2015;**116**:283-289. DOI: 10.1016/j.phytochem.2015.02.029
- [58] Ghorbanpour M, Hatami M, Kariman K, Dahaji PA. Phytochemical variations and enhanced efficiency of antioxidant and antimicrobial ingredients in *Salvia officinalis* as inoculated with different rhizobacteria. *Chemistry & Biodiversity*. 2016;**13**: 319-330. DOI: 10.1002/cbdv.201500082
- [59] Banchio E, Bogino PC, Santoro M, Torres L, Zygadlo J, Giordano W. Systemic induction of monoterpene biosynthesis in *Origanum majoricum* by soil bacteria. *Journal of Agricultural and Food Chemistry*. 2010;**58**:650-654. DOI: 10.1021/jf9030629
- [60] Banchio E, Xie X, Zhang H, Paré PW. Soil bacteria elevate essential oil accumulation and emissions in sweet basil. *Journal of Agricultural and Food Chemistry*. 2009;**57**:653-657. DOI: 10.1021/jf8020305
- [61] del Rosario Cappellari L, Santoro MV, Nievas F, Giordano W, Banchio E. Increase of secondary metabolite content in marigold by inoculation with plant growth-promoting rhizobacteria. *Applied Soil Ecology*. 2013;**70**:16-22. DOI: 10.1016/j.apsoil.2013.04.001
- [62] Liang Z, Ma Y, Xu T, Cui B, Liu Y, Guo Z, et al. Effects of abscisic acid, gibberellin, ethylene and their interactions on production of phenolic acids in *Salvia miltiorrhiza* bunge hairy roots. *PLoS One*. 2013;**8**:e72806. DOI: 10.1371/journal.pone.0072806
- [63] McSteen P, Zhao Y. Plant hormones and signaling: Common themes and new developments. *Developmental Cell*. 2008;**14**:467-473. DOI: 10.1016/j.devcel.2008.03.013
- [64] Abbasi BH, Stiles AR, Saxena PK, Liu CZ. Gibberellic acid increases secondary metabolite production in *Echinacea purpurea* hairy roots. *Applied Biochemistry and Biotechnology*. 2012;**168**:2057-2066. DOI: 10.1007/s12010-012-9917-z
- [65] Zobayed SMA, Afreen F, Kozai T. Temperature stress can alter the photosynthetic efficiency and secondary metabolite concentrations in St. John's wort. *Plant Physiology and Biochemistry*. 2005;**43**:977-984. DOI: 10.1016/j.plaphy.2005.07.013
- [66] Binder BYK, Peebles CAM, Shanks JV, San K-Y. The effects of UV-B stress on the production of terpenoid indole alkaloids in *Catharanthus roseus* hairy roots. *Biotechnology Progress*. 2009;**25**: 861-865. DOI: 10.1002/btpr.97
- [67] Tari I, Kiss G, Deér AK, Csiszár J, Erdei L, Gallé A, et al. Salicylic acid increased aldose reductase activity and

- sorbitol accumulation in tomato plants under salt stress. *Biologia Plantarum*. 2010;**54**:677-683. DOI: 10.1007/s10535-010-0120-1
- [68] Liu H, Wang X, Wang D, Zou Z, Liang Z. Effect of drought stress on growth and accumulation of active constituents in *Salvia miltiorrhiza* bunge. *Industrial Crops and Products*. 2011;**33**:84-88. DOI: 10.1016/j.indcrop.2010.09.006
- [69] Gupta P, Sharma S, Saxena S. Biomass yield and steviol glycoside production in callus and suspension culture of *Stevia rebaudiana* treated with proline and polyethylene glycol. *Applied Biochemistry and Biotechnology*. 2015; **176**:863-874. DOI: 10.1007/s12010-015-1616-0
- [70] Threlfall DR, Whitehead IM. The use of biotic and abiotic elicitors to induce the formation of secondary plant products in cell suspension cultures of solanaceous plants. *Biochemical Society Transactions*. 1988;**16**:71-75. DOI: 10.1042/bst0160071
- [71] Furze JM, Rhodes MJ, Parr AJ, Robins RJ, Whitehead IM, Threlfall DR. Abiotic factors elicit sesquiterpenoid phytoalexin production but not alkaloid production in transformed root cultures of *Datura stramonium*. *Plant Cell Reports Germany*. 1991;**10**:111-114. DOI: 10.1007/BF00232039
- [72] Shakeran Z, Keyhanfar M, Asghari G, Ghanadian M. Improvement of atropine production by different biotic and abiotic elicitors in hairy root cultures of *Datura metel*. *Turkish Journal of Biology*. 2015;**39**:111-118. DOI: 10.3906/biy-1405-25
- [73] Cai Z, Kastell A, Speiser C, Smetanska I. Enhanced resveratrol production in *Vitis vinifera* cell suspension cultures by heavy metals without loss of cell viability. *Applied Biochemistry and Biotechnology*. 2013; **171**:330-340. DOI: 10.1007/s12010-013-0354-4
- [74] Savitha BC, Thimmaraju R, Bhagyalakshmi N, Ravishankar GA. Different biotic and abiotic elicitors influence betalain production in hairy root cultures of *Beta vulgaris* in shake-flask and bioreactor. *Process Biochemistry*. 2006;**41**:50-60. DOI: 10.1016/j.procbio.2005.03.071
- [75] Jochum GM, Mudge KW, Thomas RB. Elevated temperatures increase leaf senescence and root secondary metabolite concentrations in the understory herb *Panax quinquefolius* (Araliaceae). *American Journal of Botany*. 2007;**94**:819-826. DOI: 10.3732/ajb.94.5.819
- [76] Zhao Y, Qi L-W, Wang W-M, Saxena PK, Liu C-Z. Melatonin improves the survival of cryopreserved callus of *Rhodiola crenulata*. *Journal of Pineal Research*. 2011;**50**:83-88. DOI: 10.1111/j.1600-079X.2010.00817.x
- [77] Chan LK, Koay SS, Boey PL, Bhatt A. Effects of abiotic stress on biomass and anthocyanin. *Biological Research*. 2010;**43**:127-135. DOI: /S0716-97602010000100014
- [78] Liu W, Liu C, Yang C, Wang L, Li S. Effect of grape genotype and tissue type on callus growth and production of resveratrols and their piceids after UV-C irradiation. *Food Chemistry*. 2010;**122**: 475-481. DOI: 10.1016/j.foodchem.2010.03.055
- [79] Bor M, Seckin B, Ozgur R, Yilmaz O, Ozdemir F, Turkan I. Comparative effects of drought, salt, heavy metal and heat stresses on gamma-aminobutyric acid levels of sesame (*Sesamum indicum* L.). *Acta Physiologiae Plantarum*. 2009; **31**:655-659. DOI: 10.1007/s11738-008-0255-2
- [80] Chen Y, Guo Q, Liu L, Liao L, Zhu Z. Influence of fertilization and drought

- stress on the growth and production of secondary metabolites in *Prunella vulgaris* L. The Journal of Medicinal Plants Research. 2011;5:1749-1755
- [81] Pavlík M, Vacek J, Klejdus B, Kubáň V. Hypericin and hyperforin production in *St. John's wort* *in vitro* culture: Influence of saccharose, polyethylene glycol, methyl jasmonate, and agrobacterium tumefaciens. Journal of Agricultural and Food Chemistry. 2007; 55:6147-6153. DOI: 10.1021/jf070245w
- [82] Butassi E, Svetaz LA, Zhou S, Wolfender JL, Cortés JCG, Ribas JC, et al. The antifungal activity and mechanisms of action of quantified extracts from berries, leaves and roots of *Phytolacca tetramera*. Phytomedicine. 2019;16:152884. DOI: 10.1016/j.phymed.2019.152884
- [83] Talibi I, Askarne L, Boubaker H, Boudyach EH, Msanda F, Saadi B, et al. Antifungal activity of Moroccan medicinal plants against citrus sour rot agent *Geotrichum candidum*. Letters in Applied Microbiology. 2012;55:155-161. DOI: 10.1111/j.1472-765X.2012.03273.x
- [84] Anaruma ND, Schmidt FL, Duarte MC, Figueira GM, Delarmelina C, Benato LA, et al. Control of *Colletotrichum gloeosporioides* (penz.) Sacc. In yellow passion fruit using *Cymbopogon citratus* essential oil. Brazilian Journal of Microbiology. 2010; 41:66-73. DOI: 10.1590/S1517-838220100001000012
- [85] Zhao W, Xu LL, Zhang X, Gong XW, Zhu DL, Xu XH, et al. Three new phenanthrenes with antimicrobial activities from the aerial parts of *Juncus effusus*. Fitoterapia. 2018;130:247-250. DOI: 10.1016/j.fitote.2018.09.007
- [86] Rodríguez AT, Morales D, Ramírez MA. Efecto de extractos vegetales sobre el crecimiento *in vitro* de hongos fitopatógenos. Cultivos Tropicales. 2000;21:79-82
- [87] Salhi N, Saghir SAM, Terzi V, Brahmi I, Ghedairi N, Bissati S. Antifungal activity of aqueous extracts of some dominant Algerian medicinal plants. BioMed Research International. 2017;2017:7526291. DOI: 10.1155/2017/7526291
- [88] Mongalo NI, Dikhoba PM, Soyingbe SO, Makhafole TJ. Antifungal, anti-oxidant activity and cytotoxicity of south African medicinal plants against mycotoxigenic fungi. Heliyon. 2018;4: e00973. DOI: 10.1016/j.heliyon.2018. e00973
- [89] Mahlo SM, Chauke HR, McGaw L, Eloff J. Antioxidant and antifungal activity of selected medicinal plant extracts against phytopathogenic fungi. African Journal of Traditional, Complementary, and Alternative Medicines. 2016;13:216-222. DOI: 10.21010/ajtcam.v13i4.28
- [90] Costa E, Silva J, de Sousa Carlos Mourão D, de Oliveira Lima FS, de Almeida Sarmento R, Sunti Dalcin M, et al. The efficiency of noni (*Morinda citrifolia* L.) essential oil on the control of leaf spot caused by *Exserohilum turcicum* in maize culture. Medicines. 2017;4:60. DOI: 10.3390/medicines4030060
- [91] Nkomo MM, Katerere D, Vismer H, Cruz TT, Stephane S, Balayssac SS, et al. Fusarium inhibition by wild populations of the medicinal plant *Salvia africana-lutea* L. linked to metabolomic profiling. BMC Complementary and Alternative Medicine. 2014;14(99):2-9. DOI: 10.1186/1472-6882-14-99
- [92] Elshafie HS, Grul'ová D, Baranová B, Caputo L, De Martino L, Sedlák V, et al. Antimicrobial activity and chemical composition of essential oil extracted from *Solidago canadensis* L. growing wild in Slovakia. Molecules. 2019;24:1206. DOI: 10.3390/molecules24071206

[93] Tabti L, Amine Dib ME, Gaouar N, Samira B, Tabti B. Antioxidant and antifungal activity of extracts of the aerial parts of *Thymus capitatus* (L.) Hoffmanns against four phytopathogenic fungi of *Citrus sinensis*. Jundishapur Journal of Natural Pharmaceutical Products. 2014;**9**:49-54. DOI: 10.17795/jjnpp-13972

[94] Slazak B, Kapusta M, Strömstedt AA, Słomka A, Krychowiak M, Shariatgorji M, et al. How does the sweet violet *Viola odorata* L. fight pathogens and pests-cyclotides as a comprehensive plant host defense system. Frontiers in Plant Science. 2018;**9**:1296. DOI: 10.3389/fpls.2018.01296

[95] Garcia-Mier L, Jimenez-Garcia SN, Guevara-González RG, Feregrino-Perez AA, Contreras-Medina LM, Torres-Pacheco I. Elicitor mixtures significantly increase bioactive compounds, antioxidant activity, and quality parameters in sweet bell pepper. Journal of Chemistry. 2015;**2015**:8. DOI: 10.1155/2015/269296. Article ID: 269296

[96] Vargas-Hernández M, Torres-Pacheco I, Gautier F, Álvarez-Mayorga B, Cruz-Hernández A, García-Mier L, et al. Influence of hydrogen peroxide foliar applications on *in vitro* antimicrobial activity in *Capsicum chinense* Jacq. Plant Biosystems. 2017; **151**:269-275. DOI: 10.1080/11263504.2016.1168494

[97] Zunun-Pérez AY, Guevara-Figueroa T, Jimenez-Garcia SN, Feregrino-Pérez AA, Gautier F, Guevara-González RG. Effect of foliar application of salicylic acid, hydrogen peroxide and a xyloglucan oligosaccharide on capsate content and gene expression associated with capsinoids synthesis in *Capsicum annuum* L. Journal of Biosciences. 2017; **42**:245-250. DOI: 10.1007/s12038-017-9682-9